



HHS Public Access

Author manuscript

Med Sci Sports Exerc. Author manuscript; available in PMC 2020 June 01.

Published in final edited form as:

Med Sci Sports Exerc. 2019 June ; 51(6): 1303–1313. doi:10.1249/MSS.0000000000001942.

Physical Activity, Injurious Falls, and Physical Function in Aging: An Umbrella Review

Loretta DiPietro¹, Wayne W. Campbell², David M. Buchner³, Kirk I. Erickson⁴, Kenneth E. Powell⁵, Bonny Bloodgood⁶, Timothy Hughes¹, Kelsey R. Day¹, Katrina L. Piercy⁷, Alison Vaux-Bjerke⁷, and Richard D. Olson⁷ 2018 Physical Activity Guidelines Advisory Committee

¹Milken Institute School of Public Health, The George Washington University, Washington, DC

²Department of Nutrition Science, Purdue University, West Lafayette, IN

³University of Illinois at Urbana-Champaign, Champaign, IL

⁴University of Pittsburgh, Pittsburgh, PA

⁵Centers for Disease Control and Prevention, Atlanta, GA

⁶ICF, Fairfax, VA

⁷Office of Disease Prevention and Health Promotion, Office of the Assistant Secretary for Health, U.S. Department of Health and Human Services, Rockville, MD

Abstract

Purpose: To review and update the evidence of the relationship between physical activity, risk of fall-related injury, and physical function in community-dwelling older people that was presented in the *2018 Physical Activity Guidelines Advisory Committee Scientific Report* (PAGAC Report).

Methods: Duplicate independent screenings of 1,415 systematic reviews and meta-analyses published between 2006 and 2016 identified from PubMed®, Cochrane Library, and CINAHL databases yielded 111 articles used for the PAGAC Report. The PAGAC Aging Subcommittee members graded scientific evidence strength based upon a 5-criteria rubric and assigned one of four grades: strong, moderate, limited, or not assignable. An updated search of 368 articles published between January 2017 and March 2018 yielded 35 additional pertinent articles.

Address for Correspondence: Loretta DiPietro, PhD, MPH, Milken Institute School of Public Health, 950 New Hampshire Ave, Suite 2, Washington, DC 20052, (Telephone) 202.994.4910, ldp1@gwu.edu.

*The 2018 Physical Activity Guidelines Advisory Committee includes David M. Buchner, Wayne W. Campbell, Loretta DiPietro, Kirk I. Erickson, Charles H. Hillman, John M. Jakicic, Kathleen F. Janz, Peter T. Katzmarzyk, Abby C. King, William E. Kraus, Richard F. Macko, David X. Marquez, Anne McTiernan, Russell R. Pate, Linda S. Pescatello, Kenneth E. Powell and Melicia C. Whitt-Glover.

This paper is being published as an official pronouncement of the American College of Sports Medicine. This pronouncement was reviewed for the American College of Sports Medicine by members-at-large and the Pronouncements Committee. This paper serves as an update to the topics covered in the 2009 ACSM position stand, "Exercise and Physical Activity for Older Adults" [*Med. Sci. Sports Exerc.* 2009; 41(7):1510-30]. *Disclaimer:* Care has been taken to confirm the accuracy of the information present and to describe generally accepted practices. However, the authors, editors, and publisher are not responsible for errors or omissions or for any consequences from application of the information in this publication and make no warranty, expressed or implied, with respect to the currency, completeness, or accuracy of the contents of the publication. Application of this information in a particular situation remains the professional responsibility of the practitioner; the clinical treatments described and recommended may not be considered absolute and universal recommendations.

Results: Strong evidence demonstrated that physical activity reduced the risk of fall-related injuries by 32-40%, including severe falls requiring medical care or hospitalization. Strong evidence also supported that physical activity improved physical function and reduced the risk of age-related loss of physical function in an inverse graded manner among the general aging population, and improved physical function in older people with frailty and with Parkinson's disease. Aerobic, muscle-strengthening, and/or multicomponent physical activity programs elicited the largest improvements in physical function in these same populations. Moderate evidence indicated that for older adults who sustained a hip fracture or stroke, extended exercise programs and mobility-oriented physical activity improved physical function.

Conclusions: Regular physical activity effectively helps older adults improve or delay the loss of physical function and mobility while reducing the risk of fall-related injuries. These important public health benefits underscore the importance of physical activity among older adults, especially those living with declining physical function and chronic health conditions.

Keywords

Exercise; Functional Performance; Geriatrics; Mobility

INTRODUCTION

Advances in public health and in health care are keeping people alive longer, and consequently, the proportion of older people in the global population is increasing rapidly. As of 2016, individuals ages 65 years and older comprise about 13% of the United States population, and their numbers are projected to reach 72.1 million (19% of the total population) by the year 2030 (1). This represents a two-fold increase compared with the older adult population in 2000. Moreover, the number of people 85 years and older is projected to rise to 14.6 million by 2040 (1). Due to these growing demographic trends, the prevention of chronic disease, the maintenance of functional status, and the preservation of physical independence in aging present major challenges that have substantial personal and public health implications.

Physical activity is any bodily movement that results in increased energy expenditure, and can be achieved by a variety of leisure-time, work or transportation-related activities (2). Exercise refers to physical activities that are planned, structured, repetitive, and intended to improve or maintain fitness, function, and health (2). Ample evidence now exists that regular physical activity is key to preventing and managing major chronic diseases common to older people (3). Physical activity is also important for preserving physical function and mobility, which can then delay the onset of major disability (4). Current physical activity guidelines for older people recommend at least 150 min/week of moderate-intensity aerobic activity, with muscle-strengthening activity performed on two or more days/week (5). Despite the known benefits of physical activity to health and physical function in aging, however, the proportion of older adults meeting recommended physical activity guidelines for aerobic activity remains low (27%), based on data from the 2011-2012 National Health and Nutrition Examination Survey (NHANES)(6). This low prevalence of physical activity has important implications since it is a modifiable behavior that contributes substantially to the burden of chronic disease mortality in the United States (7).

Since the *2008 Physical Activity Guidelines Advisory Committee (PAGAC) Scientific Report* (3), considerable evidence has emerged regarding the relative benefits of additional modes or combinations of physical activity to specific physical function outcomes (e.g., strength, gait speed, balance, activities of daily living [ADL] function). These additional physical activity interventions include progressive resistance training, multicomponent exercise, dual-task training, active video gaming, tai chi, yoga, and dance. In addition, the current research has begun to address the issues of the dose-response relationship between physical activity and physical function in aging. Similar to studies of pharmacologic agents, it is not only important to determine if a graded relationship exists, but also to determine the shape of the relationship for specific health outcomes, in order to establish a minimal effective dose and a maximal threshold dose for safety.

The 2018 PAGAC Scientific Report (8) expanded upon the 2008 Report by examining the relationship between physical activity and the risk of fall-related injuries, as well as the relationship between physical activity and physical function, in both the general aging population and in people living with specific chronic diseases (cardiovascular disease (CVD), chronic obstructive pulmonary disease (COPD), cognitive impairment, frailty, hip fracture, osteoporosis, Parkinson's Disease, stroke, and visual impairments). The 2018 PAGAC Scientific Report further leveraged current research in examining: 1) the dose-response relationship between exposure and outcome; 2) the mode of activity most beneficial to a specific functional outcome; and 3) whether the relationship between physical activity and physical function varied by age, race, sex, sociodemographic characteristics, or by body weight. This current article summarizes the evidence from the 2018 PAGAC Scientific Report and includes new evidence from an updated search of the effects of physical activity on fall-related injuries, and physical function in older people.

METHODS

Search Strategy, Study Selection, and Quality Assessment

Table 1 provides the specific questions and subquestions addressed by the Aging Subcommittee of the 2018 PAGAC in their report. A first search was undertaken to include publications from 2006 to 2016. The searches were conducted in PubMed®, CINAHL, and Cochrane Library and supplemented through hand-searches of reference lists of included articles and are reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (9). The Subcommittee determined that systematic reviews, meta-analyses, pooled analyses, and reports provided a wealth of quality information to answer two of its three research questions. Thus, to increase work efficiency, the searches were limited to these types of reviews. For Question 1 (What is the relationship between physical activity and risk of injury due to a fall?), the Subcommittee found that existing reviews (systematic reviews, meta-analyses, pooled analyses, and reports) covered only evidence from randomized controlled trials (RCTs), and therefore a supplementary search for cohort and case-control studies was conducted to capture the most complete literature. Search terms included physical activity and sedentary time terms combined with physical function or falls and injuries terms. The full search strategies are available at https://health.gov/paguidelines/second-edition/report/supplementary_material/pdf/

[Aging_Q1_Risk_of_Injuries_Evidence_Portfolio.pdf](#) and https://health.gov/paguidelines/second-edition/report/supplementarymaterial/pdf/Aging_Q2_Physical_Function_Evidence_Portfolio.pdf.

The inclusion criteria were pre-defined, and studies were considered potentially eligible if they were systematic reviews, meta-analyses, pooled analyses, or reports published in English from 2006 until February 2016 (also cohort studies published in English from 2006 until 2016 for risk of fall-related injuries) and investigated the association between all types and intensities of physical activity and physical function and/or risk of injuries from falls in the aging population. Studies of non-ambulatory adults, hospitalized patients, or animals were excluded. Two reviewers independently screened titles, abstracts, and full-text of the identified articles. A third reviewer helped resolve disagreement between reviewers.

In March, 2018, two updated systematic reviews were undertaken to identify additional systematic reviews and meta-analyses published between January 2017 through February 2018 that assessed the relationship between 1) any type of physical activity and fall-related injury or 2) physical function in the aging population. The searches were also conducted in PubMed®, CINAHL, and Cochrane Library and supplemented through hand-searches of reference lists of included articles. The updated review followed an established protocol that was registered as two reviews with PROSPERO [CRD42018096687 (fall-related injuries) and CRD42018095776 (physical function)].

Evidence to inform each question was graded as strong, moderate, limited, or “not assignable” based on several grading criteria, including applicability, generalizability, risk of bias/study limitations, quantity and consistency of results across studies, and magnitude and precision of effect (8) (see Table, Supplemental Digital Content 1, 2018 Physical Activity Guidelines Advisory Committee Grading Criteria). Table 2 provides a summary of the relationships and level of evidence for each health outcome examined by the 2018 PAGAC Aging Subcommittee.

RESULTS

After duplicates were removed, a total of 1,415 articles were identified from the original search process. Following full-text review, a total of four articles were deemed relevant to the question about fall-related injuries (Question 1); 38 were relevant to physical function in the general aging population (Question 2); and 63 were relevant to physical function in older people with specific chronic diseases (Question 3). Quality for each systematic review (SR), meta-analysis (MA), or article was assessed using AMSTARExBP (10). Risk of bias, or internal validity, was assessed for each original study using an adapted version of the USDA Nutrition Evidence Library Bias Assessment Tool (NEL BAT)(11).

The updated systematic search for fall-related injury risk identified 38 unique systematic reviews and meta-analyses after duplicates were removed. Of these, 32 were excluded after a review of the titles and abstracts and five more were excluded after full-text review leaving one new review (Figure 1). The updated search for physical function in the general aging population and in those with selected chronic conditions identified 330 systematic reviews

and meta-analyses, of which 288 were excluded after review of titles, six were excluded after review of the abstracts, and two more excluded after full-text review leaving 34 new reviews (Figure 2).

Physical Activity and Risk of Fall-Related Injury

The 2018 PAGAC Scientific Report cited strong and consistent evidence from RCTs demonstrating that multicomponent physical activity (i.e., combinations of aerobic, muscle-strengthening, balance, and flexibility) significantly reduced the risk of fall-related injuries by about 32-40%, including severe falls that result in bone fracture, head trauma, open wound soft tissue injury, or any other injury requiring medical care or admission to hospital (12–15). Moreover, the benefits of physical activity programs to reduce the risk of these four categories of fall-related injuries were similar between older adults identified as being at high risk of falling versus those who were at an unspecified risk (12). Also, fall prevention programs using multicomponent activity reduced the risk of fall-related bone fractures by 40-66% among older adults in community and home settings (12–15). These RCT findings were supported by data from three prospective cohort studies (16–18) and one case-control study (19).

The updated search identified one review that supported the evidence from the 2018 PAGAC Report (20). A meta-analysis of five RCTs conducted in Asian countries reported that participation in physical activity programs (primarily tai chi) by community-dwelling older adults reduced the risk of fall-related injuries by 50% (relative risk (RR) = 0.50; 95% confidence interval (CI): 0.35 to 0.71).

Dose-Response.—There was some evidence in the 2018 PAGAC Scientific Report to suggest that a dose-response relationship exists between the amount of moderate-to-vigorous physical activity or home and group exercise and risk of fall-related injury and bone fracture; however, the small number of studies available and the diverse array of physical activities studied made it difficult to describe the shape of the relationship. Consistent results from four high-quality epidemiologic studies (three cohort and one case-control) suggested that adults aged 65 years and older who participated in physical activity of at least moderate-intensity for 30 minutes per day (16) or engaging in high/very high levels of activity (i.e. a weekly physical activity index score ≥ 25)(17), reduced the risk of fall-related injury and bone fracture. Evidence also exists that even adults ages 85 years and older obtained similar benefits from 60 minutes per week of home- or group-based physical activity (18). However, it is important to note that lower amounts of moderate-intensity physical activity (16, 17) and walking (18) may not be sufficient to reduce the risk of fall-related injury and bone fracture in older age.

Physical Activity Type.—The physical activity programs that effectively reduced the risk of fall-related injuries and bone fractures contained a variety of group- and home-based activities (12, 14, 15, 18, 19). Most programs were multicomponent and included various combinations of moderate-intensity balance, strength, endurance, gait, flexibility, and “physical function” training, as well as recreational activities (e.g., dancing, cycling, gardening, sports). Although the research was limited, it does not support the use of low-

intensity walking as a primary mode of physical activity to reduce the risk of fall-related injuries and fractures among older adults (18, 19), although walking may be included in multicomponent physical activity regimens. Unfortunately, insufficient information was available from the systematic reviews to determine the effects of individual elements (e.g., strength training, balance training) of the multicomponent training programs on the risk of fall-related injuries.

Effect Modification by Sociodemographic Characteristics or Pre-existing Disability.—There was insufficient evidence available to determine whether the relationship between physical activity and risk of fall-related injuries and bone fractures varies by age, sex, race/ethnicity, socio-economic status, or weight status or whether factors such as level of physical function ability and pre-existing gait disability modify the relationship between physical activity and risk of injury due to a fall.

Physical Activity and Physical Function in the General Older Population

The 2018 PAGAC Scientific Report cited strong evidence from RCTs and cohort studies that aerobic, muscle-strengthening, balance, and/or multicomponent physical activity programs improved physical function and reduced risk of age-related loss of physical function in the general aging population (8). One high quality meta-analysis by Chase, et al. (21) analyzed data from 28 RCTs using objective composite measures of physical function, such as the short physical performance battery (SPPB), timed-up-and-go tests (TUG), the continuous scale physical performance test (CS-PPT), and the physical performance test (PPT). The summary effect size (ES) describing the magnitude of the relation between physical activity and physical function was 0.45 (95% CI: 0.27 to 0.64), with the higher quality studies reporting smaller effect sizes. The updated search identified nine SR/MAs that supported the 2018 Scientific Report by also providing strong to moderate evidence demonstrating the benefits of physical activity to physical function in this population (22–30).

Dose-Response.—The 2018 PAGAC Scientific Report found strong evidence of an inverse dose-response relationship between volume of aerobic physical activity and risk of physical functional limitations in the general aging population. One review of 24 comparisons from prospective cohort studies with covariate adjustment classified dose of aerobic activity reported in cohort studies into four ordinal categories (0=no activity; 1= low activity; 2=moderate activity; and 3 = vigorous activities and/or high activity volume) (31). With this analysis framework, virtually every study showed an inverse dose-response relationship of aerobic activity with risk of limitations in physical function (Figure 3). The 2018 Scientific Report cited limited evidence of a dose-response relationship between either balance (32) or muscle-strengthening training and physical function (21) in this same population. The updated search, however provided important additional information. Roberts et al. (29) examined the effects of different physical activity types and levels (determined from calculated MET-min and categorized as low, moderate, or high) on ADL function. Results from the pooled meta-analysis indicated that, compared with a control condition, there were beneficial effects of physical activity on ADL physical performance (standardized mean difference (SMD)=0.72; 95% CI: 0.45 to 1.00; 29 studies) with the largest effects observed for moderate activity levels (SMD=1.07; 95% CI: 0.44 to 1.70; 9

studies), compared with low levels (SMD=0.57; 95% CI:0.29 to 0.86; 20 studies). There were no studies of ADL performance that used high physical activity levels, and therefore a conclusion regarding their benefits could not be drawn. With regard to various levels of multi-task activities (those activities that combine physical and mental tasks), the greatest difference in ADL performance between the exercise and the control groups were reported for the high level multi-task group (SMD=1.36; 95% CI: 0.46 to 2.26; 6 studies), followed by the moderate level group (SMD=0.74; 95% CI: 0.41 to 1.06; 13 studies), and the low level group (SMD=0.45; 95% CI: -0.01 to 0.91; 8 studies).

Types of activity.—The 2018 PAGAC Scientific Report cited strong evidence that, specifically, aerobic, muscle-strengthening, and multicomponent physical activity improved physical function in the general aging population, and moderate evidence indicating that balance training improved physical function in a dose-response manner among these same persons (8). The evidence linking physical activities such as tai chi exercise, dance training, active video gaming, and dual-task training to improvements in physical function in healthy aging were limited, however, due to the lack of studies at this time. Similarly, insufficient evidence was available to determine the effects of flexibility activity, yoga, and qigong exercise on physical function in the general aging population.

The updated search identified a meta-analysis by Farlie and colleagues (24) that examined the effects of four different types of balance training interventions [multidimensional (activities such as “functional exercises”, tai chi, and ball games), control center of mass (COM), mobility, and reaching] on several dimensions of balance performance. Overall, the meta-analysis resulted in small to moderate effects in favor of programs that included multidimensional (SMD=0.50; 95% CI: 0.36 to 0.65), reaching (SMD=0.48; 95% CI: 0.33 to 0.64), COM (SMD=0.42; 95% CI: 0.27 to 0.56) and mobility (SMD=0.31; 95% CI: 0.20 to 0.43) balance training versus no balance training interventions. The authors noted, however, that there was substantial between-study heterogeneity among the interventions (I^2 range 50.4% to 80.6%).

Multi-task activities with high levels of physical (speed, coordination, balance), mental and social demands (e.g., dancing, team sports, handball) appeared particularly effective in improving functional performance, relative to moderate or lower levels of such activities and compared with control conditions (29). Nordic walking also has demonstrated moderate effectiveness in improving dynamic balance (effect size (ES)=0.30), functional balance (ES=0.62), muscle strength of upper (ES=0.66) and lower (ES=0.43) limbs, and aerobic capacity (ES=0.92) compared with no exercise comparison groups in healthy older people (23). A meta-analysis of 23 studies by Liu and colleagues (27) among community-dwelling older people with low physical function reported that progressive resistance exercise was effective in improving lower-body muscle strength and static standing balance; however multicomponent exercise was more effective in improving muscle strength, balance, and lower-body physical functioning, compared with progressive resistance exercise alone. Neither progressive resistance nor multicomponent exercise was effective in improving activities of daily living, however.

Howe and colleagues (25) examined the effects of active computer gaming (ACG) on physical function in 1,838 healthy people >65 years of age. Their meta-analysis of 35 RCTs reported significant moderate effects in favor of ACG over a control intervention on balance (SMD = 0.52; 95% CI: 0.24 to 0.79; 17 studies) and on functional exercise capacity when the volume of the ACG was >120 minutes per week (SMD = 0.53; 95% CI: 0.15 to 0.90; 5 studies). The authors noted, however, that the quality of the evidence for all comparisons was graded low or very low.

Finally, Weber, et al., (30) reviewed 14 studies (6 RCTs) of interventions that integrated and embedded “functional exercise” into the daily activities of older people. The most frequently evaluated intervention in this review was the Lifestyle-integrated Functional Exercise (LiFE) program (33). “Functional exercises” are designed to improve lower-body strength, balance, and motor performance, as well as for increasing daily levels of physical activity. Examples of such include postures or walking with gradual reduction in the base of support (e.g., upgrading tandem stand to one-leg stand over time) and dynamic movements that perturb the center of gravity (stepping over obstacles). “Functional exercises” for improving lower-body strength include squatting, chair stands, and toe raises with a gradual increase to more intense and challenging activities (33). Importantly, the LiFE intervention has strategies for behavior change that are based on the habit re-framing theory (34). Indeed, these exercises are linked to everyday tasks by using situational and environmental cues (e.g., tooth brushing, housework) as prompts to action, with the idea of performing the activities intentionally and consciously until they become a habit. Evidence from three RCTs in the Weber, et al. (30) review suggested that the LiFE intervention significantly improved balance, strength, and functional performance compared with either no intervention, low-intensity activities (e.g., walking), or structured exercise programs. Two of these RCTs also reported a significant reduction in incident falls in participants in the LiFE group, compared with those in either a no intervention or a low-intensity activity group. Thus, although the data are limited, the LiFE approach appears to be a promising alternative or complement to traditional structured exercise programs.

In sum, our current findings pertaining to the effectiveness of different types of physical activity for improving physical function support those from the *2018 PAGAC Scientific Report*, which cited strong evidence of the benefits of aerobic, muscle-strengthening, and multicomponent physical activity on improvements in physical function in the general aging population, and moderate evidence indicating that balance training improves physical function in these same persons. The evidence linking ACG (or active video gaming), Nordic walking, or functional exercise to improvements in physical function remains limited at this time.

Sociodemographic Characteristics and Weight Status.—Limited evidence from the 2018 PAGAC Scientific Report suggested that the relationship between physical activity and physical function did not vary by age, sex, or weight status in the general population of older adults (8). One meta-analysis reported sex and body mass index (BMI) were not significant effect modifiers of the relationship of physical activity on composite physical function scores (21). A meta-analysis of cohort studies reported the relationship between aerobic activity and ADL dependency did not differ significantly by age (75 years and

younger vs older than 75 years) (35). The available evidence was insufficient to determine whether the relationship between physical activity and physical function varied by race/ethnicity and socioeconomic status in the general population of older adults. No relevant analyses were located in the updated search. We note, however, that several findings with “limited” evidence also have high public health importance. Adults age 75 years and older have more age-related loss of physical function, are more likely to be women, and the majority have a BMI in the range of overweight-to-obese.

Effect Modification by Visual, Cognitive, or Physical Impairments.—None of the SR/MAs identified in either search examined whether visual or cognitive impairments modified the effects of physical activity on physical function. Limited evidence suggested that physical activity has a stronger effect on physical function in older adults with limitations in physical function, compared with relatively healthy older adults. One meta-analysis compared the effect size in non-frail adults (ES=0.35; 95% CI: 0.17-0.54) to that in frail adults (ES=1.09; 95% CI: 0.55-1.64) and found the effect size was significantly larger in frail adults ($P<.05$)(21). A meta-analysis identified from the updated search reported that the significant benefits of aerobic exercise training to improvements in peak aerobic capacity were observed in both healthy older people (MD=1.72; 95% CI: 0.34 to 3.10; 6 studies) and in those with existing chronic conditions (MD=1.47 (95% CI: 0.60 to 2.34; 4 studies) (22). This aerobic training involved walking, cycling, treadmill walking, walking/running on a mini-trampoline, or combinations of these activities performed 3 times/week over a range of 12 to 26 weeks.

Physical Activity and Physical Function in Older People with Specific Chronic Conditions.

This question builds upon Question 2 by addressing the relationship between physical activity and physical function in discreet populations of older people having selected chronic conditions. The chronic conditions were selected based on their prevalence in older age, as well as on the availability of published research linking physical activity to physical function within each condition (8).

Table 2 shows the level of evidence for the relationship between various types of physical activities and physical function in older people with specific chronic conditions from the 2018 PAGAC Scientific Report. The strongest evidence was observed for the benefits of multicomponent activities among people with frailty and with Parkinson’s disease. Moderate evidence indicated that for community-dwelling older adults who sustain a hip fracture, extended exercise programs (which begin after formal hip fracture rehabilitation ends) was effective for improving physical function and that mobility-oriented physical activity improved walking function for individuals after a stroke.

Cardiovascular Disease.—In the 2018 PAGAC Scientific Report, there was limited evidence suggesting that physical activities such as muscle-strengthening and alternative/complementary exercises (tai chi, qigong, Baduanjin) improved physical function among older people with cardiovascular disease. The updated search identified no additional evidence of this relationship.

Chronic Obstructive Pulmonary Disease (COPD).—Limited evidence from the 2018 Report suggested that tai chi and qigong exercise improved one aspect of physical function (walking ability) in individuals with COPD (36, 37). The updated search identified one SR/MA in people with severe COPD that supported the limited evidence reported in 2018 (38). This meta-analysis of 10 RCTs reported that exercise training improved performance on the 6 minute walk test (6MWT; SMD=3.86; 95% CI: 2.04 to 5.67), compared with a control condition. The exercise training interventions comprised primarily leg exercises, cycling, and walking, with intensity of exercise ranging from 70% to 90% of maximal velocity achieved during incremental testing at baseline.

Cognitive Impairment.—Limited evidence from the 2018 PAGAC Scientific Report suggested that for individuals with cognitive impairment, physical activity programs improved physical function, including ADL measures. Two SR/MA examining the relationship between physical activity and physical function in older people with cognitive impairment were identified in the updated search (39, 40). One of these meta-analyses (39) comprising 43 trials (N=3,988) reported significant differences between supervised exercise training and control conditions on improvements in performance on the 30-second sit-to-stand test (mean difference (MD)=2.1 repetitions; 95% CI: 0.3 to 3.9; 4 trials), step length (MD=5 cm; 95% CI: 2 to 8; 5 trials), Berg Balance Scale (MD=3.6 points; 95% CI: 0.3 to 7.0; 6 trials), functional reach (MD=3.9 cm; 95% CI: 2.2 to 5.5; 6 trials), TUG test (MD=-1 second; 95% CI: -2 to 0; 11 trials), walking speed (MD=0.13 m/s; 95% CI: 0.03 to 0.24; 7 trials), and the 6MWT (MD=50 m; 95% CI: 18 to 81; 7 trials) in this population. Importantly, about 45% of the training programs used multicomponent exercise with a resistance exercise component, while 23% relied on aerobic training.

Frailty.—All of the 15 systematic reviews/meta-analyses included in the 2018 PAGAC Scientific Report cited that physical activity improved some or all measures of physical function in older people with frailty (8). A meta-analysis (41) of 19 RCTs among community-dwelling older adults with frailty reported that overall, physical activity decreased the time needed to walk 10 meters by 1.73 seconds. This has important clinical relevance for older people with frailty, as gait speed is a strong predictor of mortality risk and there is evidence that increments in speed as small as 0.1 m/s significantly lowers that risk (42). The updated search identified five additional systematic reviews that support the strong evidence from the 2018 PAGAC Scientific Report (43–47). Most of these reviewed RCTs and experimental studies examined multicomponent exercise involving resistance training, balance, gait, or endurance training (44–47). In a review of 16 studies involving 1350 frail older adults, Lopez and colleagues (45) reported that resistance training either alone or as part of multicomponent training improved maximal muscle strength between 6.6% and 37%. Similarly, the authors report gains in muscle mass (3.4% to 7.5%), muscle power (8.2%), and functional capacity (4.7% to 58.1%). Moreover, gait speed improved between 5.9% and 14.5%, as did score on the TUG (5.5% to 20.4%).

Hip Fracture.—Moderate evidence from the 2018 PAGAC Scientific Report indicated that older people who have sustained a hip fracture also benefitted from weight-bearing, multicomponent activity (8). The updated search identified only one SR/MA that

investigated this relationship. The meta-analysis of RCTs by Lee, et al., (48) reported that progressive resistance exercise significantly improved overall physical function after hip fracture surgery compared with a control group (SMD= 0.408; 95% CI: 0.238 to 0.578; 8 studies) and it was especially effective in improving mobility (SMD=0.501; 95% CI: 0.297 to 0.705), ADLs (SMD=0.238; 95% CI: 0.040 to 0.437), balance (SMD=0.554; 95% CI: 0.310 to 0.797), lower-limb strength or power (SMD=0.421; 95% CI: 0.101 to 0.741), and performance tasks (SMD=0.841; 95% CI: 0.197 to 1.484).

Osteoporosis/Osteopenia.—Limited evidence from the 2018 PAGAC Scientific Report suggested that muscle-strengthening and agility activities performed on two or more days per week improved physical function in older people who are at risk of fragility fractures due to osteoporosis or osteopenia (8). The updated search identified one SR/MA examining this relationship that adds considerably to the level of evidence. Based on a meta-analysis of 25 RCTs, Varaha, et al, (49) reported that multicomponent exercise significantly improved timed mobility (SMD=- 0.56; 95% CI: - 0.81 to - 0.32), balance (SMD=0.50; 95% CI: 0.27 to 0.74), and self-reported functioning (SMD= - 0.69; 95% CI: - 1.04 to - 0.34) compared with a control condition in 2,113 older people (95% of whom were women). Moreover, the results for multicomponent exercise (10 studies) were more pronounced than those for gait, balance, and functional tasks (4 studies); strength/resistance training (9 studies); or tai chi (5 studies) interventions, suggesting that multicomponent exercise was more effective in improving a broad range of functional outcomes in people with osteoporosis.

Parkinson's Disease.—The 2018 PAGAC Scientific Report cited strong evidence that physical activities, such as aerobic and resistance training, tango dancing, virtual reality training, yoga, and tai chi improved a number of physical function outcomes, including walking, balance, strength, and disease-specific motor scores in older people with Parkinson's disease, with effect sizes ranging from small (ES=0.33; 95% CI: 0.17 to 0.49 for gait velocity) to moderate (ES=0.72; 95% CI:0.08 to 1.36 for the 6 min walk time) (8). Three additional systematic reviews on this relationship were identified in the updated search (50–52), two of which (51, 52) included a meta-analysis. Overall, the new papers supported the findings of the 2018 Report. One small meta-analysis (5 RCTs; N=159 participants) compared dance interventions (Tango or Irish) with other physical activity interventions (3 RCTs) or with no intervention (2 RCTs)(51). Dance practice at least 3 times/week promoted significant improvements in motor scores assessed with the Unified Parkinson's Disease Rating Scale (UPDRS) III (SMD= - 2.52; 95% CI: - 4.59 to - 0.45) and a significant decrease in TUG time (SMD= - 1.15 sec; 95% CI: - 2.03 to - 0.27 sec) compared with other interventions. Dance also improved the UPDRS III score when compared with no intervention (SMD= - 8.35; 95% CI: - 13.79 to - 2.91).

Stroke.—In the 2018 PAGAC Scientific Report, moderate evidence indicated that mobility-oriented physical activity improved walking function for individuals after a stroke and that among stroke survivors, treadmill walking (especially with cadence cuing) improved walking speed by approximately 0.23m/sec (8). Seven systematic reviews relating physical activity to improved physical function in people having had a stroke were identified from the updated search (53–59). Of these reviews, 6 included a meta-analysis (53–58). The

interventions examined included aerobic exercise (54), circuit-based training (53), dual-task balance and mobility training (56), progressive task-oriented exercise (58), and treadmill training (57, 59). Two novel meta-analyses examined the effects of circuit-based exercise (versus other types of exercise or no therapy) on various measures of physical function (53, 55) and both reported that circuit-based training was equal or superior to other forms of therapy in improving measures of gait speed, balance, and functional mobility. In their meta-analysis (N=10 studies; 835 participants), English and colleagues (55) reported that circuit training was superior to comparison interventions in improving walking capacity on the 6MWT (MD=60.86 m; 95% CI: 44.50 to 77.17 m) and gait speed (MD=0.15 m/sec; 95% CI: 0.10 to 0.19 m/sec). Interestingly, these same authors observed no difference in the effectiveness of circuit-training versus a control on walking endurance between stroke survivors who began training within 12 month of their stroke (MD=46.56; 95% CI: 21.35 to 71.77) and those who started training 12 months or more after their stroke (MD=71.15; 95% CI: 49.76 to 92.54). Additionally, superior benefits of circuit training were observed for scores on the TUG test and Activities of Balance Confidence, but not for the Berg Balance Score or the Step Test.

Visual Impairments.—Insufficient evidence was available from either search to determine the effects of physical activity on physical function in older adults with visual impairments.

DISCUSSION

This updated systematic review extends the findings of the 2018 PAGAC Scientific Report by providing new evidence that corroborates the benefits of physical activity to a lower risk of fall-related injuries, as well as to improved physical function among the general older population and among those with selected chronic conditions. The types of activities reviewed in the updated search are similar to those reviewed in the 2018 PAGAC Scientific Report with the addition of different types of balance training interventions, functional exercises that become embedded in everyday life-style activities, Nordic walking, and circuit-based training. Perhaps the most convincing evidence from the 2018 PAGAC Report, along with that from the updated search, relates to the greater benefits of multicomponent, relative to single-component, exercise to the prevention of fall-related injuries and to improvements in physical function in older age. Moreover, multicomponent and multi-task activities that are incorporated into the daily routine may be a promising alternative to structured, single-task exercise programs for older adults.

One in four individuals 65 years falls in the United States every year (60). Furthermore, falls are the leading cause of fatal injury and the most common cause of nonfatal trauma-related hospital admissions among older adults (60). Physical activity programs that emphasize combinations of moderate-intensity balance, strength, endurance, gait, and physical function training appear most effective in reducing the risk of fall-related injuries and fractures in older adults. Thus, the effectiveness of these programs (that were performed in community settings or at home) for risk reduction has significant public health relevance in older age, due to the high prevalence of falls and fall-related injuries and fractures among older adults, as well as the consequent morbidity, disability and reduced quality of life.

Age-related limitations in physical function are prevalent in older adults. The National Health Interview Survey ascertained the prevalence of physical limitations in 2001-2007, with limitations defined as great difficulty doing (or inability to do) basic tasks of life (e.g., walk a quarter of a mile, lift a 10-pound bag of groceries)(61). At that time, 22.9% of older adults ages 60-69 years and 42.9% of adults ages 80 years reported functional limitations. Older adults with lower levels of physical function generally have higher health care expenditures (60). In addition, about 80% of adults 60 years of age have at least one chronic condition, and 77% have at least two. Moreover, approximately 20 to 30% of adults older than age 65 years suffer from either mild cognitive impairment or dementia (60). Chronic diseases account for 75% of health care spending in the United States (60). Low levels of daily physical activity often co-exist with chronic disease, thereby accelerating the risk of functional decline, disability, and mortality. Ample evidence now indicates that physical inactivity is among the strongest predictors of physical disability in older people (4). Aerobic, muscle-strengthening, and multicomponent physical activity appear to have the strongest relationship to improvements in physical function in the general aging population, as well as among those with chronic conditions. Thus, such activities may delay or improve mobility disability, frailty, and loss of independence in aging.

Both the 2018 Scientific Report (8) and this umbrella review underscore the need for future research in several areas. For example, the relationship between the minimal effective dose of activity (150 min/week of moderate intensity activity) and health has been described in detail (3, 8). There still is, however, a need to examine greater volumes and intensities of physical activity to establish safety thresholds for older people –especially for those with pre-existing conditions or limitations. To accomplish this, studies need to examine several levels of activity and to monitor and report adverse events. Also, the feasibility and benefits of alternative and complimentary activities, rigorous multi-task activities, as well as novel interventions that integrate “functional exercises” into everyday tasks, need to be examined further. Moreover, the effectiveness of any new intervention needs to be examined across different socioeconomic strata to address existing disparities in prevention strategies and in health among older people . Given the rapidly increasing trends in aging demographics in the United States, preventing or delaying fall-related injuries and loss of physical function and mobility has important public health benefits, and this may be especially so for older people with already established chronic conditions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the contributions of Sarah Prowitt, MPH (HHS) and Kyle Sprow, MPH (National Cancer Institute, National Institutes of Health) for management support; Anne Brown Rodgers, HHS consultant for technical writing support; and ICF librarians, abstractors, and additional support staff.

Role of the Funder/Sponsor

HHS staff provided general administrative support to the Committee and assured that the Committee adhered to the requirements for Federal Advisory Committees. HHS also contracted with ICF, a global consulting services company, to provide technical support for the literature searches conducted by the Committee. HHS and ICF staff

collaborated with the Committee in the design and conduct of the searches by assisting with the development of the analytical frameworks, inclusion/exclusion criteria, and search terms for each primary question; using those parameters, ICF performed the literature searches.

REFERENCES

1. Administration on Aging, US Department of Health and Human Services. A profile of older Americans: 2016. Washington, DC: US Department of Health and Human Services; 2016.
2. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public health reports (Washington, DC : 1974)*. 1985;100(2):126–131.
3. Physical Activity Guidelines Advisory Committee. *Physical Activity Guidelines Advisory Committee Report*, 2008. Washington, DC: US Department of Health and Human Services; 2008 <https://health.gov/paguidelines/guidelines/report.aspx>. Published 2008. Accessed January 4, 2018.
4. Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA*. 2014;311(23):2387–2396. doi:10.1001/jama.2014.5616. [PubMed: 24866862]
5. US Department of Health and Human Services. Physical activity guidelines for Americans. 2nd Edition Washington, DC: US DHHS, 2018.
6. Keadle S, McKinnon R, Graubard BI, Troiano RP. Prevalence and trends in physical activity among older adults in the United States: A comparison across three national surveys. *Prev Med*. 8 2016;89:37–43. doi:10.1016/j.ypmed.2016.05.009. [PubMed: 27196146]
7. Lee I-M, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT, for the Lancet Physical Activity Series Working Group. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012; 7 18, 2012 10.1016/S0140-6736(12)61031-9.
8. Physical Activity Guidelines Advisory Committee. Physical Activity Guidelines Advisory Committee Report, 2018. Washington, DC: US Department of Health and Human Services; 2008 <https://health.gov/paguidelines/guidelines/report.aspx>. Published 2018. Accessed January 4, 2018.
9. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, et al. (2009) The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLOS Medicine* 6(7): e1000100 10.1371/journal.pmed.1000100 [PubMed: 19621070]
10. Johnson BT, MacDonald HV, Bruneau ML Jr, et al. Methodological quality of meta-analyses on the blood pressure response to exercise: a review. *J Hypertens*. 2014;32(4):706–723. doi:10.1097/HJH.000000000000097. [PubMed: 24463936]
11. U.S. Department of Agriculture (USDA). 2015 Dietary Guidelines Advisory Committee (DGAC) nutrition evidence library methodology. 2017 https://www.cnpp.usda.gov/sites/default/files/usda_nutrition_evidence_library/2015DGAC-SR-Methods.pdf. Accessed January 16, 2018.
12. El-Khoury F, Cassou B, Charles MA, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2013;347:f6234. doi: 10.1136/bmj.f6234. [PubMed: 24169944]
13. Gillespie LD, Robertson M, Gillespie WJ, Sherrington C, Gates S, Clemson LM, Lamb SE. Interventions for preventing falls in older people living in the community. *Cochrane Database of Systematic Reviews*. 2012;9:CD007146. doi:10.1002/14651858.CD007146.pub3.
14. Zhao R, Feng F, Wang X. Exercise interventions and prevention of fall-related fractures in older people: a meta-analysis of randomized controlled trials. *Int J Epidemiol*. 2016. doi: 10.1093/ije/dyw142.
15. Health Quality Ontario. Prevention of falls and fall-related injuries in community-dwelling seniors: an evidence-based analysis. *Ont Health Technol Assess Ser*. 2008;8:1–78.
16. Cauley JA, Harrison SL, Cawthon PM, Ensrud KE, Danielson ME, Orwoll E, Mackey DC. Objective measures of physical activity, fractures and falls: the osteoporotic fractures in men study. *J Am Geriatr Soc*. 2013;61:1080–1088. doi:10.1111/jgs.12326. [PubMed: 23855842]

17. Heesch KC, Byles JE, Brown WJ. Prospective association between physical activity and falls in community-dwelling older women. *J Epidemiol Community Health*. 2008;62:421–426. [PubMed: 18413455]
18. Iinattiniemi S, Jokelainen J, Luukinen H. Exercise and risk of injurious fall in home-dwelling elderly. *Int J Circumpolar Health*. 2008;67:235–244. [PubMed: 18767343]
19. Peel NM, McClure RJ, Hendrikz JK. Health-protective behaviours and risk of fall-related hip fractures: a population-based case-control study. *Age Ageing*. 2006;35:491–497. [PubMed: 16772364]
20. Hill KDS, Lin P, Tsang SI, Ashari WNW, Hamid A, Farrier TAA, Burton K. What works in falls prevention in Asia: a systematic review and meta-analysis of randomized controlled trials. *BMC Geriatrics*. 2018; 18: 3.2018208.
21. Chase JD, Phillips LJ, Brown M. Physical activity intervention effects on physical function among community-dwelling older adults: a systematic review and meta-analysis. *J Aging Phys Act*. 2017;25(1):149–170. doi:10.1123/japa.2016-0040. [PubMed: 27620705]
22. Bouaziz W, Kanagaratnam L, Vogel T, Schmitt E, Dramé M, Kaltenbach G, Geny B Lang PO. Effect of Aerobic Training on Peak Oxygen Uptake Among Seniors Aged 70 or Older: A Meta-Analysis of Randomized Controlled Trials. *Rejuvenation Research*. 2018; 10.1089/rej.2017.1988.2018.
23. Bullo V, Gobbo S, Vendramin B, Duregon F, Cugusi L, Di Blasio A, Bocalini DS, Zaccaria M, Bergamin M, and Ermolao A. Nordic walking can be incorporated in the exercise prescription to increase aerobic capacity, strength, and quality of life for elderly: a systematic review and meta-analysis. *Rejuvenation Research* 2018;10.1089/rej.2017.1921.2018;
24. Farlie MK, Robins L, Haas R, Keating JL, Molloy E, and Haines TP. Programme frequency, type, time and duration do not explain the effects of balance exercise in older adults: a systematic review with a meta-regression analysis. *BJSM*. 2018; 10.1136/bjsports-2016-096874.
25. Howe SC, Charles DK, Marley J, Pedlow K, McDonough SM. Gaming for health: systematic review and meta-analysis of the physical and cognitive effects of active computer gaming in older adults. *Physical Therapy*. 2017; 97: 1122–1137.017. [PubMed: 29077911]
26. Lai CC, Tu Y-K, Wang T-G, Y-T Huang Y-T, and Chien K-L. Effects of resistance training, endurance training and whole-body vibration on lean body mass, muscle strength and physical performance in older people: a systematic review and network meta-analysis. *Age and Ageing* 2018; 47: 367–373. [PubMed: 29471456]
27. Liu C-j, Chang W-P, Araujo de Carvalho I, Savage KEL, Radford LW and Thiyagarajan JA. Effects of physical exercise in older adults with reduced physical capacity: meta-analysis of resistance exercise and multimodal exercise. *International Journal of Rehabilitation Research* 2017; 40:303–314. [PubMed: 29023317]
28. Papa E, Dong X, and Hassan M. Resistance training for activity limitations in older adults with skeletal muscle function deficits: a systematic review. *Clinical Interventions in Aging* 2017;12 955–961. [PubMed: 28670114]
29. Roberts LH, Cooper CL, Gray S, Allan JL. Effect of different types of physical activity on activities of daily living in older adults: systematic review and meta-analysis. *Journal Aging Phys Act* 2017; 25: 653–670.2017.
30. Weber M, Belala N, Clemson L, Boulton E, Hawley-Hague H, Becker C, and Schwenk M. Feasibility and effectiveness of intervention programmes integrating functional exercise into daily life of older adults: a systematic review. *Gerontology* 2018;64:172–187; doi: 10.1159/000479965 2018 [PubMed: 28910814]
31. Paterson DH, Warburton DE. Physical activity and functional limitations in older adults: a systematic review related to Canada's Physical Activity Guidelines. *Int J Behav Nutr Phys Act*. 2010;7:38. doi:10.1186/1479-5868-7-38. [PubMed: 20459782]
32. Lesinski M, Hortobágyi T, Muehlbauer T, Gollhofer A, Granacher U. Effects of balance training on balance performance in healthy older adults: a systematic review and meta-analysis. *Sports Med*. 2015;45(12):1721–1738. doi:10.1007/s40279-015-0375-y. [PubMed: 26325622]

33. Clemsen L, et al.: Integration of balance and strength training into daily life activity to reduce rate of falls in older people (the LiFE study): randomised parallel trial. *BMJ* 2012;345: e4547.2012. [PubMed: 22872695]
34. Lally P, Gardner B. Promoting habit formation. *Health Psychol Rev* 2013; 7: 137–158.
35. Tak E, Kuiper R, Chorus A, Hopman-Rock M. Prevention of onset and progression of basic ADL disability by physical activity in community dwelling older adults: a meta-analysis. *Ageing Res Rev.* 2013;12(1):329–338. doi:10.1016/j.arr.2012.10.001. [PubMed: 23063488]
36. Ding M, Zhang W, Li K, Chen X. Effectiveness of t'ai chi and qigong on chronic obstructive pulmonary disease: a systematic review and meta-analysis. *J Altern Complement Med.* 2014;20(2):79–86. doi:10.1089/acm.2013.0087. [PubMed: 23961940]
37. Ngai SPC, Jones AYM, Tam W. Tai chi for chronic obstructive pulmonary disease (COPD). *Cochrane Database Syst Rev.* 2016;(6):Cd009953. doi:10.1002/14651858.CD009953.pub2. [PubMed: 27272131]
38. Paneroni M, Simonelli C, Vitacca M, and Ambrosino N. Aerobic exercise training in very severe chronic obstructive pulmonary disease. A systematic review and meta-analysis. *Am J Phys Med Rehabil* 2017;96:541–548. [PubMed: 28099192]
39. Lam FMH, Huang M-Z, Liao L-R, Chung RCK, Kwok TCY, Pang MYC. Physical exercise improves strength, balance, mobility, and endurance in people with cognitive impairment and dementia: a systematic review. *Journal of Physiotherapy* 2018; 64: 4–15. [PubMed: 29289581]
40. Goncalves A-C, Cruz J, Marques A, Demain S, Samuel D. Evaluating physical activity in dementia: a systematic review of outcomes to inform the development of a core outcome set. *Age and Ageing* 2018; 47: 34–41 2018 [PubMed: 28985262]
41. Giné-Garriga M, Roqué-Fíguls M, Coll-Planas L, Sitjà-Rabert M, Salvà A. Physical exercise interventions for improving performance-based measures of physical function in community-dwelling, frail older adults: a systematic review and meta-analysis. *Arch Phys Med Rehabil.* 2014;95(4):753–769.e3. doi:10.1016/j.apmr.2013.11.007. [PubMed: 24291597]
42. Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *JAMA.* 2011;305(1): 50–58. doi:10.1001/jama.2010.1923. [PubMed: 21205966]
43. Apostolo J, Cooke R, Bobrowicz-Campos E, Santana S, Marcucci M, Cano A, Vollenbroek-Hutten M, Germini F, D'Avanzo B, Gwyther H, Holland C. Effectiveness of interventions to prevent pre-frailty and frailty progression in older adults: a systematic review. *JBIC Database of Systematic Reviews and Implementation Reports* 2018;16: 140–232. [PubMed: 29324562]
44. Dedyne L, Deschodt M, Verschuere S, Tournoy J, Gielen E. Effects of multi-domain interventions in (pre)frail elderly on frailty, functional, and cognitive status: a systematic review. *Clinical Interventions in Aging* 2017;12: 873–896. [PubMed: 28579766]
45. Lopez P, Silveira Pinto R, Radaelli R, Rech A, Grazioli R, Izquierdo M, Lusa Cadore E. Benefits of resistance training in physically frail elderly: a systematic review. *Aging Clinical Experimental Research.* 2017 10.1007/s40520-017-0863-z.
46. Lopez P, Izquierdo M, Radaelli R, Shurzzi G, Grazioli R, Silveira Pinto R, and Lusa Cadore E. Effectiveness of multimodal training on functional capacity in frail older people: a meta-analysis of randomized controls trials. *J Aging Phys Act* 10.1123/japa.2017-01881-36
47. Lozano-Monyoya I, Correa-Pérez A, Abraha I, Soiza RL, Cherubini A, O'Mahony D, Cruz-Jentoft AJ. Nonpharmacological interventions to treat physical frailty and sarcopenia in older patients: a systematic overview – the SENATOR Project ONTOP Series. *Clinical Interventions in Aging* 2017;12: 721–740. [PubMed: 28490866]
48. Lee SY, Yoon B-H, Beom J, Ha Y-C, Lim J-Y. Effect of lower-limb progressive resistance exercise after hip fracture surgery: a systematic review and meta-analysis of randomized controlled studies. *JAMDA* 2017;18:1096.e19–1096.e26.
49. Varaha A, Rodrigues JB, MacDermid JC, Bryant D, Birmingham T. Exercise to improve functional outcomes in persons with osteoporosis: a systematic review and meta-analysis. *Osteoporosis International* 2018; 29:265–286. [PubMed: 29306984]
50. Cwiekala-Lewis KJ, Gallek M, Taylor-Piliae RE. The effects of Tai Chi on physical function and well-being among persons with Parkinson's Disease: A systematic review. *J Bodywork Movement Therapies* 2017; 21:414–421.

51. dos Santos Delabary M, Komerowski IG, Monteiro EP, Costa RR, Haas AN. Effects of dance practice on functional mobility, motor symptoms and quality of life in people with Parkinson's disease: a systematic review with meta-analysis. *Aging Clin Exp Res* 2017; DOI 10.1007/s40520-017-0836-2.
52. Mazzarin CM, Valderramas SR, de Paula Ferreira M, Tiepolo E, Guérios L, Parisotto D, Israel VL. Effects of dance and of tai chi on functional mobility, balance, and agility in Parkinson disease. a systematic review and meta-analysis. *Topics in Geriatric Rehabilitation* 2017;33:262–272.
53. Bonini-Roche AC, Souza de Andrade AL, Moraes AM, Matheus LBG, Rios Diniz L, Martins WR. Effectiveness of circuit-based exercises on gait speed, balance, and functional mobility in people affected by stroke: a meta-analysis. *PM R* 2018;10:398–409. [PubMed: 29111465]
54. Boyne P, Welge J, Kissela B, Dunning K. Factors influencing the efficacy of aerobic exercise for improving fitness and walking capacity after stroke: a meta-analysis with meta-regression. *Archives Physical Medicine Rehabilitation* 2017;98:581–595.
55. English C, Hillier S, Lynch E. Circuit class therapy for improving mobility after stroke. *Stroke* 2017;48:e275–e276. DOI: 10.1161/STROKEAHA.117.018601.
56. He Y, Yang L, Zhou J, Yao L, Pang MYC. Dual-task training effects on motor and cognitive functional abilities in individuals with stroke: a systematic review. *Clinical Rehabilitation* 2018; 10.1177/026921551875848.
57. Mehrhotz J, Pohl M, Elsner B. Treadmill training and body weight support for walking after stroke. *Cochrane Database of Systematic Reviews* 2014, Issue 1 Art. No.: CD002840. DOI: 10.1002/14651858.CD002840.pub3.
58. Stretton CM, Mudge S, Kayes NM, and McPherson KM. Interventions to improve real-world walking after stroke: a systematic review and meta-analysis. *Clinical Rehabilitation* 2017;31:310–318. [PubMed: 27056251]
59. Tally Z, Boetefuer L, Kauk C, Perez G, Schrand L, and Hoder J. The efficacy of treadmill training on balance dysfunction in individuals with chronic stroke: a systematic review. *Topics in Stroke Rehabilitation* 2017; 24:539–546; DOI: 10.1080/10749357.2017.13454452017. [PubMed: 28687056]
60. National Council on Aging. Fact sheet: healthy aging. <https://www.ncoa.org/news/resources-for-reporters/get-the-facts/healthy-aging-facts>. Accessed January 4, 2018 Arlington, VA: National Council on Aging; 2016.
61. Holmes J, Powell-Griner E, Lethbridge-Cejku M, Heyman K. Aging differently: physical limitations among adults aged 50 years and over: United States, 2001–2007. *NCHS Data Brief*. 7 2009;(20):1–8.




Figure 1. Risk of injuries from falls: systematic reviews, meta-analyses, pooled analyses, and report flow diagram of search strategy and study selection from the updated search, March 2018.




Figure 2. Physical function systematic review, meta-analyses, pooled analysis, and report flow diagram of search strategy and study selection from the updated search, March 2018.




Figure 3.

Results of a systematic review of the inverse dose-response relationship between volume of aerobic physical activity and risk of physical functional limitations in the general aging population. Categories of physical activity level were derived from prospective cohort studies with covariate adjustment: 0)=no activity; 1) = light activities only occasion walking or gardening; 2) = moderate level of activity (volume = 3-5 days/week for 30 min/day; and 3) = vigorous activities and/or high volume of systematic activity. The odds ratio is the odds of disability in the physical activity group relative to a comparison group. ADL=activities of daily living; IADL=instrumental activities of daily living; QOL=quality of life score. Figure reproduced with permission from Paterson & Warburton, 2010.

Table 1-

List of questions addressed by the Aging Subcommittee of the 2018 Physical Activity Guidelines Advisory Committee.

1	<p>What is the relationship between physical activity and risk of injury due to a fall?</p> <ul style="list-style-type: none"> a. Is there a dose-response relationship? If yes, what is the shape of the relationship? b. Does the relationship vary by age, sex, race/ethnicity, socioeconomic status, or weight status? c. What type(s) of physical activity are effective for preventing injuries due to a fall? d. What factors (e.g., level of physical function, existing gait disability) modify the relationship between physical activity and risk of injury due to a fall?
2	<p>What is the relationship between physical activity and physical function among the general aging population?</p> <ul style="list-style-type: none"> a. Is there a dose-response relationship? If yes, what is the shape of the relationship? b. Does the relationship vary by age, sex, race/ethnicity, socioeconomic status, or weight status? c. What type(s) of physical activity (single component, dual task, multicomponent) are effective for improving or maintaining physical function among the general aging population? d. What impairment(s) (e.g., visual impairment, cognitive impairment, physical impairment) modify the relationship between physical activity and physical function among the general aging population?
3	<p>What is the relationship between physical activity and physical function in older adults with selected chronic conditions? These conditions are: 1) cardiovascular disease; 2) chronic obstructive pulmonary disease (COPD); 3) cognitive impairment; 4) frailty; 5) hip fracture; 6) osteoporosis and osteopenia; 7) Parkinson's disease; 8) stroke; and 9) visual impairment.</p>

Table 2 –

Summary of the level of evidence for the relationship between physical activity, fall-related injuries, and physical function in older people: 2018 PAGAC Scientific Report

	Overall Evidence	Dose-Response	Type of Activity Reported	Effect Modification by Demographic Factors, Weight, or other Functional factors
Fall-related Injury	STRONG	LIMITED (MVPA)	MODERATE (multicomponent)	NOT ASSIGNABLE
Physical Function in the General Aging Population	STRONG	STRONG (aerobic) LIMITED (strength and balance)	STRONG (aerobic, strength, multicomponent) MODERATE (balance) LIMITED (tai chi, dance, AVG, dual-task, functional) NOT ASSIGNABLE (flexibility, yoga, qigong)	LIMITED/NOT ASSIGNABLE
Physical Function in Older People with Specific Chronic Conditions*				
• Cardiovascular Disease	LIMITED		Muscle-strengthening, tai chi, qigong, aerobic	
• COPD	LIMITED		Tia chi, qigong, walking, cycling, leg exercises	
• Cognitive Impairment	LIMITED		Supervised multicomponent	
• Frailty	STRONG		Multicomponent	
• Hip Fracture	MODERATE		Weight-bearing multicomponent	
• Osteoporosis	LIMITED		Muscle-strengthening, multicomponent	
• Parkinson's Disease	STRONG		Aerobic, resistance, dance, VRT, yoga, tai chi	
• Stroke	MODERATE		Mobility-oriented, treadmill walking	
• Visual Impairment	INSUFFICIENT			

MVPA= moderate-to-vigorous intensity physical activity; AVG=active video gaming; VRT=virtual reality training. The activities in parentheses refer to the level of evidence. For example among the general aging population, we found strong evidence linking aerobic, strength, and multicomponent activities to improvements in physical function.

* Question 3 did not examine dose-response or effect modification.