

<https://doi.org/10.1038/s41467-019-10052-z>

# Author Correction: PBX3 is targeted by multiple miRNAs and is essential for liver tumour-initiating cells

Haibo Han<sup>1</sup>, Yantao Du<sup>1</sup>, Wei Zhao<sup>1</sup>, Sheng Li<sup>1</sup>, Dongji Chen<sup>1</sup>, Jing Zhang<sup>2</sup>, Jiang Liu<sup>2</sup>, Zhenhe Suo<sup>3</sup>, Xiuwu Bian<sup>4</sup>, Baocai Xing<sup>5</sup> & Zhiqian Zhang<sup>1</sup>

Correction to: *Nature Communications* <https://doi.org/10.1038/ncomms9271>, published online 30 September 2015

This Article contains an error in Figure 3. In panel g, images representing miR-222-TuD and miR-424-TuD were both taken from the miR-222-TuD image. The correct version of Figure 3 is shown below as Fig. 1.

Published online: 21 May 2019

---

<sup>1</sup> Department of Cell Biology, Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Center for Molecular and Translational Medicine, Peking University Cancer Hospital and Institute, 52 Fucheng Road, Beijing 100142, China. <sup>2</sup> Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China. <sup>3</sup> Department of Pathology, Oslo University Hospital, Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo 0379, Norway. <sup>4</sup> Institute of Pathology and Southwest Cancer Center, Southwest Hospital, Third Military Medical University, Chongqing 400038, China. <sup>5</sup> Department of Hepatobiliary Surgery I, Peking University Cancer Hospital and Institute, Beijing 100142, China. Correspondence and requests for materials should be addressed to Z.Z. (email: [zlzqzhang@bjmu.edu.cn](mailto:zlzqzhang@bjmu.edu.cn))




Fig. 1