



# HHS Public Access

Author manuscript

*Cell Stem Cell*. Author manuscript; available in PMC 2019 May 22.

Published in final edited form as:

*Cell Stem Cell*. 2019 May 02; 24(5): 829. doi:10.1016/j.stem.2019.04.002.

## Differentiation of Human Pluripotent Stem Cells into Colonic Organoids via Transient Activation of BMP Signaling

Jorge O. Múnera, Nambirajan Sundaram, Scott A. Rankin, David Hill, Carey Watson, Maxime Mahe, Jefferson E. Vallance, Noah F. Shroyer, Katie L. Sinagoga, Adrian Zarzoso-Lacoste, Jonathan R. Hudson, Jonathan C. Howell, Praneet Chatuvedi, Jason R. Spence, John M. Shannon, Aaron M. Zorn, Michael A. Helmrath, and James M. Wells\*

---

In the STAR Methods section of our manuscript, in the last sentence of the “Generation of human gut tube cultures” subsection, we mistakenly stated, “DE was then treated with hindgut induction medium (RPMI 1640, 2 mM L-glutamine, 2% decomplexed FBS, penicillin-streptomycin, and 100 ng/mL Activin A) for 4 days with 500 ng/mL FGF4 (R&D) and 3  $\mu$ M Chiron 99021 (Tocris) to induce formation of mid-hindgut spheroids.” However, the hindgut induction medium did not include 100 ng/mL Activin A, but rather, non-essential amino acids instead. We apologize for the oversight in this description.

---

\*Correspondence: james.wells@cchmc.org.