

- Respirology* 2008; **13**(1):128-33.
14. Meijer RJ, Postma DS, Kauffman HF, Arends LR, Koeter GH, Kerstjens HA. Accuracy of eosinophils and eosinophil cationic protein to predict steroid improvement in asthma. *Clin Exp Allergy* 2002; **32**(7):1096-103. <http://dx.doi.org/10.1046/j.1365-2222.2002.01412.x>
 15. Nowak D. Management of asthma with anti-immunoglobulin E: a review of clinical trials of omalizumab. *Respir Med* 2006; **100**(11):1907-17. <http://dx.doi.org/10.1016/j.rmed.2005.10.004>
 16. Smith AD, Cowan JO, Brasset KP, et al. Exhaled nitric oxide: a predictor of steroid response. *Am J Respir Crit Care Med* 2005; **172**(4):453-9. <http://dx.doi.org/10.1164/rccm.200411-1498OC>
 17. Dahl M, Vestbo J, Lange P, Bojesen SE, Tybjaerg-Hansen A, Nordestgaard BG. C-reactive protein as a predictor of prognosis in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2007; **175**(3):250-5. <http://dx.doi.org/10.1164/rccm.200605-713OC>
 18. de Torres JP, Pinto-Plata V, Casanova C, et al. C-reactive protein levels and survival in patients with moderate to very severe COPD. *Chest* 2008; **133**(6):1336-43. <http://dx.doi.org/10.1378/chest.07-2433>
 19. Gelb AF, Flynn TC, Shinar CM, Gutierrez C, Zamel N. Role of spirometry and exhaled nitric oxide to predict exacerbations in treated asthmatics. *Chest* 2006; **129**(6):1492-9. <http://dx.doi.org/10.1378/chest.129.6.1492>
 20. Green RH, Brightling CE, McKenna S, et al. Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet* 2002; **360**(9347):1715-21. [http://dx.doi.org/10.1016/S0140-6736\(02\)11679-5](http://dx.doi.org/10.1016/S0140-6736(02)11679-5)
 21. ATS/ERS recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide, 2005. *Am J Respir Crit Care Med* 2005; **171**(8):912-30. <http://dx.doi.org/10.1164/rccm.200406-710ST>
 22. Deykin A, Halpern O, Massaro AF, Drazen JM, Israel E. Expired nitric oxide after bronchoprovocation and repeated spirometry in patients with asthma. *Am J Respir*
 23. Schneider A, Tilemann L, Schermer T, et al. Diagnosing asthma in general practice with portable exhaled nitric oxide measurement: results of a prospective diagnostic study. *Respir Res* 2009; **10**:15. <http://dx.doi.org/10.1186/1465-9921-10-15>
 24. American Association for Respiratory Care. AARC Clinical Practice Guideline. Body plethysmography: 2001 revision and update. *Respir Care* 2001; **46**:506-13.
 25. Crapo RO, Casaburi R, Coates AL, et al. Guidelines for methacholine and exercise challenge testing—1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med* 2000; **161**(1):309-29.
 26. Papi A, Romagnoli M, Baraldo S, et al. Partial reversibility of airflow limitation and increased exhaled NO and sputum eosinophilia in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000; **162**(5):1773-7.
 27. Fabbri LM, Romagnoli M, Corbetta L, et al. Differences in airway inflammation in patients with fixed airflow obstruction due to asthma or chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003; **167**(3):418-24. <http://dx.doi.org/10.1164/rccm.200203-183OC>
 28. Gan WQ, Man SF, Sin DD. The interactions between cigarette smoking and reduced lung function on systemic inflammation. *Chest* 2005; **127**(2):558-64. <http://dx.doi.org/10.1378/chest.127.2.558>
 29. Pinto-Plata VM, Mullerova H, Toso JF, et al. C-reactive protein in patients with COPD, control smokers and non-smokers. *Thorax* 2006; **61**(1):23-8. <http://dx.doi.org/10.1136/thx.2005.042200>
 30. Arif AA, Delclos GL, Colmer-Hamood J. Association between asthma, asthma symptoms and C-reactive protein in US adults: data from the National Health and Nutrition Examination Survey, 1999-2002. *Respirology* 2007; **12**(5):675-82. <http://dx.doi.org/10.1111/j.1440-1843.2007.01122.x>
 31. Price D, Berg J, Lindgren P. An economic evaluation of NIOX MINO airway inflammation monitor in the United Kingdom. *Allergy* 2009; **64**:431-8. <http://dx.doi.org/10.1111/j.1365-9953.2008.01855.x>

Available online at <http://www.thepcrj.org>

CORRIGENDUM

Corrigendum to 'Asthma incidence and accuracy of diagnosis in the Portuguese sentinel practice network' (*Prim Care Resp J* 2010; **19**(4):352-357)

Jaime Correia de Sousa, Manuel Luciano Silva, Filipa Almada Lobo, John Yaphe

The authors regret that two errors occurred on pages 354 and 356 of this article. These have already been corrected in the online version of the paper.

The first paragraph of the Discussion section on pg 354 reads:

"It also found that the accuracy of the diagnosis by family physicians/GPs was 62% using accepted diagnostic criteria."

- but it should read as follows:

"It also found that the accuracy of the diagnosis by family physicians/GPs was 84% using accepted diagnostic criteria."

The bottom of pg 356, in the section referring to diagnostic accuracy, reads:

"A incidence of 2 new cases per 1000 persons per year with a diagnostic accuracy of 62% suggests that there is considerable under-diagnosis of a common, treatable condition."

- but it should read as follows:

"An incidence of 2 new cases per 1000 persons per year with a diagnostic accuracy of 84% suggests that there is considerable under-diagnosis of a common, treatable condition."

* Corresponding author: Dr Jaime Correia de Sousa, University of Minho, School of Health Sciences, Gualtar Campus, Braga, 4710-057, Portugal
Tel: 351 253 604823 Fax: 351 253 604862 E-mail: jaimecsousa@gmail.com

DOI of original article: <http://dx.doi.org/10.4104/pcrj.2010.00022>