

Mitochondrial dysfunction and its role in tissue-specific cellular stress

David Pacheu-Grau^{1,*}, Robert Rucktäschel¹ and Markus Deckers^{1,*}

¹ Department of Cellular Biochemistry, University Medical Center Göttingen, Germany.

* Corresponding Authors:

David Pacheu-Grau, University Medical Center Göttingen, Department of Cellular Biochemistry, Humboldtallee 23, 37073 Göttingen, Germany. Phone: +49-(0)551-394571; E-mail: David.Pacheu-Grau@med.uni-goettingen.de;

Markus Deckers, University Medical Center Göttingen, Department of Cellular Biochemistry, Humboldtallee 23, 37073 Göttingen, Germany. Phone: +49-(0)551-395983; E-mail: Markus.Deckers@medizin.uni-goettingen.de

ABSTRACT Mitochondrial bioenergetics require the coordination of two different and independent genomes. Mutations in either genome will affect mitochondrial functionality and produce different sources of cellular stress. Depending on the kind of defect and stress, different tissues and organs will be affected, leading to diverse pathological conditions. There is no curative therapy for mitochondrial diseases, nevertheless, there are strategies described that fight the various stress forms caused by the malfunctioning organelles. Here, we will revise the main kinds of stress generated by mutations in mitochondrial genes and outline several ways of fighting this stress.

doi: 10.15698/cst2018.07.147

Received originally: 26.04.2018

in revised form: 13.06.2018,

Accepted 14.06.2018,

Published 13.07.2018.

Keywords: mitochondrial dysfunction, cellular stress, mitochondrial pathology, therapy.

Abbreviations:

ADOA – autosomal dominant optic atrophy,

ARO – autosomal recessive optic atrophy,

ARS – aminoacyl-tRNA synthetase,

CL – cardiolipin,

CRISPR – clustered regularly

interspaced short palindromic repeats,

LHON – Leber's hereditary optic neuropathy,

mt - mitochondrial

OXPPOS – oxidative phosphorylation,

ROS – reactive oxygen species.

MITOCHONDRIA AND CELL METABOLISM

Mitochondria play a pivotal role in eukaryotic metabolism. They catabolise redox equivalents, derived from nutrient uptake, and use them to provide the bulk of cellular energy in the form of ATP. The oxidative phosphorylation system (OXPPOS) is responsible for this energy production and it is composed of five multi-oligomeric complexes present in the inner mitochondrial membrane. Transfer of electrons through complexes I to IV reduce molecular oxygen to water. This process is coupled to proton pumping from the matrix to the intermembrane space (IMS), while the return of protons to the matrix through the F1Fo ATPase generates ATP [1]. However, an inefficient flow of electrons through the respiratory chain complexes would partially reduce oxygen and produce reactive oxygen species (ROS)

like superoxide and hydrogen peroxide. At low concentrations, these molecules act as second messengers and can activate gene transcription and trigger cellular responses, like cellular growth, production of cellular antioxidants or stimulation of mitochondrial biogenesis [2, 3]. However, once a certain threshold is exceeded, these molecules may incite oxidative damage in the form of mitochondrial DNA (mtDNA) alterations or lipid peroxidation, generating cellular stress that leads to aging or cell death.

In addition, mitochondria are involved in many other key cellular functions. Dissipation of the proton gradient by uncoupling proteins (UCPs) generates heat instead of energy and this plays an important role in exposure to cold or hibernation [4]. Calcium (Ca²⁺) uptake inside mitochondria is mediated by the mitochondrial calcium uniporter (MCU).

Although the complex has a low affinity for Ca^{2+} , the transport takes place due to the high concentration of Ca^{2+} ($>10 \mu\text{M}$) present in micro domains located in the contact sites between endoplasmic reticulum (ER) and mitochondria [5]. The mitochondrial Ca^{2+} uptake not only shapes the cytosolic Ca^{2+} dynamics, which is crucial for muscle contraction, exocytosis and gene transcription, but also modulates at least three dehydrogenases of the Krebs cycle, thus regulating energy metabolism. Finally, Ca^{2+} overload in mitochondria regulates apoptosis due to formation of the permeable transition pore (PTP) and release of cytochrome *c* from the IMS [6]. Mitochondria are involved in the biogenesis and maturation of different cofactors, like heme, biotin or iron-sulfur (Fe/S) clusters. Despite the chemical simplicity of Fe/S clusters, their biosynthesis requires more than two dozen proteins in eukaryotes and takes place both in mitochondria and the cytosol [7]. Alterations in these mechanisms are linked to severe neurodegenerative, metabolic or haematological diseases [8].

Since mitochondria take part in many different metabolic processes, mitochondrial malfunction can affect numerous aspects of the cell. As a consequence, various forms of cellular stress are generated, leading to a large variety of pathological conditions. Here, we review different forms of cellular stress caused by mitochondrial malfunction and the strategies used to fight this stress.

MITOCHONDRIAL DEFECTS

Mitochondria have retained their own genome, the mtDNA. This small, circular, double-stranded DNA is located in the mitochondrial matrix in all cell types, and can be found with copy numbers that range from several to thousands of copies. In human, the mtDNA encodes 13 polypeptides of the respiratory chain, as well as for part of the translation machinery, required for the synthesis of these polypeptides within mitochondria: two ribosomal RNAs (mt-rRNAs) and 22 transfer RNAs (mt-tRNAs) [9]. The remaining mitochondrial proteins (approx. 99%) are encoded in the nucleus, synthesized on the cytosolic ribosomes and imported into mitochondria. Therefore, we will distinguish between mitochondrial malfunction caused by mutations in the mtDNA and those caused by mutations of nuclear genes encoding mitochondrial proteins.

Alterations in the mtDNA

Some features of mtDNA make it especially sensitive to oxidative damage and mutation. Firstly, mtDNA has no introns, so every single nucleotide carries information essential for protein coding; mtDNA is naked, there are no histone proteins protecting it from damage; and although DNA repair systems do exist in mitochondria, their mechanisms and extent are poorly understood [10], therefore mutations usually remain and are transmitted to the next generation until they are removed by selection [11]. Moreover, the proximity to the respiratory chain, a ROS producing source, increases the risk of potential damage. For all these reasons, the mutational rate of the mitochondrial

genome is much higher than that of the nuclear genome [12].

Pathological changes in the mtDNA can appear as point mutations in protein coding sequences, mt-tRNAs or even mt-rRNAs. In addition, major rearrangements of mtDNA, like deletions or insertions/duplications, are a cause of disease. Due to the fact that every cell contains a variable number of mtDNA molecules, mutations can be present in homoplasmy (all copies share the same mtDNA genotype) or heteroplasmy (only a population of DNA is mutated). The level of heteroplasmy of a mutation is a critical determinant of the cellular stress of a certain tissue or organ and has a major role in the disease phenotype. Finally, a reduction of mtDNA copies (depletion syndrome) can also hamper energy production and generate cellular stress (see **Table 1**, **Figure 1**) [12].

Alterations in the mitochondrial proteins encoded in the nucleus

Due to the diverse cellular roles that mitochondria fulfill, there are many mitochondrial processes that cause a pathology when disturbed. In the last years, massive sequencing approaches have significantly increased the number of known mutations implicated in mitochondrial diseases. Examples of this are defects in: factors involved in the biogenesis or integrity of respiratory chain complexes, those that regulate mtDNA maintenance, proteins required for transcription of mt-mRNA elements involved in translation of mtDNA encoded proteins, regulators of lipid metabolism, factors involved in cellular signalling and even enzymes of the Krebs cycle (see **Table 2**).

CELLULAR EFFECTS ON DIFFERENT TISSUES

Typically, mitochondrial disorders have been divided between those presenting with multiple symptoms, usually known as syndromes, and those characterized by tissue specific phenotypes. It remains to be addressed, which factors determine the tissue-specificity of mitochondrial diseases. However, to better address the different kinds of stress caused by mitochondrial distress, we will describe them classified by tissues/organs and give some examples of alterations that cause these problems.

Sensory organs

Hearing loss is one of the most prevalent sensory disorders [61]. Genetic factors are thought to account for more than half of congenital and childhood-onset hearing loss, including mutations in mtDNA [62] and mitochondrial nuclear genes like the heme A biogenesis factor COX10 [63, 64] or the AAA protease responsible for complex III assembly BSC1L [65, 66].

Mutations in the 12S mt-rRNA (m.1555A>G and m.1494 C>T) have been associated with aminoglycoside-induced ototoxicity and mitochondrial non-syndromic hearing loss. Studies using mitochondrial cybrids derived from HeLa cells and lymphoblasts have shown that these mutations affect the integrity and fidelity of the mitochondrial ribosome, therefore causing decreased mitochondrial translation,




FIGURE 1: Mitochondrial dysfunctions are related to mutations in mtDNA and defects in nuclear encoded mitochondrial proteins. (I) Overview of mutations within the mtDNA. **(II)** The majority of mitochondrial defects based on a malfunction of OXPHOS complexes and mitochondrial translation. **(III)** Defects in other processes, like mitochondrial fusion and fission or lipid homeostasis, leads to different mitochondrial diseases. **(IV)** Different strategies to fight the diverse forms of mitochondrial stress. (a: Leigh Syndrome LS; b: Leber Hereditary Optic Neuropathy LHON; c: Neurogenic Muscle Weakness, Ataxia and Retinitis Pigmentosa NARP; d: Mitochondrial Encephalomyopathy, Lactic Acidosis and Stroke-like Episodes MELAS; e: Myoclonic Epilepsy and Ragged Red Fiber Disease MERRF; f: Sensorineural Hearing Loss SNHL; g: mitochondrial non-syndromic Hearing Loss; Δ: Kearns Sayre Syndrome KSS).

either in the presence or the absence of aminoglycosides, and resulting in a cell growth defect [67, 68]. However, a study using osteosarcoma 143B derived cybrids showed no effect on mitochondrial translation after aminoglycoside treatment [69]. This discrepancy, the phenotypic differences between asymptomatic relatives and patients all harbouring the same mutational load and the fact that only in some cases the defect arose upon antibiotic treatment, raised the search for modifying factors of aminoglycoside induced ototoxicity within the nuclear genetic background [70]. Indeed, no negative effect was observed after aminoglycoside treatment in primate cells from the Cercopithecidae family where the m.1494 C>T was fixed as the wild-type allele and cells carried a compensating mutation in mitochondrial ribosomal protein S12 (MRPS12) [71], whereas primate cells from orangutan carrying the m.1555A>G mutation and no MRP mutation showed a drastic effect after antibiotic treatment [72]. In addition, this biochemical effect has been linked to stress signalling. Cybrids carrying the m.1555A>G mutation showed hypermethylation of the mitochondrial ribosome, disturbed mi-

tochondrial translation and assembly of the respiratory chain, resulting in increased production of ROS. Enhanced superoxide levels are sensed by AMPK, which signals further to E2F1, activating pro-apoptotic signalling in the cell. This induction seems to be tissue-specific, happens mainly in the inner ear and may explain the specific hearing defect observed in the presence of this particular mutation [73] (see **Table 1**).

Eye complications are also frequently found to be associated with mitochondrial dysfunction [74] and can be divided into primary and secondary. Primary afflictions are caused by genetic defects, whereas secondary afflictions are produced by hypertensive angiopathy of the retinal arteries, or diabetic retinopathy in mitochondrial diseases with diabetes [75]. Mitochondrial optic neuropathies have been associated with mutations in mtDNA and in nuclear genes. The most frequent eye disorder due to mtDNA mutation is Leber’s hereditary optic neuropathy (LHON) [76]. LHON usually affects young male adults and is characterised by mostly bilateral subacute or acute, painless, loss of central vision, with decreased colour vision [77]. There are

three main mtDNA mutations that underly the majority of LHON cases and all of them are found in complex I genes: m.11778G > A in the ND4 gene, m.3460G > A in the ND1 gene, and m.14484T > C in the ND6 gene (see **Table 1**). In addition, these mutations are usually present in homoplasm, indicating that probably other factors are involved in the development of the disorder. The molecular mechanisms underlying LHON are not yet fully understood. There have been some risk factors proposed, like specific mitochondrial haplogroups, smoking, alcohol consumption, and the use of some antibiotics. Differences in mitochondrial

mass have been also postulated to play a role in the incomplete manifestation of the disease. LHON mutation carriers with no pathological phenotype have significantly higher mtDNA copy number in leukocytes than affected carriers. By comparing fibroblasts from unaffected and affected mutation carriers, along with controls, it was shown that unaffected carriers have increased mitochondrial transcripts, respiratory chain proteins and enzyme activities compared to controls and affected carriers. Therefore, increased mitochondrial mass may play a protective role in LHON and compensate for complex I dys-

TABLE 1. Types of mitochondrial disease caused by mitochondrial encoded genes.

| Mitochondrial defects - mitochondrial encoded (mtDNA) [60] | | | |
|--|--|--|-------------------------------------|
| Disease | Coding | Mutation | Reference |
| Kearns Sayre Syndrome (KSS) | ND3, ND4, ND4L, ND5, COX3, ATP6, ATP8, tRNA ^{Leu} , tRNA ^{Ser} , tRNA ^{His} , tRNA ^{Arg} , tRNA ^{Gly} , tRNA ^{Lys} | Δ4977 (5 kb deletion) | [13, 14] |
| Leigh Syndrome (LS) | ATP6 | m.8993T>C m.9176T>G | [15] [16] |
| | ND3 | m.10158T>C | [17-19] |
| | ND4 | m.11777C>A | [20, 21] |
| | ND5 | m.12706T>C | [22] |
| | ND6 | m.14459G>A m.14487T>C | [23, 24] [25, 26] |
| Leber Hereditary Optic Neuropathy (LHON) | ND4 | m.11778G>A | [27] |
| | ND1 | m.3460G>A | [28, 29] |
| | ND6 | m.14484T>C | [30-32] |
| Neurogenic Muscle Weakness, Ataxia and Retinitis Pigmentosa (NARP) | ATP6 | m.8993T>G | [33, 34] |
| Mitochondrial Encephalomyopathy, Lactid Acidosis and Stroke-like Episodes (MELAS) | ND1 | m.3697C>A | [35] |
| | ND5 | m.13513G>A m.13514A>G | [36] [37] |
| | tRNA ^{Phe} | m.583G>A | [38, 39] |
| | tRNA ^{Leu} (UUR) | m.3243A>G m.3256C>T m.3271T>C m.3291T>C | [40] [41, 42] [43-45] [46] |
| | tRNA ^{Gln} | m.4332G>A | [47] |
| Myoclonic Epilepsy and Ragged Red Fiber Disease (MERRF) | tRNA ^{Lys} | m.8344A>G m.8356T>C m.8363G>A | [48, 49] [50-52] [53] |
| Sensorineural Hearing Loss (SNHL) | tRNA ^{Ser} | m.7445A>G m.7511T>C | [54] [55] |
| Deafness (DEAF) | 12s rRNA | m.1494C>T m.1555A>G | [56] [57-59] |

function [78]. In addition, males seem to be more affected because of the lack of protective effects from estrogen. Indeed, a study using cybrids carrying LHON mtDNA mutations showed that the addition of estradiol increased mitochondrial biogenesis and decreased ROS production by enhancing the activity of detoxifying enzymes like SOD2, leading to a decrease in apoptosis [79] (see **Table 1**).

The most common eye afflictions associated with nDNA mutations are autosomal dominant optic atrophy (ADOA), most frequently due to mutations in the Dynamin-like GTPase OPA1, and autosomal recessive optic atrophy (AROA), which has been mainly associated with mutations in the aconitate hydratase ACO2, or the uncharacterised transmembrane protein TMEM126A (OPA7) [76]. ADOA is clinically characterised by bilaterally symmetric progressive deterioration of the central visual acuity. Approximately 60-70% of ADOA cases are caused by genetic alterations in OPA1, other genes implicated in this pathology are OPA2 [80], OPA3 [81], OPA4 [82], OPA5 [83], OPA8 [84] and WFS1 [85] (see **Table 1**). OPA1 is a protein with eight different isoforms, processed by the mitochondrial metallochaperones YME1L and OMA1 [86-88]. The best-known function of OPA1 is for inner mitochondrial fusion during mitochondrial dynamics. In addition, OPA1 is involved in the remodelling of cristae by tethering inter-cristae membranes and proper function of the protein is required for maintaining cristae structure [89]. OPA1 mutations cause defective mitochondrial fusion and altered cristae structure, leading to direct effects on mitochondrial bioenergetics, including a decreased mitochondrial membrane potential and ATP synthesis and increased ROS production [90]. Interestingly, deletion of YME1L in murine heart, which alters OPA1 processing and function in a tissue-specific way, causes dilated cardiomyopathy (**Figure 1**) [91].

AROA presents with progressive impairment of visual capacity. The defect could either be spontaneously recovered or may lead to bilateral and progressive blindness [77]. Mutations in ACO2, affect the mitochondrial tricarboxylic acid cycle and therefore mitochondrial energy supply is depleted in patients [92]. Although there have been several AROA patients with mutations in TMEM126A, the exact function of the protein and therefore the molecular mechanism underlying optic atrophy has yet to be determined [93-95].

Heart

Cardiac muscle has a high energetic demand, therefore cardiac complications are frequent among mitochondrial diseases. One of the most common cardiac afflictions present in these pathologies is cardiomyopathy, which is estimated to occur in 20-40% of children with mitochondrial disease [4, 96, 97]. However, other symptoms like arrhythmia, conduction defects, pulmonary hypertension, dilated aortic root, pericardial effusion or coronary heart disease can also be developed as consequence of mitochondrial malfunction [5, 98].

Mitochondrial cardiomyopathies are characterised by abnormal myocardial structure or function that results from genetic defects that impair the mitochondrial respira-

tory chain [6, 98]. Hypertrophic cardiomyopathy is the most common form, present in more than 50% of cases [7, 96], but other forms, like dilated, restrictive, histiocytoid and left ventricular non-compactation cardiomyopathies can also be found among these patients [8, 99].

As described before (see above), genetic defects affecting the integrity of respiratory chain complexes, mitochondrial translation, maintenance of mtDNA, lipid metabolism and other metabolic pathways inside mitochondria might lead to cardiac disease. Several important perturbations have been described in subunits or factors required for the proper assembly of respiratory chain complexes. In general, mutations in these proteins cause an impairment of respiration and ATP production, increased ROS production and finally, cellular stress derived from a bioenergetics impairment. To date, pathological mutations have been found in 26 structural subunits of complex I [9, 100], that together with mutations in assembly factors represent around 30% of childhood mitochondrial diseases [11, 101]. Complex I defects can be present with isolated cardiomyopathy or together with multi-organic failure. Mutations in subunits of complex II or III have also been associated with different types of cardiomyopathy [12, 102-104]. Of special interest are defects of the cytochrome *c* oxidase, caused by mutations in assembly factors and in nuclear-encoded structural subunits. Mutations in the complex IV assembly factors COX10 and COX15 have been associated with hypertrophic cardiomyopathy [16, 105]. Both assembly factors are involved in the biosynthesis of heme A, the prosthetic group of the cytochrome *c* oxidase. Mutations in COX6B1, have been associated with cardiomyopathy and encephalopathy and showed decreased levels of the mature cytochrome *c* oxidase complex in patient-derived tissues and cells [12, 106, 107]. Although COX6B1 was thought to be a loosely interacting structural subunit of complex IV, studies have postulated Cox12 (yeast homolog of COX6B1) to be involved in the delivery of copper to Cox2, together with other metallochaperones like Sco1, Sco2 and Coa6 [61, 108]. Indeed, mutations in human SCO2 have been mainly associated with cardioencephalopathy [62, 109-111], whereas mutations in SCO1 have been associated with hepatic failure and encephalopathy [67, 68, 112, 113], as well as cardiomyopathy [64, 114]. Both proteins contain a CXXXC that is able to coordinate copper. They bind to apo-COX2 and deliver two copper atoms to the Cu_A center. Both enzymes have different but cooperative functions and disruptions in their function impair the maturation of cytochrome *c* oxidase [70, 115]. In addition, the SCO1 and SCO2 proteins are involved in regulating cellular copper homeostasis [71, 116]. Recently, it has been shown that SCO1 keeps the copper transporter CTR1 in the plasma membrane, this function being essential for the development of adult myocardium in mice [72, 117]. Mutations in COA6 have been described in infants with hypertrophic cardiomyopathy and combined complex I and IV, or isolated complex IV deficiency in the heart [73, 118, 119]. COA6 is required for cytochrome *c* oxidase assembly [74, 120, 121]. It is involved in the insertion of copper into COX2 and it has been described to interact with SCO2 [75, 122] and SCO1

[76, 123] after the translocation of the COX2 C-terminal domain into the IMS by COX18 [124]. However, why disturbance of copper metabolism, or ultimately of the cytochrome *c* oxidase, specifically affects the heart remains unclear (see **Table 2**, **Figure 1**).

Mutations in the mitochondrial translation machinery have also been associated with cardiomyopathy. Primary defects produce a decreased synthesis of mitochondrial polypeptides, but ultimately also impair mitochondrial bioenergetics and cause cellular stress. Mutations in the 16S mt-rRNA, and the m.1555A>G mutation in the 12S mt-rRNA, have been associated with hypertrophic and restrictive cardiomyopathy [78, 125, 126]. Mutations in ribosomal proteins (MRPL3 and MRPL44) and the translation elongation factor (TSFM) can cause cardiomyopathy, together with multi-organic disease [79, 127-130]. Finally, defects in mitochondrial tRNAs can be linked to isolated cardiomyopathy or multi-organic dysfunction [131-133] (see **Table 1** and **Table 2**).

Alterations of lipid metabolism inside mitochondria can also be a determinant for cardiac disease. Barth syndrome is an x-linked autosomal recessive disease, characterized by cardiomyopathy, skeletal myopathy, neutropenia, growth retardation, and 3-methylglutaconic aciduria [80, 134-136]. This disorder is caused by mutations in the Tafazzin protein, TAZ1, a mitochondrial acyl-transferase involved in the biogenesis of cardiolipin (CL), a phospholipid almost exclusively found in the inner mitochondrial membrane [81, 137]. The adequate presence of CL is required for structural stability of many critical protein complexes in the mitochondrial membrane and it is therefore essential for many mitochondrial processes ranging from protein import, cristae morphology, function of the respiratory chain or cell stress signaling [82, 136]. Interestingly, oxidation of CL causes loss of interaction with cytochrome *c*, a pre-requisite for triggering apoptosis. Oxidized CL has been found to be involved in the opening of the mitochondrial permeability transition pore (MPTP). In addition, CL is exposed to the outer mitochondrial membrane during apoptosis, where it is used as a binding platform for pro-apoptotic factors. Therefore, CL homeostasis plays an important role in cardiomyocyte programmed death upon ischaemia or reperfusion (**Figure 1**) [83, 136].

Mutations in another lipid related enzyme, the acylglycerol kinase AGK, have been associated with hypertrophic cardiomyopathy, myopathy, cataracts, exercise intolerance and lactic acidosis (Sengers syndrome). AGK was recently described as a component of the carrier protein translocase of the inner membrane (TIM22) [84, 138, 139], meaning that a defective import of carrier proteins alters mitochondrial metabolism and may disturb the function of the heart.

Neurological disorders

Similar to previously described organs and tissues and due to the high energy demands, neurological complications are commonly linked to mitochondrial dysfunction. Indeed, some of the most known mitochondrial syndromes caused by abnormalities in the mtDNA present with drastic neuro-

logical symptoms: Kearns–Sayre syndrome (KSS), a multi-system disorder with progressive external ophthalmoplegia, pigmentary retinopathy, heart block and frequently other signs like ataxia, dementia or endocrine problems is associated with single deletions of mtDNA [140]. MELAS (Mitochondrial Myopathy, Encephalopathy, Lactic Acidosis and Stroke-like Episodes) is caused in 80% of the cases by the m.3243A>G mutation in the tRNA^{LEU(UUR)} gene, although there have been other mutations described in protein coding genes [35, 36, 40, 80, 136, 141]. MERRF (Myoclonic Epilepsy and Ragged-Red Fiber), which usually also presents with cerebellar ataxia is mainly caused by mutations in the tRNA^{lys} gene (m8344A>G, m8356T>C, m.8363G>A), being the m.8344A>G the most frequent of them [49, 51, 53, 142]. The previously described defects affect the gene expression machinery of the mitochondrial genome and will generally affect mitochondrial protein synthesis, moreover an increased ROS production has been described in cybrids carrying the MELAS m.3243A>G mutation or the KSS associated common deletion Δ 4977 [89, 143]. In addition, there are mutations described in protein coding genes or in mitochondrial nuclear genes that would only affect individual complexes of the respiratory chain: NARP (Neuropathy, Ataxia and Retinitis Pigmentosa) has been mainly associated to the m.8933T>G/C mutation in the complex V subunit mt-ATP6 [33]. NARP patient derived cells were also found to have increased ROS production and decreased levels of ATP production [143]. Leigh syndrome is a progressive neurometabolic disorder that usually presents with seizures, hypotonia, fatigue, nystagmus, poor reflexes, eating and swallowing difficulties, breathing problems, poor motor function, and ataxia. This unique mitochondrial disorder is found to be caused by both mutations in the mtDNA and the nDNA. Mutations in many different genes have been identified to be the origin of Leigh syndrome, including mtDNA subunits of the complex I, IV and V, mt-tRNAs, nuclear encoded subunits of complex I, IV and II, the pyruvate dehydrogenase complex, or some assembly factors of the cytochrome *c* oxidase (SURF1, SCO1, SCO2, COX10, COX15) or complex III (BSC1L) [144]. Cells derived from patients with 3 different complex I mutations and Leigh syndrome exhibited increased ROS production [143, 145] (see **Table 1** and **Table 2**, **Figure 1**).

As already mentioned, many more genes are being identified as the reason behind mitochondrial dysfunction. Defects in mtDNA maintenance may result into defective mtDNA replication and lead to quantitative loss of mtDNA (mtDNA depletion) or qualitative one (mtDNA deletion). Downstream perturbation of mitochondrial protein synthesis will finally lead to a bioenergetics defect. MPV17 is a mitochondrial inner membrane protein involved in maintenance of mtDNA. It is believed to be involved in the import of deoxynucleotides into mitochondria. Pathogenic variants in MPV17 have been reported to cause hepatocerebral mtDNA depletion syndrome with liver failure, development delay and other neurological manifestations [146, 147]. In addition, infantile Navajo neuropathy (NNH), a neurohepatological disorder prevalently present among

Navajo children in the southwestern of USA has been found to be caused by mutations in MPV17 [148]. A recent report analysing new pathological variants in MPV17

showed that most patients exhibited a single or combined respiratory chain complex activity decrease [149] (see **Table 2**).

TABLE 2. Mitochondrial defects caused by nuclear encoded genes.

| Mitochondrial defects - nuclear encoded (nDNA) [60] | | | | | | | | | |
|---|----------|------------|--------|-------------|---------|------------|--|-----------|--|
| OXPHOS (structural proteins and assembly factors) | | | | | | | | | |
| Complex I | | Complex II | | Complex III | | Complex IV | | Complex V | |
| NDUFS1 | NDUFS2 | SDH-A | CYC1 | COX4I1 | COX4I2 | ATP5E | | | |
| NDUFS3 | NDUFS4 | SDH-B | UQCRC2 | COX5A | COX6B1 | ATP5A1 | | | |
| NDUFS6 | NDUFS7 | SDH-C | UQCRB | COX6A1 | COX8A | ATP8A2 | | | |
| NDUFS8 | NDUFB3 | SDH-D | UQCRQ | COX7B | COX15 | ATPAF2 | | | |
| NDUFB9 | NDUFB10 | SDHAF1 | BCS1L | SURF1 | COX20 | TMEM70 | | | |
| NDUFB11 | NDUFV1 | SDHF2 | LYRM7 | SCO1 | COA3 | | | | |
| NDUFA2 | NDUFA9 | | UQCC2 | SCO2 | COA5 | | | | |
| NDUFA10 | NDUFA11 | | UQCC3 | COX10 | COA6 | | | | |
| NDUFA12 | NDUFA13 | | | COX14 | COA7 | | | | |
| NDUFAF1 | NDUFAF2 | | | PET100 | LRPPRC | | | | |
| NDUFAF3 | NDUFAF4 | | | APOPT1 | FASTKD2 | | | | |
| NDUFAF5 | NDUFAF6 | | | | TACO1 | | | | |
| NUBPL | FOXRED | | | | | | | | |
| ACAD9 | | | | | | | | | |
| mtDNA maintenance | | | | | | | | | |
| POLG | POLG2 | ANT1 | MPV17 | OPA1 | MFN2 | C10ORF2 | | | |
| FBXL4 | DGUOK | RRM2B | SUCLA2 | SUCLG1 | TK2 | TFAM | | | |
| MGME1 | | | | | | | | | |
| Mitochondrial Import | | | | | | | | | |
| DDP | DNAJC19 | | | | | | | | |
| Mitochondrial Protein Synthesis | | | | | | | | | |
| AARS2 | CARS2 | DARS2 | EARS2 | FARS2 | GARS2 | HARS2 | | | |
| IARS2 | KARS | LARS | LARS2 | NARS2 | PARS2 | RARS2 | | | |
| SARS2 | TARS2 | VAR2 | YARS2 | EFG1 | TSFM | TUFM | | | |
| GTPBP3 | MTFMT | MTO1 | TRMT5 | TRMT10C | TRMU | GFM1 | | | |
| GFM2 | C12orf65 | RMND1 | MRPL3 | MRPS7 | MRPL12 | MRPS16 | | | |
| MRPS22 | MRPL44 | PUS1 | | | | | | | |
| Iron Homeostasis | | | | | | | | | |
| FRDA | ABCB7 | GLRX5 | ISCU | BOLA3 | NFU1 | ISCA2 | | | |
| IBA57 | LYRM4 | LYRM7 | FDXL1 | | | | | | |
| Coenzyme Q10 biogenesis | | | | | | | | | |
| COQ2 | COQ4 | COQ5 | COQ6 | COQ7 | COQ9 | APTX | | | |
| PDSS1 | PDSS2 | CABC1 | | | | | | | |
| Mitochondrial quality control | | | | | | | | | |
| SPG7 | AFG3L2 | | | | | | | | |
| Mitochondrial Integrity | | | | | | | | | |
| DLP1 | TAZ1 | RMRP | | | | | | | |
| Mitochondrial Metabolism | | | | | | | | | |
| PDHA1 | ETHE1 | ATAD3 | | | | | | | |

Aminoacyl-tRNA synthetases (ARS), are a family of proteins encoded in the nucleus and present in either the cytosol or mitochondria that ensure the proper conjugation of an amino acid with its cognate tRNA molecules. All mt-ARS are synthesized in the cytosol, imported to mitochondria due to an N-terminal targeting sequence (presequence) which is cleaved upon translocation to the matrix. Pathogenic variants of mt-ARS will affect mitochondrial translation and have been implicated in human neurological disorders of the brain, spinal cord and motor neurons in addition to other symptoms. Some of the most typical presentations are leukoencephalopathy with involvement of the brainstem and spinal cord and high lactate due to mt-aspartyl-tRNA synthetase (DARS2) mutations [150], leukoencephalopathy with thalamus and brainstem involvement and high lactate, caused by mt-glutamyl-tRNA synthetase (EARS2 [151]). However, there are other mt-ARS mutations which may also produce white matter lesions. The similar symptoms shown by ARS mutations may imply a shared mechanism of disease, however such a mechanism has not been yet demonstrated. Among the possible molecular reasons are: a reduced aminoacylation activity, altered dimerization, mislocalization, gain of function though pathogenic interactions and loss of noncanonical function [152] (see **Table 2**).

NEW STRATEGIES TO FIGHT MITOCHONDRIAL DERIVED STRESS

Nowadays there is no actual treatment for mitochondrial diseases. Nevertheless, in the last years a number of therapeutic strategies have been proposed, mainly in animal models. They can be classified into those acting on common pathways, and therefore applicable to different diseases, and those which aim to ameliorate a particular disorder (**Figure 1**) [153].

Those tissues or organs affected by decreased ATP production, and therefore impaired bioenergetics, can benefit from increased mitochondrial mass and activity. The transcriptional co-activator peroxisome proliferator activated receptor-1alpha (PGC1alpha) is the master regulator of mitochondrial biogenesis. It increases the activity of several transcription factors, like the nuclear respiratory factors (NR1 and NR2), thereby controlling the expression of OXPHOS related genes. In addition, PGC1alpha interacts with the peroxisomal proliferator activator receptors (PPARs), which regulate the expression of fatty acid oxidation genes [154]. PGC1alpha is activated either by deacetylation by Sirt1, or phosphorylation by AMPK, both of which can be modulated pharmacologically [155]. Under physiological conditions, PGC1alpha shows its highest expression levels in the heart, and mouse models lacking this protein have shown a normal cardiac function in unstimulated conditions. However, an impaired cardiac function was observed during certain stress conditions, like intense exercise or aortic constriction. Thus, the physiological role of PGC1alpha seems to be in fighting cellular stress [156].

Another possible strategy is to bypass the block in the respiratory chain from specific complex defects. In such a

way, electrons would flow again and reduce ROS production. Concomitantly, unaffected complexes would pump protons across the inner membrane and increase ATP production. The yeast *Saccharomyces cerevisiae* NADH reductase (Ndi1), which transfers electrons from NADH to coenzyme Q (CoQ), has been used to bypass CI defects [157]. In a similar approach, the alternative oxidase (AOX), which transfers electrons from CoQ to molecular oxygen in different organisms, has been used to bypass CIII and IV defects in cell culture [158] and to ameliorate to different extent respiratory defects in fly models [159, 160]. The enzyme has been successfully expressed in murine models [161], however correction of respiratory chain defects has not been shown yet *in vivo* in mammals.

As previously described (see above), the dynamin-like GTPase OPA1 is required for proper mitochondrial shaping. Regulating fission and fusion helps fight mitochondrial malfunction. Increasing the expression of long isoforms of OPA1 improves respiration efficiency by enhancing super-complex assembly and protects *in vivo* from many insults, such as ischemia/reperfusion, denervation/induced muscle atrophy, and OXPHOS deficiency [89, 162, 163].

In order to cope with increased oxidative damage generated in damaged mitochondria, different small molecules with antioxidant properties have been tested. Some examples, like Idebenone, lipoic acid, or Coenzyme Q₁₀, directly transfer electrons to the respiratory chain and bypass defective complexes. Others, like EPI-743 and RP103, enhance the biogenesis of glutathione, an important cellular antioxidant. KH176, can reduce altered cellular ROS levels and protect OXPHOS deficient cells against redox stress by targeting the Thioredoxin/Peroxiredoxin system [164]. MTP-131 is a member of the Szeto-Schiller (SS) peptide family and binds to CL. It increases OXPHOS capacity and improves the way mitochondria respond to metabolic changes. L-Arginine, a donor of nitric oxide, which thus regulates vascular tone, was shown to induce an improvement in aerobic capacity and muscle metabolism in models for mitochondrial disease [165].

Finally, genetic approaches can be used to correct mutations at a genomic level. Mitochondrially targeted restriction endonucleases have been used to shift heteroplasmy levels in cell lines with mutations in mtDNA and in heteroplasmic mice. Introduction of TALE and zinc finger nucleases (TALEN and ZFN) enabled the addition of specificity to the nucleases so that mutant DNA molecules could be selected for by directing unspecific restriction enzyme (FokI) to appropriate specific sequence assembling ZFN or TALE modules [166, 167]. However, this approach requires very large constructs that do not so easily fit into adeno-associated viruses (AAV) vectors. The Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) system is a bacterial immune system that has been modified for genome engineering. Due to the simplicity and adaptability of this technique, CRISPR has quickly displaced the previously established TALENs or ZFNs for genome engineering. CRISPR consists of two elements: a guide RNA (gRNA) and a non-specific CRISPR-associated endonuclease (Cas9). The gRNA is a short synthetic RNA

composed of a "scaffold", necessary for Cas9 binding, and a 20 nt targeting sequence that is specific to the gene of interest [168]. CRISPR was originally employed to knock-out target genes, but it has also been used to chromosomally modify or tag proteins, and to activate or repress target genes. However, the viability of this approach to target mitochondrial genes, mainly because of the requirement of a reliable nucleotide import system into mitochondria is not yet clear [169].

CONCLUDING REMARKS

Mitochondrial diseases show very complex and various clinical presentations. Because of the dual genetic origin of mitochondrial proteins, the number of genes susceptible of causing a mitochondrial pathology is large. Thus, the diagnostic process of mitochondrial diseases is usually complicated and very long, and in many cases although there is a clear suspicion of a mitochondrial defect, the final defect behind the phenotype remains undercover [170].

Despite the heterogeneity of the diseases and the genetic defects the final kind of cellular stresses are similar. In the most cases, there is a bioenergetic defect or an increased production of ROS. It remains unclear why although many genetic defects are present in the whole body, only certain tissues are affected. There may be different mechanisms to cope with mitochondrial stress, which may be tissue specific [73]. Indeed, disease models of mtDNA replication machinery failure have been linked to imbalance of the cellular dNTPs pool and consequently to increased glutathione biogenesis through *de novo* serine biogenesis. This metabolic switch was proposed to be a specific and rapid response to cellular stress/mtDNA damage in skeletal muscle and heart [171]. Together with this pathway, transcriptional response and mitochondrial unfolded protein response constitute the integrated mitochondrial stress response (ISRmt), which is controlled by the metabolic signalling regulator mTORC1 in muscle. However, long-term activation of cellular stress responses may be detrimental since chronic upregulation of anabolism contributes to mitochondrial myopathy pathogenesis

[172]. In addition, there is a certain multifactorial component in mitochondrial diseases. In some cases, ancient mt-rRNAs mutations rendered no adverse phenotype unless some environmental factors were used [72]. Therefore, the different incorporation and tolerance of tissues and organs for different xenobiotics may be different. Population variation plays also an important role in enhancing/ diminishing mitochondrial-related phenotypes, like population polymorphisms in mt-rRNAs and side effects derived from antibiotic treatment [173]. Nevertheless, a deeper investigation will be required to understand tissue specificity of mitochondrial diseases.

There have been several approaches proposed to treat mitochondrial diseases, however their application to the clinics is still a challenge [153]. Nevertheless, the discovery of new genome editing tools and the development of stem cell technologies will provide open new avenues of possibilities for the treatment of mitochondrial diseases.

ACKNOWLEDGMENTS

We are indebted to Sylvie Callegari for critical reading of the manuscript and to Peter Rehling and Alexander Schenzlielorz for fruitful discussion.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

COPYRIGHT

© 2018 Pacheu-Grau *et al.* This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: David Pacheu-Grau, Robert Rucktäschel and Markus Deckers (2018). Mitochondrial dysfunction and its role in tissue-specific cellular stress. *Cell Stress* 2(8): 184-199. doi: 10.15698/cst2018.07.147

REFERENCES

- Mick DU, Dennerlein S, Wiese H, Reinhold R, Pacheu-Grau D, Lorenzi I, Sasarman F, Weraarpachai W, Shoubridge EA, Warscheid B, and Rehling P (2012). MITRAC links mitochondrial protein translocation to respiratory-chain assembly and translational regulation. *Cell* 151(7): 1528–1541. doi: 10.1016/j.cell.2012.11.053
- Yoboue ED, and Devin A (2012). Reactive oxygen species-mediated control of mitochondrial biogenesis. *Int J Cell Biol* 2012:403870. doi: 10.1155/2012/403870
- Di Meo S, Reed TT, Venditti P, and Victor VM (2016). Role of ROS and RNS Sources in Physiological and Pathological Conditions. *Oxid Med Cell Longev* 2016: 1245049. doi: 10.1155/2016/1245049
- Busiello RA, Savarese S, and Lombardi A (2015). Mitochondrial uncoupling proteins and energy metabolism. *Front Physiol* 6: 36. doi: 10.3389/fphys.2015.00036
- Penna E, Espino J, De Stefani D, and Rizzuto R (2018). The MCU complex in cell death. *Cell calcium* 69: 73–80. doi: 10.1016/j.ceca.2017.08.008
- Raffaello A, Mammucari C, Gherardi G, and Rizzuto R (2016). Calcium at the Center of Cell Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes. *Trends Biochem Sci* 41(12): 1035–1049. doi: 10.1016/j.tibs.2016.09.001
- Lill R (2009). Function and biogenesis of iron-sulphur proteins. *Nature* 460(7257): 831–838. doi: 10.1038/nature08301
- Cardenas-Rodriguez M, Chatzi A, and Tokatlidis K (2018). Iron-sulfur clusters: from metals through mitochondria biogenesis to disease. *J Biol Inorg Chem* 23(4):509-520. doi: 10.1007/s00775-018-1548-6
- Montoya J, López-Pérez MJ, and Ruiz-Pesini E (2006). Mitochondrial DNA transcription and diseases: past, present and future. *Biochim*

- Biophys Acta** 1757(9-10): 1179–1189. doi: 10.1016/j.bbabi.2006.03.023
10. Zinovkina LA (2018). Mechanisms of Mitochondrial DNA Repair in Mammals. **Biochemistry** 83(3): 233–249. doi: 10.1134/S0006297918030045
11. Ruiz-Pesini E, Mishmar D, Brandon M, Procaccio V, and Wallace DC (2004). Effects of purifying and adaptive selection on regional variation in human mtDNA. **Science** 303(5655): 223–226. doi: 10.1126/science.1088434
12. Montoya J, López-Gallardo E, Herrero-Martín MD, Martínez-Romero I, Gómez-Durán A, Pacheu D, Carreras M, Díez-Sánchez C, López-Pérez MJ, and Ruiz-Pesini E (2009). Diseases of the human mitochondrial oxidative phosphorylation system. **Adv Exp Med Biol** 652(Chapter 5): 47–67. doi: 10.1007/978-90-481-2813-6_5
13. Zeviani M, Moraes CT, DiMauro S, Nakase H, Bonilla E, Schon EA, and Rowland LP (1988). Deletions of mitochondrial DNA in Kearns-Sayre syndrome. **Neurology** 38(8): 1339–1339. doi: 10.1212/WNL.38.8.1339
14. Lestienne P, and Ponsot G (1988). KEARNS-SAYRE SYNDROME WITH MUSCLE MITOCHONDRIAL DNA DELETION. **The Lancet** 331(8590): 885. doi: 10.1016/S0140-6736(88)91632-7
15. de Vries DD, van Engelen BGM, Gabreëls FJM, Ruitenbeek W, and van Oost BA (2004). A second missense mutation in the mitochondrial ATPase 6 gene in Leigh's syndrome. **Ann Neurol** 34(3): 410–412. doi: 10.1002/ana.410340319
16. Carrozzo R, Murray J, Santorelli FM, and Capaldi RA (2000). The T9176G mutation of human mtDNA gives a fully assembled but inactive ATP synthase when modeled in *Escherichia coli*. **FEBS Lett** 486(3): 297–299. doi: 10.1016/S0014-5793(00)02244-4
17. Crimi M, Papadimitriou A, Galbiati S, Palamidou P, Fortunato F, Bordonni A, Papandreou U, Papadimitriou D, Hadjigeorgiou GM, Drogari E, Bresolin N, and Comi GP (2004). A New Mitochondrial DNA Mutation in ND3 Gene Causing Severe Leigh Syndrome with Early Lethality. **Pediatric Res** 55(5): 842–846. doi: 10.1203/01.PDR.0000117844.73436.68
18. McFarland R, Kirby DM, Fowler KJ, Ohtake A, Ryan MT, Amor DJ, Fletcher JM, Dixon JW, Collins FA, Turnbull DM, Taylor RW, and Thorburn DR (2004). De novo mutations in the mitochondrial ND3 gene as a cause of infantile mitochondrial encephalopathy and complex I deficiency. **Ann Neurol** 55(1): 58–64. doi: 10.1002/ana.10787
19. Lebon S, Chol M, Benit P, Mugnier C, Chretien D, Giurgea I, Kern I, Girardin E, Hertz-Pannier L, de Lonlay P, Rötig A, Rustin P, and Munnich A (2003). Recurrent de novo mitochondrial DNA mutations in respiratory chain deficiency. **J Med Genet** 40(12): 896–899. doi: 10.1136/jmg.40.12.896
20. Deschauer M, Bamberg C, Claus D, Zierz S, Turnbull DM, and Taylor RW (2003). Late-onset encephalopathy associated with a C11777A mutation of mitochondrial DNA. **Neurology** 60(8): 1357–1359. doi: 10.1212/01.WNL.0000055869.99975.4B
21. Komaki H, Akanuma J, Iwata H, Takahashi T, Mashima Y, Nonaka I, and Goto Y-I (2003). A novel mtDNA C11777A mutation in Leigh syndrome. **Mitochondrion** 2(4): 293–304. doi: 10.1016/S1567-7249(03)00003-5
22. Taylor RW, Morris AA, Hutchinson M, and Turnbull DM (2002). Leigh disease associated with a novel mitochondrial DNA ND5 mutation. **Eur J Hum Genet** 10(2): 141–144. doi: 10.1038/sj.ejhg.5200773
23. Kirby DM, Kahler SG, Freckmann ML, Reddihough D, and Thorburn DR (2000). Leigh disease caused by the mitochondrial DNA G14459A mutation in unrelated families. **Ann Neurol** 48(1): 102–104. doi: 10.1002/1531-8249(200007)48:1<102::AID-ANA15>3.3.CO;2-D
24. Jun AS, Brown MD, and Wallace DC (1994). A mitochondrial DNA mutation at nucleotide pair 14459 of the NADH dehydrogenase subunit 6 gene associated with maternally inherited Leber hereditary optic neuropathy and dystonia. **Proc Natl Acad Sci U S A** 91(13): 6206–6210. doi: 10.1073/pnas.91.13.6206
25. Solano A, Roig M, Vives-Bauza C, Hernandez-Peña J, Garcia-Arumi E, Playan A, López-Pérez MJ, Andreu AL, and Montoya J (2003). Bilateral striatal necrosis associated with a novel mutation in the mitochondrial ND6 gene. **Ann Neurol** 54(4): 527–530. doi: 10.1002/ana.10682
26. Ugalde C, Triepels RH, Coenen MJH, Van Den Heuvel LP, Smeets R, Uusimaa J, Briones P, Campistol J, Majamaa K, Smeitink JAM, and Nijtmans LGJ (2003). Impaired complex I assembly in a Leigh syndrome patient with a novel missense mutation in the ND6 gene. **Ann Neurol** 54(5): 665–669. doi: 10.1002/ana.10734
27. Wallace D, Singh G, Lott M, Hodge J, Schurr T, Lezza A, Elsas L, and Nikoskelainen E (1988). Mitochondrial DNA mutation associated with Leber's hereditary optic neuropathy. **Science** 242(4884): 1427–1430. doi: 10.1126/science.3201231
28. Howell N, Bindoff LA, McCullough DA, Kubacka I, Poulton J, Mackey D, Taylor L, and Turnbull DM (1991). Leber hereditary optic neuropathy: identification of the same mitochondrial ND1 mutation in six pedigrees. **Am J Hum Genet** 49(5): 939–950. PMID: 1928099
29. Huoponen K, Vilkkij J, Aula P, Nikoskelainen EK, and Savontaus ML (1991). A new mtDNA mutation associated with Leber hereditary optic neuropathy. **Am J Hum Genet** 48(6): 1147–1153. PMID: 1674640
30. Brown MD, Voljavec AS, Lott MT, MacDonald I, and Wallace DC (1992). Leber's hereditary optic neuropathy: a model for mitochondrial neurodegenerative diseases. **FASEB J** 6(10): 2791–2799. PMID: 1634041
31. Howell N, Kubacka I, Xu M, and McCullough DA (1991). Leber hereditary optic neuropathy: involvement of the mitochondrial ND1 gene and evidence for an intragenic suppressor mutation. **Am J Hum Genet** 48(5): 935–942. PMID: 2018041
32. Johns DR, Neufeld MJ, and Park RD (1992). An ND-6 mitochondrial DNA mutation associated with Leber hereditary optic neuropathy. **Biochem Biophys Res Commun** 187(3): 1551–1557. doi: 10.1016/0006-291x(92)90479-5
33. Holt IJ, Harding AE, Petty RK, and Morgan-Hughes JA (1990). A new mitochondrial disease associated with mitochondrial DNA heteroplasmy. **Am J Hum Genet** 46(3): 428–433. PMID: 2137962
34. Harding AE, Holt IJ, Sweeney MG, Brockington M, and Davis MB (1992). Prenatal diagnosis of mitochondrial DNA8993 T→G disease. **Am J Hum Genet** 50(3): 629–633. PMID: 1539598
35. Kirby DM, McFarland R, Ohtake A, Dunning C, Ryan MT, Wilson C, Ketteridge D, Turnbull DM, Thorburn DR, and Taylor RW (2004). Mutations of the mitochondrial ND1 gene as a cause of MELAS. **J Med Genet** 41(10): 784–789. doi: 10.1136/jmg.2004.020537
36. Santorelli FM, Tanji K, Kulikova R, Shanske S, Vilarinho L, Hays AP, and DiMauro S (1997). Identification of a novel mutation in the mtDNA ND5 gene associated with MELAS. **Biochem Biophys Res Commun** 238(2): 326–328. doi: 10.1006/bbrc.1997.7167
37. Corona P, Antozzi C, Carrara F, D'Incerti L, Lamantea E, Tiranti V, and Zeviani M (2001). A novel mtDNA mutation in the ND5 subunit of complex I in two MELAS patients. **Ann Neurol** 49(1): 106–110. doi: 10.1002/1531-8249(200101)49:1<106::aid-ana16>3.0.co;2-t
38. Darin N, Kollberg G, Moslemi A-R, Tulinius M, Holme E, Grönlund MA, Andersson S, and Oldfors A (2006). Mitochondrial myopathy with exercise intolerance and retinal dystrophy in a sporadic patient with a G583A mutation in the mt tRNA(phe) gene. **Neuromuscul Disord**

16(8): 504–506. doi: 10.1016/j.nmd.2006.05.010

39. Hanna MG, Nelson IP, Morgan-Hughes JA, and Wood NW (1998). MELAS: a new disease associated mitochondrial DNA mutation and evidence for further genetic heterogeneity. *J Neurol Neurosurg Psychiatry* 65(4): 512–517. doi: 10.1136/jnnp.65.4.512

40. Goto Y, Nonaka I, and Horai S (1990). A mutation in the tRNA(Leu)(UUR) gene associated with the MELAS subgroup of mitochondrial encephalomyopathies. *Nature* 348(6302): 651–653. doi: 10.1038/348651a0

41. Sato W, Hayasaka K, Shoji Y, Takahashi T, Takada G, Saito M, Fukawa O, and Wachi E (1994). A mitochondrial tRNA(Leu)(UUR) mutation at 3,256 associated with mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes (MELAS). *Biochem Mol Biol Int* 33(6): 1055–1061. PMID: 7804130

42. Moraes CT, Ciacci F, Bonilla E, Jansen C, Hirano M, Rao N, Lovelace RE, Rowland LP, Schon EA, and DiMauro S (1993). Two novel pathogenic mitochondrial DNA mutations affecting organelle number and protein synthesis. Is the tRNA(Leu)(UUR) gene an etiologic hot spot? *J Clin Invest* 92(6): 2906–2915. doi: 10.1172/JCI116913

43. Hayashi J, Ohta S, Takai D, Miyabayashi S, Sakuta R, Goto Y, and Nonaka I (1993). Accumulation of mtDNA with a mutation at position 3271 in tRNA(Leu)(UUR) gene introduced from a MELAS patient to HeLa cells lacking mtDNA results in progressive inhibition of mitochondrial respiratory function. *Biochem Biophys Res Commun* 197(3): 1049–1055. doi: 10.1006/bbrc.1993.2584

44. Sakuta R, Goto Y, Horai S, and Nonaka I (1993). Mitochondrial DNA mutations at nucleotide positions 3243 and 3271 in mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes: a comparative study. *J Neurol Sci* 115(2): 158–160. doi: 10.1016/0022-510x(93)90219-o

45. Goto Y, Nonaka I, and Horai S (1991). A new mtDNA mutation associated with mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like episodes (MELAS). *Biochim Biophys Acta* 1097(3): 238–240. doi: 10.1016/0925-4439(91)90042-8

46. Goto YI, Tsugane K, Tanabe Y, Nonaka I, and Horai S (1994). A New Point Mutation at Nucleotide Pair-3291 of the Mitochondrial Transfer-RNA(Leu)(Uur) Gene in a Patient with Mitochondrial Myopathy, Encephalopathy, Lactic-Acidosis, and Stroke-Like Episodes (MELAS). *Biochem Biophys Res Commun* 202(3): 1624–1630. doi: 10.1006/bbrc.1994.2119

47. Bataillard M, Chatzoglou E, Rumbach L, Sternberg D, Tournade A, Laforet P, Jardel C, Maisonnobe T, and Lombes A (2001). Atypical MELAS syndrome associated with a new mitochondrial tRNA glutamine point mutation. *Neurology* 56(3): 405–407. doi: 10.1212/WNL.56.3.405

48. Shoffner JM, Lott MT, Lezza AM, Seibel P, Ballinger SW, and Wallace DC (1990). Myoclonic epilepsy and ragged-red fiber disease (MERRF) is associated with a mitochondrial DNA tRNA(Lys) mutation. *Cell* 61(6): 931–937. doi: 10.1016/0092-8674(90)90059-n

49. Wallace DC, Zheng XX, Lott MT, Shoffner JM, Hodge JA, Kelley RI, Epstein CM, and Hopkins LC (1988). Familial mitochondrial encephalomyopathy (MERRF): genetic, pathophysiological, and biochemical characterization of a mitochondrial DNA disease. *Cell* 55(4): 601–610. doi: 10.1016/0092-8674(88)90218-8

50. Masucci JP, Davidson M, Koga Y, Schon EA, and King MP (1995). In vitro analysis of mutations causing myoclonus epilepsy with ragged-red fibers in the mitochondrial tRNA(Lys) gene: two genotypes produce similar phenotypes. *Mol Cell Biol* 15(5): 2872–2881. doi: 10.1128/MCB.15.5.2872

51. Zeviani M, Muntoni F, Savarese N, Serra G, Tiranti V, Carrara F, Mariotti C, and DiDonato S (1993). A MERRF/MELAS overlap syndrome

associated with a new point mutation in the mitochondrial DNA tRNA(Lys) gene. *Eur J Hum Genet* 1(1): 80–87. doi: 10.1159/000472390

52. Silvestri G, Moraes CT, Shanske S, Oh SJ, and DiMauro S (1992). A new mtDNA mutation in the tRNA(Lys) gene associated with myoclonic epilepsy and ragged-red fibers (MERRF). *Am J Hum Genet* 51(6): 1213–1217. PMID: 1361099

53. Ozawa M, Nishino I, Horai S, Nonaka I, and Goto Y-I (1997). Myoclonus epilepsy associated with ragged-red fibers: A G-to-A mutation at nucleotide pair 8363 in mitochondrial tRNA(Lys) in two families. *Muscle Nerve* 20(3): 271–278. doi: 10.1002/(SICI)1097-4598(199703)20:3<271::AID-MUS2>3.0.CO;2-8

54. Reid FM, Vernham GA, and Jacobs HT (1994). A novel mitochondrial point mutation in a maternal pedigree with sensorineural deafness. *Human Mut* 3(3): 243–247. doi: 10.1002/humu.1380030311

55. Sue CM, Tanji K, Hadjigeorgiou G, Andreu AL, Nishino I, Krishna S, Bruno C, Hirano M, Shanske S, Bonilla E, Fischel-Ghodsian N, DiMauro S, and Friedman R (1999). Maternally inherited hearing loss in a large kindred with a novel T7511C mutation in the mitochondrial DNA tRNA(Ser)(UCN) gene. *Neurology* 52(9): 1905–1908. doi: 10.1212/WNL.52.9.1905

56. Hema Bindu L, and Reddy PP (2009). Genetics of aminoglycoside-induced and prelingual non-syndromic mitochondrial hearing impairment: A review. *Int J Audiol* 47(11): 702–707. doi: 10.1080/14992020802215862

57. Fischel-Ghodsian N, Prezant TR, Bu X, and Öztas S (1993). Mitochondrial ribosomal RNA gene mutation in a patient with sporadic aminoglycoside ototoxicity. *Am J Otolaryngol* 14(6): 399–403. doi: 10.1016/0196-0709(93)90113-L

58. Hutchin T, Haworth I, Higashi K, Fischel-Ghodsian N, Stoneking M, Saha N, Arnos C, and Cortopassi G (1993). A molecular basis for human hypersensitivity to aminoglycoside antibiotics. *Nucleic Acids Research* 21(18): 4174–4179. PMID: 8414970

59. Prezant TR, Agapian JV, Bohlman MC, Bu X, Öztas S, Qiu WQ, Arnos KS, Cortopassi GA, Jaber L, and Rotter JI (1993). Mitochondrial ribosomal RNA mutation associated with both antibiotic-induced and non-syndromic deafness. *Nat Genet* 4(3): 289–294. doi: 10.1038/ng0793-289

60. Lott MT, Leipzig JN, Derbeneva O, Xie HM, Chalkia D, Sarmady M, Procaccio V, and Wallace DC (2013). mtDNA Variation and Analysis Using Mitomap and Mitomaster. *Curr Protoc Bioinformatics* 44(1): 1.23.1–26. doi: 10.1002/0471250953.bi012344

61. Kral A, and O'Donoghue GM (2010). Profound deafness in childhood. *N Engl J Med* 363(15): 1438–1450. doi: 10.1056/NEJMra0911225

62. Mutai H, Watabe T, Kosaki K, Ogawa K, and Matsunaga T (2017). Mitochondrial mutations in maternally inherited hearing loss. *BMC Med Genet* 18(1): 32. doi: 10.1186/s12881-017-0389-4

63. Pitceathly RDS, Taanman J-W, Rahman S, Meunier B, Sadowski M, Cirak S, Hargreaves I, Land JM, Nanji T, Polke JM, Woodward CE, Sweeney MG, Solanki S, Foley AR, Hurles ME, Stalker J, Blake J, Holton JL, Phadke R, Muntoni F, Reilly MM, Hanna MG, UK10K Consortium (2013). COX10 mutations resulting in complex multisystem mitochondrial disease that remains stable into adulthood. *JAMA Neurol* 70(12): 1556–1561. doi: 10.1001/jamaneurol.2013.3242

64. Antonicka H, Leary SC, Guercin G-H, Agar JN, Horvath R, Kennaway NG, Harding CO, Jaksch M, and Shoubridge EA (2003). Mutations in COX10 result in a defect in mitochondrial heme A biosynthesis and account for multiple, early-onset clinical phenotypes associated with isolated COX deficiency. *Hum Mol Genet* 12(20): 2693–2702. doi: 10.1093/hmg/ddg284

65. Hinson JT, Fantin VR, Schönberger J, Breivik N, Siem G, McDonough B, Sharma P, Keogh I, Godinho R, Santos F, Esparza A, Nicolau Y, Selvaag E, Cohen BH, Hoppel CL, Tranebjaerg L, Eavey RD, Seidman JG, and Seidman CE (2007). Missense mutations in the BCS1L gene as a cause of the Björnstad syndrome. **N Engl J Med** 356(8): 809–819. doi: 10.1056/NEJMoa055262
66. Fernandez-Vizarra E, Bugiani M, Goffrini P, Carrara F, Farina L, Procopio E, Donati A, Uziel G, Ferrero I, and Zeviani M (2007). Impaired complex III assembly associated with BCS1L gene mutations in isolated mitochondrial encephalopathy. **Hum Mol Genet** 16(10): 1241–1252. doi: 10.1093/hmg/ddm072
67. Guan MX, Fischel-Ghodsian N, and Attardi G (2000). A biochemical basis for the inherited susceptibility to aminoglycoside ototoxicity. **Hum Mol Genet** 9(12): 1787–1793. PMID: 10915767
68. Iglesias E, Llobet L, Pacheu-Grau D, Gómez-Durán A, and Ruiz-Pesini E (2012). Cybrids for Mitochondrial DNA Pharmacogenomics. **Drug Development Research** 73(8): 453–460. doi: 10.1002/ddr.21037
69. Giordano C, Pallotti F, Walker WF, Checcharelli N, Musumeci O, Santorelli F, d'Amati G, Schon EA, DiMauro S, Hirano M, and Davidson MM (2002). Pathogenesis of the deafness-associated A1555G mitochondrial DNA mutation. **Biochem Biophys Res Commun** 293(1): 521–529. doi: 10.1016/S0006-291X(02)00256-5
70. Pacheu-Grau D, Gómez-Durán A, López-Pérez MJ, Montoya J, and Ruiz-Pesini E (2010). Mitochondrial pharmacogenomics: barcode for antibiotic therapy. **Drug Discov Today** 15(1-2): 33–39. doi: 10.1016/j.drudis.2009.10.008
71. Emperador S, Pacheu-Grau D, Bayona-Bafaluy MP, Garrido-Pérez N, Martín-Navarro A, López-Pérez MJ, Montoya J, and Ruiz-Pesini E (2014). An MRPS12 mutation modifies aminoglycoside sensitivity caused by 12S rRNA mutations. **Front Genet** 5: 469. doi: 10.3389/fgene.2014.00469
72. Pacheu-Grau D, Gómez-Durán A, López-Gallardo E, Pinós T, Andreu AL, López-Pérez MJ, Montoya J, and Ruiz-Pesini E (2011). “Progress” renders detrimental an ancient mitochondrial DNA genetic variant. **Hum Mol Genet** 20(21): 4224–4231. doi: 10.1093/hmg/ddr350
73. Raimundo N, Song L, Shutt TE, McKay SE, Cotney J, Guan M-X, Gilliland TC, Hohuan D, Santos-Sacchi J, and Shadel GS (2012). Mitochondrial stress engages E2F1 apoptotic signaling to cause deafness. **Cell** 148(4): 716–726. doi: 10.1016/j.cell.2011.12.027
74. Finsterer J, Zarrouk-Mahjoub S, and Daruich A (2016). The Eye on Mitochondrial Disorders. **J Child Neurol** 31(5): 652–662. doi: 10.1177/0883073815599263
75. Zhu Y, Gu X, and Xu C (2016). A Mitochondrial DNA A8701G Mutation Partly Associated with Maternally Inherited Hypertension and Dilated Cardiomyopathy in a Chinese Pedigree. **Chin Med J** 129(15): 1890. doi: 10.4103/0366-6999.186656
76. Leruez S, Amati-Bonneau P, VERNY C, Reynier P, Procaccio V, Bonneau D, and Milea D (2014). Mitochondrial dysfunction affecting visual pathways. **Rev Neurol** 170(5): 344–354. doi: 10.1016/j.neuro.2014.03.009
77. Finsterer J, Mancuso M, Pareyson D, Burgunder J-M, and Klopstock T (2017). Mitochondrial disorders of the retinal ganglion cells and the optic nerve. **Mitochondrion**. doi: 10.1016/j.mito.2017.10.003
78. Giordano C et al. (2014). Efficient mitochondrial biogenesis drives incomplete penetrance in Leber's hereditary optic neuropathy. **Brain** 137(Pt 2): 335–353. doi: 10.1093/brain/awt343
79. Giordano C, Montopoli M, Perli E, Orlandi M, Fantin M, Ross-Cisneros FN, Caparrotta L, Martinuzzi A, Ragazzi E, Ghelli A, Sadun AA, d'Amati G, and Carelli V (2011). Oestrogens ameliorate mitochondrial dysfunction in Leber's hereditary optic neuropathy. **Brain** 134(Pt 1): 220–234. doi: 10.1093/brain/awq276
80. Katz BJ, Zhao Y, Warner JEA, Tong Z, Yang Z, and Zhang K (2006). A family with X-linked optic atrophy linked to the OPA2 locus Xp11.4-Xp11.2. **Am J Med Genet A** 140(20): 2207–2211. doi: 10.1002/ajmg.a.31455
81. Sergouniotis PI, Perveen R, Thiselton DL, Giannopoulos K, Sarros M, Davies JR, Biswas S, Ansons AM, Ashworth JL, Lloyd IC, Black GC, and Votruba M (2015). Clinical and molecular genetic findings in autosomal dominant OPA3-related optic neuropathy. **Neurogenetics** 16(1): 69–75. doi: 10.1007/s10048-014-0416-y
82. Kerrison JB, Arnould VJ, Ferraz Sallum JM, Vagefi MR, Barmada MM, Li Y, Zhu D, and Maumenee IH (1999). Genetic heterogeneity of dominant optic atrophy, Kjer type: Identification of a second locus on chromosome 18q12.2-12.3. **Arch Ophthalmol** 117(6): 805–810. PMID: 10369594
83. Barbet F, Hakiki S, Orssaud C, Gerber S, Perrault I, Hanein S, Ducrocq D, Dufier J-L, Munnich A, Kaplan J, and Rozet J-M (2005). A third locus for dominant optic atrophy on chromosome 22q. **J Med Genet** 42(1): e1. doi: 10.1136/jmg.2004.025502
84. Carelli V, Schimpf S, Fuhrmann N, Valentino ML, Zanna C, Iommarini L, Papke M, Schaich S, Tippmann S, Baumann B, Barboni P, Longanesi L, Rugolo M, Ghelli A, Alavi MV, Youle RJ, Bucchi L, Carroccia R, Giannoccaro MP, Tonon C, Lodi R, Cenacchi G, Montagna P, Liguori R, and Wissinger B (2011). A clinically complex form of dominant optic atrophy (OPA8) maps on chromosome 16. **Hum Mol Genet** 20(10): 1893–1905. doi: 10.1093/hmg/ddr071
85. Eiberg H, Hansen L, Kjer B, Hansen T, Pedersen O, Bille M, Rosenberg T, and Tranebjaerg L (2006). Autosomal dominant optic atrophy associated with hearing impairment and impaired glucose regulation caused by a missense mutation in the WFS1 gene. **J Med Genet** 43(5): 435–440. doi: 10.1136/jmg.2005.034892
86. Anand R, Wai T, Baker MJ, Kladt N, Schauss AC, Rugarli E, and Langer T (2014). The i-AAA protease YME1L and OMA1 cleave OPA1 to balance mitochondrial fusion and fission. **J Cell Biol** 204(6): 919–929. doi: 10.1083/jcb.201308006
87. Baker MJ, Lampe PA, Stojanovski D, Korwitz A, Anand R, Tatsuta T, and Langer T (2014). Stress-induced OMA1 activation and autocatalytic turnover regulate OPA1-dependent mitochondrial dynamics. **EMBO J** 33(6): 578–593. doi: 10.1002/emboj.201386474
88. Ehses S, Raschke I, Mancuso G, Bernacchia A, Geimer S, Tondera D, Martinou J-C, Westermann B, Rugarli E, and Langer T (2009). Regulation of OPA1 processing and mitochondrial fusion by m-AAA protease isoenzymes and OMA1. **J Cell Biol** 187(7): 1023–1036. doi: 10.1083/jcb.200906084
89. Pernas L, and Scorrano L (2016). Mito-Morphosis: Mitochondrial Fusion, Fission, and Cristae Remodeling as Key Mediators of Cellular Function. **Annu Rev Physiol** 78: 505–531. doi: 10.1146/annurev-physiol-021115-105011
90. Kao S-H, Yen M-Y, Wang A-G, Yeh Y-L, and Lin A-L (2015). Changes in Mitochondrial Morphology and Bioenergetics in Human Lymphoblastoid Cells With Four Novel OPA1 Mutations. **Invest Ophthalmol Vis Sci** 56(4): 2269–2278. doi: 10.1167/iovs.14-16288
91. Wai T, García-Prieto J, Baker MJ, Merkwirth C, Benit P, Rustin P, Rupérez FJ, Barbas C, Ibañez B, and Langer T (2015). Imbalanced OPA1 processing and mitochondrial fragmentation cause heart failure in mice. **Science** 350(6265): aad0116. doi: 10.1126/science.aad0116
92. Metodiev MD, Gerber S, Hubert L, Delahodde A, Chretien D, Gérard X, Amati-Bonneau P, Giacomotto M-C, Boddart N, Kaminska A, Desguerre I, Amiel J, Rio M, Kaplan J, Munnich A, Rötig A, Rozet J-M, and Besmond C (2014). Mutations in the tricarboxylic acid cycle enzyme, aconitase 2, cause either isolated or syndromic optic neuropathy with encephalopathy and cerebellar atrophy. **J Med Genet** 51(12): 834–838. doi: 10.1136/jmedgenet-2014-102532

93. Hanein S, Perrault I, Roche O, Gerber S, Khadom N, Rio M, Boddart N, Jean-Pierre M, Brahimi N, Serre V, Chretien D, Delphin N, Fares-Taie L, Lachheb S, Rötig A, Meire F, Munnich A, Dufier J-L, Kaplan J, and Rozet J-M (2009). TMEM126A, encoding a mitochondrial protein, is mutated in autosomal-recessive nonsyndromic optic atrophy. **Am J Hum Genet** 84(4): 493–498. doi: 10.1016/j.ajhg.2009.03.003
94. Désir J, Coppieters F, Van Regemorter N, De Baere E, Abramowicz M, and Cordonnier M (2012). TMEM126A mutation in a Moroccan family with autosomal recessive optic atrophy. **Mol Vis** 18: 1849–1857. PMID: 22815638
95. Meyer E, Michaelides M, Tee LJ, Robson AG, Rahman F, Pasha S, Luxon LM, Moore AT, and Maher ER (2010). Nonsense mutation in TMEM126A causing autosomal recessive optic atrophy and auditory neuropathy. **Mol Vis** 16: 650–664. PMID: 20405026
96. Scaglia F, Towbin JA, Craigen WJ, Belmont JW, Smith EO, Neish SR, Ware SM, Hunter JV, Fernbach SD, Vladutiu GD, Wong L-JC, and Vogel H (2004). Clinical spectrum, morbidity, and mortality in 113 pediatric patients with mitochondrial disease. **Pediatrics** 114(4): 925–931. doi: 10.1542/peds.2004-0718
97. Holmgren D, Wåhlander H, Eriksson BO, Oldfors A, Holme E, and Tulinius M (2003). Cardiomyopathy in children with mitochondrial disease; clinical course and cardiological findings. **Eur Heart J** 24(3): 280–288. doi: 10.1016/s0195-668x(02)00387-1
98. El-Hattab AW, and Scaglia F (2016). Mitochondrial Cardiomyopathies. **Front Cardiovasc Med** 3: 25. doi: 10.3389/fcvm.2016.00025
99. Finsterer J, and Kothari S (2014). Cardiac manifestations of primary mitochondrial disorders. **Int J Cardiol** 177(3): 754–763. doi: 10.1016/j.ijcard.2014.11.014
100. Rodenburg RJ (2016). Mitochondrial complex I-linked disease. **Biochim Biophys Acta** 1857(7): 938–945. doi: 10.1016/j.bbabi.2016.02.012
101. Brunel-Guitton C, Levtova A, and Sasarman F (2015). Mitochondrial Diseases and Cardiomyopathies. **Can J Cardiol** 31(11): 1360–1376. doi: 10.1016/j.cjca.2015.08.017
102. Alston CL, Ceccatelli Berti C, Blakely EL, Oláhová M, He L, McMahon CJ, Olpin SE, Hargreaves IP, Noll C, McFarland R, Goffrini P, O'Sullivan MJ, and Taylor RW (2015). A recessive homozygous p.Asp92Gly SDHD mutation causes prenatal cardiomyopathy and a severe mitochondrial complex II deficiency. **Hum Genet** 134(8): 869–879. doi: 10.1007/s00439-015-1568-z
103. Andreu AL, Checcarelli N, Iwata S, Shanske S, and DiMauro S (2000). A missense mutation in the mitochondrial cytochrome b gene in a revisited case with histiocytoid cardiomyopathy. **Pediatr Res** 48(3): 311–314. doi: 10.1203/00006450-200009000-00008
104. Carossa V, Ghelli A, Tropeano CV, Valentino ML, Iommarini L, Maresca A, Caporali L, La Morgia C, Liguori R, Barboni P, Carbonelli M, Rizzo G, Tonon C, Lodi R, Martinuzzi A, De Nardo V, Rugolo M, Ferretti L, Gandini F, Pala M, Achilli A, Olivieri A, Torroni A, and Carelli V (2014). A novel in-frame 18-bp microdeletion in MT-CYB causes a multisystem disorder with prominent exercise intolerance. **Hum Mutat** 35(8): 954–958. doi: 10.1002/humu.22596
105. Antonicka H, Mattman A, Carlson CG, Glerum DM, Hoffbuhr KC, Leary SC, Kennaway NG, and Shoubridge EA (2003). Mutations in COX15 produce a defect in the mitochondrial heme biosynthetic pathway, causing early-onset fatal hypertrophic cardiomyopathy. **Am J Hum Genet** 72(1): 101–114. doi: 10.1086/345489
106. Abdulhag UN, Soiferman D, Schueler-Furman O, Miller C, Shaag A, Elpeleg O, Edvardson S, and Saada A (2015). Mitochondrial complex IV deficiency, caused by mutated COX6B1, is associated with encephalomyopathy, hydrocephalus and cardiomyopathy. **Eur J Hum Genet** 23(2): 159–164. doi: 10.1038/ejhg.2014.85
107. Massa V, Fernandez-Vizarrá E, Alshahwan S, Bakhsh E, Goffrini P, Ferrero I, Mereghetti P, D'Adamo P, Gasparini P, and Zeviani M (2008). Severe infantile encephalomyopathy caused by a mutation in COX6B1, a nucleus-encoded subunit of cytochrome c oxidase. **Am J Hum Genet** 82(6): 1281–1289. doi: 10.1016/j.ajhg.2008.05.002
108. Ghosh A, Pratt AT, Soma S, Theriault SG, Griffin AT, Trivedi PP, and Gohil VM (2016). Mitochondrial disease genes COA6, COX6B and SCO2 have overlapping roles in COX2 biogenesis. **Hum Mol Genet** 25(4): 660–671. doi: 10.1093/hmg/ddv503
109. Jaksch M, Horvath R, Horn N, Auer DP, Macmillan C, Peters J, Gerbitz KD, Kraegeloh-Mann I, Muntau A, Karcagi V, Kalmachey R, Lochmuller H, Shoubridge EA, and Freisinger P (2001). Homozygosity (E140K) in SCO2 causes delayed infantile onset of cardiomyopathy and neuropathy. **Neurology** 57(8): 1440–1446. PMID: 11673586
110. Mobley BC, Enns GM, Wong L-J, and Vogel H (2009). A novel homozygous SCO2 mutation, p.G193S, causing fatal infantile cardioencephalomyopathy. **Clin Neuropathol** 28(2): 143–149. doi: 10.5414/npp28143
111. Papadopoulou LC, Sue CM, Davidson MM, Tanji K, Nishino I, Sadlock JE, Krishna S, Walker W, Selby J, Glerum DM, Coster RV, Lyon G, Scalais E, Lebel R, Kaplan P, Shanske S, De Vivo DC, Bonilla E, Hirano M, DiMauro S, and Schon EA (1999). Fatal infantile cardioencephalomyopathy with COX deficiency and mutations in SCO2, a COX assembly gene. **Nat Genet** 23(3): 333–337. doi: 10.1038/15513
112. Leary SC, Antonicka H, Sasarman F, Weraarpachai W, Cobine PA, Pan M, Brown GK, Brown R, Majewski J, Ha KCH, Rahman S, and Shoubridge EA (2013). Novel mutations in SCO1 as a cause of fatal infantile encephalopathy and lactic acidosis. **Hum Mutat** 34(10): 1366–1370. doi: 10.1002/humu.22385
113. Valnot I, Osmond S, Gigarel N, Mehaye B, Amiel J, Cormier-Daire V, Munnich A, Bonnefont JP, Rustin P, and Rötig A (2000). Mutations of the SCO1 gene in mitochondrial cytochrome c oxidase deficiency with neonatal-onset hepatic failure and encephalopathy. **Am J Hum Genet** 67(5): 1104–1109. doi: 10.1016/S0002-9297(07)62940-1
114. Stiburek L, Vesela K, Hansikova H, Hulkova H, and Zeman J (2009). Loss of function of Sco1 and its interaction with cytochrome c oxidase. **Am J Physiol Cell Physiol** 296(5): C1218–26. doi: 10.1152/ajpcell.00564.2008
115. Leary SC, Kaufman BA, Pellicchia G, Guercin G-H, Mattman A, Jaksch M, and Shoubridge EA (2004). Human SCO1 and SCO2 have independent, cooperative functions in copper delivery to cytochrome c oxidase. **Hum Mol Genet** 13(17): 1839–1848. doi: 10.1093/hmg/ddh197
116. Leary SC, Cobine PA, Kaufman BA, Guercin G-H, Mattman A, Palaty J, Lockitch G, Winge DR, Rustin P, Horvath R, and Shoubridge EA (2007). The human cytochrome c oxidase assembly factors SCO1 and SCO2 have regulatory roles in the maintenance of cellular copper homeostasis. **Cell Metab** 5(1): 9–20. doi: 10.1016/j.cmet.2006.12.001
117. Baker ZN, Jett K, Boulet A, Hossain A, Cobine PA, Kim B-E, Zawily EI AM, Lee L, Tibbits GF, Petris MJ, and Leary SC (2017). The mitochondrial metallochaperone SCO1 maintains CTR1 at the plasma membrane to preserve copper homeostasis in the murine heart. **Hum Mol Genet** 26(23): 4617–4628. doi: 10.1093/hmg/ddx344
118. Calvo SE, Compton AG, Hershman SG, Lim SC, Lieber DS, Tucker EJ, Laskowski A, Garone C, Liu S, Jaffe DB, Christodoulou J, Fletcher JM, Bruno DL, Goldblatt J, DiMauro S, Thorburn DR, and Mootha VK (2012). Molecular Diagnosis of Infantile Mitochondrial Disease with Targeted Next-Generation Sequencing. **Sci Transl Med** 4(118): 118ra10–118ra10. doi: 10.1126/scitranslmed.3003310
119. Baertling F, A M van den Brand M, Hertecant JL, Al-Shamsi A, P

- van den Heuvel L, Distelmaier F, Mayatepek E, Smeitink JA, Nijtmans LGJ, and Rodenburg RJT (2015). Mutations in COA6 cause cytochrome c oxidase deficiency and neonatal hypertrophic cardiomyopathy. *Hum Mutat* 36(1): 34–38. doi: 10.1002/humu.22715
120. Ghosh A, Trivedi PP, Timbalia SA, Griffin AT, Rahn JJ, Chan SSL, and Gohil VM (2014). Copper supplementation restores cytochrome c oxidase assembly defect in a mitochondrial disease model of COA6 deficiency. *Hum Mol Genet* 23(13): 3596–3606. doi: 10.1093/hmg/ddu069
121. Vögtle F-N, Burkhart JM, Rao S, Gerbeth C, Hinrichs J, Martinou J-C, Chacinska A, Sickmann A, Zahedi RP, and Meisinger C (2012). Inter-membrane space proteome of yeast mitochondria. *Mol Cell Proteomics* 11(12): 1840–1852. doi: 10.1074/mcp.M112.021105
122. Pacheu-Grau D, Bareth B, Dudek J, Juris L, Vögtle F-N, Wissel M, Leary SC, Dennerlein S, Rehling P, and Deckers M (2015). Cooperation between COA6 and SCO2 in COX2 maturation during cytochrome c oxidase assembly links two mitochondrial cardiomyopathies. *Cell Metab* 21(6): 823–833. doi: 10.1016/j.cmet.2015.04.012
123. Stroud DA, Maher MJ, Lindau C, Vögtle F-N, Frazier AE, Surgenor E, Mountford H, Singh AP, Bonas M, Oeljeklaus S, Warscheid B, Meisinger C, Thorburn DR, and Ryan MT (2015). COA6 is a mitochondrial complex IV assembly factor critical for biogenesis of mtDNA-encoded COX2. *Hum Mol Genet* 24(19): 5404–5415. doi: 10.1093/hmg/ddv265
124. Bourens M, and Barrientos A (2017). Human mitochondrial cytochrome c oxidase assembly factor COX18 acts transiently as a membrane insertase within the subunit 2 maturation module. *J Biol Chem* 292(19): 7774–7783. doi: 10.1074/jbc.M117.778514
125. Liu Z, Song Y, Li D, He X, Li S, Wu B, Wang W, Gu S, Zhu X, Wang X, Zhou Q, Dai Y, and Yan Q (2014). The novel mitochondrial 16S rRNA 2336T>C mutation is associated with hypertrophic cardiomyopathy. *J Med Genet* 51(3): 176–184. doi: 10.1136/jmedgenet-2013-101818
126. Santorelli FM, Tanji K, Manta P, Casali C, Krishna S, Hays AP, Mancini DM, DiMauro S, and Hirano M (1999). Maternally inherited cardiomyopathy: an atypical presentation of the mtDNA 12S rRNA gene A1555G mutation. *Am J Hum Genet* 64(1): 295–300. doi: 10.1086/302188
127. Ahola S, Isohanni P, Euro L, Brillhante V, Palotie A, Pihko H, Lönnqvist T, Lehtonen T, Laine J, Tyynismaa H, and Suomalainen A (2014). Mitochondrial EFTs defects in juvenile-onset Leigh disease, ataxia, neuropathy, and optic atrophy. *Neurology* 83(8): 743–751. doi: 10.1212/WNL.0000000000000716
128. Distelmaier F, Haack TB, Catarino CB, Gallenmüller C, Rodenburg RJ, Strom TM, Baertling F, Meitinger T, Mayatepek E, Prokisch H, and Klopstock T (2015). MRPL44 mutations cause a slowly progressive multisystem disease with childhood-onset hypertrophic cardiomyopathy. *Neurogenetics* 16(4): 319–323. doi: 10.1007/s10048-015-0444-2
129. Galmiche L, Serre V, Beinat M, Assouline Z, Lebre A-S, Chretien D, Nietschke P, Benes V, Boddaert N, Sidi D, Brunelle F, Rio M, Munnich A, and Rötig A (2011). Exome sequencing identifies MRPL3 mutation in mitochondrial cardiomyopathy. *Human Mut* 32(11): 1225–1231. doi: 10.1002/humu.21562
130. Emperador S, Bayona-Bafaluy MP, Fernández-Marmiesse A, Pineda M, Felgueroso B, López-Gallardo E, Artuch R, Roca I, Ruiz-Pesini E, Couce ML, and Montoya J (2016). Molecular-genetic characterization and rescue of a TSFM mutation causing childhood-onset ataxia and nonobstructive cardiomyopathy. *Eur J Hum Genet* 25(1): 153–156. doi: 10.1038/ejhg.2016.124
131. Giordano C, Perli E, Orlandi M, Pisano A, Tuppen HA, He L, Ierinò R, Petruzzello L, Terzi A, Autore C, Petrozza V, Gallo P, Taylor RW, and d'Amati G (2013). Cardiomyopathies due to homoplasmic mitochondrial tRNA mutations: morphologic and molecular features. *Hum Pathol* 44(7): 1262–1270. doi: 10.1016/j.humpath.2012.10.011
132. Goldstein JD, Shanske S, Bruno C, and Perszyk AA (1999). Maternally inherited mitochondrial cardiomyopathy associated with a C-to-T transition at nucleotide 3303 of mitochondrial DNA in the tRNA(Leu(UUR)) gene. *Pediatr Dev Pathol* 2(1): 78–85. doi: 10.1007/s100249900094
133. Alila-Fersi O, Tabei M, Maalej M, Belguith N, Keskes L, Mkaouer-Rebai E, and Fakhfakh F (2018). First description of a novel mitochondrial mutation in the MT-TI gene associated with multiple mitochondrial DNA deletion and depletion in family with severe dilated mitochondrial cardiomyopathy. *Biochem Biophys Res Commun* 497(4): 1049–1054. doi: 10.1016/j.bbrc.2018.02.173
134. Barth PG, Valianpour F, Bowen VM, Lam J, Duran M, Vaz FM, and Wanders RJA (2004). X-linked cardioskeletal myopathy and neutropenia (Barth syndrome): an update. *Am J Med Genet A* 126A(4): 349–354. doi: 10.1002/ajmg.a.20660
135. Barth PG, Van den Bogert C, Bolhuis PA, Scholte HR, van Gennip AH, Schutgens RB, and Ketel AG (1996). X-linked cardioskeletal myopathy and neutropenia (Barth syndrome): respiratory-chain abnormalities in cultured fibroblasts. *J Inherit Metab Dis* 19(2): 157–160. doi: 10.1007/bf01799418
135. Dudek J, and Maack C (2017). Barth syndrome cardiomyopathy. *Cardiovasc Res*. doi: 10.1093/cvr/cvx014
137. Xu Y, Malhotra A, Ren M, and Schlame M (2006). The enzymatic function of tafazzin. *J Biol Chem* 281(51): 39217–39224. doi: 10.1074/jbc.M606100200
138. Vukotic M, Nolte H, König T, Saita S, Ananjew M, Krüger M, Tatsuta T, and Langer T (2017). Acylglycerol Kinase Mutated in Sengers Syndrome Is a Subunit of the TIM22 Protein Translocase in Mitochondria. *Mol Cell* 67(3): 471–483. doi: 10.1016/j.molcel.2017.06.013
139. Kang Y, Stroud DA, Baker MJ, De Souza DP, Frazier AE, Liem M, Tull D, Mathivanan S, McConville MJ, Thorburn DR, Ryan MT, and Stojanovski D (2017). Sengers Syndrome-Associated Mitochondrial Acylglycerol Kinase Is a Subunit of the Human TIM22 Protein Import Complex. *Mol Cell* 67(3): 457–470. doi: 10.1016/j.molcel.2017.06.014
140. Moraes CT, DiMauro S, Zeviani M, Lombes A, Shanske S, Miranda AF, Nakase H, Bonilla E, Werneck LC, and Servidei S (1989). Mitochondrial DNA deletions in progressive external ophthalmoplegia and Kearns-Sayre syndrome. *N Engl J Med* 320(20): 1293–1299. doi: 10.1056/NEJM198905183202001
141. Manfredi G, Schon EA, Moraes CT, Bonilla E, Berry GT, Sladky JT, and DiMauro S (1995). A new mutation associated with MELAS is located in a mitochondrial DNA polypeptide-coding gene. *Neuromuscul Disord* 5(5): 391–398. doi: 10.1016/0960-8966(94)00079-o
142. Shoffner JM, Lott MT, Lezza AM, Seibel P, Ballinger SW, and Wallace DC (1990). Myoclonic epilepsy and ragged-red fiber disease (MERRF) is associated with a mitochondrial DNA tRNA(Lys) mutation. *Cell* 61(6): 931–937. doi: 10.1201/b15310-58
143. Nissanka N, and Moraes CT (2018). Mitochondrial DNA damage and reactive oxygen species in neurodegenerative disease. *FEBS Lett* 592(5): 728–742. doi: 10.1002/1873-3468.12956
144. Lake NJ, Compton AG, Rahman S, and Thorburn DR (2016). Leigh syndrome: One disorder, more than 75 monogenic causes. *Ann Neurol* 79(2): 190–203. doi: 10.1002/ana.24551
145. Wojtala A, Karkucinska-Wieckowska A, Sardao VA, Szczepanowska J, Kowalski P, Pronicki M, Duszyński J, and Wieckowski MR (2017). Modulation of mitochondrial dysfunction-related oxidative stress in fibroblasts of patients with Leigh syndrome by inhibition of prooxidative p66Shc pathway. *Mitochondrion* 37: 62–79. doi: 10.1016/j.mito.2017.07.002

146. Spinazzola A, Viscomi C, Fernandez-Vizarra E, Carrara F, D'Adamo P, Calvo S, Marsano RM, Donnini C, Weiher H, Strisciuglio P, Parini R, Sarzi E, Chan A, DiMauro S, Rötig A, Gasparini P, Ferrero I, Mootha VK, Tiranti V, and Zeviani M (2006). MPV17 encodes an inner mitochondrial membrane protein and is mutated in infantile hepatic mitochondrial DNA depletion. *Nat Genet* 38(5): 570–575. doi: 10.1038/ng1765
147. Spinazzola A, Santer R, Akman OH, Tsiakas K, Schaefer H, Ding X, Karadimas CL, Shanske S, Ganesh J, Di Mauro S, and Zeviani M (2008). Hepatocerebral form of mitochondrial DNA depletion syndrome: novel MPV17 mutations. *Arch Neurol* 65(8): 1108–1113. doi: 10.1001/archneur.65.8.1108
148. Spinazzola A, Massa V, Hirano M, and Zeviani M (2008). Lack of founder effect for an identical mtDNA depletion syndrome (MDS)-associated MPV17 mutation shared by Navajos and Italians. *Neuromuscul Disord* 18(4): 315–318. doi: 10.1016/j.nmd.2007.12.007
149. El-Hattab AW, Wang J, Dai H, Almannai M, Stauffer C, Alfadhel M, Gambello MJ, Prasun P, Raza S, Lyons HJ, Afqi M, Saleh MAM, Faqeh EA, Alzaidan HI, Alshenqiti A, Flore LA, Hertecant J, Sacharow S, Barbooth DS, Murayama K, Shah AA, Lin HC, and Wong L-JC (2018). MPV17-related mitochondrial DNA maintenance defect: New cases and review of clinical, biochemical, and molecular aspects. *Human Mut* 39(4): 461–470. doi: 10.1002/humu.23387
150. Miyake N, Yamashita S, Kurosawa K, Miyatake S, Tsurusaki Y, Doi H, Saito H, and Matsumoto N (2011). A novel homozygous mutation of DARS2 may cause a severe LBSL variant. *Clin Genet* 80(3): 293–296. doi: 10.1111/j.1399-0004.2011.01644.x
151. S ahin S, Cansu A, Kalay E, Dinçer T, Kul S, Çakır ISM, Kamaşak T, and Budak GYGL (2016). Leukoencephalopathy with thalamus and brainstem involvement and high lactate caused by novel mutations in the EARS2 gene in two siblings. *J Neurol Sci* 365: 54–58. doi: 10.1016/j.jns.2016.04.008
152. Boczonadi V, Jennings MJ, and Horvath R (2018). The role of tRNA synthetases in neurological and neuromuscular disorders. *FEBS Lett* 592(5): 703–717. doi: 10.1002/1873-3468.12962
153. Viscomi C (2016). Toward a therapy for mitochondrial disease. *Biochem Soc Trans* 44(5): 1483–1490. doi: 10.1042/BST20160085
154. Scarpulla RC (2008). Transcriptional paradigms in mammalian mitochondrial biogenesis and function. *Physiol Rev* 88(2): 611–638. doi: 10.1152/physrev.00025.2007
155. Puigserver P, and Spiegelman BM (2003). Peroxisome proliferator-activated receptor-gamma coactivator 1 alpha (PGC-1 alpha): transcriptional coactivator and metabolic regulator. *Endocr Rev* 24(1): 78–90. doi: 10.1210/er.2002-0012
156. Villena JA (2015). New insights into PGC-1 coactivators: redefining their role in the regulation of mitochondrial function and beyond. *FEBS J* 282(4): 647–672. doi: 10.1111/febs.13175
157. Perales-Clemente E, Bayona-Bafaluy MP, Pérez-Martos A, Barrientos A, Fernández-Silva P, and Enríquez JA (2008). Restoration of electron transport without proton pumping in mammalian mitochondria. *Proc Natl Acad Sci U S A* 105(48): 18735–18739. doi: 10.1073/pnas.0810518105
158. Dassa EP, Dufour E, Gonçalves S, Paupe V, Hakkaert GAJ, Jacobs HT, and Rustin P (2009). Expression of the alternative oxidase complements cytochrome c oxidase deficiency in human cells. *EMBO Mol Med* 1(1): 30–36. doi: 10.1002/emmm.200900001
159. Kempainen KK, Rinne J, Sriram A, Lakanmaa M, Zeb A, Tuomela T, Popplestone A, Singh S, Sanz A, Rustin P, and Jacobs HT (2014). Expression of alternative oxidase in *Drosophila* ameliorates diverse phenotypes due to cytochrome oxidase deficiency. *Hum Mol Genet* 23(8): 2078–2093. doi: 10.1093/hmg/ddt601
160. Camargo AF, Chioda MM, Rodrigues APC, Garcia GS, McKinney EA, Jacobs HT, and Oliveira MT (2018). Xenotopic expression of alternative electron transport enzymes in animal mitochondria and their impact in health and disease. *Cell Biol Int* 42(6):664–669. doi: 10.1002/cbin.10943
161. Szibor M, Dhandapani PK, Dufour E, Holmström KM, Zhuang Y, Salwig I, Wittig I, Heidler J, Gizatullina Z, Gainutdinov T, German Mouse Clinic Consortium, Fuchs H, Gailus-Durner V, de Angelis MH, Nandania J, Velagapudi V, Wietelmann A, Rustin P, Gellerich FN, Jacobs HT, and Braun T (2017). Broad AOX expression in a genetically tractable mouse model does not disturb normal physiology. *Dis Model Mech* 10(2): 163–171. doi: 10.1242/dmm.027839
162. Civileto G, Varanita T, Cerutti R, Gorletta T, Barbaro S, Marchet S, Lamperti C, Viscomi C, Scorrano L, and Zeviani M (2015). Opa1 overexpression ameliorates the phenotype of two mitochondrial disease mouse models. *Cell Metab* 21(6): 845–854. doi: 10.1016/j.cmet.2015.04.016
163. Varanita T, Soriano ME, Romanello V, Zaglia T, Quintana-Cabrera R, Semenzato M, Menabò R, Costa V, Civileto G, Pesce P, Viscomi C, Zeviani M, Di Lisa F, Mongillo M, Sandri M, and Scorrano L (2015). The OPA1-dependent mitochondrial cristae remodeling pathway controls atrophic, apoptotic, and ischemic tissue damage. *Cell Metab* 21(6): 834–844. doi: 10.1016/j.cmet.2015.05.007
164. Beyrath J, Pellegrini M, Renkema H, Houben L, Pecheritsyna S, van Zandvoort P, van den Broek P, Bekel A, Eftekhari P, and Smeitink JAM (2018). KH176 Safeguards Mitochondrial Diseased Cells from Redox Stress-Induced Cell Death by Interacting with the Thioredoxin System/Peroxisome Enzyme Machinery. *Sci Rep* 8(1): 6577. doi: 10.1038/s41598-018-24900-3
165. Koopman WJ, Beyrath J, Fung C-W, Koene S, Rodenburg RJ, Willems PH, and Smeitink JA (2016). Mitochondrial disorders in children: toward development of small-molecule treatment strategies. *EMBO Mol Med* 8(4): 311–327. doi: 10.15252/emmm.201506131
166. Bacman SR, Williams SL, Pinto M, Peralta S, and Moraes CT (2013). Specific elimination of mutant mitochondrial genomes in patient-derived cells by mitoTALENs. *Nat Med* 19(9): 1111–1113. doi: 10.1038/nm.3261
167. Gammage PA, Rorbach J, Vincent AI, Rebar EJ, and Minczuk M (2014). Mitochondrially targeted ZFNs for selective degradation of pathogenic mitochondrial genomes bearing large-scale deletions or point mutations. *EMBO Mol Med* 6(4): 458–466. doi: 10.1002/emmm.201303672
168. Wang G, Shimada E, Zhang J, Hong JS, Smith GM, Teitell MA, and Koehler CM (2012). Correcting human mitochondrial mutations with targeted RNA import. *Proc Natl Acad Sci U S A* 109(13): 4840–4845. doi: 10.1073/pnas.1116792109
169. Gammage PA, Moraes CT, and Minczuk M (2017). Mitochondrial Genome Engineering: The Revolution May Not Be CRISPR-ized. *Trends Genet* 9525(17): 30191–9. doi: 10.1016/j.tig.2017.11.001
170. Frazier AE, Thorburn DR, and Compton AG (2017). Mitochondrial energy generation disorders: genes, mechanisms and clues to pathology. *J Biol Chem*. doi: 10.1074/jbc.R117.809194
171. Nikkanen J, Forsström S, Euro L, Paetau I, Kohnz RA, Wang L, Chilov D, Viinämäki J, Roivainen A, Marjamäki P, Liljenbäck H, Ahola S, Buzkova J, Terzioglu M, Khan NA, Pirnes-Karhu S, Paetau A, Lönnqvist T, Sajantila A, Isohanni P, Tyynismaa H, Nomura DK, Battersby BJ, Velagapudi V, Carroll CJ, and Suomalainen A (2016). Mitochondrial DNA Replication Defects Disturb Cellular dNTP Pools and Remodel One-Carbon Metabolism. *Cell Metab* 23(4): 635–648. doi: 10.1016/j.cmet.2016.01.019
172. Khan NA, Nikkanen J, Yatsuga S, Jackson C, Wang L, Pradhan S, Kivelä R, Pessia A, Velagapudi V, and Suomalainen A (2017). mTORC1 Regulates Mitochondrial Integrated Stress Response and Mitochon-

drial Myopathy Progression. **Cell Metab** 26(2): 419–428. doi: 10.1016/j.cmet.2017.07.007

173. Pacheu-Grau D, Gómez-Durán A, Iglesias E, López-Gallardo E, Montoya J, and Ruiz-Pesini E (2013). Mitochondrial antibiograms in personalized medicine. **Hum Mol Genet** 22(6): 1132–1139. doi: 10.1093/hmg/dds517