

Original Article



Protease-Activated Receptors 2-Antagonist Suppresses Asthma by Inhibiting Reactive Oxygen Species-Thymic Stromal Lymphopoietin Inflammation and Epithelial Tight Junction Degradation

OPEN ACCESS

Received: Sep 25, 2018

Revised: Feb 10, 2019

Accepted: Feb 12, 2019

Correspondence to

Soo-Jong Hong, MD, PhD

Department of Pediatrics, Childhood Asthma Atopy Center, Asan Medical Center, University of Ulsan College of Medicine, 88 Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea.

Tel: +82-2-3010-3379

Fax: +82-2-473-3725

E-mail: sjhong@amc.seoul.kr

Copyright © 2019 The Korean Academy of Asthma, Allergy and Clinical Immunology · The Korean Academy of Pediatric Allergy and Respiratory Disease

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Ha-Jung Kim

<https://orcid.org/0000-0002-2699-0263>

Seung-Hwa Lee

<https://orcid.org/0000-0001-9359-0318>

Sekyoo Jeong

<https://orcid.org/0000-0002-9201-0227>

Soo-Jong Hong

<https://orcid.org/0000-0003-1409-2113>

Disclosure

There are no financial or other issues that might lead to conflict of interest.

Ha-Jung Kim ¹, **Seung-Hwa Lee** ², **Sekyoo Jeong** ³, **Soo-Jong Hong** ^{4,5*}

¹Department of Internal Medicine, Chonnam National University College of Veterinary Medicine, Gwangju, Korea

²Asan Institute for Life Sciences, University of Ulsan College of Medicine, Seoul, Korea

³CRID Center, NeoPharm Co., Ltd., Daejeon, Korea

⁴Department of Pediatrics, Childhood Asthma Atopy Center, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

⁵Environmental Health Center, Asan Medical Center, Seoul, Korea

ABSTRACT

Purpose: Protease-activated receptor 2 (PAR2) reportedly triggers the immune response in allergic asthma. We aimed to investigate the mechanism on allergic inflammation mediated by PAR2.

Methods: Human lung epithelial cells (A549 cells) were used for *in vitro*, and the German cockroach extract (GCE)-induced mouse model was developed for *in vivo* studies.

Results: In A549 cells, the levels of reactive oxygen species (ROS) and thymic stromal lymphopoietin (TSLP) were significantly increased by GCE treatment, but were suppressed by PAR2-antagonist (PAR2-ant) or N-acetylcysteine (NAC) treatment. Claudin-1 was degraded by GCE, and was restored by PAR2-ant or NAC in the cells. In the mouse model, the clinical appearance including bronchial hyperresponsiveness, bronchoalveolar lavage fluid analysis and total immunoglobulin E were significantly suppressed by PAR2-ant or NAC. Moreover, TSLP levels in the lung were suppressed by the same treatments in the lung. Claudin-1 was also degraded by GCE, and was restored by PAR2-ant or NAC.

Conclusions: ROS generation and epidermal tight junction degradation are triggered by protease, followed by the induction of TSLP in allergic asthma. Our findings could suggest that PAR2-ant or anti-oxidants could be considered for allergic diseases as preventive alternatives.

Keywords: Asthma; protease-activated receptor 2; thymic stromal lymphopoietin; tight junction; reactive oxygen species

INTRODUCTION

More than 3 billion people suffer from allergies (*i.e.*, asthma, atopic dermatitis and rhinitis) worldwide. Several protein allergens in the environment could induce allergic immune responses through activation of their functional components, including proteases.¹⁻³

Protease-activated receptors (PARs) are activated by proteases. Although PAR2 reportedly mediates the T helper 2 (Th2)-immune response in allergic diseases, the exact underlying mechanism is poorly understood.^{4,5} In addition, thymic stromal lymphopoietin (TSLP) is produced by epithelial cells and triggers Th2-type inflammation by proteases.⁵ Moreover, PAR2 induces TSLP activation in response to serine proteases.⁵ The body is protected against the inflammation caused by proteases on epithelial cells due to the mucosal barrier (*e.g.*, lung and skin). However, it is unclear whether these mechanisms are involved in allergic human diseases.

Reactive oxygen species (ROS) have recently been reported as an important factor influencing the development of allergic diseases.⁶⁻⁸ ROS also induce the Th2 response through the oxidization of lipids on epithelial cells, which in turn is mediated by the toll-like receptor 4 (TLR4) and the adaptor protein TIR-domain-containing adapter-inducing interferon (IFN)- β .⁹ Although the relationship between ROS and PAR2 has been addressed in a study on allergic asthma,¹⁰ the exact mechanism remains unclear.

In the present study, we aimed to assess the mechanism underlying the induction of allergic asthma by a cockroach allergen extract that includes protease. First, we evaluated the PAR2-ROS-TSLP-related initiation of inflammation and mucosal tight junction degradation caused by protease in the mouse airway *in vivo* and in human bronchial epithelial cells (A549 cells) *in vitro*. Secondly, the PAR2-antagonist (PAR2-ant) and anti-oxidants were used to block the initiation of inflammation in a mouse model of asthma and in A549 cells. Our findings could suggest whether alternative new treatments with PAR2-ant or anti-oxidants could be considered for allergic diseases.

MATERIALS AND METHODS

German cockroach extract (GCE) preparation

GCE was prepared as previously described.⁹ Fifty grams of frozen German cockroaches were homogenized in liquid nitrogen and defatted in 200 mL of ethyl ether and 200 mL of ethyl acetate. Extractions were performed with slow stirring at 4°C overnight in phosphate buffer saline (PBS) containing 6 mmol/L 2-mercaptoethanol and 1 mg/mL 1-phenyl-3-(2-thiazolyl)-2-thiourea to prevent melanization. The extract was then centrifuged at 10,000 \times g for 30 minutes at 4°C, and the supernatant was finally filtered through 0.2- μ m filters. The endotoxin from the supernatant was removed by the Detoxi-Gel™ Endotoxin Removing Gel (Pierce Biotechnology, Rockford, IL, USA). The endotoxin from the GCE was removed and measured to be below 0.01 EU/mL by the chromogenic Limulus Amebocyte Lysate test (Lonza, Walkersville, MD, USA). The GCE contained 0.11 U/mg Bla g1.

ROS levels

The ROS levels were measured in human bronchial epithelial cells (A549) (Costar, Cambridge, MA, USA) *in vitro* and in the mouse lung *in vivo*. By using the OxiSelect ROS Assay Kit (Cell Biolabs, San Diego, CA, USA), ROS levels on A549 cells were measured according to the manufacturer's guidelines. In brief, the attached cells seeded on a black 96-well plate (1×10^5 cells/well) were pretreated with washing buffer and then treated with 100 μ L of the cell permeable fluorogenic probe 2', 7' dichlorodihydrofluorescein diacetate (DCFH-DA, Sigma-Aldrich, St Louis, MO, USA) in Dulbecco's modified eagle medium (DMEM, Sigma-Aldrich) media for 1 hour at 37°C. After the cells were washed 3 times with Dulbecco's PBS, GCE (100 μ g/mL) with or without PAR2-ant (NPS-1577, 200 μ M; NeoPharm Pharmacy, Daejeon, Korea) was added to the

DMEM media. Pretreatment with PAR2-ant was ensured for 30 minutes prior to GCE treatment. To quench ROS generation, 2 mM of N-acetyl-L-cysteine (NAC; Sigma A9165; Sigma-Aldrich, St Louis, MO, USA) was added. The plates were read at excitation and emission wavelengths of 450 and 520 nm, respectively, every 30 minutes for 2 hours by using a VICTOR™ X3 Multilabel Plate Reader (PerkinElmer, Santa Clara, CA, USA).

By using immunohistochemistry (IHC), the expression of ROS-related enzyme glutathione S-transferase 1 (GSTP1) was measured in the mouse lung *in vivo* (described in IHC section).

Cell viability assay

Cell viability was determined by using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. In brief, A549 cells at the logarithmic growth phase were randomly seeded onto 96-well culture plates at a density of 1×10^3 cells/mL, and were cultured with 100 μ L DMEM media (supplemented with 10% fetal bovine serum [FBS]) in each well. After cell adhesion was achieved, the cells were incubated with various concentrations of GCE (0, 50, 100, 200 and 400 μ g/mL) for 1, 2, 4 and 8 hours with or without PAR2-ant (200 μ M) and NAC (2 mM). For the cell viability assay, 10 μ L of MTT solution (Sigma-Aldrich; 5 mg/mL) was added to each well, and incubation was continued at 37°C for 4 hours. Following centrifugation at 3,000 rpm for 10 minutes, the supernatant was removed to obtain the formazan pellet. The pellet was then dissolved completely with 100 μ L of dimethyl sulfoxide (DMSO). An enzyme linked immunosorbent assay (ELISA) plate reader was used to measure the absorbance at a wavelength of 570 nm, in order to determine the amount of pellet.

Real-time reverse transcription polymerase chain reaction (PCR) in the A549 cells

A549 cells were cultured in DMEM (Sigma-Aldrich) media and supplemented with 10% FBS (Sigma-Aldrich) and 1% antibiotics (penicillin/streptomycin; Invitrogen, Carlsbad, CA, USA) under standard tissue culture conditions. A total of 6×10^4 cells were seeded per well on 12-well plates, which were exposed to 100 μ g/mL or 150 μ g/mL (claudin-1) of GCE with or without 200 μ M PAR2-ant and 2mM NAC for 2 hours.

Pretreatment with PAR2-ant was ensured for 30 minutes prior to GCE treatment. To confirm the PAR2 agonist (PAR2-ago) function of GCE, a positive control for the PAR2-ago (100 μ M, SLIGKV-NH₂; R&D Systems, Minneapolis, MN, USA) was also examined. To determine the optimal duration for each mediator (TSLP/claudin-1) on exposure to GCE, the expressions were measured at different time points (0, 2, 6, 8, 24, 36 and 48 hours) by using real-time PCR. At the optimal duration for each mediator, the other material (PAR2-ant, NAC and PAR2-ago) was added to the GCE-treated cells. Real-time PCR was performed with specific primers and probes using TaqMan Gene Expression Assays and TaqMan Gene Expression Master Mix (Applied Biosystems, Foster City, CA, USA) on an ABI 7500 system (Applied Biosystems, Piscataway, NJ, USA). The signals were normalized using glyceraldehyde 3-phosphate dehydrogenase (GAPDH).

Mouse model and treatment

Female BALB/c mice weighing 20–25 g (4 weeks of age) were purchased from OrientBio (Seongnam, Korea) and cared for and used in accordance with the guidelines of the Institutional Animal Care and Use Committee at Asan Medical Center and Ulsan University College of Medicine. The mice were intranasally sensitized to GCE (100 μ g in PBS) on days 1, 3, 5, 8, 10 and 12, whereas PAR2-ant (0.02% in 1% DMSO, intranasally) or NAC (100 mg/kg in PBS, intraperitoneally) was administered 30 minutes before GCE treatment on days 1, 3 and 5. Five mice were included in each group.

Clinical evaluations and analysis *in vivo*

The bronchial hyperresponsiveness (BHR) in response to inhaled methacholine (MeCh; Sigma-Aldrich) was measured in conscious, unrestrained mice by using a barometric whole-body plethysmograph (Buxco; EMKA Technologies, Paris, France). For bronchoalveolar lavage (BAL) fluid analysis, the airways were lavaged 2 times through a tracheal cannula with 1-mL aliquots of pyrogen-free saline warmed to 37°C. The recovered lavage fluid was pooled, and the cells were collected by centrifugation (5,000 rpm, 4°C and 5 minutes) and resuspended in 100 mL of cold PBS. The cells were stained with trypan blue to determine viability, and the total number of nucleated cells were counted using a hemocytometer. For differential BAL cell counts, cytospin preparations were made and stained with Diff-Quik (Sysmex, Takatsukadai, Japan). After the samples were coded, all cytospin preparations were evaluated by a single observer using an oil immersion microscope (magnification, × 1,000). At least 200 cells were counted per preparation, and the absolute number of each cell type was calculated.

Serum was obtained after BHR measurement, and 100 µL (1/10 dilution in carbonate-bicarbonate buffer) was added to each well of a 96-well plate. An immunoglobulin E (IgE)-specific ELISA was used to determine the total IgE levels in the serum by using matching antibody pairs (eBioscience, San Diego, CA, USA) according to the manufacturer's instructions.

For the histological evaluation of lung tissue, the left lung of each mouse was embedded in paraffin, sectioned to a thickness of 5 µm, and stained with hematoxylin and eosin to assess eosinophilic infiltration. Inflammation was scored by 2 independent investigators in a blind manner. The degree of peribronchial and perivascular inflammation was evaluated on a subjective scale of 0–3.

To measure the levels of inflammatory cytokines (interleukin [IL]-4, IL-13, and IFN-γ), whole-lung RNA was extracted using RNeasy (Qiagen, Valencia, CA, USA), and real-time PCR was performed with specific primers and probes using TaqMan Gene Expression Assays and TaqMan Gene Expression Master Mix (Applied Biosystems) on an ABI 7500 system (Applied Biosystems). Signals were normalized using GAPDH.

IHC

Sections from formalin-fixed, paraffin-embedded tissue blocks of lung were cut (5-µm thickness) and mounted on glass slides. IHC was conducted using an automated slide preparation system Benchmark XT (Ventana Medical systems Inc, Tucson, AZ, USA). Deparaffinization, epitope retrieval and immunostaining were performed according to the manufacturer's instructions by using cell conditioning solutions and the BMK ultraVIEW diaminobenzidine detection system (Ventana Medical Systems). Lung sections were stained with TSLP (1:700 dilution, ab115700; Abcam, Cambridge, United Kingdom), GSTP1 (1:800 dilution, GTX112695; GeneTex, Irvine, CA, USA), and claudin-1 (1:25 dilution, 71-7800; Invitrogen). Positive signals were amplified by using ultraVIEW copper, and sections were counterstained with hematoxylin and bluing reagent.

Determination of immunohistopathological staining intensity

Immunohistopathological staining was objectively analyzed by using a computer imaging program (ImageJ® 1.41; <http://rsb.info.nih.gov/ij/>).

Statistical analysis

The statistical differences of each parameter among the groups were calculated (GraphPad Prism 5.0, San Diego, CA, USA) by using 1-way analysis of variance (Tukey's multiple comparison test). A *P* value of < 0.05 was considered to indicate statistical significance.

RESULTS

GCE induces inflammation and degrades claudin-1 via a PAR2- and ROS-mediated mechanism in human bronchial epithelial cells (A549 cells)

DCFH-DA analysis indicated that GCE treatment generated ROS after 2 hours in A549 cells, whereas PAR2-ant and NAC treatment significantly suppressed ROS generation (Fig. 1A and B). This suggests that GCE induces ROS generation, which is mediated via PAR2.




Fig. 1. GCE induces inflammation and degrades tight junctions in human bronchial epithelial cells (A549 cells). The 2', 7'-DCF intensity is significantly increased following GCE exposure and suppressed following PAR2-ant and NAC treatment in A549 cells (1 hour (A) and 2 hours (B)). The expression of TSLP is significantly increased following GCE and PAR2-ago treatment and suppressed following PAR2-ant and NAC treatment in A549 cells after 2 hours (C). The expression of claudin-1 is significantly decreased by GCE and PAR2-ago treatment, but is significantly restored by PAR2-ant and NAC treatment in A549 cells (D). (A, B) Fluorescence intensity measurement at wavelengths of 450 nm and 520 nm; (C, D) real-time polymerase chain reaction (Taqman method).

GCE, German cockroach extract; NAC, N-acetylcysteine; DCF, dichlorofluorescein; PAR2-ant, protease-activated receptor 2-antagonist; PAR2-ago, protease-activated receptor 2 agonist; TSLP, thymic stromal lymphopoietin.

* $P < 0.01$ and † $P < 0.001$. The experiments were repeated 6 times.

Effects of PAR2 Antagonist in a Model of Asthma




Fig. 2. GCE induces allergic inflammation in the mouse model of asthma. (A) BHR is significantly increased following GCE treatment and suppressed following PAR2-ant and NAC treatment. $^{\ddagger}P < 0.001$ compared to saline. (B) The total number of cells in bronchoalveolar lavage fluid is also significantly increased following GCE treatment and suppressed following PAR2-ant and NAC treatment. (C) The numbers of macrophages, eosinophils and neutrophils are significantly increased following GCE treatment and suppressed following PAR2-ant and NAC treatment. (D) The total immunoglobulin E level is also significantly increased following GCE treatment and suppressed following PAR2-ant and NAC treatment. (E) Lung inflammation (hematoxylin and eosin staining 200 \times magnification) and (F) peribronchovascular inflammation is significantly increased following GCE treatment and suppressed following PAR2-ant and NAC treatment. BHR, bronchial hypersensitivity; GCE, German cockroach extract; CE, cockroach extract; NAC, N-acetylcysteine; PAR2-ant, protease-activated receptor 2-antagonist. $^*P < 0.05$, $^{\dagger}P < 0.01$, and $^{\ddagger}P < 0.001$. The experiments were repeated 5 times.

To evaluate the ROS- and PAR2-related inflammatory mechanism in the A549 cells following exposure to GCE, we measured the levels of TSLP and claudin-1 using real-time PCR. The expression of TSLP was significantly increased by GCE or PAR2-ago treatment after 2 hours (**Fig. 1C**). However, the expression of claudin-1 was significantly suppressed by GCE or PAR2-ago treatment after 6 hours (**Fig. 1D**). Thus, it is likely that the expression of inflammatory cytokine TSLP and tight junction protein claudin-1 is mediated by PAR2 following exposure to GCE in A549 cells.

The expression of TSLP was significantly suppressed by PAR2-ant and NAC treatment (**Fig. 1C**), whereas the expression of claudin-1 was significantly increased following PAR2-ant and NAC treatment (**Fig. 1D**). Hence, the induction of inflammatory mechanisms involving the TSLP and tight junction protein degradation by GCE may be mediated by PAR2 and ROS.

GCE induces allergic inflammation, which is mediated by a PAR2 and ROS mechanism, in a mouse model of asthma

To confirm whether the PAR2 and ROS mechanism was involved in the response to GCE *in vivo* in a similar manner to that noted *in vitro*, a mouse model of asthma was established using GCE. The phenotypes of allergic asthma including BHR, total cell count in BAL fluid, total IgE levels, histopathology and lung inflammatory score were significantly increased following GCE treatment (**Fig. 2**). Differential cell counts indicated eosinophilic and neutrophilic inflammation in the BAL fluid (**Fig. 2C**). The phenotypes were significantly suppressed by PAR2-ant and NAC treatment. This suggests that the allergic inflammation in the mouse model was mediated via the PAR2 and ROS mechanism.

GCE induces Th2 inflammation in the lung, which is mediated by the PAR2 and ROS mechanism, in a mouse model of asthma

The level of Th2 cytokines IL-4 and IL-13 were significantly increased in the lung following GCE treatment, and were suppressed following PAR2-ant and NAC treatment (**Fig. 3A and C**). In contrast, the expression of Th1 cytokine IFN- γ was significantly suppressed by GCE treatment and increased following PAR2-ant and NAC treatment (**Fig. 3B**). This suggests that GCE induces Th2 inflammation in the lung, which is mediated by the PAR2 and ROS mechanism, in a mouse model of asthma.

GCE induces ROS-TSLP-related inflammation in the lung, which is mediated by PAR2, in a mouse model of asthma

The expression of ROS-related enzyme GSTP1 in the lung was significantly decreased following GCE treatment. IHC of the lung indicated that the decreased GSTP1 levels had recovered following PAR2-ant and NAC treatment (**Fig. 4**). Moreover, the expression of TSLP in the lung appeared to be regulated via the PAR2-ant and NAC mechanism, as per the findings of IHC examination (**Fig. 5**). Thus, the ROS generation by GCE treatment was followed by an increase in TSLP expression, which was mediated by PAR2, in the lung of a mouse model of asthma.

GCE causes disruption of the tight junction protein in the lung, which is mediated by a PAR2- and ROS-related mechanism, in a mouse model of asthma

Claudin-1 expression was disrupted by GCE treatment in the lung, but recovered following PAR2-ant and NAC treatment, as noted on IHC examination (**Fig. 6**). This suggests that GCE induces tight junction degradation, which is mediated by a PAR2- and ROS-related inflammatory mechanism.

Effects of PAR2 Antagonist in a Model of Asthma




Fig. 3. GCE induces T helper 2-allergic inflammation in a mouse model of asthma. The levels of IL-4 and IL-13 are significantly increased following GCE treatment and suppressed following PAR2-ant and NAC treatment in the lung (A and C). Moreover, the level of IFN- γ is significantly decreased following GCE treatment and increased following PAR2-ant and NAC treatment (B). Data were measured using real-time polymerase chain reaction. GCE, German cockroach extract; IL, interleukin; IFN, interferon; PAR2-ant, protease-activated receptor 2-antagonist; NAC, N-acetylcysteine. * $P < 0.05$; † $P < 0.01$. The experiments were repeated 5 times.




Fig. 4. GCE significantly suppresses GSTP1 expression in a mouse model. GSTP1 expression is significantly increased by PAR2-ant and NAC treatment in the lung, as indicated by immunohistochemistry (immunohistochemistry, 200 \times magnification) (A) and intensity quantification (B). GCE, German cockroach extract; GSTP1, glutathione S-transferase 1; NAC, N-acetylcysteine; PAR2-ant, protease-activated receptor 2-antagonist. * $P < 0.01$; † $P < 0.001$. The experiments were repeated 5 times.




Fig. 5. GCE significantly increases TSLP expression in a mouse model. TSLP expression is significantly suppressed by PAR2-ant and NAC treatment in the lung, as indicated by immunohistochemistry (200 × magnification) (A) and intensity quantification (B).

GCE, German cockroach extract; TSLP, thymic stromal lymphopoietin; NAC, N-acetylcysteine; PAR2-ant, protease-activated receptor 2-antagonist.

* $P < 0.001$. The experiments were repeated 5 times.




Fig. 6. GCE significantly disturbs claudin-1 expression in a mouse model. Claudin-1 expression is significantly restored by PAR2-ant and NAC treatment in the lung, as indicated by immunohistochemistry (200 × magnification) (A) and intensity quantification (B).

GCE, German cockroach extract; PAR2-ant, protease-activated receptor 2-antagonist; NAC, N-acetylcysteine.

* $P < 0.01$; † $P < 0.001$. The experiments were repeated 5 times.

DISCUSSION

In the present study, we found that ROS-TSLP-related inflammation was induced by GCE exposure via PAR2 in human bronchial epithelial cells and a mouse model of asthma. Moreover, we observed that tight junction proteins could be affected in allergic inflammation via PAR2 in a mouse model of asthma. Although asthma is regarded as an airway inflammatory disease that is driven by a skewed Th2 mechanism, the process of initiation of the disease on the surface of respiratory epithelial cells remains unclear. Protease allergens reportedly elicit airway reactions by triggering innate immunity receptors, such as PARs, and thus activating epithelial cells, mast cells and dendritic cells (DCs), which in turn induce the additional release of mediators.¹¹ In patients with bronchial asthma, PAR2 expression is increased on the surface of respiratory epithelial cells.¹² Hence, blocking of PAR2 could be considered as an option for the treatment of asthma. Although some studies have indicated the presence of a PAR2-related mechanism in the mouse model,^{10,13} no studies have shown any specific association among the mediators that could play an important role in the development of asthma.

The present study showed that the protease-containing allergen GCE first activated PAR2, and then induced ROS generation in A549 cells and in the mouse model. The present study also showed PAR2 could be one of the possible mechanism of allergic inflammation. ROS play a central role in orchestrating Th2 inflammation, by inducing the formation of oxidized lipids that trigger the TLR4 and toll or IL-1 receptor domain-containing adaptor-inducing IFN- β -mediated induction of TSLP in epithelial cells.⁹ In the present study, the expression of TSLP was regulated by PAR2-ant or anti-oxidants, and ROS generation was mediated by PAR2. Hence, TSLP appears to be induced by the ROS generated by PAR2 activation, consistent with the findings of a previous study.⁹

Mutation of the barrier protein *filaggrin* is a critical factor in the pathogenesis of atopic dermatitis.^{14,15} Although *filaggrin* is not expressed in the lower airway respiratory epithelium,¹⁶ the barrier function of the airway epithelium is impaired in asthma, thus indicating a shared common underlying pathogenic mechanism.¹⁷ In the present study, the expression of claudin-1 was significantly decreased following exposure to GCE or PAR2-ago in A549 cells, and was also decreased in the lung of a mouse model sensitized to GCE. Thus, the findings confirm that the tight junction protein claudin-1 is closely related to the development of asthma. The expression was also regulated by PAR2 and ROS *in vitro* and *in vivo*, suggesting that tight junction proteins could be degraded through the activation of PAR by proteases in allergens and via the generation of ROS.

The initiation of sensitization to allergens in the airway is preceded by their uptake and processing by a subpopulation of mucosal DCs, followed by the presentation of specific peptide epitopes to naïve T cells in association with major histocompatibility class II.¹⁸ Protease enzymatic activity of allergens that could promote intraepithelial DCs functions for allergic reactions by cleaving the tight-junction proteins.¹⁹ The activation of DCs could induce Th2 cell responses via epithelial cell-derived factors such as TSLP, which conditions DCs to induce Th2 cell responses.^{20,21} Thus, allergen source-derived proteases compromise epithelial barrier function by degrading tight junction proteins, thus facilitating allergen delivery across the epithelial layers.¹⁷ The above mentioned theory is summarized in **Supplementary Fig. S1**.

In conclusion, GCE activates PAR2 and generates ROS, which degrade tight junction proteins and facilitate the penetration of allergens to induce TSLP-Th2 inflammation in airway epithelial cells *in vitro* and in the lung of a mouse model. Thus, the blocking of PAR2 is an alternative treatment option for allergic inflammation.

ACKNOWLEDGMENTS

This study was supported by a grant of the Korean Health Technology R&D Project, Ministry of Health & Welfare, Republic of Korea (HI13C1674).

SUPPLEMENTARY MATERIAL

Supplementary Fig. S1

A summary of the suggested mechanism underlying asthma. PAR2, activated by a protease-containing allergen, induces the generation of ROS, which then degrades tight

junction proteins and facilitates TSLP-Th2 inflammation via DC modulation in bronchial epithelial cells.

[Click here to view](#)

REFERENCES

1. Milovanovic M, Heine G, Zuberbier T, Worm M. Allergen extract-induced interleukin-10 in human memory B cells inhibits immunoglobulin E production. *Clin Exp Allergy* 2009;39:671-8.
[PUBMED](#) | [CROSSREF](#)
2. Traidl-Hoffmann C, Jakob T, Behrendt H. Determinants of allergenicity. *J Allergy Clin Immunol* 2009;123:558-66.
[PUBMED](#) | [CROSSREF](#)
3. Chapman MD, Pomés A, Breiteneder H, Ferreira F. Nomenclature and structural biology of allergens. *J Allergy Clin Immunol* 2007;119:414-20.
[PUBMED](#) | [CROSSREF](#)
4. Ramachandran R, Noorbakhsh F, Defea K, Hollenberg MD. Targeting proteinase-activated receptors: therapeutic potential and challenges. *Nat Rev Drug Discov* 2012;11:69-86.
[PUBMED](#) | [CROSSREF](#)
5. Kouzaki H, O'Grady SM, Lawrence CB, Kita H. Proteases induce production of thymic stromal lymphopoietin by airway epithelial cells through protease-activated receptor-2. *J Immunol* 2009;183:1427-34.
[PUBMED](#) | [CROSSREF](#)
6. Kim HJ, Lee E, Lee SH, Kang MJ, Hong SJ. Mold elicits atopic dermatitis by reactive oxygen species: epidemiology and mechanism studies. *Clin Immunol* 2015;161:384-90.
[PUBMED](#) | [CROSSREF](#)
7. Qu J, Li Y, Zhong W, Gao P, Hu C. Recent developments in the role of reactive oxygen species in allergic asthma. *J Thorac Dis* 2017;9:E32-43.
[PUBMED](#) | [CROSSREF](#)
8. Lin TJ, Karmaus WJJ, Chen ML, Hsu JC, Wang JJ. Interactions between bisphenol A exposure and GSTP1 polymorphisms in childhood asthma. *Allergy Asthma Immunol Res* 2018;10:172-9.
[PUBMED](#) | [CROSSREF](#)
9. Tang H, Cao W, Kasturi SP, Ravindran R, Nakaya HI, Kundu K, et al. The T helper type 2 response to cysteine proteases requires dendritic cell-basophil cooperation via ROS-mediated signaling. *Nat Immunol* 2010;11:608-17.
[PUBMED](#) | [CROSSREF](#)
10. Nadeem A, Alharbi NO, Vliagoftis H, Tyagi M, Ahmad SF, Sayed-Ahmed MM. Proteinase activated receptor-2-mediated dual oxidase-2 up-regulation is involved in enhanced airway reactivity and inflammation in a mouse model of allergic asthma. *Immunology* 2015;145:391-403.
[PUBMED](#) | [CROSSREF](#)
11. Kauffman HF. Innate immune responses to environmental allergens. *Clin Rev Allergy Immunol* 2006;30:129-40.
[PUBMED](#) | [CROSSREF](#)
12. Roche N, Stirling RG, Lim S, Oliver BG, Oates T, Jazrawi E, et al. Effect of acute and chronic inflammatory stimuli on expression of protease-activated receptors 1 and 2 in alveolar macrophages. *J Allergy Clin Immunol* 2003;111:367-73.
[PUBMED](#) | [CROSSREF](#)
13. Shin YS, Sohn JH, Kim JY, Lee JH, Cho SH, Hong SJ, et al. Endotoxin is not essential for the development of cockroach induced allergic airway inflammation. *Yonsei Med J* 2012;53:593-602.
[PUBMED](#) | [CROSSREF](#)
14. Palmer CN, Irvine AD, Terron-Kwiatkowski A, Zhao Y, Liao H, Lee SP, et al. Common loss-of-function variants of the epidermal barrier protein filaggrin are a major predisposing factor for atopic dermatitis. *Nat Genet* 2006;38:441-6.
[PUBMED](#) | [CROSSREF](#)
15. Moniaga CS, Jeong SK, Egawa G, Nakajima S, Hara-Chikuma M, Jeon JE, et al. Protease activity enhances production of thymic stromal lymphopoietin and basophil accumulation in flaky tail mice. *Am J Pathol* 2013;182:841-51.
[PUBMED](#) | [CROSSREF](#)

16. Ying S, Meng Q, Corrigan CJ, Lee TH. Lack of filaggrin expression in the human bronchial mucosa. *J Allergy Clin Immunol* 2006;118:1386-8.
[PUBMED](#) | [CROSSREF](#)
17. Matsumura Y. Role of allergen source-derived proteases in sensitization via airway epithelial cells. *J Allergy (Cairo)* 2012;2012:903659.
[PUBMED](#) | [CROSSREF](#)
18. Jahnsen FL, Strickland DH, Thomas JA, Tobagus IT, Napoli S, Zosky GR, et al. Accelerated antigen sampling and transport by airway mucosal dendritic cells following inhalation of a bacterial stimulus. *J Immunol* 2006;177:5861-7.
[PUBMED](#) | [CROSSREF](#)
19. Hammad H, Lambrecht BN. Dendritic cells and epithelial cells: linking innate and adaptive immunity in asthma. *Nat Rev Immunol* 2008;8:193-204.
[PUBMED](#) | [CROSSREF](#)
20. Paul WE, Zhu J. How are T(H)2-type immune responses initiated and amplified? *Nat Rev Immunol* 2010;10:225-35.
[PUBMED](#) | [CROSSREF](#)
21. Zhu J, Yamane H, Paul WE. Differentiation of effector CD4 T cell populations (*). *Annu Rev Immunol* 2010;28:445-89.
[PUBMED](#) | [CROSSREF](#)