

Unraveling ancestry, kinship, and violence in a Late Neolithic mass grave

Hannes Schroeder^{a,1,2}, Ashot Margaryan^{b,c,1}, Marzena Szmyt^{d,e}, Bertrand Theulot^b, Piotr Włodarczyk^f, Simon Rasmussen^{g,h}, Shyam Gopalakrishnan^a, Anita Szczepanek^{f,i}, Tomasz Konopka^j, Theis Z. T. Jensen^{a,k}, Barbara Witkowska^{d,e,l}, Stanisław Wilk^j, Marcin M. Przybyła^m, Łukasz Pospieszny^{n,o}, Karl-Göran Sjögren^p, Zdzisław Belka^q, Jesper Olsen^r, Kristian Kristiansen^p, Eske Willerslev^{b,s,t}, Karin M. Frei^u, Martin Sikora^b, Niels N. Johannsen^{v,2}, and Morten E. Allentoft^{b,2}

^aSection for Evolutionary Genomics, Department of Biology, University of Copenhagen, 1353 Copenhagen K, Denmark; ^bLundbeck Foundation GeoGenetics Centre, Department of Biology, University of Copenhagen, 1350 Copenhagen K, Denmark; ^cInstitute of Molecular Biology, National Academy of Sciences of the Republic of Armenia, 0014 Yerevan, Armenia; ^dInstitute of Eastern Studies, Adam Mickiewicz University, 61-614 Poznań, Poland; ^eArchaeological Museum, 61-781 Poznań, Poland; ^fInstitute of Archaeology and Ethnology, Polish Academy of Sciences, Centre for Mountains and Uplands Archaeology, 31-016 Kraków, Poland; ^gNovo Nordisk Foundation Center for Protein Research, University of Copenhagen, 2200 Copenhagen, Denmark; ^hFaculty of Health and Medical Sciences, University of Copenhagen, 2200 Copenhagen, Denmark; ⁱDepartment of Anatomy, Jagiellonian University Medical College, 31-034 Kraków, Poland; ^jDepartment of Forensic Medicine, Jagiellonian University Collegium Medicum, 31-008 Kraków, Poland; ^kBioArCh, Department of Archaeology, University of York, York YO10 5NG, United Kingdom; ^lInstitute of Archaeology, Jagiellonian University, 31-007 Kraków, Poland; ^mDolmen S.C., 30-512 Kraków, Poland; ⁿInstitute of Archaeology and Ethnology, Centre for Studies into Late Antiquity and Early Medieval Times, Polish Academy of Sciences, 61-612 Poznań, Poland; ^oDepartment of Anthropology and Archaeology, University of Bristol, B58 1UU Bristol, United Kingdom; ^pDepartment of Historical Studies, University of Gothenburg, 405 30 Gothenburg, Sweden; ^qIsotope Laboratory, Adam Mickiewicz University, 61-680 Poznań, Poland; ^rAarhus AMS Centre, Department of Physics and Astronomy, Aarhus University, 8000 Aarhus C, Denmark; ^sDepartment of Zoology, University of Cambridge, CB2 3EJ Cambridge, United Kingdom; ^tWellcome Trust Sanger Institute, CB10 1SA Hinxton, United Kingdom; ^uUnit for Environmental Archaeology and Materials Science, National Museum of Denmark, 1220 Copenhagen K, Denmark; and ^vDepartment of Archaeology and Heritage Studies, Aarhus University, 8270 Højbjerg, Denmark

Edited by Clark Spencer Larsen, The Ohio State University, Columbus, OH, and approved March 25, 2019 (received for review November 27, 2018)

The third millennium BCE was a period of major cultural and demographic changes in Europe that signaled the beginning of the Bronze Age. People from the Pontic steppe expanded westward, leading to the formation of the Corded Ware complex and transforming the genetic landscape of Europe. At the time, the Globular Amphora culture (3300–2700 BCE) existed over large parts of Central and Eastern Europe, but little is known about their interaction with neighboring Corded Ware groups and steppe societies. Here we present a detailed study of a Late Neolithic mass grave from southern Poland belonging to the Globular Amphora culture and containing the remains of 15 men, women, and children, all killed by blows to the head. We sequenced their genomes to between 1.1- and 3.9-fold coverage and performed kinship analyses that demonstrate that the individuals belonged to a large extended family. The bodies had been carefully laid out according to kin relationships by someone who evidently knew the deceased. From a population genetic viewpoint, the people from Koszyce are clearly distinct from neighboring Corded Ware groups because of their lack of steppe-related ancestry. Although the reason for the massacre is unknown, it is possible that it was connected with the expansion of Corded Ware groups, which may have resulted in competition for resources and violent conflict. Together with the archaeological evidence, these analyses provide an unprecedented level of insight into the kinship structure and social behavior of a Late Neolithic community.

ancient DNA | archaeology | kinship | migration | violence

In 2011, archaeological excavations near the village of Koszyce in southern Poland uncovered a ca. 5,000-y-old mass grave (Fig. 1) associated with the Globular Amphora culture and containing the remains of 15 men, women, and children who had been killed, but carefully buried with rich grave goods (1). Closer study of the skeletons (2) revealed that the individuals had all been killed by blows to the head, possibly during a raid on their settlement. To shed light on this Late Neolithic community and the events that unfolded at Koszyce 5,000 y ago, we sequenced their genomes to between 1.1- and 3.9-fold coverage (Table 1) and performed genome-wide analyses to explore their genetic ancestry and kinship relations. In addition, we obtained 16 radiocarbon dates (*SI Appendix*, section 4 and *Dataset S1*) to narrow

down the date of the massacre to 2880–2776 BCE (*SI Appendix*, Fig. S5). We also provide a detailed description of the injuries (*SI Appendix*, section 3), and strontium isotope measurements of dental enamel provide information on mobility and residence patterns (*SI Appendix*, section 5). Together, the analyses enable us to draw up a remarkably detailed picture of this Late Neolithic community, including their genetic ancestry, physical appearance, kinship structure, and social organization.

Significance

We sequenced the genomes of 15 skeletons from a 5,000-y-old mass grave in Poland associated with the Globular Amphora culture. All individuals had been brutally killed by blows to the head, but buried with great care. Genome-wide analyses demonstrate that this was a large extended family and that the people who buried them knew them well: mothers are buried with their children, and siblings next to each other. From a population genetic viewpoint, the individuals are clearly distinct from neighboring Corded Ware groups because of their lack of steppe-related ancestry. Although the reason for the massacre is unknown, it is possible that it was connected with the expansion of Corded Ware groups, which may have resulted in violent conflict.

Author contributions: N.N.J. and M.E.A. designed and led the study; M. Szmyt, P.W., A.S., T.K., T.Z.T.J., B.W., S.W., M.M.P., Ł.P., and K.-G.S. provided samples for analysis and/or archaeological and osteological background information; A.M., Z.B., J.O., and K.M.F. generated the data; H.S., B.T., S.R., S.G., Z.B., K.M.F., M. Sikora, and M.E.A. analyzed the data; H.S., M. Szmyt, Z.B., J.O., K.K., E.W., K.M.F., M. Sikora, N.N.J., and M.E.A. interpreted the data; and H.S., N.N.J., and M.E.A. wrote the paper with input from all the other authors.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Data deposition: The aligned reads reported in this paper have been deposited in the European Nucleotide Archive and are available for download under accession no. [PRJEB28451](https://www.ebi.ac.uk/ena/record/PRJEB28451).

¹H.S. and A.M. contributed equally to this work.

²To whom correspondence may be addressed. Email: hshroeder@bio.ku.dk, nnj@cas.au.dk, or meallentoft@bio.ku.dk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1820210116/-DCSupplemental.

Published online May 6, 2019.




Fig. 1. The mass grave at Koszyce, southern Poland. (A) Photograph of the 15 skeletons and grave goods buried at Koszyce site 3 (reproduced with permission from ref. 2). (B) Map of Poland showing the location of Koszyce and four other Globular Amphora/Złota group sites included in this study.

Results and Discussion

DNA was isolated from teeth and petrous bones, using established protocols, and the libraries were sequenced on Illumina HiSeq 2500 platforms. The sequence data showed all the hallmarks of damaged ancient DNA, and modern contamination was estimated to be very low (Table 1). The human endogenous DNA contents ranged between 13% and 75% (Dataset S3). According to the sequencing data, eight of the individuals in the grave were males and seven were females (Table 1). This is consistent with previously published results based on skeletal traits, with the exception of one male (individual 4) who had previously been identified as a probable female and five juveniles who could not be sexed previously (2). Data on phenotypic traits based on imputed genotypes (Dataset S5) revealed that the individuals had mostly brown eyes, dark or dark-blond hair, and intermediate to dark skin.

Genetic Affinities. To investigate their genetic ancestry, we merged the 15 Koszyce genomes with the Human Origins data set (3), as well as 168 previously published ancient genomes (Dataset S6). In addition, we included genome-wide data for nine individuals from

four contemporary, neighboring sites in southern Poland belonging to the Globular Amphora culture and its Złota group variant that we sequenced to between 0.2- and 1-fold coverage (Dataset S3). We then performed a principal component analysis and found that all 24 Globular Amphora/Złota group individuals clustered with other previously sequenced Globular Amphora individuals (4, 5) and other Neolithic groups (Fig. 24). This confirms earlier suggestions that the Globular Amphora people belonged to the Neolithic gene pool of Europe, as typified by early Anatolian farmers (4, 5).

To further investigate the ancestry of the Globular Amphora individuals, we performed a supervised ADMIXTURE (6) analysis, specifying typical western European hunter-gatherers (Loschbour), early Neolithic Anatolian farmers (Barçın), and early Bronze Age steppe populations (Yamnaya) as ancestral source populations (Fig. 2B). The results indicate that the Globular Amphora/Złota group individuals harbor ca. 30% western hunter-gatherer and 70% Neolithic farmer ancestry, but lack steppe ancestry. To formally test different admixture models and estimate mixture proportions, we then used *qpAdm* (7) and find that the Polish Globular

Table 1. Sequencing results for the Koszyce individuals

Individual no.	Age at death, y	Average depth of coverage	Genome coverage (%)	Chromosomal sex	5' C-T (%)	Contamination rate (%)	mtDNA haplogroup	ChrY haplogroup
1	25–30	2.8x	82	XX	19	0.3	T2b	—
2	1.5–2	1.9x	75	XY	24	0.7	T2b	I2a-L801
3	30–35	1.3x	67	XX	24	0.1	H27+16093	—
4	16–17	3.0x	83	XY	21	0.1	K1a1b1e	I2a-L801
5	20–25	3.9x	85	XY	19	0.0	HV0a	I2a-L801
6	13–14	1.1x	60	XX	20	0.4	K1a1b1e	—
7	2–2.5	2.1x	77	XY	21	0.0	HV16	I2a-L801
8	30–35	3.0x	83	XX	22	0.2	J1c3f	—
9	15–16	2.5x	81	XX	21	0.2	J1c3f	—
10	18–20	2.4x	80	XY	20	0.1	HV0a	I2a-L801
11	40–50	1.1x	60	XY	26	0.0	HV0a	I2a-L801
12	30–40	3.7x	85	XX	22	0.4	K1a1b1e	—
13	5–6	2.6x	81	XY	19	0.0	J1c3f	I2a-L801
14	50–60	3.0x	83	XX	24	0.3	HV0a	—
15	40–50	2.9x	82	XY	21	0.0	HV0a	I2a-L801

Age at death was estimated based on standard osteological methods (47, 48). Genetic sex estimates were obtained by assessing the reads mapping to the Y versus X chromosome (49). Deamination rates (5' C-T) were estimated using MapDamage (34). Contamination rates are based on Schmutzi (35). For more information, see *Materials and Methods*.




Fig. 2. Genetic affinities of the Koszyce individuals and other GAC groups (here including Złota) analyzed in this study. (A) Principal component analysis of previously published and newly sequenced ancient individuals. Ancient genomes were projected onto modern reference populations, shown in gray. (B) Ancestry proportions based on supervised ADMIXTURE analysis ($K = 3$), specifying Western hunter-gatherers, Anatolian Neolithic farmers, and early Bronze Age steppe populations as ancestral source populations. LP, Late Paleolithic; M, Mesolithic; EN, Early Neolithic; MN, Middle Neolithic; LN, Late Neolithic; EBA, Early Bronze Age; PWC, Pitted Ware culture; TRB, Trichterbecherkultur/Funnelbeaker culture; LBK, Linearbandkeramik/Linear Pottery culture; GAC, Globular Amphora culture; Złota, Złota culture.

Amphora/Złota group individuals can be modeled as a mix of western European hunter-gatherer (17%) and Anatolian Neolithic farmer (83%) ancestry (*SI Appendix, Table S2*), mirroring the results of previous studies (4, 5).

Kinship and Consanguinity. Analyses of ancient genomes can provide detailed information on the kinship structures and social organization of past communities (8–10). At Koszyce, mitochondrial DNA (mtDNA) analysis revealed the presence of six different maternal lineages, whereas analysis of the non-recombining region of the Y chromosome showed that all males carried the same Y chromosome haplotype: I2a-L801 (Table 1). We then estimated genomic runs of homozygosity (ROH) and found that the Koszyce individuals were not particularly inbred. Although a slightly larger section of the Koszyce genomes is contained within ROH compared with typical modern European populations (*SI Appendix, Fig. S7*), this signal is mainly driven by an increased fraction of short ROH (<2 Mb), which is indicative of ancestral restrictions in population size rather than recent inbreeding. On the basis of genome-wide patterns of allelic identity-by-state (IBS), we computed kinship coefficients between all pairs of individuals and applied established cutoff values for possible kinship categories (*Materials and Methods*). We find that the Koszyce burial represents a large extended family connected via several first- and second-degree relationships (Fig. 3 and *SI Appendix, Fig. S9*).

Overall, we identified four nuclear families in the grave, which are for the most part represented by mothers and their children (Fig. 3). Closely related kin were buried next to each other: a mother was buried cradling her child, and siblings were placed side by side. Evidently, these individuals were buried by people who knew them well and who carefully placed them in the grave according to familial relationships. For example, individual 14, the oldest individual in the grave, was buried close to her two sons (individuals 5 and 15), whereas individual 8, a 30–35-y-old woman, was buried with her teenage daughter (individual 9) and

5-y-old son (individual 13). Using genome-wide patterns of IBS, we were also able to reconstruct more complex relationships: individuals 5, 10, 11, and 15 all appear to be brothers, and yet they do not have the same mother (individual 14 is the mother of individuals 5 and 15, but not 10 and 11), suggesting that they might be half-brothers. However, all four of them share the same mitochondrial DNA haplotype, suggesting that their mothers might also have been related.

Interestingly, the older males/fathers are mostly missing from the grave, suggesting that it might have been them who buried their kin. The only father present in the grave is individual 10, whose partner and son are placed together opposite him in the grave. In addition, there is a young boy (individual 7), aged 2–2.5 y, whose parents are not in the grave, but he is placed next to other individuals to whom he is closely related through various second-degree relationships. Finally, there is individual 3, an adult female, who does not seem to be genetically related to anyone in the group. However, her position in the grave close to individual 4, a young man, suggests that she may have been as close to him in life as she was in death. These biological data and burial arrangements show that the social relationships held to be most significant in these societies were identical with genetic and reproductive relationships. However, they also demonstrate that nuclear families were nested in larger, extended family groups, either permanently or for parts of the year.

Social Organization, Residence Patterns, and Subsistence Strategies.

The presence of unrelated females and related males in the grave is interesting because it suggests that the community at Koszyce was organized along patrilineal lines of descent, adding to the mounting evidence that this was the dominant form of social organization among Late Neolithic communities in Central Europe (11, 12). Usually, patrilineal forms of social organization go hand in hand with female exogamy (i.e., the practice of women marrying outside their social group). Indeed, several studies (11, 12) have shown that patrilocal residence patterns and female

aimed specifically at males from competing groups (21). The former strategy presumably aims at eradicating a competing group from a given territory, whereas in the latter case, it may be assumed that the women and/or children are taken as captives, a practice that is well documented ethnographically and historically for prestate societies (23). Although alternative scenarios (e.g., ritualistic violence or familicide) cannot be ruled out, it seems most plausible that the massacre at Koszyce falls in the former category. The fact that most adult males of the group are missing from the grave most likely reflects that they were away (or fled) when the raid occurred, leaving the remaining group vulnerable.

Although it is impossible to identify the culprits of the massacre that took place at Koszyce around 2880–2776 BCE, it is interesting to note that it occurred right around the time when the Corded Ware complex started to spread rapidly across large parts of Central Europe, and it seems plausible that the group from Koszyce fell victim to some violent intergroup conflict related to the territorial expansion of Corded Ware groups or another competing group in the area. If the general interaction between Globular Amphora people and neighboring, steppe-related cultures (including early Corded Ware) was primarily hostile, it would explain why Globular Amphora individuals carry no steppe ancestry and, in part, why Europe experienced such a dramatic reduction in Neolithic genomic ancestry at this time (7, 17).

Concluding Remarks

Brutal events such as the family massacre documented in the Koszyce burial may have been all too common in the unstable, tumultuous centuries at the beginning of the third millennium BCE. However, along with all the violence and aggression illustrated by the Koszyce find, our study also demonstrates the strong sense of family affiliation and cohesion that prevailed among this group of people. From the careful positioning of the bodies in the grave, it is clear that both nuclear and extended family relations were key to how people organized their lives, and that these relations represented major, normative values in Globular Amphora communities of this period. Although it has often been suggested that nuclear and/or extended family structures were important in many prehistoric societies, the archaeological and genomic data we have presented here provide actual proof that this was indeed the case. Furthermore, as we start to better understand the mating patterns and social structures of past populations, we can start to move beyond our often all too simplistic assumptions used to model population history and culture change (24). Ultimately, it will be through the integration of archaeological, anthropological, and biological data and theory that we will be able to produce more accurate representations of past social realities and population histories.

Materials and Methods

Samples. The samples analyzed in this study (21 petrous bones and three teeth) stem from five different archaeological sites in southern Poland, all associated with the Globular Amphora culture or its Złota group variant (*SI Appendix, section 1*). For detailed site descriptions, see *SI Appendix, section 2*.

DNA Extraction, Library Preparation, and Sequencing. DNA was extracted from petrous bones and tooth roots, using established protocols specifically designed for ancient DNA (*SI Appendix, section 6*). Double-stranded DNA libraries were built, using Illumina-specific adapters, and amplified and indexed using a dual-indexing protocol (25). The optimal number of PCR cycles was determined using qPCR. The indexed and amplified libraries were purified and quantified on an Agilent 2200 TapeStation before being pooled in equimolar amounts. The pooled libraries were sequenced (80 bp, single read) on Illumina HiSeq 2500 platforms at the Danish National High-throughput DNA Sequencing Centre.

Raw Read Processing and Alignment. Base-calling was performed using CASAVA 1.8.2. Adapter sequences and leading/trailing stretches of Ns were trimmed using AdapterRemoval 2.1.3 (26). Reads of at least 30 bp were mapped to the human reference genome GRCh37 using bwa 0.7.10 (27) with the seed disabled. Duplicates and reads with MAPQ <30 were removed using

samtools 1.3.1 (28). All reads were then merged to sample level and realigned using GATK-3.3.0 (29). The realigned reads had the md-tag updated and extended base alignment qualities calculated using samtools calmd. Read depth and coverage were determined using BEDtools 2.27.0 genomcov (30). For full details, see *SI Appendix, section 6*.

Genotype Likelihoods and Imputation. Genotype likelihoods were determined using GATK 3.7.0 (31). We called biallelic sites present in the 1000 Genomes (–genotyping_mode GENOTYPE_GIVEN_ALLELES) filtering for transitions by setting the genotype likelihoods to 0 for all three genotypes (e.g., hom ref, het, and hom alt). Subsequently, the individual VCF (variant call format) files were merged using BCftools 1.3.1. The combined VCF was then split into separate files containing 200,000 markers each and imputed separately using Beagle 4.0 (r1399) (32), using the 1000 Genomes (33) phase 3 map included with Beagle (*.phase3.v5a.snps.vcf.gz and plink.chr*.GRCh37.map) with input through the genotype likelihood option.

MapDamage Analysis. We used mapDamage 2.0 (34) to assess the extent of DNA damage in our samples and found that all of them showed the features characteristic for ancient DNA, including short average fragment lengths, an increased occurrence of purines before strand breaks, and an increased frequency of apparent cytosine (C) to thymine (T) substitutions at 5'-ends (*Dataset S3*), which likely results from the deamination of cytosine residues that occur primarily in the single-stranded overhangs of DNA fragments.

Mitochondrial DNA. We used Schmutzi (35) to determine the endogenous consensus mtDNA sequences. Reads were mapped to the Cambridge reference sequence and filtered for MAPQ \geq 30. Haploid variants were called using the endoCaller program implemented in Schmutzi (35), and only variants with a posterior probability exceeding 50 on the PHRED scale (probability of error: 1/100,000) were retained. We then used Haplogrep 2.2 (36) to determine the mtDNA haplogroups, specifying PhyloTree (build 17) as the reference phylogeny (37).

Y Chromosome DNA. To determine the Y chromosome haplogroups, we first called haploid genotypes for reads mapping (MAPQ \geq 30) to the Y chromosome, using samtools/bcftools (27, 38). We then extracted genotypes for all individuals at SNPs included in the Y-DNA haplogroup tree version 13.57 from the International Society of Genetic Genealogy (<https://isogg.org>). The full list of observed alleles defining subhaplogroups of I2a is listed in *Dataset S4* and plotted in *SI Appendix, Fig. S8*.

Principal Component Analysis. Principal component analysis was performed using smartPCA (39), using around 1,000 modern Eurasian individuals from the HO dataset (3) and 168 previously published ancient genomes (*Dataset S6*). The ancient genomes were projected onto the modern variation, using the option lsq project. Individuals with too much missing data (mind 0.1) were excluded from the analysis.

Model-Based Clustering. We ran ADMIXTURE (6) in supervised mode on the same merged dataset, specifying Western hunter-gatherers (Loschbour), Neolithic farmers (Barcın), and Yamnaya as ancestral source populations. The resulting admixture proportions are plotted in Fig. 2B. The plot shows population averages, and to help visualize the results, only a selected number of ancient populations are shown.

qpAdm Analysis. Admixture proportions were modeled using qpAdm (7), specifying Mesolithic Western European hunter-gatherers, Anatolian Neolithic farmers (Barcın), and either Yamnaya or Eastern hunter-gatherers as possible source populations. In most cases, a two-way model without steppe ancestry provides the best fit (*SI Appendix, Table S2*). The only exception is Książnice, for which a small amount of additional gene-flow from some steppe-related population provides a better fit than a simple two-way model. To determine the source of this additional gene-flow, we modeled the Książnice genomes as a mix of Western European hunter-gatherers, Neolithic farmers, and either Yamnaya or Eastern hunter-gatherer ancestry, and found that both provide an equally good fit, suggesting that this additional gene-flow is not necessarily Yamnaya related.

Phenotype. To reconstruct the physical appearance of the Koszyce individuals, we first imputed missing genotypes for our low-coverage genomes and then selected 39 SNPs (40) to determine phenotypic traits, including skin, hair, and eye color. The allele counts were uploaded to the HliisPlex-5 website to obtain the phenotype predictions (*Dataset S5*).

Runs of Homozygosity. We identified runs of homozygosity using IBDseq (41) on a merge of the imputed genotypes of the Koszyce individuals and 214 individuals from two populations (IBS, TSI) of the 1000 Genomes Project (33). Genotypes for the Koszyce individuals with imputed genotype probability <0.99 were set to missing for this analysis. We ran IBDseq on the merged and filtered dataset with default parameters, except allowing for a higher genotype error rate (errormax = 0.005). Inferred tracts were filtered for LOD score ≥ 2 , and then further processed by splitting tracts that spanned centromeres or those that spanned large assembly gaps with less than 1 Mb on both sides of the gaps. For each individual, we inferred the total length contained in short (<2 Mb) and long (>8 Mb) ROH by summing up individual tract lengths within those length categories, as done in ref. 42 (SI Appendix, Fig. S7).

Kinship Analysis. Kinship analysis was performed using a recently described method based on pairwise sharing of alleles IBS (10, 43). Because of the low sequencing coverage, we first computed a matrix of expected pairwise IBS sharing for all pairs of individuals as the 2D site frequency spectrum, using the program realSFS from the ANGSD package (44). For each pair, the 2D-site frequency spectrum with the best likelihood across 10 replicate runs was retained and was used to calculate a number of relatedness estimators: the kinship estimator introduced by Manichaikul et al. (45), implemented in the KING package [equation (9) in ref. 45], and the “R0” and “R1” ratios described in Waples et al. (43)

We used the kinship coefficient ranges described in Manichaikul et al. (45) to classify pairs of individuals into first-, second-, and third-degree relatives. First-degree relatives were further distinguished using the R0 ratio, into parent–offspring ($R0 \leq 10$) or sibling relationships ($R0 > 10$ –2; SI Appendix, Fig. S9 and Dataset S7). The resulting relatedness networks of first- and second-degree relatives were visualized using Cytoscape version 3.4.0 (46), including annotations of individual sex and mitochondrial haplogroups.

ACKNOWLEDGMENTS. We thank Jesper Stenderup and the staff at the Danish National High-throughput Sequencing Centre for technical assistance. We also thank Agata Ulanowska, Eva Andersson Strand, Małgorzata Siennicka, and Ulla Mannering for helpful discussion. The project was funded by a Villum Foundation Young Investigator grant (Grant 10120 to M.E.A.). The Centre for GeoGenetics is funded by the National Research Foundation of Denmark and the Lundbeck Foundation. H.S. was supported by HERA (Humanities in the European Research Area) and the European Union’s Horizon 2020 Research and Innovation Programme (Grant 649307). S.R. was supported by the Novo Nordisk Foundation (Grant NNF14CC0001). T.Z.T.J. is supported by the European Union’s EU Framework Programme for Research and Innovation Horizon 2020 under Grant Agreement 676154 (ArchSci2020). Work on the site’s radiocarbon chronology was funded by a grant from the Aarhus University Research Foundation (N.N.J.). The strontium isotope measurements were funded by the Carlsberg Foundation “Tales of Bronze Age Women” project (Grant CF15-0878 to K.M.F.). Publication fees were covered by the Materials, Culture and Heritage research program at Aarhus University.

- Przybyła MM, Szczepanek A, Włodarczak P, eds (2013) *Koszyce stanowisko 3. Przemoc i rytuał u schyłku neolitu* (Profil-Archeo, Kraków-Pekowice, Poland).
- Konopka T, Szczepanek A, Przybyła MM, Włodarczak P (2016) Evidence of interpersonal violence or a special funeral rite in the Neolithic multiple burial from Koszyce in southern Poland—a forensic analysis. *Anthropol Rev* 79:69–85.
- Lazaridis I, et al. (2014) Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature* 513:409–413.
- Tassi F, et al. (2017) Genome diversity in the Neolithic Globular Amphora culture and the spread of Indo-European languages. *Proc Biol Sci* 284:20171540.
- Mathieson I, et al. (2018) The genomic history of southeastern Europe. *Nature* 555:197–203.
- Alexander DH, Novembre J, Lange K (2009) Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* 19:1655–1664.
- Haak W, et al. (2015) Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* 522:207–211.
- Kennett DJ, et al. (2017) Archaeogenomic evidence reveals prehistoric matrilineal dynasty. *Nat Commun* 8:14115.
- O’Sullivan N, et al. (2018) Ancient genome-wide analyses infer kinship structure in an Early Medieval Alemannic graveyard. *Sci Adv* 4:eaa01262.
- Sikora M, et al. (2017) Ancient genomes show social and reproductive behavior of early Upper Paleolithic foragers. *Science* 358:659–662.
- Haak W, et al. (2008) Ancient DNA, strontium isotopes, and osteological analyses shed light on social and kinship organization of the later stone age. *Proc Natl Acad Sci USA* 105:18226–18231.
- Knipper C, et al. (2017) Female exogamy and gene pool diversification at the transition from the final Neolithic to the early Bronze Age in central Europe. *Proc Natl Acad Sci USA* 114:10083–10088.
- Szmyt M (1999) *Between West and East: People of the Globular Amphora Culture in Eastern Europe; 2950-2350 BC* (Institute of Prehistory, Adam Mickiewicz University, Poznan, Poland).
- Woidich M (2014) The Western Globular Amphora culture. A new model for its emergence and expansion. *eTopoi* 3:67–85.
- Johannsen N, Laursen S (2010) Routes and wheeled transport in late 4th–early 3rd millennium funerary customs of the Jutland Peninsula: Regional evidence and European context. *Præhist Z* 85:15–58.
- Czebreszuk J, Szmyt M (2011) Identities, differentiation and interactions on the Central European Plain in the 3rd millennium BC. *Sozialarchäologische Perspektiven: Gesellschaftlicher Wandel 5000-1500 v. Chr. Zwischen Atlantik Und Kaukasus*, Archäologie in Eurasien, eds Hansen S, Müller J (Philipp von Zabern, Mainz, Germany), pp 247–269.
- Allentoft ME, et al. (2015) Population genomics of Bronze Age Eurasia. *Nature* 522:167–172.
- Kelly RC (1985) *The Nuer Conquest: The Structure and Development of an Expansionist System* (University of Michigan Press, Ann Arbor, MI).
- Schulting RJ, Fibiger L (2012) *Sticks, Stones, and Broken Bones: Neolithic Violence in a European Perspective* (Oxford Univ Press, Oxford).
- Chenal F, Perrin B, Barrand-Emam H, Boulestin B (2015) A farewell to arms: A deposit of human limbs and bodies at Bergheim, France, c. 4000 BC. *Antiquity* 89:1313–1330.
- Meyer C, et al. (2018) Early Neolithic executions indicated by clustered cranial trauma in the mass grave of Halberstadt. *Nat Commun* 9:2472.
- Milner GR, Anderson E, Smith VG (1991) Warfare in late prehistoric west-central Illinois. *Am Antiq* 56:581–603.
- Cameron CM (2016) *Captives: How Stolen People Changed the World* (University of Nebraska Press, Lincoln, NE).
- Johannsen NN, Larson G, Meltzer DJ, Vander Linden M (2017) A composite window into human history. *Science* 356:1118–1120.
- Kircher M, Sawyer S, Meyer M (2012) Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res* 40:e3.
- Shubert M, Lindgreen S, Orlando L (2016) AdapterRemoval v2: Rapid adapter trimming, identification, and read merging. *BMC Res Notes* 9:88.
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.
- Li H, et al.; 1000 Genome Project Data Processing Subgroup (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079.
- DePristo MA, et al. (2011) A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 43:491–498.
- Quinlan AR, Hall IM (2010) BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* 26:841–842.
- McKenna A, et al. (2010) The genome analysis toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20:1297–1303.
- Browning BL, Browning SR (2013) Improving the accuracy and efficiency of identity-by-descent detection in population data. *Genetics* 194:459–471.
- Auton A, et al.; 1000 Genomes Project Consortium (2015) A global reference for human genetic variation. *Nature* 526:68–74.
- Jönsson H, Ginolhac A, Schubert M, Johnson PLF, Orlando L (2013) mapDamage2.0: Fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* 29:1682–1684.
- Renaud G, Slon V, Duggan AT, Kelso J (2015) Schmutzi: Estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol* 16:224.
- Weissensteiner H, et al. (2016) HaploGrep 2: Mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Res* 44:W58–63.
- van Oven M (2015) PhyloTree Build 17: Growing the human mitochondrial DNA tree. *Forensic Sci Int Genet Suppl Ser* 5:e392–e394.
- Li H (2011) A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* 27:2987–2993.
- Patterson N, Price AL, Reich D (2006) Population structure and eigenanalysis. *PLoS Genet* 2:e190.
- Chaitanya L, et al. (2018) The HIRisPlex-S system for eye, hair and skin colour prediction from DNA: Introduction and forensic developmental validation. *Forensic Sci Int Genet* 35:123–135.
- Browning BL, Browning SR (2013) Detecting identity by descent and estimating genotype error rates in sequence data. *Am J Hum Genet* 93:840–851.
- Schroeder H, et al. (2018) Origins and genetic legacies of the Caribbean Taino. *Proc Natl Acad Sci USA* 115:2341–2346.
- Waples RK, Albrechtsen A, Moltke I (2019) Allele frequency-free inference of close familial relationships from genotypes or low-depth sequencing data. *Mol Ecol* 28:35–48.
- Kornelissen TS, Albrechtsen A, Nielsen R (2014) ANGSD: Analysis of next generation sequencing data. *BMC Bioinformatics* 15:356.
- Manichaikul A, et al. (2010) Robust relationship inference in genome-wide association studies. *Bioinformatics* 26:2867–2873.
- Shannon P, et al. (2003) Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13:2498–2504.
- Cunningham C, Scheuer L, Black S (2016) *Developmental Juvenile Osteology* (Elsevier Academic Press, Burlington, MA).
- Ubelaker DH (1989) *Human Skeletal Remains: Excavation, Analysis, Interpretation* (Taraxacum, Washington, DC).
- Skoglund P, Storå J, Götherström A, Jakobsson M (2013) Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J Arch Sci* 40:4477–4482.