

CORRECTION

Open Access



Correction to: Prognostic significance of preoperative plasma D-dimer level in patients with surgically resected clinical stage I non-small cell lung cancer: a retrospective cohort study

Kaoru Kaseda^{1*}, Keisuke Asakura¹, Akio Kazama² and Yukihiro Ozawa³

Correction to: J Cardiothorac Surg

<https://doi.org/10.1186/s13019-017-0676-3>

The original version of this article [1] did not cite the following sources [2–5], which were used to write the article.

The authors apologise for any inconvenience that this might have caused.

harbouring epidermal growth factor receptor and K-ras mutation. *Thorac Cancer*. 2017;8:229–37.

Author details

¹Department of Thoracic Surgery, Sagamihara Kyodo Hospital, 2-8-18 Hashimoto, Midori-ku, Sagamihara, Kanagawa 252-5188, Japan. ²Department of Pathology, Sagamihara Kyodo Hospital, 2-8-18 Hashimoto, Midori-ku, Sagamihara, Kanagawa 252-5188, Japan. ³Yuai Clinic, 1-6-2 Kitashinyokohama, Kohoku-Ku, Yokohama, Kanagawa 223-0059, Japan.

Received: 13 June 2019 Accepted: 13 June 2019

Published online: 20 June 2019

References

1. Kaseda K, Asakura K, Kazama A, Ozawa Y. Prognostic significance of preoperative plasma D-dimer level in patients with surgically resected clinical stage I non-small cell lung cancer: a retrospective cohort study. *J Cardiothorac Surg*. 2017;12(1):102.
2. Fukumoto K, Taniguchi T, Usami N, et al. The preoperative plasma D-dimer level is an independent prognostic factor in patients with completely resected non-small cell lung cancer. *Surg Today*. 2015;45:63.
3. Jiang HG, Li J, Shi SB, et al. Value of fibrinogen and D-dimer in predicting recurrence and metastasis after radical surgery for non-small cell lung cancer. *Med Oncol*. 2014;31:22.
4. Kaseda K, Watanabe K, Asakura K, Kazama A, Ozawa Y. Identification of false-negative and false-positive diagnoses of lymph node metastases in non-small cell lung cancer patients staged by integrated 18F-fluorodeoxyglucose positron emission tomography/computed tomography: A retrospective cohort study. *Thorac Cancer*. 2016;7:473–80.
5. Kaseda K, Asakura K, Kazama A, Ozawa Y. Clinicopathological and prognostic features of surgically resected pathological stage I lung adenocarcinoma

* Correspondence: kaseda@wb4.so-net.ne.jp

¹Department of Thoracic Surgery, Sagamihara Kyodo Hospital, 2-8-18 Hashimoto, Midori-ku, Sagamihara, Kanagawa 252-5188, Japan
Full list of author information is available at the end of the article

