



# Many rice genes are differentially spliced between roots and shoots but cytokinin has minimal effect on splicing

Nowlan H. Freese<sup>1\*</sup> | April R. Estrada<sup>1\*</sup> | Ivory C. Blakley<sup>1</sup> | Jinjie Duan<sup>2</sup> |  
Ann E. Loraine<sup>1</sup>

<sup>1</sup>Department of Bioinformatics and Genomics, University of North Carolina at Charlotte, Charlotte, North Carolina

<sup>2</sup>Department of Biomedicine, Aarhus University, Aarhus C, Denmark

## Correspondence

Ann E. Loraine, Department of Bioinformatics and Genomics, University of North Carolina at Charlotte, Charlotte, NC.  
Email: aloraine@uncc.edu

## Funding information

National Science Foundation, Grant/Award Number: 1523814; NIH, Grant/Award Number: 2R01GM103463-06A1

## Abstract

Alternatively spliced genes produce multiple spliced isoforms, called transcript variants. In differential alternative splicing, transcript variant abundance differs across sample types. Differential alternative splicing is common in animal systems and influences cellular development in many processes, but its extent and significance is not as well known in plants. To investigate differential alternative splicing in plants, we examined RNA-Seq data from rice seedlings. The data included three biological replicates per sample type, approximately 30 million sequence alignments per replicate, and four sample types: roots and shoots treated with exogenous cytokinin delivered hydroponically or a mock treatment. Cytokinin treatment triggered expression changes in thousands of genes but had negligible effect on splicing patterns. However, many genes were differentially spliced between mock-treated roots and shoots, indicating that our methods were sufficiently sensitive to detect differential splicing between data sets. Quantitative fragment analysis of reverse transcriptase-PCR products made from newly prepared rice samples confirmed 9 of 10 differential splicing events between rice roots and shoots. Differential alternative splicing typically changed the relative abundance of splice variants that co-occurred in a data set. Analysis of a similar (but less deeply sequenced) RNA-Seq data set from *Arabidopsis* showed the same pattern. In both the *Arabidopsis* and rice RNA-Seq data sets, most genes annotated as alternatively spliced had small minor variant frequencies. Of splicing choices with abundant support for minor forms, most alternative splicing events were located within the protein-coding sequence and maintained the annotated reading frame. A tool for visualizing protein annotations in the context of genomic sequence (ProtAnnot) together with a genome browser (Integrated Genome Browser) were used to visualize and assess effects of differential splicing on gene function. In general, differentially spliced regions coincided with conserved protein domains, indicating that differential alternative splicing is likely to affect protein function between root and shoot tissue in rice.

\*These authors are contributed equally.

This manuscript was previously deposited as a preprint at <https://www.biorxiv.org/content/10.1101/293217v5>

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Plant Direct* published by American Society of Plant Biologists, Society for Experimental Biology and John Wiley & Sons Ltd.

**KEYWORDS**

alternative splicing, cytokinin, rice, RNA-Seq, root, shoot

**1 | INTRODUCTION**

Alternative splicing of pre-mRNA transcripts enables one gene to produce multiple transcript variants encoding different functions. Alternative splicing is an almost universal phenomenon in higher eukaryotes, occurring to varying degrees in every animal and plant genome examined to date (Kalsotra & Cooper, 2011; Reddy, Marquez, Kalyna, & Barta, 2013). In animals, differential expression of splice variants has been recruited as a regulatory mechanism in multiple processes, such as sex determination in invertebrates and neuronal differentiation in mammals (Barbosa-Morais et al., 2012; Kalsotra & Cooper, 2011; Salz, 2011).

In plants, less is known about the functional significance and patterns of alternative splicing. However, several trends are apparent. Genes involved in circadian regulation are highly alternatively spliced, often producing multiple splice variants that fluctuate in concert with day/night cycling along with overall transcript abundance (Filichkin, Cumbie, et al., 2015). The serine and arginine-rich (SR) family of RNA-binding, splicing regulatory proteins is greatly expanded compared to mammals and includes many plant-specific forms (Barbosa-Morais, Carmo-Fonseca, & Aparício, 2006; Filichkin, Priest, Megraw, & Mockler, 2015; Kalyna & Barta, 2004; Plass, Agirre, Reyes, Camara, & Eyra, 2008). SR transcripts themselves are also highly alternatively spliced in plants, with the relative abundance of these alternative transcripts varying according to environmental stresses and hormones (Filichkin, Priest, et al., 2015; Gullledge, Roberts, Vora, Patel, & Loraine, 2012; Keller et al., 2016; Mei, Boatwright, Feng, Schnable, & Barbazuk, 2017; Palusa, Ali, & Reddy, 2007).

A growing body of evidence indicates that cell and tissue specific regulation of alternative splicing occurs in plants, but its significance and extent is not well established (Li, Yamada, Han, Ohler, & Benfey, 2016; Sun et al., 2018; Vitulo et al., 2014). We previously found through analysis of RNA-Seq data from *Arabidopsis* pollen that the relative abundance of splice variants was similar between leaves and pollen, despite the differences between the two tissues (Loraine, McCormick, Estrada, Patel, & Qin, 2013). However, this latter analysis was limited by having just one biological replicate for pollen and only two biological replicates for leaves. A more comprehensive analysis of multiple *Arabidopsis* data sets found a high incidence of isoform switching, in which the identity of the most prevalent variant differs between sample types (Vanechoutte, Estrada, Lin, Loraine, & Vandepoele, 2017). This splicing diversity may have arisen in part from the heterogeneity of the data sets used, which were produced using rapidly changing (and improving) sequencing technologies at different times by different groups.

In this study, we used a well-replicated RNA-Seq data set from rice to re-examine prevalence of alternative splicing between tissues and hormone (cytokinin) treatment. This data set was previously generated

to investigate cytokinin regulation of gene expression in roots and shoots from 10-day-old rice seedlings (Raines et al., 2016). A parallel study produced an analogous data set from *Arabidopsis* for comparison, but was less deeply sequenced (Zubo et al., 2017). Both the rice and *Arabidopsis* RNA-Seq data sets included three biological replicates per sample type and four sample types—roots and shoots treated with exogenous cytokinin or a mock, vehicle-only treatment. In both data sets, the treatment triggered differential expression of thousands of genes, with roots affected to a greater degree than shoots.

For most alternatively spliced genes, regardless of whether or not they were differentially spliced, the relative abundance of splicing forms was highly skewed, with most alternatively spliced genes producing one major form. Nonetheless, there were many alternatively spliced genes where minor forms were abundant and therefore seemed likely to affect gene function. We found that the relative abundance of transcript variants for most alternatively spliced genes was remarkably stable, with very few differentially spliced genes between cytokinin treated and control samples. By contrast, many more genes were differentially spliced between roots and shoot, and most differential splicing occurred within the protein-coding sequence. Moreover, nearly every differential splicing event detected changed the relative abundance of splice variants that co-occurred in the same sample. These results suggest differential alternative splicing likely contributes to gene function diversification between roots and shoots by moderating the relative abundance of co-expressed splice variants, but plays little role in cytokinin signaling.

**2 | MATERIALS AND METHODS****2.1 | RNA-Seq data**

RNA-Seq data from rice and *Arabidopsis* shoot and root tissue were from our previous studies on the effect of cytokinin on gene expression (Raines et al., 2016; Zubo et al., 2017). Rice sequence data are available from the Sequence Read Archive under accession SRP049054. Aligned, processed data are available from October 2011 rice genome assembly IgbQuickload directories at <http://lorainelab-quickload.scidas.org/rnaseq/>. *Arabidopsis* sequence data are available from the Sequence Read Archive under accession SRP059384. Aligned, processed data are available from the TAIR10 (June 2009) *Arabidopsis* genome assembly IgbQuickload directories at <http://lorainelab-quickload.scidas.org/rnaseq/>.

**2.2 | Data processing**

Rice sequences were aligned onto the *Oryza sativa japonica* genome assembly Os-Nipponbare-Reference-IRGSP-1.0 (Kawahara et al., 2013) and *Arabidopsis* sequences were aligned onto the TAIR10 June 2009

release of the *Arabidopsis* genome (Lamesch et al., 2012) using TopHat (Kim et al., 2013) and BowTie2 (Langmead & Salzberg, 2012) with maximum intron size set to 5,000 bases. A command-line, Java program called “FindJunctions” was used to identify exon-exon junctions from gapped read alignments in the RNA-Seq data. FindJunctions produces BED format files containing junction features, and the score field of the BED file lists the number of read alignments that supported the junction. Only reads that aligned to a unique location in the genome were considered. Source code and compiled versions of FindJunctions are available from <https://bitbucket.org/lorainelab/findjunctions>.


### 2.3 | Identification of alternative splicing events and differential splicing

To date, there have been two major releases of *O. sativa japonica* gene models: the MSU7 gene set (Kawahara et al., 2013) and the RAP-Db gene set (Sakai et al., 2013). The two gene model sets contain mostly the same data, but the MSU7 gene models appear to be the most heavily used and annotated with Gene Ontology terms. For simplicity, and to take advantage of available functional annotations, we used the MSU7 annotations here. For analysis of *Arabidopsis* data, we used TAIR10 (Lamesch et al., 2012) and Araport11 (Cheng et al., 2017) gene models.

Annotated alternative splicing events and the number of reads supporting each alternative were identified using the exon-intron overlap method introduced in English, Patel, and Loraine (2010) and further developed here for use with RNA-Seq data. Exons and introns from pairs of gene models from the same locus were compared to identify alternatively spliced regions. Regions where an intron in one model overlapped an exon in another model on the same strand were identified and used to define mutually exclusive splicing choices (Figure 1). Gene models that included an alternatively spliced region were designated the “L” form (for “Long”) with respect to the splicing choice. Likewise, models that lacked an alternatively spliced region were designated “S” (for “Short”). The major form was defined as the transcript variant with the highest expression when comparing pairs of gene models. Thus, the minor form had lower expression in comparison to the major form.

Alternatively spliced regions were labeled according to the type of alternative splicing, as follows. Regions flanked by alternative donor sites were designated “DS” for alternative donor site. Regions flanked by alternative acceptor sites were labeled “AS” for alternative acceptor site. AS and DS events that coincided with exon skipping were labeled “AS/ES” and “DS/ES”. Alternatively spliced regions arising from introns that the spliceosome sometimes failed to excise were designated “RI” for retained intron.

For each alternatively spliced region representing two mutually exclusive splicing choices, RNA-Seq read alignments that unambiguously supported one or the other splicing choice were counted. For AS and DS events, only gapped reads that aligned across intron junctions were counted. For RI events, gapped reads that aligned across the retained intron were counted as support for the intron-removed (S) form, and un-gapped reads that overlapped at least 20 bases



**FIGURE 1** Alternative splicing annotation. The overlap between an intron in one gene model and an exon in another gene model defines an alternatively spliced region. Arrows indicate splice sites, named AS for acceptor site and DS for donor site. Use of sites named AS-L or DS-L causes inclusion of the differentially spliced region, generating the longer (L) form. Similarly, DS-S and AS-S refer to sites that exclude the differentially spliced region and generate the shorter (S) form. (a) Alternative donor sites, in which the U2 snRNP complex forms at alternative locations on the 5’ end of introns. (b) Alternative acceptor sites, in which the U1 snRNP complex forms at alternative sites near the 3’ end of alternatively spliced introns. (c) Alternatively spliced intron, in which a donor/acceptor site pairing can either be used or not used, forming a retained intron (RI). (d) Alternatively spliced, skipped exon. In exon skipping, alternative splicing involves four sites, indicated by DS-S/L, AS-L, DS-L, and SD-S/L. Exon inclusion requires assembly of two spliceosome complexes linking DS-S/L with AS-L and DS-L with AS-S/L, while exon skipping requires linking DS-S/L and AS-S/L only

within the intron were counted as support for the intron-retained (L) form.

For each alternatively spliced region in each biological replicate, the number of reads supporting L or S, were used to calculate percent-spliced-in (PSI) as  $100 * L / (S + L)$ , where L was the number of reads supporting only the L form and S + L the number of reads that supported S or L but not both. This is the same as the splicing index described in Katz, Wang, Airolidi, and Burge (2010). To identify differentially spliced regions, a two-sided t test was used to compare PSI between sample

types. Because PSI variance was large for events with small *M* (very few informative reads), only alternatively spliced regions where *M* was 10 or more in at least three replicate libraries were tested. A false discovery rate (FDR) was calculated for each test using the method of Benjamini and Hochberg (Benjamini & Hochberg, 1995), as implemented in the R programming language “p.adjust” method. Alternative splicing events with  $FDR \leq 0.1$  were considered differentially alternatively spliced.

Software used to identify and quantify alternative events is available from <https://bitbucket.org/lorainelab/altspliceanalysis>. Data analysis code used to analyze RNA-Seq data is available from <https://bitbucket.org/lorainelab/ricealtsplice>. Data analysis code is implemented as R Markdown files designed to be run in the RStudio development environment. Readers interested in experimenting with different analysis parameters can clone the repository, modify the code, and re-run analyses as desired. RNA-Seq alignments, coverage graphs, and junctions data are available for visualization in Integrated Genome Browser (Freese, Norris, & Loraine, 2016).

## 2.4 | RT-PCR and capillary gel electrophoresis analysis of alternative splicing

Differential alternative splicing detected by analysis of RNA-Seq was re-tested using the reverse transcriptase, PCR-based fragment analysis method described in (Stamm, Smith, & Lührmann, 2012). Differentially spliced regions identified computationally were PCR-amplified using fluorescently labeled primers and quantified using capillary gel electrophoresis. One benefit of the method is that the results are expressed as relative abundances of splice variants within a sample, thus eliminating the need to normalize using reference genes as in traditional qRT-PCR experiments aimed at measuring overall gene expression.

For splicing validation, new rice seedlings equivalent to the mock-treated (control) samples from the RNA-Seq experiment were grown and harvested. Seedlings were grown hydroponically in pots containing either liquid media only or calcined clay granules watered with liquid media as recommended in (Eddy, Hahn, & Moulton, 2012). After 12 days, plants were removed from the pots and roots and shoots were collected separately. Roots and shoots from the same pot were combined to form paired biological replicates. Samples were flash frozen in nitrogen and stored at  $-80^{\circ}\text{C}$  prior to RNA extraction.

RNA was extracted using the RNeasy Plant Mini Kit from Qiagen following the manufacturer's instructions, including the optional DNase I digestion to eliminate DNA contamination. First strand cDNA was synthesized using oligo dT primers and 1  $\mu\text{g}$  of total RNA per 20- $\mu\text{l}$  reaction. PCR amplification of cDNA was performed using primers flanking differentially spliced regions, including one primer labeled with 6-carboxyfluorescein (6-FAM) for amplicon detection during fragment analysis. Cycle parameters included denaturation at  $94^{\circ}\text{C}$  for 2 min, followed by 24 cycles of  $94^{\circ}\text{C}$  for 15 s,  $50^{\circ}\text{C}$  for 30 s, and  $70^{\circ}\text{C}$  for 1 min, with a final elongation step of  $72^{\circ}\text{C}$  for 10 min. This was essentially the same regime described in (Stamm et al., 2012) but with fewer cycles to ensure reactions were stopped before exiting the logarithmic phase. PCR

products were combined with size standards and separated on a 3730 Genetic Analyzer (Life Technologies). Amplicons were quantified using manufacturer-provided software by calculating the area under each amplicon peak. The percentage of the variant containing the alternatively spliced region (%L, see above) was calculated by dividing the long form area by the total area for both long and short forms. Spreadsheets with data exported from the instrument, along with PSI calculations, are available in the project git repository (<https://bitbucket.org/lorainelab/ricealtsplice>) in a subfolder named “Experimental Testing.”

## 3 | RESULTS

### 3.1 | Most genes annotated as alternatively spliced favored one dominant form in rice

RNA-Seq read alignments from our previously generated rice libraries were used to assess alternative splicing in four sample types: roots and shoots from seedlings treated with the cytokinin compound benzyladenine (BA) or with a mock, control treatment (Raines et al., 2016). Using the exon-intron overlap method described previously (English et al., 2010), alternative splicing events within each gene were identified and annotated (Figure 1). Gene models that included an alternatively spliced region were designated the “L-long” alternative form and models that lacked an alternatively spliced region were designated “S-short”. For each alternative splicing event, the number of sequence alignments unambiguously supporting each alternative (L or S) was counted. These counts were then used to calculate percent-spliced-in (PSI), the percentage of read alignments supporting the longer (L) form.

In the combined data from all libraries from the rice data set there were 11,192 AS events. Of these, 77% had at least one read supporting each of the two splicing choices, and 19.8% had support for just one splicing choice. Only 2.8% of AS events has no reads supporting either form; these corresponded to genes with low or no expression in any of the sample types tested.

We further analyzed the 77% of AS events that had at least one read supporting each of the two splicing choices (L and S). Most genes annotated as alternatively spliced favored a single transcript, that is, 70% of the splicing events had a PSI  $<20\%$  or  $>80\%$  (Figure 2). Nevertheless, 29% of the splicing events (representing 1,861 genes) had a PSI between 20% and 80%, indicating that many genes produce multiple transcript variants (Figure 2).

### 3.2 | Rice genes with abundant support for both alternative splicing choices perform many diverse functions

We used Gene Ontology term enrichment to determine if the subset of genes in rice for which alternatively spliced forms were unusually abundant exhibited enrichment with specific functions or processes, for example, circadian cycling, in which alternative splicing might play a prominent regulatory role. We asked if some Gene Ontology terms were significantly enriched with genes containing alternative splicing

events with a PSI between 20% and 80%, corresponding to the central region of Figure 2. Interestingly, we found that these genes exhibited a diversity of gene functions, with no significant enrichment of functional categories. Thus, alternative splicing in which both transcripts are prevalent appears to affect genes with many functions in rice.


### 3.3 | Many rice genes are differentially spliced between roots and shoots but cytokinin hormone application has minimal effect on splicing

In animals, differential splicing between cell or tissue types contributes to cellular differentiation, especially in the nervous system (Naftelberg, Schor, Ast, & Kornblihtt, 2015). Less is known about the role of differential splicing in regulating cellular differentiation and other processes in plants. Rice shoots and roots are profoundly different tissues, but our previous analysis of rice RNA-Seq data found that many of the same genes were expressed in both tissues (Raines et al., 2016). This raises the question of how these two different tissues are able to carry out their specialized roles, and suggest the hypothesis that differential splicing could enable differential functions in genes expressed in both tissues, as proposed in Reddy et al. (2013).


Our previous analysis of the effects of cytokinin treatment on rice identified many thousands of genes that were differentially expressed in response to cytokinin (Raines et al., 2016). However, little is known about the role of alternative splicing during cytokinin response, except for one study in *Arabidopsis* that reported a shift in splicing of SR protein genes following cytokinin hormone treatment (Palusa et al., 2007). Therefore, we examined differential splicing in the rice RNA-Seq data set comparing root and shoot tissue with or without cytokinin.

First, we asked: When an alternatively spliced gene was expressed in two different sample types, was the relative abundance of splice variants the same or different? To address this, we examined the correlation of PSI between roots and shoots or between

BA-treated and mock-treated samples (Figure 3). We found that PSI was similar between BA-treated and untreated samples, as revealed by the tighter clustering of scatter plot points (Figure 3a,b). This indicated that genes that were alternatively spliced in BA-treated samples were also alternatively spliced in the controls, and that the relative abundance of splice variants was similar. Thus, the cytokinin hormone treatment had minimal effect on splicing. By contrast, there were many genes where the relative abundance of splice variants was different between roots and shoots (Figure 3c). Consistent with Figure 3, statistical testing of PSI differences between sample types identified 90 genes, where PSI was significantly different between roots and shoots ( $FDR \leq 0.1$ ), but only four and two genes where PSI was different between BA-treated samples and controls



**FIGURE 2** Distribution of percent-spliced-in (PSI) for annotated splicing events in rice where each choice was supported by at least one RNA-Seq alignment. The y axis is the frequency of splicing events, and the x axis the PSI between 0 and 100%. PSI was calculated as  $100 \cdot L / (S + L)$ , where L and S were the number of reads that supported the splicing choice that included (L) or excluded (S) the differentially spliced region. Read alignment counts from all 12 libraries were combined to obtain a global view of alternative splicing occurrence in rice seedlings. The U-shaped character of the distribution persisted whether lower or higher thresholds of informative reads were used



**FIGURE 3** Scatter plots comparing percent-spliced-in (PSI) between sample types in rice for annotated splicing events. PSI was calculated from RNA-Seq reads obtained from sequencing rice-seedling shoots and roots grown hydroponically and subjected to a 2-hr treatment with BA, a cytokinin analog, or a mock-treatment (control). PSI is the average of three biological replicates. Only events with at least 15 informative read alignments in all six samples being compared were included. (a) BA-treated rice roots (y axis) compared to mock roots (x axis). (b) BA-treated rice shoots (y axis) compared to mock shoots (x axis). (c) Mock shoots (y axis) compared to mock roots (x axis)

in roots and shoots, respectively (Table S1). Thus, we observed limited but non-trivial levels of differential alternative splicing between roots and shoots but minimal differential alternative splicing between control and BA-treated samples.

### 3.4 | Comparison of *Arabidopsis* differential splicing shows similar patterns to rice

To determine whether the observed patterns of differential splicing are similar in other plants, we analyzed splicing in *Arabidopsis* roots and shoots that had also been treated with the cytokinin compound benzyladenine in our previous study (Zubo et al., 2017). Due to the *Arabidopsis* libraries not being sequenced to the same depth as the rice libraries, many more splicing events had little or no support. Using the same FDR threshold as with the rice data set ( $FDR \leq 0.1$ ), we identified few differentially spliced regions between shoots and roots (3) and none in the control to treatment comparisons (Table S2). However, PSI was distributed similarly to rice in that most alternatively spliced genes expressed one major form (Figure S1a). In addition, scatter plots showing average PSI in treated versus untreated samples showed a much tighter clustering of points as compared to scatter plots comparing roots and shoots (Figure S1b–d). Statistical testing of PSI differences confirmed the cytokinin hormone treatment had minimal effect on splicing in *Arabidopsis*. Thus, the general pattern of more differential splicing between tissue types as compared to treatment with exogenous cytokinin appears conserved between rice and *Arabidopsis*.

### 3.5 | Alternative splicing remodeled protein-coding sequence more often than disrupting it in rice

Alternative splicing can occur anywhere in a gene, including UTR and protein-coding regions. Because three bases encode one amino acid, the lengths of spliced coding regions in a transcript are multiples of three. Thus, when alternatively spliced regions occur in coding regions and are not multiples of three, then inclusion of these regions in transcripts is likely to introduce a frame shift,

resulting in a premature stop codon and a truncated protein product. As shown in Table 1, there was an enrichment of alternatively spliced regions in rice that were evenly divisible by three in coding regions versus non-coding in all subsets of the data. These subsets included all annotated alternatively spliced regions, regions where the minor form was prevalent (the region between 20% and 80% PSI in Figure 2), and differentially spliced regions. Thus, alternative splicing within the coding regions of genes was biased against introducing frame shifts and promoted protein remodeling rather than truncation.

Most differential splicing between roots and shoots (67%) occurred within protein-coding regions (Table S1), suggesting that differential splicing is likely to affect gene function at the level of the protein product. In every instance of differential splicing, major and minor forms were both detected, with differential splicing observed as a change in the relative abundance of the two forms.

To further understand the effects of splicing on protein-coding sequences, we visualized differentially spliced regions together with RNA-Seq alignments, coverage graphs, and inferred junctions using genome browsers. Two genome browsers were used to visualize the data—Integrated Genome Browser (Freese et al., 2016) and ProtAnnot (Mall et al., 2016). Integrated Genome Browser (IGB) was used to examine RNA-Seq read alignments and compare alignments to the annotated gene structures. ProtAnnot, an IGB App, was used to search the InterPro database of conserved protein domains to find out how (or if) splicing inferred from RNA-Seq data was likely to affect gene function through remodeling of protein domains as detected by the InterProScan Web service (Finn et al., 2017).

Of the 105 differentially spliced regions between rice roots and shoots, 71 overlapped protein-coding sequence regions, suggesting that in these cases, splicing affected protein function. All but one (70/71) of the differentially spliced regions embedded in coding regions overlapped a predicted functional domain (e.g., a predicted transmembrane helix) or a region found by protein classification systems (e.g., Pfam (Finn et al., 2016) or PANTHER (Thomas et al., 2003)) to be conserved among members of the same protein family (Table S1 and Figure 4).

**TABLE 1** Counts of alternative splicing choices in rice where the length in nucleotides of the splicing event is evenly divisible by three or with remainder of 1 or 2. A remainder of 1 or 2 leads to a shift in frame in translation. A  $p$ -value  $< 0.05$  is considered significant

Alternative splicing					
Location	Event	Divisible by 3	Remainder of 1	Remainder of 2	$p$ -Value <sup>a</sup>
Coding region	Annotated as alternatively spliced	3,248	2,466	2,411	3e-36
UTR	Annotated as alternatively spliced	1,152	1,127	1,113	1
Coding region	Minor form is expressed	173	149	153	8e-6
UTR	Minor form is expressed	34	20	13	0.03
Coding region	Differentially spliced	33	24	18	0.03
UTR	Differentially spliced	6	5	11	0.79

<sup>a</sup> $p$ -Value obtained from binomial test of the null hypothesis that the true probability of a differentially spliced region having a length divisible by three is 1 in 3 and an alternative hypothesis that the probability is greater than 1 in 3.

Figure 4a shows LOC\_Os06g05110, encoding a putative chloroplast superoxide dismutase enzyme. Superoxide dismutase enzymes protect the plant from reactive oxygen species by catalyzing toxic superoxide radicals to molecular oxygen (Alscher, Erturk, & Heath, 2002). Transcript LOC\_Os06g05110.3 contained a retained intron, which caused a frame shift, a premature stop codon, and a truncated match to a manganese/iron superoxide dismutase C-terminal profile (Figure 4a). The lower panel of 4A shows RNA-Seq coverage graphs for shoots and root samples. Roots had a higher relative proportion of read support for the retained intron (long) form (38.5% in root, 13.6% in shoot) (Table 2). Average expression in roots was lower than in shoots, 12.5 versus 39.7 Reads Per Kilobase per Million mapped reads (RPKM) (Table 2). Thus, roots had lower overall expression of LOC\_Os06g05110 and an increased relative proportion of the retained intron transcript with a shortened truncated match to a manganese/iron superoxide dismutase profile.

Figure 4b shows LOC\_Os02g05830, which codes for ribulose biphosphate carboxylase (RuBisCO) small chain chloroplast precursor. Retained intron event in transcript LOC\_Os02g05830.2 caused a frame shift, a premature stop codon, and alternative carboxy terminal regions. This resulted in a shorter match to a RuBisCO small subunit profile from the Interpro database, as shown in the upper panel of 4B. The bottom panel of 4B shows that roots had an increased proportion of reads supporting the retained intron (long) form (88.2% in root, 10.1% in shoot) as well as lower overall expression compared to shoots, 4.8 versus 12.3 RPKM (Table 2).

Rice contains five RuBisCO small subunit genes with varying expression levels, but only LOC\_Os02g05830 was differentially alternatively spliced between roots and shoots. The relatively low levels of expression of LOC\_Os02g05830 contrasted with the expression of the paralogous RuBisCO small subunit gene LOC\_Os12g19470 (80.2 and 942 RPKM in roots and shoots). LOC\_Os12g19470 also contained an alternative spliced, retained intron form that introduced a frameshift and premature stop codon. However, we did not find statistically significant differential splicing between roots and shoots for LOC\_Os12g19470. Its proportion of long to short form splicing was stable across tissues and expression levels. Understanding how these splicing ratios were maintained is beyond the scope of this study, but the existence of differentially spliced paralogs may provide an opportunity to identify sequence features that stabilize splicing.

Figure 4c shows LOC\_Os12g08260, encoding dehydrogenase E1 component domain containing protein. In InterPro, the dehydrogenase E1 component domain is annotated as belonging to many dehydrogenases of diverse biological function. Thus, there is no known function for LOC\_Os12g08260 apart from its putative enzymatic activity. The top panel of Figure 4c shows that transcript variant LOC\_Os12g08260.5 contained a retained intron event that introduced a frame shift. The frame shift changed the carboxy terminus of the protein and disrupted a profile match to the dehydrogenase E1 component domain. The bottom panel of Figure 4c shows that the root sample had an increased proportion of reads supporting the retained intron (long) form (55.7% in root, 2.9% in shoot), and gene expression was lower overall in roots (8.0 vs. 30.3 RPKM) (Table 2).

### 3.6 | RT-PCR with capillary gel electrophoresis confirmed differential splicing between rice roots and shoots for 9 of 10 genes tested







We used a method based on capillary gel electrophoresis of fluorescently tagged PCR products to assay splicing of 10 genes detected as differentially spliced between rice roots and shoots (Stamm et al., 2012). New rice seedlings were grown under a close-to-identical replication of the RNA-Seq experiment. Primers were designed to amplify differentially spliced regions, including one primer that was conjugated to a fluorescent tag. Following PCR amplification of cDNA prepared from the new rice samples, products were resolved on a capillary-based sequencer and PSI calculated (Table 2). In 9 out of 10 genes, differential alternative splicing was confirmed. In the one case where differential alternative splicing was not confirmed, there were very few RNA-Seq read alignments covering the differentially spliced region, suggesting this was likely a false positive result. The FDR cutoff used to detect differential splicing in the RNA-Seq data was 0.1, corresponding to 1 in 10 false discoveries, in line with results from the microcapillary-based analysis.

## 4 | DISCUSSION

This study profiled alternative splicing using a high coverage RNA-Seq data from 10-day old, hydroponically grown rice seedlings treated with a cytokinin hormone. A less deeply sequenced data set from cytokinin-treated *Arabidopsis* seedlings provided a comparison between monocot versus eudicot. We found that cytokinin treatment induced very few splicing changes between treated and untreated controls for both rice and *Arabidopsis*. However, there were many differences in splicing between untreated rice roots and shoots, and most of these changed the protein-coding region of genes.

Palusa et al. (2007) found that BA-treatment of *Arabidopsis* seedlings triggered splicing changes in multiple SR genes, encoding RNA-binding proteins whose counterparts in metazoans regulated alternative splice site choice. Their study used PCR amplification of cDNA followed by agarose gel electrophoresis to detect changes in splicing and focused on SR protein genes only. Thus, we expected to observe changes in SR gene splicing due to the cytokinin treatment, leading to changes in splicing for many downstream genes. However, close examination of SR splicing genes within our RNA-Seq datasets revealed no statistically significant differential splicing due to BA-treatment. It is possible that the differences in methodology used to measure splicing changes between the two studies (RNA-Seq vs. visualization of PCR amplification of cDNA) could account for the differences in observations.

One possible explanation for why the cytokinin treatment had minimal effect on splicing was that the treatment itself was ineffective. However, differential expression analysis showed that many genes were up- or downregulated by the treatment in the two data sets tested—rice and *Arabidopsis* (Raines et al., 2016; Zubo et al., 2017). Fewer genes were detected as differentially expressed in the





**FIGURE 4** ProtAnnot and IGB images showing differences in splicing between rice shoot and root. ProtAnnot (upper panels) shows coding region exons color-coded by frame, with regions matching InterPro profiles indicated by green, linked rectangles. Asterisk characters highlight differences in profile matches. Integrated Genome Browser (lower panels) shows a zoomed-in view of RNA-Seq coverage graphs from rice root (blue) and shoot (green). Y axis is the number of RNA-Seq aligned sequences with MSU7 gene models in black below. (a) LOC\_Os06g05110 isoforms .3 and .1 encoding superoxide dismutase, matching InterPro profile IPR019831 and IPR019832. (b) LOC\_Os02g05830 isoforms .2 and .1 encoding RuBisCO small subunit, matching InterPro profile IPR000894. (c) LOC\_Os12g08260 isoforms .5 and .1 encoding dehydrogenase E1 component, matching InterPro profile PTHR11516

*Arabidopsis* data set, most likely reflecting higher variability between biological replicates combined with lower sequencing depth as compared to the rice data set. Nevertheless, known cytokinin response genes were differentially regulated in both data sets, showing the cytokinin treatment penetrated plant cells and induced stereotypical cytokinin signaling without also triggering changes in splicing.

The relative lack of differential splicing between cytokinin-treated and mock-treated samples suggests that cytokinin signaling does not employ alternative splicing as a regulatory mechanism to the same degree as with other plant hormones, notably abscisic acid (ABA). ABA plays a role in perception and response to stresses, especially desiccation stress (Maia, Dekkers, Dolle, Ligterink, & Hilhorst, 2014). ABA also plays a role in regulating splicing of SR45, an SR-like protein, and SR45 plays a role in regulating downstream splicing of multiple genes (Cruz, Carvalho, Richardson, & Duque, 2014). Thus, there is a clear connection between stress signaling and the plant stress hormone ABA.

By contrast, cytokinin signaling involves transfer of phosphate groups between successive elements of a phosphorelay signaling pathway culminating in phosphorylation-dependent activation of Myb-type transcription factor proteins called type B ARR. Cytokinin treatment has no or little effect on transcription of type B ARRs, the key regulators of cytokinin signaling (Argyros et al., 2008; Kieber &

Schaller, 2018). In addition, type B transcriptional regulators are not highly alternatively spliced. By contrast, a closely related family of similar genes encoding so-called “pseudo-response regulators” have similar sequence to type B ARRs and are highly alternatively spliced (James et al., 2012). These genes are involved in regulating the circadian clock and have nothing to do with cytokinin signaling.

Using the same methods and data set, we identified a relatively large number of genes in rice (90) that were differentially spliced genes between shoots and roots, and we validated 9 of 10 using fragment analysis of independently produced rice samples (Tables S1 and 2). This observation of differential splicing between roots and shoots is important for two key reasons. First, it shows that our data analysis methods can identify differential alternative splicing in a data set. In other words, the roots versus shoots comparison provided a positive control for differential splicing detection. Second, the detection of differential splicing between roots and shoots illuminates the function of alternative splicing in plant cells. Our data supports the growing body of evidence that alternative splicing is cell, tissue, and stage specific in plants (Gupta et al., 2015; Sun et al., 2018; Vitulo et al., 2014), including in roots (Li et al., 2016).

Interestingly, in the three examples highlighted in Figure 4, the proportion of retained intron form was anti-correlated with transcript

**TABLE 2** Differential splicing of 10 genes in rice. Genes were first detected using RNA-Seq and re-tested using capillary gel electrophoresis (CGE). A  $p$ -value < 0.05 is considered significant

Gene	AS type	Avg. RPKM expression		RNA-seq PSI (%)		CGE PSI (%)		$p$ -Value <sup>a</sup>
		Root	Shoot	Root	Shoot	Root	Shoot	
LOC_Os01g25484, ferredoxin nitrite reductase	RI	300	131	74.2	31.7	29.9	13	5e-05
LOC_Os01g35580, unknown	AS	55.7	34.9	44.0	66.3	49.9	66.9	2.47e-04
LOC_Os01g45274, carbonic anhydrase	ES	171	1,380	96.8	24.3	97.4	15.4	3e-09
LOC_Os01g51290, protein kinase	RI	49.6	49.9	88.4	95.1	13.3	17.2	0.03466
LOC_Os03g05390, citrate transporter	RI	219	174	86.0	96.3	86	95.5	1.2e-04
LOC_Os12g08260, dehydrogenase E1	RI	8.03	30.3	55.7	2.9	4.1	0.87	3e-05
LOC_Os01g61670, ureidoglycolate hydrolase	DS	59	37.8	78.0	31.0	59	37.8	1.6e-09
LOC_Os05g48040, MATE efflux family protein	DS	7.05	6.72	88.1	100.0	89.4	88.27	0.632
LOC_Os02g05830, ribulose biphosphate carboxylase	RI	4.79	12.3	88.2	10.1	23.9	3.06	8e-04
LOC_Os06g05110, superoxide dismutase	RI	12.5	39.7	38.5	13.6	22.7	6.5	1.3e-07

<sup>a</sup> $p$ -Value obtained from comparing roots and shoots PSI from CGE.



levels overall. In all three cases, the retained intron led to a premature stop codon and truncation of the predicted domain. In general, superoxide dismutase and RuBisCO are important in photosynthetic tissue, and LOC\_Os06g05110 and LOC\_Os02g05830 had higher expression in shoots than roots (Alscher et al., 2002; Andersson & Backlund, 2008). Although we cannot say to what extent alternative splicing affected the protein function, we presume that the truncated domains would have a functional difference. As expression is already decreased in roots compared to shoots for LOC\_Os06g05110 and LOC\_Os02g05830, the increased proportion of retained intron in roots (and thus truncated domain) may act to further downregulate the number of fully functional proteins. It is likely that alternative splicing plays a role in fine-tuning gene function to meet the needs of different plant tissues or cell types where a gene is expressed.

We also examined the prevalence of alternative splicing, independent from differential splicing. That is, we used RNA-Seq read alignments to assess how often annotated alternative splice sites were used in our RNA-Seq data sets. For most genes annotated as alternatively spliced, the minor form frequency was typically low, accounting for <20% of sequence read alignments across the differentially spliced region. Genes where minor form frequency exceeded 20% exhibited a diversity of functions. Thus, many diverse processes in rice involved alternatively spliced genes in which splice variants were expressed at levels likely to affect gene function in different ways.

A major limitation of this study was that we limited our analysis to annotated splice forms and did not attempt to form new transcript models based on the RNA-Seq data. This was done mainly because the libraries used were not strand-specific and attempts to assemble transcripts using transcript assembly tools led to incorrect fusion of neighboring genes and other artifacts (not shown). Future studies will therefore benefit greatly from using better library preparation protocols to simplify and streamline data analysis. Nonetheless, this analysis provides new insight into the role of alternative splicing in plant tissues and hormone response.

In conclusion, by analyzing the number of reads that supported different splice variants, we identified examples of differential splicing with confirmation by RT-PCR with capillary gel electrophoresis. There were 90 genes differentially spliced between rice root and shoot tissues, but only four between cytokinin-treated and non-treated samples. For most differential splicing events, the protein-coding regions were affected, strongly suggesting that differential splicing is playing a role in modulating gene function between roots and shoots.

## ACKNOWLEDGMENTS

The authors would like to acknowledge Dr. Alex Feltus and the Renaissance Computing Institute for providing access to computational resources. This material is based upon work supported by the National Science Foundation through the Postdoctoral Research Fellowship in Biology under Grant No. 1523814 awarded to NHF and the Plant Genome Research Program award number IOS-1238051. NIH award 2R01GM103463-06A1 to AEL supports development of the IGB software. Computational resources used in this work

were supported by NSF Grant No. 1659300 “CC\*Data: National Cyberinfrastructure for Scientific Data Analysis at Scale (SciDAS).”

## CONFLICT OF INTEREST

The authors declare no conflict of interest associated with the work described in this manuscript.

## AUTHOR CONTRIBUTIONS

AEL designed the research. ARE and ICB performed the experiments. NHF, ARE, ICB, JD, and AEL analyzed the data. NHF, ARE, and AEL drafted the manuscript with contributions from ICB and JD.

## ORCID

Nowlan H. Freese  <https://orcid.org/0000-0002-5616-9230>

Ann E. Loraine  <https://orcid.org/0000-0002-8365-0177>

## REFERENCES

- Alscher, R. G., Erturk, N., & Heath, L. S. (2002). Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *Journal of Experimental Botany*, *53*, 1331–1341.
- Andersson, I., & Backlund, A. (2008). Structure and function of Rubisco. *Plant Physiology and Biochemistry*, *46*, 275–291.
- Argyros, R. D., Mathews, D. E., Chiang, Y.-H., Palmer, C. M., Thibault, D. M., Etheridge, N., ... Schaller, G. E. (2008). Type B response regulators of *Arabidopsis* play key roles in cytokinin signaling and plant development. *The Plant Cell*, *20*, 2102–2116.
- Barbosa-Morais, N. L., Carmo-Fonseca, M., & Aparicio, S. (2006). Systematic genome-wide annotation of spliceosomal proteins reveals differential gene family expansion. *Genome Research*, *16*, 66–77.
- Barbosa-Morais, N. L., Irimia, M., Pan, Q., Xiong, H. Y., Gueroussov, S., Lee, L. J., ... Blencowe, B. J. (2012). The evolutionary landscape of alternative splicing in vertebrate species. *Science*, *338*, 1587–1593.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society*, *57*, 289–300.
- Cheng, C. Y., Krishnakumar, V., Chan, A. P., Thibaud-Nissen, F., Schobel, S., & Town, C. D. (2017). Araport11: A complete reannotation of the *Arabidopsis thaliana* reference genome. *The Plant Journal*, *89*, 789–804.
- Cruz, T. M., Carvalho, R. F., Richardson, D. N., & Duque, P. (2014). Abscisic acid (ABA) regulation of *Arabidopsis* SR protein gene expression. *International Journal of Molecular Sciences*, *15*, 17541–17564.
- Eddy, R., Hahn, D. T., & Moulton, T. (2012). Optimizing greenhouse rice production: What is the best root medium? *Purdue Methods for Rice Growth*. Retrieved from <https://docs.lib.purdue.edu/pmrg/1>
- English, A. C., Patel, K. S., & Loraine, A. E. (2010). Prevalence of alternative splicing choices in *Arabidopsis thaliana*. *BMC Plant Biology*, *10*, 102.
- Filichkin, S. A., Cumbie, J. S., Dharmawardhana, P., Jaiswal, P., Chang, J. H., Palusa, S. G., ... Mockler, Todd. C. (2015). Environmental stresses modulate abundance and timing of alternatively spliced circadian transcripts in *Arabidopsis*. *Molecular Plant*, *8*, 207–227.
- Filichkin, S., Priest, H. D., Megraw, M., & Mockler, T. C. (2015). Alternative splicing in plants: Directing traffic at the crossroads of adaptation and environmental stress. *Current Opinion in Plant Biology*, *24*, 125–135.

- Finn, R. D., Attwood, T. K., Babbitt, P. C., Bateman, A., Bork, P., Bridge, A. J., ... Mitchell, A. L. (2017). InterPro in 2017—Beyond protein family and domain annotations. *Nucleic Acids Research*, *45*, D190–D199.
- Finn, R. D., Coghill, P., Eberhardt, R. Y., Eddy, S. R., Mistry, J., Mitchell, A. L., ... Bateman, A. (2016). The Pfam protein families database: Towards a more sustainable future. *Nucleic Acids Research*, *44*, D279–D285.
- Freese, N. H., Norris, D. C., & Loraine, A. E. (2016). Integrated genome browser: Visual analytics platform for genomics. *Bioinformatics*, *32*, 2089–2095.
- Gulledge, A. A., Roberts, A. D., Vora, H., Patel, K., & Loraine, A. E. (2012). Mining *Arabidopsis thaliana* RNA-seq data with integrated genome browser reveals stress-induced alternative splicing of the putative splicing regulator SR45a. *American Journal of Botany*, *99*, 219–231.
- Gupta, V., Estrada, A. D., Blakley, I., Reid, R., Patel, K., Meyer, M. D., ... Loraine, A. E. (2015). RNA-Seq analysis and annotation of a draft blueberry genome assembly identifies candidate genes involved in fruit ripening, biosynthesis of bioactive compounds, and stage-specific alternative splicing. *Gigascience*, *4*, 5.
- James, A. B., Syed, N. H., Bordage, S., Marshall, J., Nimmo, G. A., Jenkins, G. I., ... Nimmo, H. G. (2012). Alternative splicing mediates responses of the *Arabidopsis* circadian clock to temperature changes. *Plant Cell*, *24*, 961–981.
- Kalsotra, A., & Cooper, T. A. (2011). Functional consequences of developmentally regulated alternative splicing. *Nature Reviews Genetics*, *12*, 715–729.
- Kalyna, M., & Barta, A. (2004). A plethora of plant serine/arginine-rich proteins: Redundancy or evolution of novel gene functions? *Biochemical Society transactions*, *32*, 561–564.
- Katz, Y., Wang, E. T., Airoidi, E. M., & Burge, C. B. (2010). Analysis and design of RNA sequencing experiments for identifying isoform regulation. *Nature Methods*, *7*, 1009–1015.
- Kawahara, Y., de la Bastide, M., Hamilton, J. P., Kanamori, H., McCombie, W. R., Ouyang, S., ... Matsumoto, T. (2013). Improvement of the *Oryza sativa* Nipponbare reference genome using next generation sequence and optical map data. *Rice*, *6*, 1–10.
- Keller, M., Hu, Y., Meshovic, A., Fragkostefanakis, S., Schleiff, E., & Simm, S. (2016). Alternative splicing in tomato pollen in response to heat stress. *DNA research*, *24*, 205–217.
- Kieber, J. J., & Schaller, G. E. (2018). Cytokinin signaling in plant development. *Development*, *145*, 1–7.
- Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., & Salzberg, S. L. (2013). TopHat2: Accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biology*, *14*, R36.
- Lamesch, P., Berardini, T. Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., ... Huala, E. (2012). The Arabidopsis Information Resource (TAIR): Improved gene annotation and new tools. *Nucleic Acids Research*, *40*, D1202–D1210.
- Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, *9*, 357.
- Li, S., Yamada, M., Han, X., Ohler, U., & Benfey, P. N. (2016). High resolution expression map of the *Arabidopsis* root reveals alternative splicing and lincRNA regulation. *Developmental Cell*, *39*, 508–522.
- Loraine, A. E., McCormick, S., Estrada, A., Patel, K., & Qin, P. (2013). RNA-seq of *Arabidopsis* pollen uncovers novel transcription and alternative splicing. *Plant Physiology*, *162*, 1092–1109.
- Maia, J., Dekkers, B. J., Dolle, M. J., Ligterink, W., & Hilhorst, H. W. (2014). Abscisic acid (ABA) sensitivity regulates desiccation tolerance in germinated *Arabidopsis* seeds. *New Phytologist*, *203*, 81–93.
- Mall, T., Eckstein, J., Norris, D., Vora, H., Freese, N. H., & Loraine, A. E. (2016). ProtAnnot: An App for Integrated Genome Browser to display how alternative splicing and transcription affect proteins. *Bioinformatics*, *32*, 2499–2501.
- Mei, W., Boatwright, L., Feng, G., Schnable, J. C., & Barbazuk, W. B. (2017). Evolutionarily conserved alternative splicing across monocots. *Genetics*, *207*, 465–480.
- Naftelberg, S., Schor, I. E., Ast, G., & Kornblihtt, A. R. (2015). Regulation of alternative splicing through coupling with transcription and chromatin structure. *Annual Review of Biochemistry*, *84*, 165–198.
- Palusa, S. G., Ali, G. S., & Reddy, A. S. N. (2007). Alternative splicing of pre-mRNAs of Arabidopsis serine/arginine-rich proteins: Regulation by hormones and stresses. *The Plant Journal*, *49*, 1091–1107.
- Plass, M., Agirre, E., Reyes, D., Camara, F., & Eyras, E. (2008). Co-evolution of the branch site and SR proteins in eukaryotes. *Trends in Genetics*, *24*, 590–594.
- Raines, T., Blakley, I. C., Tsai, Y.-C., Worthen, J. M., Franco-Zorrilla, J. M., Solano, R., ... Kieber, J. J. (2016). Characterization of the cytokinin-responsive transcriptome in rice. *BMC Plant Biology*, *16*, 260.
- Reddy, A. S. N., Marquez, Y., Kalyna, M., & Barta, A. (2013). Complexity of the alternative splicing landscape in plants. *The Plant Cell*, *25*, 3657–3683.
- Sakai, H., Lee, S. S., Tanaka, T., Numa, H., Kim, J., Kawahara, Y., ... Itoh, T. (2013). Rice Annotation Project Database (RAP-DB): An integrative and interactive database for rice genomics. *Plant and Cell Physiology*, *54*, e6.
- Salz, H. K. (2011). Sex Determination in Insects: A binary decision based on alternative splicing. *Current Opinion in Genetics & Development*, *21*, 395–400.
- Stamm, S., Smith, C., & Lührmann, R. (2012). *Alternative pre-mRNA splicing: Theory and protocols*. Weinheim, Germany: Wiley-Blackwell.
- Sun, Y., Hou, H., Song, H., Lin, K., Zhang, Z., Hu, J., & Pang, E. (2018). The comparison of alternative splicing among the multiple tissues in cucumber. *BMC Plant Biology*, *18*, 5.
- Thomas, P. D., Campbell, M. J., Kejarawal, A., Mi, H., Karlak, B., Daverman, R., ... Narechania, A. (2003). PANTHER: A Library of Protein Families and Subfamilies Indexed by Function. *Genome Research*, *13*, 2129–2141.
- Vanechoutte, D., Estrada, A. R., Lin, Y. C., Loraine, A. E., & Vandepoele, K. (2017). Genome-wide characterization of differential transcript usage in *Arabidopsis thaliana*. *The Plant Journal*, *92*, 1218–1231.
- Vitulo, N., Forcato, C., Carpinelli, E. C., Telatin, A., Campagna, D., D'Angelo, M., ... Valle, G. (2014). A deep survey of alternative splicing in grape reveals changes in the splicing machinery related to tissue, stress condition and genotype. *BMC Plant Biology*, *14*, 99–99.
- Zubo, Y. O., Blakley, I. C., Yamburenko, M. V., Worthen, J. M., Street, I. H., Franco-Zorrilla, J. M., ... Schaller, G. E. (2017). Cytokinin induces genome-wide binding of the type-B response regulator ARR10 to regulate growth and development in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, *114*, E5995–E6004.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Freese NH, Estrada AR, Blakley IC, Duan J, Loraine AE. Many rice genes are differentially spliced between roots and shoots but cytokinin has minimal effect on splicing. *Plant Direct*. 2019;3:1–11. <https://doi.org/10.1002/pld3.136>