

RESEARCH ARTICLE

$\gamma\delta$ T cells shape memory-phenotype $\alpha\beta$ T cell populations in non-immunized mice

Swati Popat Phalke^{1‡}, Yafei Huang², Kira Rubtsova¹, Andrew Getahun³, Deming Sun⁴, Richard L. Reinhardt^{1,3}, Rebecca L. O'Brien^{1,3}, Willi K. Born^{1,3*}

1 Department of Biomedical Research, National Jewish Health, Denver, CO, United States of America, **2** Joint Laboratory for Stem Cell Engineering and Technology Transfer, School of Basic Medicine, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, PR China, **3** Department of Immunology and Microbiology, University of Colorado Health Sciences Center, Aurora, CO, United States of America, **4** Doheny Eye Institute and Department of Ophthalmology, David Geffen School of Medicine at UCLA, Los Angeles, CA, United States of America

‡ Current address: David Vermijlen Lab, Department of Pharmacotherapy and Pharmaceuticals—Institute of Medical Immunology (IMI), University Libre de Bruxelles, Brussels, Belgium.

* bornw@njhealth.org



OPEN ACCESS

Citation: Phalke SP, Huang Y, Rubtsova K, Getahun A, Sun D, Reinhardt RL, et al. (2019) $\gamma\delta$ T cells shape memory-phenotype $\alpha\beta$ T cell populations in non-immunized mice. PLoS ONE 14 (6): e0218827. <https://doi.org/10.1371/journal.pone.0218827>

Editor: Moriya Tsuji, Rockefeller University, UNITED STATES

Received: April 16, 2019

Accepted: June 11, 2019

Published: June 25, 2019

Copyright: © 2019 Phalke et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: This research was funded by NIH/NIAID 1R21 AI122135-01A1 to WKB. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Size and composition of $\gamma\delta$ T cell populations change dramatically with tissue location, during development, and in disease. Given the functional differentiation of $\gamma\delta$ T cell subsets, such shifts might alter the impact of $\gamma\delta$ T cells on the immune system. To test this concept, and to determine if $\gamma\delta$ T cells can affect other immune cells prior to an immune response, we examined non-immunized mice derived from strains with different genetically induced deficiencies in $\gamma\delta$ T cells, for secondary changes in their immune system. We previously saw extensive changes in pre-immune antibodies and B cell populations. Here, we report effects on $\alpha\beta$ T cells. Similarly to the B cells, $\alpha\beta$ T cells evidently experience the influence of $\gamma\delta$ T cells at late stages of their pre-immune differentiation, as single-positive heat stable antigen-low thymocytes. Changes in these and in mature $\alpha\beta$ T cells were most prominent with memory-phenotype cells, including both CD8+ and CD4+ populations. As previously observed with B cells, most of the effects on $\alpha\beta$ T cells were dependent on IL-4. Unexpectedly, IL-4 seemed to be produced mainly by $\alpha\beta$ T cells in the non-immunized mice, albeit strongly regulated by $\gamma\delta$ T cells. Similarly to our findings with B cells, changes of $\alpha\beta$ T cells were less pronounced in mice lacking all $\gamma\delta$ T cells than in mice lacking only some, suggesting that the composition of the $\gamma\delta$ T cell population determines the nature of the $\gamma\delta$ -influence on the other pre-immune lymphocytes.

Introduction

$\gamma\delta$ T cell populations change in numbers and in cellular composition with human age [1], but also with infectious and autoimmune diseases [2]. In primary immune-deficiencies, the genetic background of an individual can be associated with such variations as well [2], and hematopoietic transplantation induces changes because $\gamma\delta$ T cells are only slowly and

incompletely restored [3, 4]. Likewise, mouse ontogeny is associated with an ordered appearance of $\gamma\delta$ T cells expressing different TCR-V γ genes (subsets), causing compositional shifts in the population over time. Furthermore, throughout life, different mouse tissues contain $\gamma\delta$ T cell populations with distinct subset compositions [5, 6]. It was found that the TCR-V γ -defined cell subsets have different functional potentials [7–12], suggesting a connection between innate TCR-ligand-specificities and functional differentiation [13–15], and overall changes of $\gamma\delta$ T cell functions during ontogeny and between the tissues.

Albeit in small numbers, $\gamma\delta$ T cells can exert a strong regulatory influence on inflammatory and immune responses [16, 17]. They also affect the immune system in the steady state [18]. Because of their functional differences, compositional shifts in the $\gamma\delta$ T cell populations ought to alter the net-effect $\gamma\delta$ T cells might have on other immune cells, but this influence has been difficult to demonstrate because external conditions that lead to shifts in $\gamma\delta$ T cell populations often also directly affect other cell-types. Therefore, to specifically investigate the effect of compositional changes in $\gamma\delta$ T cell populations on other cells, we assembled a collection of background-matched mice with genetic mutations in TCR genes causing the loss of some or all $\gamma\delta$ T cells [19–21], as a model for the natural shifts in $\gamma\delta$ T cell populations introduced above. These mice were not immunized, challenged, or treated in other ways. Nevertheless, their comparison revealed numerous secondary changes of the immune system, including differences in levels of immunoglobulin (Ig) subclasses [18, 22], in the composition of antibody specificities (repertoire) [18], in serum levels of cytokines and the capability of T cells to produce them [18], as well as shifts in granulocytes and B cells [18, 23].

We initially focused our investigation on the secondary changes in B cells, and confirmed that they were the result of interactions with the altered $\gamma\delta$ T cell populations [18, 23]. The current study extends the investigation to secondary changes in $\alpha\beta$ T cells. In the course of exploring underlying mechanisms, we compared effects on $\alpha\beta$ T cells and B cells. As with B cells [18, 23], we found that changes in specific $\alpha\beta$ T cell populations depend on the particular deficiency in $\gamma\delta$ T cells. With either lymphocyte-type, the $\gamma\delta$ -influence begins at late immature developmental stages, and most of the effects apparently are mediated by $\gamma\delta$ T cell-regulated and $\alpha\beta$ T cell-sourced interleukin 4 (IL-4).

Materials and methods

Mice

C57BL/6J mice and $\gamma\delta$ T cell-deficient mice of the same genetic background (B6.TCR δ KO mice) were originally obtained from the Jackson Laboratory (Bar Harbor, ME) and bred at National Jewish Health (Denver, CO). TCR-V γ 4 $^{-/-}$ /V γ 6 $^{-/-}$ mice [20], which were a gift from Dr. Koichi Ikuta (Kyoto University, Kyoto, Japan) were then backcrossed onto the C57BL/6J genetic background and re-established as homozygous line after 12 backcross generations (B6.TCR-V γ 4/6KO mice). B6.TCR-V γ 1 $^{-/-}$ mice [24] were a gift from Dr. Simon Carding (Norwich Medical School, Norwich, U.K.), and provided by Dr. C. Wayne Smith (Baylor College of Medicine, Houston, TX). These mice were still further backcrossed onto the C57BL/6J genetic background and re-established as homozygous line after >10 backcross generations (B6.TCR-V γ 1KO mice). IL-4 $^{-/-}$ mice (C57BL/6 $^{-Il4tm1Nnt/J}$) were obtained from the Jackson Laboratory and were a gift from Dr. P. Murrack at National Jewish Health. IL-4 $^{-/-}$ mice were crossed with B6.TCR-V γ 4/6KO mice, double KO mice selected from the F2 generation and bred as a new homozygous line (B6.TCR-V γ 4/6KO/IL-4KO mice). All mice were cared for at National Jewish Health (NJH) following guidelines for normal and immune-deficient animals, and euthanized by CO2 inhalation following current guidelines of the American Veterinary Medical Association (AVMA). All efforts were made to minimize suffering. The experiments

were conducted under a protocol approved by the National Jewish Health Institutional Animal Care and Use Committee (Protocol number AS2484).

Although the KO mice were backcrossed $>10x$, this approach cannot completely rule out an influence of genes closely linked to the locus of interest. Therefore, we sought to complement the genetic evidence with alternative ways of changing the same $\gamma\delta$ T cell populations, using antibody treatment for $\gamma\delta$ T cell-inactivation and adoptive cell-transfer for $\gamma\delta$ T cell-reconstitution. Such experiments in our preceding studies confirmed that the altered $\gamma\delta$ T cells mediate the changes in antibodies and B cells seen in the $\gamma\delta$ -gene mutated mice [18, 22, 23, 25]. The similarity of $\gamma\delta$ -effects on the $\alpha\beta$ T-cells in this study, including changes seen with specific $\gamma\delta$ -deficiencies, the critical role of IL-4, and the developmental timing of the $\gamma\delta$ -effect on $\alpha\beta$ T cells, suggest that similar mechanisms are at work, and that here as well interactions with $\gamma\delta$ T cells are responsible for the effects.

Flow cytometric analysis

Cells obtained from single cell suspensions (5×10^5 /well) were stained in 96-well plates (Falcon, BD Biosciences) for the cell surface markers shown in the figures, using the specific mAbs listed below. Lymphocytes were gated based on their forward (FSC) and side scatter (SC), and aggregates (doublets and other conjugates) were excluded by using forward scatter area (FSC-A) and forward scatter height (FSC-H). Samples were analyzed using LSR II and LSR Fortessa flow cytometers (Becton Dickinson). At least 1×10^5 events were acquired per gated region, and the data were analyzed using FlowJo 10.5.2 software (TreeStar).

Antibody reagents

Antibodies for flow cytometry included the following clones and conjugates: anti B220 (clone RA3-6B2, BV421, Biolegend, and APCCy7, eBiosciences); anti CD93 (clone AA4.1, APC, Invitrogen); anti CD23 (clone B3B4, PECy7, eBiosciences); anti IgM (clone B76, AF488, home-made); anti Fas (clone Jo2, PE, BD Pharmingen); anti IL-4R α (clone mIL4R-M1, PE, BD Biosciences); anti CD3 (clone 17A2, PB, Biolegend); anti CD4 (clone RM4.5, APC, BD Biosciences and BV510, Biolegend); anti CD8 β .2 (clone 53–5.8, FITC, BD Biosciences); anti Fas-L (clone mFL3, PE, eBiosciences); anti TCR- β (clone H57-597, PB, Biolegend and PercpCy5.5, eBiosciences); anti CD24 (clone M1/69, BV711, BD Biosciences); anti CD8 α (clone 53–6.7, APCeF780, eBiosciences); anti CD122 (clone 5H4, PE, eBiosciences); anti CD44 (clone IM7, BV605, BD Biosciences); anti CD49d (clone R1-2, APC, Biolegend); anti CXCR3 (clone CXCR3-173, PECy7, Biolegend); anti T-bet (clone 4B10, BV421, Biolegend); anti Eomes (clone 1219A, AF488, R&D Systems).

In vivo transient inactivation of T cells

In order to selectively inactivate T cells and subsets of T cells *in vivo*, we i.v.-injected affinity-purified mAbs specific for TCR- β (clone H57.597.2), TCR- δ (clone GL3), TCR-V γ 1 (clone 2.11) and TCR-V γ 4 (clone UC3) at 200 μ g/mouse as previously described in detail [26]. This treatment does not deplete [27] but efficiently and specifically inactivates the targeted T cells (S1–S3 Figs) for up to two weeks [26]. The antibody-treated mice were used 7 days post injection for *in vivo* capture of IL-4.

In vivo production of IL-4

In vivo production of IL-4 was measured using the Mouse IL-4 *In Vivo* Capture Assay Set (BD Pharmingen), following the method of Finkelman and Morris [28]. Unlike our earlier study

[18], where we used an *in vivo* capture period of 8.5 hrs to demonstrate the difference in IL-4 production between C57BL/6 (wt) and B6.TCR-V γ 4/6KO mice, we extended the *in vivo* capture period here to 24 hrs. This modification increased the IL-4 signal roughly 2x to ~ 400 pg IL-4/ml serum but abolished the difference between the mouse strains, presumably because the injected capture antibody (10 μ g/mouse) becomes limiting.

Nomenclature

We used the nomenclature for murine TCR-V γ genes introduced by Heilig and Tonegawa [29].

Statistical analysis

Each experiment was carried out at least 3 times independently. Data are presented as means \pm SD. The One Way Nonparametric ANOVA (Kruskal Wallis) test was used for group comparisons. All data generated with $\gamma\delta$ T cell-deficient mice were compared to the data generated with wt mice. A p value <0.05 was considered statistically significant.

Results

Splenic lymphopenia and loss of $\alpha\beta$ T cells in mice lacking certain $\gamma\delta$ T cells

$\gamma\delta$ T cells encompass several subpopulations with different functional capabilities [8]. For example, the V γ 1+ $\gamma\delta$ T cell population includes most of the $\gamma\delta$ T cells capable of producing IL-4 [30], whereas the V γ 4+ and V γ 6+ populations harbor the vast majority of IL-17-producing $\gamma\delta$ T cells [31, 32]. Because of these functional differences, shifts in the composition of total $\gamma\delta$ T cells might be expected to change their overall functional “output” and consequentially, other cells and tissues.

We compared C57BL/6 mice (wt) with background-matched mice deficient in all $\gamma\delta$ T cells (B6.TCR δ KO) or in only some $\gamma\delta$ T cells, lacking V γ 1+ cells (B6.TCR-V γ 1KO) or lacking V γ 4+ and V γ 6+ cells (B6.TCR-V γ 4/6KO). In young adult mice (8–12 weeks old, male and female), the absence of all $\gamma\delta$ T cells had little if any effect on splenic lymphocyte numbers (Fig 1A), or splenic $\alpha\beta$ T cells (Fig 1B and 1C). However, the absence of V γ 1+ $\gamma\delta$ T cells only was associated with significantly diminished numbers of CD8+ $\alpha\beta$ T cells (Fig 1C). In contrast, the absence of V γ 4+ and V γ 6+ $\gamma\delta$ T cells was associated with splenic lymphopenia (Fig 1A), and a pronounced loss of CD4+ but not of CD8+ $\alpha\beta$ T cells (Fig 1B and 1C). These changes in splenic $\alpha\beta$ T cell populations also affected their relative frequencies (S4 Fig). Of note, in terms of absolute numbers, the loss of $\alpha\beta$ T cells in the spleen was much smaller than the previously reported loss of splenic B cells [23].

We previously investigated IL-4-production in non-immunized B6.TCR-V γ 4/6KO mice [18]. An *in vivo* cytokine capture assay [28], employed for a period of 8.5 hrs, indicated that these mice have higher IL-4 serum levels when compared to wt mice [18]. Their splenic T cells exhibited elevated IL-4 production following *in vitro* activation, and on a per cell basis, $\gamma\delta$ T cells produced more IL-4 than $\alpha\beta$ T cells. But $\gamma\delta$ T cells in blood and spleen are less frequent than $\alpha\beta$ T cells, and we did not resolve the relative contributions of the two T cell-types to overall IL-4 serum levels. To address this question, we again used the *in vivo* cytokine capture assay, but this time in combination with *in vivo* T cell inactivation by antibody treatment [26]. Firstly, we pre-treated the mice with i.v. injected TCR-V-specific mAbs, at doses previously shown to inactivate the targeted T cells for approximately two weeks [26]. All of the antibodies effectively targeted T cells (assessed on day 8 post injection), with no discernible effects on B




Fig 1. Deficiency in $\gamma\delta$ T cells can affect splenic lymphocyte numbers and the size of splenic $\alpha\beta$ T cell populations. A) Comparison of total numbers of lymphocytes/spleen in C57BL/6 (B6), B6.TCR δ KO, B6.TCR-V γ 1KO, B6.TCR-V γ 4/6KO and B6.TCR-V γ 4/6KO/IL-4KO mice. n equal or greater than 10 mice/group. B) Numbers of CD4+ $\alpha\beta$ T cells (TCR- β +CD4+ lymphocytes) in the spleen, in the same mouse strains as in A. n equal or greater than 8 mice/group. C) Numbers of CD8+ $\alpha\beta$ T cells (TCR- β +CD8 α + lymphocytes) in the spleen, in the same mouse strains as in A. n equal or greater than 8 mice/group. Female and male mice ages 8–12 wks were included in the comparisons shown in Fig 1. * p <0.05, ** p <0.01, *** p <0.001.

<https://doi.org/10.1371/journal.pone.0218827.g001>

cells (S1–S3 Figs). At one week into the window of T-inactivation, we secondly employed the *in vivo* cytokine capture assay. To achieve maximum sensitivity, we used a longer *in vivo* capture period (24 hrs), achieving roughly 2x greater accumulation of captured IL-4 by comparison with our earlier study [18]. However, we lost the ability to differentiate between wt control and B6.TCR-V γ 4/6KO mice, presumably because the injected capture antibody becomes limiting over the extended capture period. Indeed, the total (increased) amount of IL-4 captured in untreated wt and B6.TCR-V γ 4/6KO mice was almost the same (~ 400 pg/ml serum) (Fig 2). Importantly, a large decrease of captured IL-4 in the mice pre-treated with anti TCR- β mAbs, but not in those treated with the other mAbs, revealed that the serum IL-4 in the non-immunized mice is mainly $\alpha\beta$ T cell-derived. This bias was similar for both wt and B6.TCR-V γ 4/6KO mice (Fig 2).

We previously showed that marginal zone and follicular B cells are deficient in B6.TCR-V γ 4/6KO mice, and found that removing IL-4 (B6.TCR-V γ 4/6KO/IL-4KO) restored the numbers of these B cells [23]. Consistently, we show here that removing IL-4 abrogated their splenic lymphopenia (Fig 1A). Importantly, the same manipulation also restored splenic CD4+ and CD8+ $\alpha\beta$ T cell populations, albeit with much variation between individual mice (Fig 1B and 1C).

In all strains, individual mice varied in their lymphocyte numbers. In part, this might have been due to residual genetic heterogeneity in the backcrossed mice (see Methods), or age-related, as mice were still growing at the included ages of 8–12 weeks. Female mice consistently had lower body weights (S5 Fig), but we included mice of both genders in the groups because splenic lymphocyte numbers (S6 Fig) and effects of $\gamma\delta$ -deficiencies on splenic $\alpha\beta$ T cells (S7 Fig) were indistinguishable.

$\gamma\delta$ T cells shape memory-phenotype CD8+ $\alpha\beta$ T cell populations in non-immunized mice

Mice harbor substantial numbers of memory-phenotype (MP) CD8+ $\alpha\beta$ T cells, even if they have not been exposed to foreign antigens. These cells are distinguished by their high level of expression of CD122 and CD44 [33]. Homeostatic expansion under lymphopenic conditions, and certain cytokines including IL-7, IL-4 and IL-15, can further increase this population [34–36]. Like naïve $\alpha\beta$ T cells, MP cells have a diverse TCR repertoire, but they respond to inflammation and infections more rapidly and strongly, and provide better host protection during primary immune responses [37].

Having found lymphopenia and higher-than wt IL-4 serum levels and production by T cells in certain $\gamma\delta$ T cell-deficient mouse strains [18, 23] (and current study), we examined $\alpha\beta$ T cells in the spleen focusing on MP cells (Fig 3) (gating strategy in S8 Fig). Representative cytograms (Fig 3A, top row) show that the percentage of CD8+ $\alpha\beta$ T cells with a memory phenotype was much increased in B6.TCR-V γ 4/6KO mice, but not in B6.TCR-V γ 4/6KO/IL-4KO mice. Consistently [38], a subset of the CD8+ $\alpha\beta$ T cells in B6.TCR-V γ 4/6KO mice, but not in B6.TCR-V γ 4/6KO/IL-4KO mice, expressed IL-4R α at increased levels (S9 Fig). Because overall numbers of CD8+ $\alpha\beta$ T cells in the spleens of wt and B6.TCR-V γ 4/6KO mice did not




Fig 2. Contribution of T cells to serum IL-4 levels in non-immunized mice. Effect of *in vivo* treatment of C57BL/6 (wt) and B6.TCR-V $\gamma 4/6$ KO mice with anti TCR mAbs on serum IL-4 levels: Mice were pre-treated with i.v. injections of saline or of the different TCR-specific mAbs specified in the figure, and serum IL-4 was later captured *in vivo* during a 24 hr period using an i.v. injected anti IL-4 mAb and subsequently detected by ELISA (see [Methods](#)). Top: C57BL/6 mice, n = 5; bottom: B6.TCR-V $\gamma 4/6$ KO mice, n = 3. *p<0.05.

<https://doi.org/10.1371/journal.pone.0218827.g002>

significantly differ ([Fig 1](#)), the increased frequency of MP CD8+ $\alpha\beta$ T cells in B6.TCR-V $\gamma 4/6$ KO mice reflects a selective expansion of these cells. In contrast to B6.TCR-V $\gamma 4/6$ KO mice, no substantial changes of MP CD8+ $\alpha\beta$ T cells were seen in B6.TCR δ KO or B6.TCR-V $\gamma 1$ KO mice ([S10 Fig](#)). [Fig 3B](#) shows a comparison of all mouse strains of this study with regard to the frequencies of splenic CD8+ $\alpha\beta$ T cells and MP subsets. In B6.TCR-V $\gamma 4/6$ KO mice but not in




Fig 3. Changes of peripheral memory phenotype αβ T cell populations in γδ T cell-deficient mice. A) Comparison of splenic CD8+ or CD4+ memory-phenotype αβ T cells in C57BL/6 (wt), B6.TCR-Vγ4/6KO and B6.TCR-Vγ4/6KO/IL-4KO mice. The representative flow cytograms shows a >3fold relative increase in the larger population of CD8+ memory-phenotype αβ T cells in a B6.TCR-Vγ4/6KO mouse. CD4+ αβ T cells expressing CD44 at high levels were also increased but only few of these expressed CD122 at high levels. Removal of IL-4 (B6.TCR-Vγ4/6KO/IL-4KO) restored near normal (wt) frequencies of all populations. B,C) Hierarchical comparison of splenic αβ T cells in wt and γδ T cell deficient mice (same strains as in Fig 1), including first CD8+ or CD4+ cells among TCR-β+ T cells, second CD122+CD44+ cells among either type of single-positive αβ T cells, third CD49d low cells among the memory phenotype single-positive αβ T cells, and fourth CXCR3 high cells among CD49d low memory phenotype single-positive αβ T cells. The hierarchical comparison is extended further in Table 1. Female and male mice ages 8–12 wks were included in the comparisons of Fig 2B. n equal or greater than 9 mice/group. *p<0.05, **p<0.01, ***p<0.001.

<https://doi.org/10.1371/journal.pone.0218827.g003>

the other mice, CD8+ αβ T cells were increased, with a predominance of CD122+CD44+ MP cells. B6.TCR-Vγ4/6KO/IL-4KO mice had normal frequencies (and numbers) of such cells. As with wt mice and the other mouse strains, almost all of the expanded MP CD8+ αβ T cells in B6.TCR-Vγ4/6KO mice expressed CD49d at low levels. This is typical for antigen-inexperienced (AI) MP αβ T cells (which would not, presumably, have been previously stimulated via their TCRs), because CD49d is up-regulated following stimulation via the TCR [33]. Therefore, the data suggest that the Vγ4+ and/or Vγ6+ γδ T cells present in wt mice function to control the size of the population of AIMP CD8+ αβ T cells and that IL-4 plays a role in this regulation.

AIMP CD8+ αβ T cells that arise as a result of lymphopenia-induced peripheral homeostatic proliferation [34], “virtual memory” (VM) MP T cells in the periphery, and IL-4-dependent innate MP αβ T cells in thymus and periphery, all differ in their gene expression [34, 37]. In order to determine which might be affected by γδ T cells, we examined products of such differentially expressed genes including CXCR3, a CXC chemokine receptor involved in sensing and responding to inflammation, and the transcription factors eomesodermin (EOMES) and T-bet. Nearly all of the CD122+CD44+CD49d^{lo} AIMP CD8+ splenic αβ T cells detected in the mouse strains of this study highly expressed CXCR3 (Fig 3B), and among these CXCR3^{high} cells, almost all expressed T-bet and EOMES (Tables 1 and 2). The nearly uniform transcription factor expression pattern among the expanded CD8+ MP αβ T cells in B6.TCR-Vγ4/6KO mice resembles that of the previously described VM CD8+ αβ T cells [34, 37, 39, 40].

γδ T cells in non-immunized mice also affect memory-phenotype CD4+ αβ T cells

Whereas CD8+ MP αβ T cells in non-immunized mice have been examined extensively [34, 37, 41], not much attention has been given to the less frequent CD4+ MP αβ T cells [42, 43]. In

Table 1. T-bet and EOMES expression among antigen-inexperienced memory-phenotype αβ T cells, cell frequencies.

	%	B6	B6.Vγ4/6KO	B6.Vγ4/6KO/IL-4KO
TCRβ ⁺ CD8 ⁺ CD122 ⁺ CD44 ⁺ CD49d ^{lo} CXCR3 ^{hi}	T-bet ⁺ EOMES ⁺	97.4 ± 0.3	98.4 ± 0.5	95.1 ± 0.5
	T-bet ⁻ EOMES ⁺	0.1 ± 0.0	0.0 ± 0.0	0.1 ± 0.0
	T-bet ⁻ EOMES ⁻	0.1 ± 0.0	0.0 ± 0.0	0.1 ± 0.1
TCRβ ⁺ CD4 ⁺ CD122 ⁺ CD44 ⁺ CD49d ^{lo} CXCR3 ^{hi}	T-bet ⁺ EOMES ⁻	98.6 ± 0.3	85.7 ± 3.8	95.5 ± 1.6
	T-bet ⁺ EOMES ⁺	0.9 ± 0.3	13.9 ± 3.9	4.0 ± 1.6
	T-bet ⁻ EOMES ⁺	0.0 ± 0.0	0.4 ± 0.4	0.0 ± 0.0
	T-bet ⁻ EOMES ⁻	0.4 ± 0.2	0.0 ± 0.0	0.5 ± 0.2

Data are presented as means +/- SD; groups were compared using the nonparametric ANOVA (Kruskal Wallis) test. C57BL/6 (B6) mice (n = 8), B6. Vγ4/6KO mice (n = 3), and B6. Vγ4/6KO/IL-4KO mice (n = 7)

<https://doi.org/10.1371/journal.pone.0218827.t001>

Table 2. T-bet and EOMES expression among antigen-inexperienced memory-phenotype $\alpha\beta$ T cells, cell numbers.

	numbers (x 10 ⁴)	B6	B6.V γ 4/6KO	B6.V γ 4/6KO/ IL-4KO
TCR β ⁺ CD8 ⁺ CD122 ⁺ CD44 ⁺ CD49d ^{lo} CXCR3 ^{hi}	Tbet ⁺ EOMES ⁻	3.1 ± 0.3	5.0 ± 2.2	10.9 ± 3.9
	Tbet ⁺ EOMES ⁺	138.4 ± 17.6	314.5 ± 79.2	228.8 ± 88.5
	Tbet ⁻ EOMES ⁺	0.2 ± 0.1	0.1 ± 0.1	0.3 ± 0.1
	Tbet ⁻ EOMES ⁻	0.2 ± 0.0	0.0 ± 0.0	0.2 ± 0.1
TCR β ⁺ CD4 ⁺ CD122 ⁺ CD44 ⁺ CD49d ^{lo} CXCR3 ^{hi}	Tbet ⁺ EOMES ⁻	23.5 ± 2.8	3.7 ± 0.9	25.3 ± 6.0
	Tbet ⁺ EOMES ⁺	0.4 ± 0.1	0.5 ± 0.0	1.1 ± 0.6
	Tbet ⁻ EOMES ⁺	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
	Tbet ⁻ EOMES ⁻	0.1 ± 0.1	0.0 ± 0.0	0.2 ± 0.1

Data are presented as means +/- SD; groups were compared using the nonparametric ANOVA (Kruskal Wallis) test. C57BL/6 (B6) mice (n = 8), B6.V γ 4/6KO mice (n = 3), and B6.V γ 4/6KO/IL-4KO mice (n = 7)

<https://doi.org/10.1371/journal.pone.0218827.t002>

the wt mice of this study, only 2–4% of CD4⁺ splenic $\alpha\beta$ T cells fell into this category (Fig 3A, bottom row, Fig 3C) compared to 13–15% of CD8⁺ splenic $\alpha\beta$ T cells (Fig 3A, top row, Fig 3B). In B6.TCR δ KO mice (Fig 3C, S10 Fig), the frequency of MP cells among CD4⁺ $\alpha\beta$ T cells was similarly low. In most but not all B6.TCR-V γ 1KO mice (Fig 3C, S10 Fig), MP cells among CD4⁺ $\alpha\beta$ T cells were even further diminished. In some B6.TCR-V γ 4/6KO mice, on the other hand, MP cells seemed to be slightly increased (Fig 3A, bottom row), but overall, these mice did not significantly differ from wt mice (Fig 3C). Whereas nearly all MP CD8⁺ $\alpha\beta$ T cells in the mouse strains examined were CD49d^{low} (Fig 3B), only about one half of MP CD4⁺ $\alpha\beta$ T cells displayed this “antigen-inexperienced” phenotype in wt mice, and in B6.TCR δ KO and B6.TCR-V γ 1KO mice (Fig 3C). The other MP CD4⁺ $\alpha\beta$ T cells expressed CD49d at high levels, perhaps reflecting better chances of MP CD4⁺ $\alpha\beta$ T cells in non-immunized mice (compared to MP CD8⁺ $\alpha\beta$ T cells) to experience TCR-stimulation. Quite distinctly, even fewer of these cells (~20%) were CD49d^{low} in B6.TCR-V γ 4/6KO mice (Fig 3C). In contrast, B6.TCR-V γ 4/6KO/IL-4KO mice resembled wt mice, suggesting that (when compared to MP CD8⁺ $\alpha\beta$ T cells) IL-4 might have the opposite effect on MP CD4⁺ $\alpha\beta$ T cells in non-immunized mice, and actually favor antigen-experienced cells in this population. Only about one half of the few MP CD4⁺ $\alpha\beta$ T cells that did express CD49d at low levels expressed CXCR3 highly, and most of these cells were Tbet⁺ EOMES⁻, as opposed to CXCR3^{hi} CD49d^{lo} MP CD8⁺ $\alpha\beta$ T cells, which were mostly Tbet⁺ EOMES⁺ (Tables 1 and 2).

The influence of $\gamma\delta$ T cells on developing $\alpha\beta$ T cells becomes evident late during intra-thymic maturation

In contrast to the spleen, there was no indication of lymphopenia in the thymi of $\gamma\delta$ T cell-deficient mouse strains (Fig 4, gating strategy in S11 Fig). Numbers of non-gated thymocytes and of total thymic lymphocytes (Fig 4A) were mostly unchanged or perhaps slightly higher compared to those in wt mice. We did not analyze the earliest precursors of $\alpha\beta$ T cells (CD4/CD8 double-negative thymocytes), but numbers of their immediate progeny, double-positive “immature” thymocytes were essentially unchanged. Likewise, CD4-single-positive CD24^{low} “mature” thymic lymphocytes (Fig 4B) were unchanged, and only CD8 single-positive “mature” thymic lymphocytes in B6.TCR-V γ 4/6KO mice were significantly increased relative to those in wt mice. The numeric changes in “mature” CD8 single-positive thymocytes increased the relative frequency of these cells and slightly decreased the frequency of CD4⁺ thymocytes (S12 Fig). Similarly to CD8⁺ $\alpha\beta$ T cells in the spleen, a much-increased fraction of




Fig 4. Thymocytes in $\gamma\delta$ T cell-deficient mice. A) No lymphopenia in the thymus of $\gamma\delta$ T cell-deficient mice. Total thymocytes and thymic lymphocytes (identified by their scatter profile) were counted in the mouse strains of this study (same strains as in Fig 1). Male and female mice, ages 8–12 wks were included. n equal or greater than 7 mice/group. B) No deficiency of double- or single-positive thymocytes in $\gamma\delta$ T cell-deficient mice. Double (CD4+CD8+) and single (CD4+ or CD8+) positive thymic lymphocytes were counted in the mouse strains of this study (same strains as in Fig 1). Male and female mice, ages 8–12 wks were included. n equal or greater than 7 mice/group. * $p < 0.05$. C) Comparison of CD8+ or CD4+ single-positive CD122+CD44+ TCR- β + thymocytes in C57BL/6 (wt), B6.TCR-V γ 4/6KO and B6.TCR-V γ 4/6KO/IL-4KO mice. The representative flow cytograms show an increase in the relative frequency of CD8-single-positive CD122+CD44+ TCR- β + thymocytes in B6.TCR-V γ 4/6KO mice, and a relative decrease of CD4-single-positive CD122+CD44+ TCR- β + thymocytes. Both cell-types were present at normal (wt) frequencies in B6.TCR-V γ 4/6KO/IL-4KO mice. D, E) Hierarchical comparison of mature TCR- β + thymocytes in wt and $\gamma\delta$ T cell deficient mice (same strains as in Fig 1), including first CD8- or CD4-single-positive cells among thymic lymphocytes, second CD122+CD44+ cells among either type of single-positive $\alpha\beta$ T thymocytes, third CD49d low cells among the memory phenotype single-positive $\alpha\beta$ thymocytes and fourth, CD49d high cells among memory phenotype single-positive $\alpha\beta$ thymocytes. Female and male mice ages 8–12 wks were included in the comparisons of Fig 1A. n equal or greater than 7 mice/group. * $p < 0.05$, ** $p < 0.01$.

<https://doi.org/10.1371/journal.pone.0218827.g004>

these expanded CD8 single-positive mature thymocytes in B6.TCR-V γ 4/6KO but not in B6.TCR-V γ 4/6KO/IL-4KO mice expressed the activation/memory markers CD122 and CD44 (representative cytograms in Fig 4C, top row, comparison with other mouse strains Fig 4D). Among the mouse strains tested, only the CD8+CD122+CD44+ thymocyte population in B6.TCR-V γ 4/6KO mice contained CD49d high cells at a lower frequency when compared to the wt mice, but the difference was small (Fig 4D). These data appear to reflect a limited role for IL-4 in the expansion of MP CD8-single positive thymic lymphocytes, and might indicate that AIMP CD8+ $\alpha\beta$ T cells originate in the thymus (at least in B6.TCR-V γ 4/6KO mice).

In stark contrast to the increased MP CD8+ mature thymic lymphocytes, MP CD4+ mature thymic lymphocytes were much decreased in B6.TCR-V γ 4/6KO mice when compared to wt mice (Fig 4C, bottom row, Fig 4E), a difference we had not observed with $\alpha\beta$ T cells in the spleen of B6.TCR-V γ 4/6KO mice (Fig 3C). Among these thymic lymphocytes, CD49d expression was skewed towards high levels when compared to the other mouse strains (consistent with stimulation via the TCR) (Fig 4E), far more prominently so than the pattern observed in the spleen (Fig 3). B6.TCR-V γ 4/6KO/IL-4KO mice had normal (low) frequencies of CD49d high cells among their MP CD4+ thymic lymphocytes, again linking the expression pattern to IL-4.

$\gamma\delta$ T cells appear to impact $\alpha\beta$ T cells and B cells at similar stages in their development

We had previously seen large differences of mature and immature transitional B cell populations in the spleens of non-immunized $\gamma\delta$ T cell-deficient mice (same strains as in the current study) but not of immature B cells represented by Hardy fractions [44] in bone marrow [23]. To ascertain the stage in B cell development at which $\gamma\delta$ T cells engage with B cells, we compared transitional B cells in the spleen of these mice with the small transitional-like bone marrow B cell population [45], which marks B cells about to exit from bone marrow (Fig 5). As noted before [23], numbers of mature (B220+IgM+CD93-) splenic B cells were much decreased in B6.TCR-V γ 4/6KO mice (Fig 5A). Because mature B cells in the spleen are more frequent than T cells, their loss accounts for most of the splenic lymphopenia in these mice. Total immature (B220+IgM+CD93+) splenic B cells were also decreased in these mice, and so were early (T1, B220+IgM+CD93+CD23-) and late (T2+T3, B220+IgM+CD93+CD23+) transitional B cells. B6.TCR-V γ 4/6KO/IL-4KO mice showed none of the changes in B cells seen with B6.TCR-V γ 4/6KO mice. By contrast, in bone marrow of B6.TCR-V γ 4/6KO mice, we failed to detect significant losses of immature B cells and their transitional-like subtypes (Fig 5B).




Fig 5. Changes of transitional B cells in spleen and transitional-like B cells in bone marrow of $\gamma\delta$ T cell-deficient mice. A) Comparison of splenic B cells (total number of cells/spleen) including mature (B220+IgM+CD93⁻), immature (B220+IgM+CD93⁺), transitional 1 (T1, B220+IgM+CD93+CD23⁻) and transitional 2+3 (T2+T3, B220+IgM+CD93+CD23⁺) B cells, same mouse strains as in Fig 1. B) Comparison of bone marrow B cells (total number of B cells/right femur) including immature (B220+IgM+CD93⁺), transitional 1-like (T1, B220+IgM+CD93

+CD23-) and transitional 2+3-like (T2+T3, B220+IgM+CD93+CD23+) B cells, same mouse strains as in Fig 1. Female and male mice were included in the comparisons shown in Fig 5, ages 8–13 wks, n equal or greater 8 mice/group. * $p < 0.05$, *** $p < 0.001$.

<https://doi.org/10.1371/journal.pone.0218827.g005>

Expression of the death receptor Fas can serve as a marker of activated/memory B cells [46]. We previously reported increased Fas+ germinal center B cells in the spleens of B6.TCR-V γ 4/6KO mice [18]. Here, we broadly assessed Fas receptor expression by B cells in the spleen, in order to compare frequencies of activated/memory B cells in the non-immunized $\gamma\delta$ T cell-deficient mice of this study (Fig 6). Similarly to the MP CD8+ $\alpha\beta$ T cells, Fas+ immature and mature B cells of B6.TCR-V γ 4/6KO but not B6.TCR-V γ 4/6KO/IL-4KO mice were increased in frequency although—unlike the T cells, absolute numbers of these B cells were not increased. Inversely, B6.TCR-V γ 1KO mice seemed to suffer an absolute loss of (mostly mature) Fas+ B cells. A comparison of these data with those on $\alpha\beta$ T cells suggests that with either lymphocyte-type, the $\gamma\delta$ -influence becomes manifest late during pre-immune maturation.

Discussion

The possibility that $\gamma\delta$ T cells affect the immune system prior to its responses has hardly been explored. To address this question, we compared immune cells in a set of genetic background-matched $\gamma\delta$ T cell-deficient mouse strains. Between these mouse strains, without immunization, we found differences in overall leucocyte numbers, including granulocytes and B cells [18, 23]. We also observed large differences in antibody levels [18], suggesting that $\gamma\delta$ T cells influence the production of natural antibodies [47], and more broadly influence the shape of the native immune system [48]. The current study extends this investigation to $\alpha\beta$ T cells. The results suggest that $\gamma\delta$ T cells begin to influence the development of $\alpha\beta$ T cells late during their intra-thymic maturation, thus shaping mature T cell populations. This pre-immune interaction was most noticeable with memory phenotype (MP) $\alpha\beta$ T cells, which might be important because MP $\alpha\beta$ T cells—even if they are not specific for relevant antigens—reportedly contribute more to host protection than do naïve T cells [34, 37].

Several shared features connect the current study of $\alpha\beta$ T cells in non-immunized mice to the previous work on B cells [18, 23]. First, similarly to the findings with B cells and antibodies, complete absence of $\gamma\delta$ T cells seemed to have little effect on $\alpha\beta$ T cells in non-immunized mice, whereas selective deficiencies of $\gamma\delta$ T cell-subsets were associated with large changes in $\alpha\beta$ T cells. This particular feature seems to reflect some interactive balance between functionally differentiated $\gamma\delta$ T cell-populations [18, 48–50]. Perturbation of the balance drastically changes the influence of the remaining $\gamma\delta$ T cells on other cell-types. Interestingly, the balance seems to be maintained in part through internal crosstalk between $\gamma\delta$ T cells. We believe this to be true because we previously observed that loss of certain $\gamma\delta$ T cell-types (in B6.TCR-V γ 4/6KO mice) changed population size and functional activity of the remaining $\gamma\delta$ T cells (i.e. V γ 1+ and V γ 7+ cell populations in B6.TCR-V γ 4/6KO mice) [18, 48]. Second, and again similarly to the findings with B cells, $\gamma\delta$ T cell-regulated IL-4 seems to be a critical mediator of the $\gamma\delta$ -effect on pre-immune $\alpha\beta$ T cells in the $\gamma\delta$ T cell-deficient mice. In earlier studies, we found increased numbers of IL-4-producing T cells (both $\alpha\beta$ and $\gamma\delta$ T cells) and elevated levels of circulating IL-4 in B6.TCR-V γ 4/6KO mice [18, 23], along with numerous cellular and molecular changes expected to occur with increased IL-4 activity [51–54]. These changes were not present in B6.TCR-V γ 4/6KO/IL-4KO mice. Likewise, in the current study, non-immunized B6.TCR-V γ 4/6KO but not B6.TCR-V γ 4/6KO/IL-4KO mice harbored enlarged populations of memory-phenotype (MP) CD8+ $\alpha\beta$ T cells, some of which are known to depend on IL-4 [34, 37]. In this study, we confirmed that most of the elevated IL-4 in B6.TCR-V γ 4/6KO is T cell-




Fig 6. Changes in Fas+ B cells in the spleen of $\gamma\delta$ T cell-deficient mice. Comparison of Fas-expressing B cells in $\gamma\delta$ T cell-deficient mouse strains for relative frequencies and absolute numbers (same strains as in Fig 1). B cells (B220+IgM+), mature B cells (B220+IgM+CD93-), immature B cells (B220+IgM+CD93+). n equal or greater than 9 mice/group. * $p < 0.05$, ** $p < 0.01$.

<https://doi.org/10.1371/journal.pone.0218827.g006>

derived. Moreover, in both non-immunized B6.TCR-V γ 4/6KO mice and C57BL/6 (wt) mice, $\alpha\beta$ T cells appear to be the major source of the circulating IL-4. It remains to be determined exactly how certain $\gamma\delta$ T cells control pre-immune IL-4-production by other cell-types. In any case, the effect of $\gamma\delta$ T cell-controlled IL-4 on B and T cells is mostly evident with their activated/memory states. Thus, in B6.TCR-V γ 4/6KO mice but not in B6.TCR-V γ 4/6KO/IL-4KO mice, we found both immature and mature, activated Fas+ B cells [46] to be much increased in frequency (though not in numbers) in the presence of huge losses of resting B cells. Likewise, CD122+CD44+ MP $\alpha\beta$ T cells were increased (in frequency and numbers) while other $\alpha\beta$ T cells were unchanged or lost. Secondly, the data of this study suggest that $\gamma\delta$ T cells begin to exert their influence on $\alpha\beta$ T cells already in the thymus, at a late single-positive (medullary?) stage of intra-thymic maturation. Of note, we cannot exclude an influence of $\gamma\delta$ T cells on DN thymocyte precursors of $\alpha\beta$ T cells but this seems unlikely given that we did not find significant changes in DP thymocytes. Again, this is reminiscent of their influence on B cells, which becomes evident at the transitional stages of B cell-development [23] (and this study). Similarly to the effects on developing B cells, the $\gamma\delta$ -dependent changes of developing CD8+ MP thymocytes correlate well with changes of mature CD8+ MP $\alpha\beta$ T cells.

The parallels between changes of B cells and $\alpha\beta$ T cells in the $\gamma\delta$ T cell-deficient mice suggest that similar mechanisms are at work. This is informative because unlike B cells, $\alpha\beta$ T cells can express TCR- γ and, sometimes, $-\delta$ genes, and thus potentially could be intrinsically affected by the mutations of the $\gamma\delta$ T cell-deficient mouse strains examined here. However, with B cells, non-genetic manipulations of $\gamma\delta$ T cells supported the notion of cellular interactions between $\gamma\delta$ T cells and B cells as cause for the changes in B cells and antibodies [18, 22, 23, 25]. On the basis of the findings with B cells, and because of the similarity of the $\gamma\delta$ -effects on B and $\alpha\beta$ T cells, we would expect that cellular interactions (rather than intrinsic effects of the mutations) also mediate the changes in $\alpha\beta$ T cells.

Consistently, the data imply that the $\gamma\delta$ -influence on B cells and $\alpha\beta$ T cells is mainly mediated by IL-4. IL-4 directly supports T and B cell-development, during the immune responses and before [51, 55]. Early studies with fetal thymocytes characterized IL-4 as a growth and differentiation factor for intra-thymic T cell precursors [56, 57], but inhibitory effects of IL-4 on thymocyte development were seen as well [58]. In the first IL-4 transgenic mice, overexpression of IL-4 lead to lymphopenia and death, so that transgenic IL-4 expression had to be attenuated in order to establish transgenic mouse strains [59]. Direct cytotoxicity and IL-4-induced apoptosis has been described with human B-lineage acute lymphoblastic leukemia cells as well as with a proportion of normal B cell progenitors [60]. In addition, we observed increased IL-21 production in B6.TCR-V γ 4/6KO mice but not in B6.TCR-V γ 4/6KO/IL-4KO mice, linking IL-4 and IL-21 (Kira Rubtsova, unpublished data). IL-21 can mediate growth arrest and apoptosis of certain B cells [61, 62]. Regarding T cells, a cytokine receptor consisting of IL-4R α and IL-13R α 1 (heteroreceptor, HR) was found to transduce IL-4 (and IL-13) signals that drive apoptosis of Th1 cells and skew neonatal immunity towards Th2 cells [63], and earlier in development, IL-4/IL-13 signaling through the HR can affect the potential of thymic progenitors to commit to the T cell lineage [64]. Therefore, it is conceivable that $\gamma\delta$ T cells, through their control of IL-4, promote the loss of (naïve?) B and T lymphocytes, while simultaneously protecting and expanding activated B and T cell populations.

Lastly, our data indicate that $\gamma\delta$ T cells affect CD4+ and CD8+ MP $\alpha\beta$ T cells differently. The expansion of CD8+ MP $\alpha\beta$ T cells in B6.TCR-V γ 4/6KO mice suggests that $\gamma\delta$ T cells deficient in these mice normally repress CD8+ MP $\alpha\beta$ T cells, in thymus and spleen. The repression could be direct or via a second, otherwise supportive cell-type, e.g. V γ 1+ $\gamma\delta$ T cells. The expanded CD8+ $\alpha\beta$ T cells in B6.TCR-V γ 4/6KO mice closely resemble antigen-inexperienced memory-phenotype (AIMP) CD8+ $\alpha\beta$ T cells previously described by others [34, 37]. In the

same mice, CD4⁺ (CD49d low and high) MP $\alpha\beta$ T cells were diminished, albeit only in the thymus. Here, the developing $\alpha\beta$ T cells might be normally supported by the $\gamma\delta$ T cells deficient in B6.TCR-V γ 4/6KO mice, or suppressed by some of those still present. Specifically, V γ 1⁺ $\gamma\delta$ T cells, when freed from the inhibition by V γ 4⁺ and/or V γ 6⁺ $\gamma\delta$ T cells, might prevent the development of CD4⁺ MP $\alpha\beta$ T cells. In apparent contradiction, a contraction of CD4⁺ MP $\alpha\beta$ T cells in the spleen (but not in the thymus) of B6.TCR-V γ 1KO mice suggests that V γ 1⁺ $\gamma\delta$ T cells normally support CD4⁺ MP $\alpha\beta$ T cells, at least in the periphery. A resolution of this apparent contradiction might lie in our earlier observation of functional differences between V γ 1⁺ $\gamma\delta$ T cells developing in different environments [18, 23]. Those developing in B6.TCR-V γ 4/6KO mice are type 2-biased, hence better equipped to support IL-4-dependent CD8⁺ AIMP $\alpha\beta$ T cells and perhaps more likely to inhibit type-1 IL-4-independent CD4⁺ MP $\alpha\beta$ T cells. In contrast, V γ 1⁺ $\gamma\delta$ T cells in wt mice, developing in the presence of all other $\gamma\delta$ T cells, tend to be type 1-biased, and thus may support primarily type-1 IL-4-independent CD4⁺ MP $\alpha\beta$ T cells. Further studies will be required to fully assess and delineate the influence of $\gamma\delta$ T cells on pre-immune $\alpha\beta$ T cells and B cells.

Supporting information

S1 Fig. Effect of *in vivo* treatment with anti TCR mAbs on splenic T cells in C57BL/6 (wt) mice. Adult mice were treated with i.v. injected anti TCR mAbs or saline alone, and analyzed by flow cytometry as detailed in the Methods.
(PDF)

S2 Fig. Effect of *in vivo* treatment with anti TCR mAbs on splenic T cells in B6.TCR-V γ 4/6 KO mice. Adult mice were treated with i.v. injected anti TCR mAbs or saline alone, and analyzed by flow cytometry as detailed in the Methods.
(PDF)

S3 Fig. No significant effect of *in vivo* treatment with anti TCR mAbs on splenic B cells in C57BL/6 (wt) and B6.TCR-V γ 4/6 KO mice. Adult mice were treated with i.v. injected anti TCR mAbs or saline alone, and analyzed by flow cytometry as detailed in the Methods.
(PDF)

S4 Fig. Relative frequencies of CD4⁺ and CD8⁺ $\alpha\beta$ T cells in the spleen of $\gamma\delta$ T cell-deficient mice. Female and male mice ages 8–12 wks were included in the comparisons shown (same mice as in Fig 1). *** $p < 0.001$.
(PDF)

S5 Fig. Growth of C57BL/6 mice and background-matched $\gamma\delta$ T cell deficient mice. A) Comparison of live body weight (grams) of female C57BL/6 (B6), B6.TCR δ KO, B6.TCR-V γ 1KO, B6.TCR-V γ 4/6KO, and B6.TCR-V γ 4/6KO/IL-4KO mice, at 4, 8 and 12 wks of age.
B) Same comparison as in A but with male mice.
n equal or greater 5 mice/group, except for B6.TCR-V γ 1KO mice (females: 8 wks n = 1, 12 wks n = 2, males: 12 wks n = 2) and C57BL/6 mice (males: 12 wks n = 3).
(PDF)

S6 Fig. Similar splenic lymphocyte numbers in female and male $\gamma\delta$ T cell deficient mice. Comparison of age-matched female and male mice for splenic lymphocyte numbers, including C57BL/6 (B6), B6.TCR δ KO, B6.TCR-V γ 1KO, B6.TCR-V γ 4/6KO, and B6.TCR-V γ 4/6KO/IL-4KO mice. Female and male mice ages 8–12 wks were included in the comparison shown in

S5 Fig. n equal or greater than 5 mice/group.
(PDF)

S7 Fig. Similar effect of $\gamma\delta$ T cell-deficiencies on total splenic $\alpha\beta$ T cells in female and male mice. Comparison of total numbers of TCR- β + cells in the spleen of C57BL/6 (B6), B6.TCR δ KO, B6.TCR-V γ 1KO, B6.TCR-V γ 4/6KO, and B6.TCR-V γ 4/6KO/IL-4KO mice. Female and male mice ages 8–12 wks were included in the comparison shown in [S6 Fig.](#) n equal or greater than 10 mice/group.
(PDF)

S8 Fig. Gating strategy for CD4+ and CD8+ memory-phenotype $\alpha\beta$ T cells in the spleen of $\gamma\delta$ T cell deficient mice.
(PDF)

S9 Fig. IL4R α Expression on CD4 and CD8 T cells. Splenocytes were stained and analyzed by flow cytometry as described in the Methods.
(PDF)

S10 Fig. CD4+ and CD8+ memory-phenotype $\alpha\beta$ T cells in the spleen of C57BL/6 (wt), B6.TCR δ KO and B6.TCR-V γ 1KO mice. Unlike B6.TCR-V γ 4/6KO mice, no substantial changes in the frequencies of memory-phenotype $\alpha\beta$ T cells were found in the spleens of B6.TCR δ KO and B6.TCR-V γ 1KO mice.
(PDF)

S11 Fig. Gating strategy for memory-phenotype CD4 and CD8 single-positive thymocytes.
(PDF)

S12 Fig. Relative frequencies of CD4+/CD8+, CD4+ and CD8+ thymocytes in $\gamma\delta$ T cell-deficient mice. Male and female mice, ages 8–12 wks were included (same mice as in [Fig 4](#)). n equal or greater than 7 mice/group. * $p < 0.05$, ** $p < 0.01$.
(PDF)

Acknowledgments

The authors would like to thank Drs. Laurent Gapin, Hua Huang, Mindy Miller and Habib Zaghoulani, and Harsha Krovi for providing perspective, helpful discussion, and experiments that were ultimately not included in the manuscript. We also thank Dr. Fred Finkelman for the generous gift of anti IL-4 mAbs, and Dr. Philippa Marrack for reading the manuscript and providing editorial advice.

Author Contributions

Conceptualization: Yafei Huang, Deming Sun, Richard L. Reinhardt, Rebecca L. O'Brien, Willi K. Born.

Data curation: Swati Popat Phalke, Kira Rubtsova, Andrew Getahun.

Formal analysis: Swati Popat Phalke, Kira Rubtsova, Andrew Getahun, Richard L. Reinhardt, Willi K. Born.

Funding acquisition: Willi K. Born.

Investigation: Swati Popat Phalke, Kira Rubtsova, Willi K. Born.

Methodology: Swati Popat Phalke, Yafei Huang, Rebecca L. O'Brien.

Supervision: Rebecca L. O'Brien.

Validation: Swati Popat Phalke.

Visualization: Swati Popat Phalke, Kira Rubtsova, Andrew Getahun.

Writing – original draft: Willi K. Born.

Writing – review & editing: Swati Popat Phalke, Richard L. Reinhardt, Rebecca L. O'Brien.

References

1. Parker CM, Groh V, Band H, Porcelli SA, Morita C, Fabbi M, et al. Evidence for extrathymic changes in the T cell receptor g/d repertoire. *J Exp Med.* 1990; 171:1597–612. <https://doi.org/10.1084/jem.171.5.1597> PMID: 2185330
2. Bank I, Marcu-Malina V. Quantitative peripheral blood perturbations of gammadelta T cells in human disease and their clinical implications. *Clinic Rev Allerg Immunol.* 2013; <https://doi.org/10.1007/s12016-013-8391-x> PMID: 24126758
3. Airoidi I, Bertaina A, Prigione I, Zorzoli A, Pagliara D, Cocco C, et al. Gammadelta T cell reconstitution after HLA-haploidentical hematopoietic transplantation depleted of TCR-alpha-beta+/CD19+ lymphocytes. *Blood.* 2015; 125:2349–58. <https://doi.org/10.1182/blood-2014-09-599423> PMID: 25612623
4. Kelly KA, O'Brien R, Born W. Reconstitution of SCID mice with haemopoietic precursors: a detailed analysis of gd T-cell reconstitution. *Immunology.* 1997; 91:65–72. <https://doi.org/10.1046/j.1365-2567.1997.00233.x> PMID: 9203967
5. Havran W, Allison JP. Developmentally ordered appearance of thymocytes expressing different T cell antigen receptors. *Nature.* 1988; 335:443–5. <https://doi.org/10.1038/335443a0> PMID: 2458531
6. Haas W, Pereira P, Tonegawa S. Gamma/delta T cells. *Annu Rev Immunol.* 1993; 11:637–85. <https://doi.org/10.1146/annurev.iy.11.040193.003225> PMID: 8476575
7. Born WK, Yin Z, Hahn YS, Sun D, O'Brien RL. Analysis of gamma delta T cell functions in the mouse. *J Immunol.* 2010; 184:4055–61. <https://doi.org/10.4049/jimmunol.0903679> PMID: 20368285
8. Bonneville M, O'Brien RL, Born WK. Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity. *Nature Reviews Immunology.* 2010; 10:467–78. <https://doi.org/10.1038/nri2781> PMID: 20539306
9. Huber SA, Graveline D, Newell MK, Born WK, O'Brien RL. Vg1+ T cells suppress and Vg4+ T cells promote susceptibility to coxsackievirus B3-induced myocarditis in mice. *J Immunology.* 2000; 165:4174–81.
10. O'Brien RL, Xiang Y, Huber S, Ikuta K, Born WK. Depletion of a gd T cell subset can increase host resistance to a bacterial infection. *J Immunol.* 2000; 165:6472–9.
11. Narayan K, Sylvia KE, Malhotra N, Yin CC, Martens G, Vallerskog T, et al. Intrathymic programming of effector fates in three molecularly distinct gammadelta T cell subtypes. *Nature Immunol.* 2012; published online April 1, 2012; <https://doi.org/10.1038/ni.2247> PMID: 22473038
12. Dong P, Zhang S, Cai M, Kang N, Hu Y, Cui L, et al. Global characterization of differential gene expression profiles in mouse Vgamma1+ and Vgamma4+ gammadelta T cells. *PLOS One.* 2014; 9(11): e112964. <https://doi.org/10.1371/journal.pone.0112964> PMID: 25405356
13. O'Brien RL, Lahn M, Born W, Huber SA. T cell receptor and function cosegregate in gamma-delta T cell subsets. *Chemical Immunology.* 2001; 79:1–28. PMID: 11478152
14. O'Brien RL, Roark CL, Jin N, Aydintug MK, French JD, Chain JL, et al. Gammadelta T cell receptors: functional correlations. *Immunol Rev.* 2007; 215:77–88. <https://doi.org/10.1111/j.1600-065X.2006.00477.x> PMID: 17291280
15. Melandri D, Zlatareva I, Chaleil RAG, Dart RJ, Chancellor A, Nussbaumer O, et al. The gammadelta TCR combines innate immunity with adaptive immunity by utilizing spatially distinct regions for agonist selection and antigen responsiveness. *Nature Immunology.* 2018; 19:1352–65. <https://doi.org/10.1038/s41590-018-0253-5> PMID: 30420626
16. Born W, Cady C, Jones-Carson J, Mukasa A, Lahn M, O'Brien R. Immunoregulatory functions of gammadelta T cells. *Adv Immunol.* 1999; 71:77–144. PMID: 9917911
17. Vantourout P, Hayday A. Six-of-the-best: unique contributions of gammadelta T cells to immunology. *Nature Reviews Immunology.* 2013; 13:88–100. <https://doi.org/10.1038/nri3384> PMID: 23348415
18. Huang Y, Heiser RA, Detanico TO, Getahun A, Kirchenbaum GA, Casper TL, et al. Gammadelta T cells affect IL-4 production and B-cell tolerance. *Proceedings of the National Academy of Sciences of the*

- United States of America. 2015; 112:E39–E48. <https://doi.org/10.1073/pnas.1415107111> PMID: 25535377
19. Mombaerts P, Arnoldi J, Russ F, Tonegawa S, Kaufmann SHE. Different roles of α B and γ δ T cells in immunity against an intracellular bacterial pathogen. *Nature*. 1993; 365:53–6. <https://doi.org/10.1038/365053a0> PMID: 8361537
 20. Sunaga S, Maki K, Komagata Y, Miyazaki J-I, Ikuta K. Developmentally ordered V-J recombination in mouse T cell receptor γ locus is not perturbed by targeted deletion of the Vg4 gene. *J Immunol*. 1997; 158:4223–8. PMID: 9126983
 21. Dalton JE, Pearson J, Scott P, Carding S. The interaction of gamma delta T cells with activated macrophages is a property of the Vg1 subset. *J Immunol*. 2003; 171:6488–94. <https://doi.org/10.4049/jimmunol.171.12.6488> PMID: 14662848
 22. Huang Y, Jin N, Roark CL, Aydintug MK, Wands JM, Huang H, et al. The influence of IgE-enhancing and IgE-suppressive gammadelta T cells changes with exposure to inhaled ovalbumin. *J Immunol*. 2009; 183:849–55.
 23. Huang Y, Getahun A, Heiser RA, Detanico T, Aviszus K, Kirchenbaum G, et al. Gammadelta T cells shape preimmune peripheral B cell populations. *J Immunol*. 2016; 196:217–31. <https://doi.org/10.4049/jimmunol.1501064> PMID: 26582947
 24. Egan CE, Dalton JE, Andrew EM, Smith JE, Gubbels MJ, Striepen B, et al. A requirement for the Vgamma1+ subset of peripheral gammadelta T cells in the control of the systemic growth of *Toxoplasma gondii* and infection-induced pathology. *J Immunol*. 2005; 175:8191–9. <https://doi.org/10.4049/jimmunol.175.12.8191> PMID: 16339558
 25. Huang Y, Aydintug MK, Loomis J, Macleod MK, McKee AS, Kirchenbaum G, et al. Antigen-Specific Regulation of IgE Antibodies by Non-Antigen-Specific gammadelta T Cells. *J Immunol*. 2013; 190(3):913–21. Epub 2013/01/01. <https://doi.org/10.4049/jimmunol.1202230> PMID: 23275606; PubMed Central PMCID: PMC3552125.
 26. Hahn Y-S, Taube C, Jin N, Sharp L, Wands JM, Kemal Aydintug M, et al. Different potentials of γ δ T cell subsets in regulating airway responsiveness: Vg1+ cells, but not Vg4+ cells, promote airway hyperreactivity, TH2 cytokines, and airway inflammation. *J Immunol*. 2004; 172:2894–902.
 27. Koenecke C, Chennupati V, Schmitz S, Malissen B, Forster R, Prinz I. In vivo application of mAb directed against the gammadelta TCR does not deplete but generates "invisible" gammadelta T cells. *European Journal of Immunology*. 2009; 39:372–9. <https://doi.org/10.1002/eji.200838741> PMID: 19130484
 28. Finkelman FD, Morris SC. Development of an assay to measure in vivo cytokine production in the mouse. *International Immunology*. 1999; 11:1811–8. <https://doi.org/10.1093/intimm/11.11.1811> PMID: 10545485
 29. Heilig JS, Tonegawa S. Diversity of murine gamma genes and expression in fetal and adult T lymphocytes. *Nature*. 1986; 322:836–40. <https://doi.org/10.1038/322836a0> PMID: 2943999
 30. Gerber DJ, Azuara V, Levraud J-P, Huang SY, Lembezat M-P, Pereira P. IL-4-producing γ δ T cells that express a very restricted TCR repertoire are preferentially localized in liver and spleen. *J Immunol*. 1999; 163:3076–82. PMID: 10477572
 31. Roark CL, French JD, Taylor MA, Bendele AM, Born WK, O'Brien RL. Exacerbation of collagen-induced arthritis by oligoclonal, iL-17-producing gammadelta T cells. *J Immunol*. 2007; 179:5576–83.
 32. Simonian PL, Wehrmann F, Roark CL, Born WK, O'Brien RL, Fontenot AP. gammadelta T cells protect against lung fibrosis via IL-22. *Journal of Experimental Medicine*. 2010; 207:2239–53. <https://doi.org/10.1084/jem.20100061> PMID: 20855496
 33. Haluszczak C, Akue AD, Hamilton SE, Johnson LDS, Pujanauski L, Teodorovic L, et al. The antigen-specific CD8+ T cell repertoire in unimmunized mice includes memory phenotype cells bearing markers of homeostatic expansion. *Journal of Experimental Medicine*. 2009; Epub.
 34. Jameson SC, Lee YJ, Hogquist KA. Innate memory T cells. *Adv Immunol*. 2015; 126:173–213. <https://doi.org/10.1016/bs.ai.2014.12.001> PMID: 25727290
 35. Goldrath AW, Bevan MJ. Low-affinity ligands for the TCR drive proliferation of mature CD8+ T cells in lymphopenic hosts. *Immunity*. 1999; 11:183–90. PMID: 10485653
 36. Kurzweil V, LaRoche A, Oliver PM. Increased peripheral IL-4 leads to an expanded virtual memory CD8+ population. *J Immunol*. 2014; 192:5643–51.
 37. White JT, Cross EW, Kedl RM. Antigen-inexperienced memory CD8+ T cells: where they come from and why we need them. *Nat Rev Immunol*. 2017; 17:391–400. <https://doi.org/10.1038/nri.2017.34> PMID: 28480897
 38. Renz H, Domenico J, Gelfand EW. IL-4-dependent up-regulation of IL-4 receptor expression in murine T and B cells. *J Immunol*. 1991; 146:3049–55.

39. Sosinowski T, White JT, Cross EW, Haluszczak C, Marrack P, Gapin L, et al. CD8alpha+ dendritic cell transpresentation of IL-15 to naive CD8+ T cells produces antigen-inexperienced T cells in the periphery with memory phenotype and functions. *J Immunol*. 2013; 190:1936–47. <https://doi.org/10.4049/jimmunol.1203149> PMID: 23355737
40. Akue AD, L J.Y., Jameson SC. Derivation and maintenance of virtual memory T cells. *J Immunol*. 2012; 188:2516–23. <https://doi.org/10.4049/jimmunol.1102213> PMID: 22308307
41. Goldrath AW, Bogatzki LY, Bevan MJ. Naive T cells transiently acquire a memory-like phenotype during homeostasis-driven proliferation. *Journal of Experimental Medicine*. 2000; 192:557–64. <https://doi.org/10.1084/jem.192.4.557> PMID: 10952725
42. Kawabe T, Zhu J, Sher A. Correspondence: Foreign antigen-independent memory-phenotype CD4+ T cells: a new player in innate immunity? *Nature Reviews Immunology*. 2018; <https://doi.org/10.1038/nri.2018.12> published online 26 Feb 2018.
43. Kedl RM, White JT. Correspondence: Foreign antigen-independent memory-phenotype CD4+ T cells: a new player in innate immunity? *Nature Rev Immunol*. 2018; <https://doi.org/10.1038/nri.2018.13> published online 26 Feb 2018.
44. Hardy RR, Hayakawa K. B cell development pathways. *Annu Rev Immunol*. 2001; 19:595–621. <https://doi.org/10.1146/annurev.immunol.19.1.595> PMID: 11244048
45. Chung JB, Silverman M, Monroe JG. Transitional B cells: step by step towards immune competence. *Trends in Immunol*. 2003; 24:342–8.
46. Hao Z, Duncan GS, Seagal J, Su Y-W, Hong C, Haight J, et al. Fas receptor expression in germinal-center B cells is essential for T and B lymphocyte homeostasis. *Immunity*. 2008; 29:615–27. <https://doi.org/10.1016/j.immuni.2008.07.016> PMID: 18835195
47. Born WK, Huang Y, Zeng W, Torres RM, O'Brien RL. A special connection between gammadelta T cells and natural antibodies? *Arch Immunol Ther Exp (Warsz)*. 2016; <https://doi.org/10.1007/s00005-016-0403-0>. PMID: 27235134
48. Born WK, Huang Y, Reinhardt RL, Huang H, Sun D, O'Brien RL. Gamma delta T cells and B cells. *Advances in Immunology*. 2017; 134:1–45. <https://doi.org/10.1016/bs.ai.2017.01.002> Epub Feb 14. Review. PMID: 28413020
49. Born WK, Huang Y, Jin N, Huang H, O'Brien RL. Balanced approach of gammadelta T cells to type 2 immunity. *Immunology and Cell biology*. 2010; 88:269–74. <https://doi.org/10.1038/icb.2009.105> PMID: 20101253
50. Hao J, Dong S, Xia S, He W, Jia H, Zhang S, et al. Regulatory role of Vgamma1 gammadelta T cells in tumor immunity through IL-4 production. *J Immunol*. 2011; 187:4979–86. <https://doi.org/10.4049/jimmunol.1101389> PMID: 21987661
51. Paul WE, Ohara J. B-cell stimulatory factor-1/Interleukin 4. *Annu Rev Immunol*. 1987; 5:429–59. <https://doi.org/10.1146/annurev.iy.05.040187.002241> PMID: 3297106
52. Erb K, Ruger B, von Brevern M, Ryyfel B, Schimpl A, Rivett K. Constitutive expression of interleukin (IL)-4 in vivo causes autoimmune-type disorders in mice. *Journal of Experimental Medicine*. 1997; 185:329–39. <https://doi.org/10.1084/jem.185.2.329> PMID: 9016881
53. Rudge EU, Cutler AJ, Pritchard NR, Smith KGC. Interleukin 4 reduces expression of inhibitory receptors on B cells and abolishes CD22 and FcgammaRII-mediated B cell suppression. *Journal of Experimental Medicine*. 2002; 195:1079–85. <https://doi.org/10.1084/jem.20011435> PMID: 11956299
54. Hidaka T, Kitani A, Hara M, Harigai M, Suzuki K, Kawaguchi Y, et al. IL-4 down-regulates the surface expression of CD5 on B cells and inhibits spontaneous immunoglobulin and IgM-rheumatoid factor production in patients with rheumatoid arthritis. *Clin Exp Immunol*. 1992; 89:223–9. <https://doi.org/10.1111/j.1365-2249.1992.tb06936.x> PMID: 1379131
55. Paul WE. Interleukin-4: a prototypic immunoregulatory lymphokine. *Blood*. 1991; 77:1859–65. PMID: 2018830
56. Palacios R, Sideras P, von Boehmer H. Recombinant interleukin 4/BSF-1 promotes growth and differentiation of intrathymic T cell precursors from fetal mice in vitro. *EMBO Journal*. 1987; 6:91–5. PMID: 3495433
57. Zlotnik A, Ransom J, Frank G, Fischer M, Howard M. Interleukin 4 is a growth factor for activated thymocytes: possible role in T-cell ontogeny. *Proceedings of the National Academy of Sciences of the United States of America*. 1987; 84:3856–60. <https://doi.org/10.1073/pnas.84.11.3856> PMID: 3495799
58. Plum J, De Smedt M, Leclercq G, Tison B. Inhibitory effect of murine recombinant IL-4 on thymocyte development in fetal thymus organ cultures. *J Immunol*. 1990; 145:1066–73. PMID: 2116475
59. Tepper RI, Levinson DA, Stanger BZ, Campos-Torres J, Abbas AK, Leder P. IL-4 induces allergic-like inflammatory disease and alters T cell development in transgenic mice. *Cell*. 1990; 62:457–67. PMID: 2116236

60. Manabe A, Coustan-Smith E, Kumagai M, Behm FG, Raimondi SC, Pui C-H, et al. Interleukin-4 induces programmed cell death (apoptosis) in cases of high-risk acute lymphoblastic leukemia. *Blood*. 1994;1731–7.
61. Jin H, Carrio R, Yu A, Malek TR. Distinct activation signals determine whether IL-21 induces B cell costimulation, growth arrest, or Bim-dependent apoptosis. *J Immunol*. 2004; 173:657–65.
62. Tortola L, Yadava K, Bachmann MF, Muller C, Kisielow J, Kopf M. IL-21 induces death of marginal zone B cells during chronic inflammation. *Blood*. 2010; 116:5200–7. <https://doi.org/10.1182/blood-2010-05-284547> PMID: 20826721
63. Li L, Lee H-H, Bell JJ, Gregg RK, Ellis JS, Gessner A, et al. IL-4 utilizes an alternative receptor to drive apoptosis of Th1 cells and skews neonatal immunity toward Th2. *Immunity*. 2004; 20:429–40. PMID: 15084272
64. Barik S, Miller MM, Cattin-Roy AN, Ukah TK, Chen W, Zaghouani H. IL-4/IL-13 signaling inhibits the potential of early thymic progenitors to commit to the T cell lineage. *J Immunol*. 2017; 199:2767–76. <https://doi.org/10.4049/jimmunol.1700498> PMID: 28893952