

## Distribution and evaluation of bone and soft tissue tumors operated in a tertiary care center

Recep Öztürk<sup>a,\*</sup>, Şefik Murat Arıkan<sup>b</sup>, Emin Kürşat Bulut<sup>a</sup>, Ahmet Fevzi Kekeç<sup>c</sup>, Fener Çelebi<sup>d</sup>, Bedii Şafak Güngör<sup>a</sup>

<sup>a</sup> Department of Orthopaedics and Traumatology, Dr Abdurrahman Yurtaslan Ankara Oncology Training and Research Hospital, Ankara, Turkey

<sup>b</sup> Department of Orthopaedics and Traumatology, Gazi University Hospital, Ankara, Turkey

<sup>c</sup> Department of Orthopaedics and Traumatology, Necmettin Erbakan University Hospital, Konya, Turkey

<sup>d</sup> Department of Orthopaedics and Traumatology, Memorial Dicle Hospital, Diyarbakır, Turkey

### ARTICLE INFO

#### Article history:

Received 27 November 2016

Received in revised form

23 December 2017

Accepted 17 March 2019

Available online 11 April 2019

#### Keywords:

Epidemiological features

Epidemiology

Cancer

Turkey

Malignant tumors

Musculoskeletal tumors

### ABSTRACT

**Objective:** The aim of this study was to retrospectively evaluate the patients who were operated in our orthopedics and traumatology clinic with the suspicion of bone and soft tissue tumors.

**Methods:** A total of 3133 patients (1146 (46.5%) female and 1318 (53.5%) male) who presented to our tertiary clinic from different regions of Turkey between January 2002 and July 2013 with the presumed diagnosis of bone and soft tissue tumors were analyzed according to age, gender, bone/soft tissue localization, tumoral localization, histopathological diagnosis, tumor size and incidence.

**Results:** Of all operated patients, 2464 (78%) were diagnosed with tumor, while non-tumoral causes were found in 669 (22%) patients. Of the cases diagnosed with tumor, 1139 were bone localized, 1004 soft tissue localized, and 321 metastasis. The most common benign bone tumors were osteochondroma (130, 20%), enchondroma (96, 15%), and simple bone cysts (90, 14%), while the most common malignant bone tumors were osteosarcoma (241, 44%), ewing's sarcoma (89, 16%), and chondrosarcoma (77, 14%); respectively. The most common benign soft tissue tumors were lipoma (141, 22%), giant cell tumors (108, 16%) and ganglion (107, 16%), while the most common malignant soft tissue tumors were liposarcoma (55, 16%), synovial sarcoma (53, 16%) and malignant mesenchymal tumors (45, 13%); respectively.

**Conclusion:** Musculoskeletal tumors are rare, but descriptive data in any region are important in order to reduce mortality and improve treatment. No significant difference was found between the data of our hospital regarding epidemiology of the musculoskeletal system tumors and those from the other regions around the world.

**Level of evidence:** Level IV, Therapeutic study.

© 2019 Turkish Association of Orthopaedics and Traumatology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Introduction

Musculoskeletal system tumors are rare, and these tumors must be kept in mind in differential diagnosis in patients with suspected and persistent symptoms.<sup>1</sup> The algorithm should always follow the pattern of medical history, physical examination, radiological investigations, systemic screening, and histopathological diagnosis.<sup>1,2</sup>

The distribution of musculoskeletal system tumors may show regional variations. In history and physical examination; under

standing the region specific age, gender, localization and incidence of tumors is helpful for the diagnosis.

In this study, approximately 11,5-year data of the orthopedics and traumatology of our hospital which is a tertiary center delivering healthcare service to patients who are referred from different provinces of Turkey, and especially from central Anatolia and eastern of Turkey were analyzed. In addition, a brief literature screening was carried out on the studies conducted about incidence and distribution of bone and soft tissue localized tumors in Turkey.

### Patients & methods

This study includes data from January 2002, the time since electronic archive system available, to July 2013. Pathological

\* Corresponding author. Polatlı Public Hospital, Ankara, Turkey. Tel.: +90 5054634794.

E-mail address: [ozturk\\_recep@windowslive.com](mailto:ozturk_recep@windowslive.com) (R. Öztürk).

Peer review under responsibility of Turkish Association of Orthopaedics and Traumatology.

outcomes of our orthopedics and traumatology clinic were obtained from the hospital registry database. Therefore, these data do not include patients with only radiological diagnosis. All patients' age, gender, date of operation, tumor localization and tumor size were analyzed. Patients were first divided into two groups as tumoral diseases and non-tumoral causes. Tumoral diseases were then further divided as primary tumors and metastatic tumors. The primary tumors were classified and examined in separate groups according to localization in bones and soft tissues, and benign and malignant status.

All different tumors and non-tumoral causes were tabulated and figured according to the incidence, localization and age ranges (Tables 1–10 and Figs. 1–3).

All statistical analyses were performed using SPSS 22.0 statistical software (SPSS, Chicaco, IL, USA). Descriptive statistics are expressed as mean, frequency and percentage.

## Results

In histopathological examination of the 3133 patients, 2464 were diagnosed with tumor. Of all patients, 1146 (46.5%) were female and 1318 (53.5%) were male.

Of the tumoral lesions, 1496 (47.7%) were localized in the low extremity involving hips and distal, 771 (24.6%) in the upper extremity with shoulder joint and distal, 154 (4.9%) in the pelvic region and pelvic bones, and 43 (1.4%) in the thoraco-lumbar-sacral region and its bones.

Of all cases, 1178 (37.5%) were primary bone tumors, 965 (30.8%) primary soft tissue tumors, and 321 (10.2%) metastasis. Whereas 669 (21.3%) cases had non-tumoral causes. Excluding the non-tumoral causes, 1282 (52%) of bone and soft tissue tumors were benign and 1182 (48%) were malignant.

### Bone tumors

The mean age was found as 28.7 years in 1182 patients with primary bone tumors. Of these cases, 662 (58%) were seen in males and 477 (42%) in females. The tumors were localized in lower

**Table 1**  
Benign bone tumors.

|  | Number | Percentage |
|--|--------|------------|
| Aneurysmal bone cyst                               | 58     | 9.0        |
| Benign fibrous histiocytoma                        | 5      | 0.7        |
| Bizarre parosteal osteochondromatous proliferation | 1      | 0.1        |
| Bone island  | 1      | 0.1        |
| Chondroblastoma                                    | 30     | 4.6        |
| Chondromyxoid fibroma                              | 7      | 1.0        |
| Desmoid tumor (bone)                               | 3      | 0.4        |
| Enchondroma  | 96     | 15.0       |
| Eosinophilic granuloma                             | 13     | 2.0        |
| Fibroma (bone)                                     | 2      | 0.3        |
| Fibrous dysplasia                                  | 49     | 7.6        |
| Giant cell tumor (bone)                            | 74     | 11.5       |
| Hemangioendotelyoma                                | 2      | 0.3        |
| Haemangiopericytoma                                | 1      | 0.1        |
| Intraosseous ganglion                              | 4      | 0.6        |
| Intraosseous hemangioma                            | 4      | 0.6        |
| Intraosseous lipoma                                | 8      | 1.2        |
| Non-ossifying fibroma                              | 30     | 4.6        |
| Osteoblastoma                                      | 4      | 0.6        |
| Osteoid osteoma                                    | 26     | 4.0        |
| Osteochondroma                                     | 130    | 20.3       |
| Periosteal chondroma                               | 1      | 0.1        |
| Schwannoma (bone)                                  | 1      | 0.1        |
| Simple bone cyst                                   | 90     | 14.0       |

**Table 2**  
Malignant bone tumors.

|                               | Number | Percentage |
|-------------------------------|--------|------------|
| Adamantinoma                  | 2      | 0.3        |
| Angiosarcoma (bone)           | 4      | 0.7        |
| Chondrosarcoma                | 77     | 14.3       |
| Chordoma                      | 7      | 1.3        |
| Epithelioid sarcoma           | 2      | 0.3        |
| Ewing's sarcoma               | 89     | 16.5       |
| Fibrosarcoma                  | 2      | 0.3        |
| Leiomyosarcoma (bone)         | 6      | 1.1        |
| Lymphoma                      | 20     | 3.7        |
| Leukemia                      | 3      | 0.5        |
| Malignant giant cell tumor    | 2      | 0.3        |
| Malignant mesenchymal tumor   | 22     | 4.0        |
| Osteosarcoma                  | 241    | 44.7       |
| Parachordoma                  | 1      | 0.1        |
| Plasmacytoma/multiple myeloma | 49     | 9.1        |
| Pleomorphic sarcoma           | 11     | 2.0        |

extremity in 714, upper extremity in 307, pelvis in 92, and vertebral region in 26 patients.

The mean age was found as 26.5 and 31.1 in patients with benign and malignant tumors; respectively.

### Benign bone tumors

Of the primary benign bone tumors, 194 (30.3%) were localized in the femur, 109 (17.0%) in the humerus, and 98 (15.3%) in the tibia (Table 7).

**Table 3**  
Benign soft tissue tumors.

|   | Number | Percentage |
|---|--------|------------|
| Arterio-venous malformation                   | 4      | 0.6        |
| Baker's cyst                                  | 6      | 0.9        |
| Benign fibrous histiocytoma                   | 2      | 0.3        |
| Calcinosis                                    | 1      | 0.1        |
| Chondromyxoid fibroma (soft tissue)           | 1      | 0.1        |
| Desmoid tumor                                 | 11     | 1.7        |
| Elastofibroma dorsi                           | 2      | 0.3        |
| Epidermal inclusion cyst                      | 22     | 3.4        |
| Fibroma                                       | 14     | 2.2        |
| Ganglion cyst                                 | 107    | 16.9       |
| Giant cell tumor (soft tissue)                | 71     | 11.2       |
| Glomus tumor                                  | 3      | 0.4        |
| Hamartoma                                     | 2      | 0.3        |
| Hemangioendotelyoma                           | 1      | 0.1        |
| Hemangioma                                    | 103    | 16.2       |
| Hemangiopericytoma                            | 4      | 0.6        |
| Inflammatory myofibroblastic tumor            | 2      | 0.3        |
| Leiomyoma                                     | 1      | 0.1        |
| Lymphangioma                                  | 6      | 0.9        |
| Lipom   | 141    | 22.3       |
| Myxoid tumor                                  | 1      | 0.1        |
| Myoma   | 1      | 0.1        |
| Myxoma  | 10     | 1.5        |
| Neurofibroma                                  | 12     | 1.8        |
| Neuroma                                       | 3      | 0.4        |
| Ossification fibromyxoid tumor                | 1      | 0.1        |
| Palmar/plantar fibromatosis                   | 4      | 0.6        |
| Papilloma                                     | 2      | 0.3        |
| Pilomatricoma                                 | 4      | 0.6        |
| Pyogenic granuloma                            | 1      | 0.1        |
| Reactive follicular hyperplasia (Kimura dis.) | 1      | 0.1        |
| Schwannoma                                    | 28     | 4.4        |
| Synovial chondromatosis                       | 5      | 0.7        |
| Synovial cyst                                 | 6      | 0.9        |
| Villonodular synovitis                        | 48     | 7.5        |
| Xantoma                                       | 1      | 0.1        |

**Table 4**  
Malignant soft tissue tumors.

|   | Number | Percentage |
|---|--------|------------|
| Angiosarcoma                            | 2      | 0.6        |
| Clear cell sarcoma                      | 3      | 0.9        |
| Dermatofibrosarcoma protuberans         | 3      | 0.9        |
| Epitheloid sarcoma                      | 3      | 0.9        |
| Extraskeletal chondrosarcoma            | 7      | 2.1        |
| Extraskeletal Ewing's sarcoma           | 11     | 3.3        |
| Extraskeletal osteosarcoma              | 7      | 2.1        |
| Fibrosarcoma                            | 4      | 1.2        |
| Histiocytic sarcoma                     | 1      | 0.3        |
| Kaposi sarcoma                          | 1      | 0.3        |
| Leiomyosarcoma                          | 13     | 4.0        |
| Lymphoma                                | 15     | 4.6        |
| Liposarcoma                             | 55     | 16.9       |
| Malignant mesenchymal tumor             | 45     | 13.8       |
| Malignant melanoma                      | 9      | 2.7        |
| Malignant peripheral nerve sheath tumor | 22     | 6.7        |
| Plasmacytoma (soft tissue)              | 1      | 0.3        |
| Pleomorphic sarcoma                     | 35     | 10.7       |
| Rhabdomyosarcoma                        | 16     | 4.9        |
| Spindle cell sarcoma                    | 3      | 0.9        |
| Squamous cell carcinoma                 | 15     | 4.6        |
| Synovial sarcoma                        | 53     | 16.3       |

The mean age of the patients with osteochondroma, which was the most common benign bone tumor was found 24.1 (6–69) years. The most common localizations were femur distal (38, 29%), tibia proximal (16, 12%) and humerus proximal (15, 11%); respectively.

#### Malignant bone tumors

Of the primary malignant bone tumors, 214 (x%) were localized in the femur, 86 (x%) in the tibia, and 69 (x%) in the humerus (Table 7).

The mean age of the patients with osteosarcoma, which was the most common primary malignant bone tumor was found 22.4 (6–79) years. The most common localizations were femur distal (109, 45%), tibia proximal (48, 19%), and humerus proximal (19, 7%), respectively.

#### Soft tissue tumors

The mean age of 965 patients with primary soft tissue tumors was 40.7 years. Of these cases, 453 (47%) were seen in male and 512 (53%) were seen in female patients. The tumors were localized in the lower extremity in 606 (60,3%), in the upper extremity in 376

**Table 5**  
Metastatic tumors.

|                      | Number | Percentage |
|----------------------|--------|------------|
| Breast               | 59     | 18.3       |
| Bladder              | 10     | 3.1        |
| Endometrium          | 4      | 1.2        |
| Gastrointestinal     | 14     | 4.3        |
| Kidney               | 48     | 14.9       |
| Lungs                | 57     | 17.7       |
| Lymphoma             | 2      | 0.6        |
| Medulloblastoma      | 1      | 0.3        |
| Mesothelioma         | 1      | 0.3        |
| Neuroendocrine tumor | 2      | 0.6        |
| Ovary                | 2      | 0.6        |
| Pancreas             | 3      | 0.9        |
| Primer unknown       | 89     | 27.7       |
| Prostate             | 15     | 4.6        |
| Thyroid              | 12     | 3.7        |
| Urinary system       | 2      | 0.6        |

**Table 6**  
Non-tumor causes.

|                           | Number | Percentage |
|---------------------------|--------|------------|
| Bursitis                  | 22     | 3.2        |
| Callus tissue             | 15     | 2.2        |
| Dystrophic calcification  | 1      | 0.1        |
| Fat necrosis              | 12     | 1.7        |
| Gout tofus                | 6      | 0.8        |
| Hematoma                  | 3      | 0.4        |
| Hydatid cyst              | 6      | 0.8        |
| Inflammatory changes      | 3      | 0.4        |
| Inflammatory event        | 155    | 22.8       |
| Inflammatory pseudotumor  | 1      | 0.1        |
| Lymphadenitis             | 2      | 0.2        |
| Metabolic disease         | 2      | 0.2        |
| Myositis                  | 4      | 0.5        |
| Myositis ossificans       | 9      | 1.3        |
| Nodular fasciitis         | 6      | 0.8        |
| Normal tissue             | 116    | 17.1       |
| Osteoarthritis            | 7      | 1.0        |
| Osteochondritis dissecans | 2      | 0.2        |
| Osteomyelitis             | 64     | 9.4        |
| Osteonecrosis             | 18     | 2.6        |
| Other                     | 79     | 11.6       |
| Paget's disease           | 4      | 0.5        |
| Synovial hypertrophy      | 16     | 2.3        |
| Synovitis                 | 121    | 17.8       |
| Tuberculosis              | 4      | 0.5        |

**Table 7**  
Placement of bone tumors.

|                 | Number | Percentage |
|-----------------|--------|------------|
| Clavicle        | 18     | 1.5        |
| Femur diaphysis | 29     | 2.5        |
| Femur distal    | 246    | 21.2       |
| Femur proximal  | 131    | 11.3       |
| Fibula          | 68     | 5.8        |
| Foot - ankle    | 54     | 4.6        |
| Hand -wrist     | 46     | 3.9        |
| Humerus         | 178    | 15.4       |
| Pelvic bones    | 92     | 7.9        |
| Radius          | 21     | 1.8        |
| Scapula         | 29     | 2.5        |
| Tibia diaphysis | 16     | 1.3        |
| Tibia distal    | 40     | 3.4        |
| Tibia proximal  | 146    | 12.3       |
| Ulna            | 15     | 1.2        |
| Vertebra        | 26     | 2.2        |

**Table 8**  
Placement of soft tissue tumors.

|                  | Number | Percentage |
|------------------|--------|------------|
| Ankle            | 34     | 3.5        |
| Arm              | 58     | 6.0        |
| Cruris           | 82     | 8.5        |
| Elbow            | 29     | 3.0        |
| Foot             | 79     | 8.2        |
| Forearm          | 49     | 5.1        |
| Hand             | 130    | 13.5       |
| Hip              | 43     | 4.4        |
| Inginal          | 6      | 0.6        |
| Knee             | 99     | 10.3       |
| Pelvis           | 4      | 0.4        |
| Shoulder         | 48     | 5.0        |
| Thigh            | 250    | 26.0       |
| Vertebral region | 3      | 0.3        |
| Wrist            | 51     | 5.3        |

**Table 9**  
Placement of metastatic tumors.

|                   | Number | Percentage |
|-------------------|--------|------------|
| Clavicle          | 4      | 1.2        |
| Cruris            | 1      | 0.3        |
| Femur diaphysis   | 8      | 2.4        |
| Femur distal      | 20     | 6.2        |
| Femur proximal    | 121    | 37.6       |
| Fibula distal     | 1      | 0.3        |
| Fibula proximal   | 2      | 0.6        |
| Foot              | 2      | 0.6        |
| Forearm-radius    | 4      | 1.2        |
| Gluteal           | 2      | 0.6        |
| Hand              | 4      | 15.5       |
| Humerus diaphysis | 5      | 1.5        |
| Humerus distal    | 5      | 1.5        |
| Humerus proximal  | 50     | 15.5       |
| Knee              | 1      | 0.3        |
| Pelvis            | 45     | 14.0       |
| Scapula           | 9      | 2.8        |
| Shoulder          | 7      | 2.1        |
| Scapula           | 9      | 2.8        |
| Thigh             | 9      | 2.8        |
| Tibia             | 9      | 2.8        |
| Vertebral region  | 12     | 3.7        |

(37,4%), in the pelvis in 17 (1,6%), and in the vertebral region (0,5%) in 5 patients.

The mean age was found as 38.0 (1–86) and 45.3 (2–87) years in patients with benign and malignant soft tissue tumors; respectively.

#### Benign soft tissue tumors

Of the primary benign soft tissue tumors, 171 (27,0%) were localized in the hand-wrist, 101 (15,9%) in the thigh, and 89 (14,0%) in the foot-ankle (Table 8).

The mean age of patients with lipoma, which was the most common benign soft tissue tumor was found as 44.5 (1–81) years. The most common localizations were thigh (32, 22,6%), shoulder (23, 16,3%), and arm (22, 15,6%).

**Table 10**  
Location of non-tumoral causes.

|                  | Number | Percentage |
|------------------|--------|------------|
| Ankle            | 10     | 1.4        |
| Arm              | 13     | 1.9        |
| Clavicle         | 9      | 1.3        |
| Cruris           | 14     | 2.0        |
| Elbow            | 24     | 3.5        |
| Femur            | 124    | 18.2       |
| Fibula           | 8      | 1.1        |
| Foot             | 44     | 6.4        |
| Forearm          | 6      | 0.8        |
| Hand             | 27     | 3.9        |
| Hip              | 16     | 2.3        |
| Humerus          | 40     | 5.8        |
| Knee             | 149    | 21.9       |
| Pelvis           | 47     | 6.9        |
| Radius           | 5      | 0.7        |
| Scapula          | 4      | 0.5        |
| Shoulder         | 8      | 1.1        |
| Thigh            | 43     | 6.3        |
| Tibia            | 60     | 8.8        |
| Ulna             | 4      | 0.5        |
| Vertebral region | 17     | 2.4        |
| Wrist            | 6      | 0.8        |

#### Malignant soft tissue tumors

Of the primary malignant soft tissue tumors, 149 (45,9%) were localized in the thigh, 39 (12,0%) in the cruris, and 24 (7,4%) in the foot-ankle (Table 8).

The mean age of patients with liposarcoma, which was the most common malignant soft tissue tumor was found as 51.4 (24–80) years. The most common localizations were the thigh (33, x%), gluteal (6, x%), and cruris (5, x%).

#### Metastases and non-tumoral causes

The mean age of 321 patients with metastatic disease was found as 57.4 (4–85). Of these cases, 187 were seen in male and 134 in female patients.

Of the metastatic tumors, 59 (18,3%) were originated from the breast, and 57 (17,7%) from the lung, while the primary of 89 (27,7%) tumors was unknown (Table 5).

The mean age of 677 patients with non-tumoral causes was found as 44.4 (1–96) years. Of these cases, 327 were seen in male and 351 in female patients.

#### Discussion

Understanding the region specific age, gender, localization and incidence of tumors is helpful for the diagnosis, and the distribution of these tumors may differ. Therefore, it is necessary to know the distribution and features of bone and soft tissue tumors in Turkey.

Although activities on cancer registry have become concentrated in Turkey recently, there is no sufficient resource including a large series that show the incidence and diagnostic distribution of bone and soft tissue tumors.<sup>3</sup>

There are many regional epidemiological study records in the literature.<sup>4,5</sup> There are limited studies from Turkey to demonstrate the distribution of musculoskeletal system tumors. Whereas some of these studies include local hospital data,<sup>6–8</sup> the others are in form of the analysis of the cases evaluated in tumor councils.<sup>9</sup> In addition, there are also studies of more specific patient groups such as childhood tumors and bone tumors, vs.<sup>10–12</sup>

In 1993, Gür et al published 11-year data of Ankara Gülhane Hospital including 420 cases, and reported that all data were consistent with the classical literature information and no any regional difference was found.<sup>6</sup> In 2000, Kösem et al reported 6-year data of bone and soft tissue tumors of 250 from the region of Van.<sup>7</sup> Differently, they reported that unlike obvious predominance of the lower extremity in the classical information, they found an equal distribution in the localization of soft tissue tumors.

In 2005, Solakoğlu et al published their 10-year data about age, gender and tumor localization of 937 cases with only bone tumors, and unlike classical literature they found giant cell tumor as the most common bone tumor in adults. Whereas Güngör et al published evaluation of 200 cases with only childhood benign tumors in 2008.<sup>10,11</sup>

In 2011, Yüçetürk et al published 20-year data of the Ege University musculoskeletal tumor council including 5658 cases.<sup>9</sup> In that study, malignant musculoskeletal tumors accounted for 40% of all tumors. The authors reported that these high data were not resulted from the high regional incidence of these tumors, and was caused by the was resulted from the University tumor council being one of the limited reference centers. In our study, the ratio of malignant tumors was found even higher and was 48%. We attributed this high incidence to that our study did not include the population of benign tumors, which were radiologically diagnosed and followed-up with non-surgical methods, and to the gradually




Fig. 1. Age distribution of bone tumors (horizontal column: ages, vertical column: numbers).




Fig. 2. Age distribution of soft tissue tumors (horizontal column: ages, vertical column: numbers).




Fig. 3. Age distribution of metastatic tumors (horizontal column: ages, vertical column: numbers).

increased number of centers where benign tumors are followed-up and treated.

In 2014, Dabak et al published 25-year data of bone and soft tissue tumors from the central Blacksea region which contained 1925 cases.<sup>8</sup> The authors reported that, soft tissue tumors were more common among women, and the ages with the benign and malignant soft tissue tumors encountered were 10 years older than the literature, and that the most common localization of soft tissue tumors was the wrist. In addition, since their

study included only the cases that they performed surgery, the tumors which treatment is often non-surgical were less frequent. Similarly to the literature, in our study the most common localization of soft tissue tumors was found as the thigh.

In the study by Sevimli in 2017 which examined musculoskeletal tumors in Malatya province and surrounding (6 year, 710 cases), the most common malignant bone tumor was found as chondrosarcoma.<sup>13</sup> Whereas in the present study, the most

common malignant bone tumor was osteosarcoma, consistently with the literature.<sup>4,5,7,9,11</sup>

The most common benign bone tumors are osteochondromes and the most common location was proximal humerus and around the knee.<sup>1</sup> In our study, as well as many other studies, was consistent with this. Differently, Solakoglu et al found that osteochondroma was the most common tumor in children, giant cell tumor was the most common tumor in adults; Yucetürk et al found that the enchondroma was the most common tumor.<sup>6,8,11,14</sup> Actually, it is known that the number of osteochondromas occur much more from indicated, whereas due to require less operation, osteochondromas were seen less in studies.<sup>1,14</sup>

Whereas the most common soft tissue sarcomas have been reported as pleomorphic sarcoma and liposarcoma, in our study the most common tumors were found as liposarcoma, synovial sarcoma and malign mesenchymal tumor. Pleomorphic sarcoma ranked four. We think that this difference was resulted from the lack of a subclassification in patients with the pathological outcome of malignant mesenchymal tumor.

In our study, the most common localizations of metastasis were found as the femur proximal, humerus proximal, and pelvis. Although the most common localization of metastasis has been reported as the vertebra in the literature, less common incidence of this region in our study could be explained by that these patients can be treated in orthopedics and neurosurgery clinics of our hospital, and this study included only the orthopedics archive and that most vertebral metastases could be treated with non-operative methods.

When the relationship between tumoral diseases and age is examined; both benign and malignant bone tumors are most commonly seen between 11 and 20 years of age, and our results are consistent with this information (Fig. 1). For soft tissue tumors, a narrow age range with an evidently high incidence of tumors can not be said as in bone tumors.<sup>8,10,15</sup> It can be said that, in our study benign soft tissue tumors were more common in 11–60 age range, and malignant soft tissue tumors in 21–60 age range (Fig. 2). In their study, Dabak et al found the most common age range of metastatic tumors as 51–60 years. In our study the most common age range was found as 61–70 years followed by 51–60 years with a little difference (Fig. 3).

Similar to other studies, bone tumors were more common in male patients (58%), while soft tissue tumors were more common in female patients (53%).<sup>8,10</sup> However, there are studies reported a higher incidence of soft tissue tumors among men in Turkey.<sup>9,12</sup>

The most common metastatic bone tumors are originated from the breast and prostate followed by lungs, kidneys and thyroid.<sup>16</sup> In our study, and in the study by Dabak et al the most common diagnoses were the tumors with unknown primary, and breast and

lung localized tumors. The possible reason for lower incidence of prostate ca metastasis in these studies is that, the mentioned studies included the patients who underwent operation and prostate ca requires less surgery due to blastic metastases.

In conclusion; musculoskeletal tumors are rare, but descriptive data in any region are important in order to reduce mortality and improve treatment. No significant difference was found between the data of our hospital regarding epidemiology of the musculoskeletal system tumors and those from the other regions around the world. Further multicenter studies are needed in order to reach the information, which better reflects the demographics of musculoskeletal tumors in Turkey.

## References

- Ozturk R. Kemik ve yumuşak doku Tümörleri. In: Atay T, ed. *Ortopedi ve spor Yaralanmaları Asistan Kitabı*. Ankara, Derman Tıbbi Yayıncılık; 2015:635–704. <https://doi.org/10.4328/DERMAN.3774>.
- Yıldız Y, Sağlık Y. Kemik ve yumuşak doku tümörlerinde biopsi. *TOTBİD dergisi*. 2003;2(1-2):37–43.
- Türkiye Halk Sağlığı Kurumu. Türkiye Kanser Kontrol Programı. Available from: [https://hsgm.saglik.gov.tr/depo/birimler/kanser-db/yayinlar/raporlar/Ulusal\\_Kanser\\_Kontrol\\_Plani\\_2013\\_2018.pdf](https://hsgm.saglik.gov.tr/depo/birimler/kanser-db/yayinlar/raporlar/Ulusal_Kanser_Kontrol_Plani_2013_2018.pdf). Accessed December 31, 2016.
- Solooki S, Vosoughi AR, Masoomi V. Epidemiology of musculoskeletal tumors in Shiraz, south of Iran. *Indian J Med Paediatr Oncol*. 2011;32(4):187–191. <https://doi.org/10.4103/0971-5851.95138>.
- Bergovec M, Kubat O, Smerdelj M, Seiwerth S, Bonevski A, Orlic D. Epidemiology of musculoskeletal tumors in a national referral orthopedic department. A study of 3482 case. *Cancer Epidemiol*. 2015;39(3):298–302. <https://doi.org/10.1016/j.canep.2015.01.015>.
- Gür E, Başbozkurt M, Ateşalp AS, Kırdemir V, Baydar ML, Erler K. Kemik ve yumuşak doku tümörlerinin 11 yıllık retrospektif analizi. *Acta Orthop Traumatol Turc*. 1993;27(2):82–86.
- Kosem M, Bayram I. Van Yöresindeki Ekstremitelerde Kemik ve Yumuşak Doku Tümörlerinin Dağılımı ve Değerlendirilmesi. *Van Med J*. 2001;8(3):93–96.
- Dabak N, Çıraklı A, Gülman B, Selçuk MB, Barış S. Distribution and evaluation of bone and soft tissue tumors in the middle Black Sea Region. *Acta Orthop Traumatol Turc*. 2014;48(1):17–24. <https://doi.org/10.3944/AOTT.2014.3013>.
- Yucetürk G, Sabah D, Keçeci B, Kara AD, Yalçınkaya S. Prevalence of bone and soft tissue tumors. *Acta Orthop Traumatol Turc*. 2011;45(3):135–143. <https://doi.org/10.3944/AOTT.2011.2504>.
- Solakoglu D, Benzer E. Distribution of bone tumors according to age, sex and tumor site. *Acta Oncol Turc*. 2005;38(1):38–43.
- Güngör Ş, Karakoç Y, Arıkan M, Metin E, Gürler D. Demographic features of childhood benign bone tumors and evaluation of clinical approach. *Acta Oncol Turc*. 2008;41(1):7–13.
- Ozturk R, Arıkan ŞM, Şimşek MA, Özanağan E, Güngör BŞ. Management of solitary fibrous tumors localized in extremity: case series and a review of the literature. *Eklemler Hastalıkları Cerrahisi*. 2017;28(2):121–127. <https://doi.org/10.5606/ehc.2017.52092>.
- Sevimli R. Distribution and evaluation of primary bone and soft tissue tumors admitted from Malatya province and surrounding provinces. *Med Sci*. 2017;6(3):546–550. <https://doi.org/10.5455/medscience.2017.06.8619>.
- Rodop O. Osteokondroma. In: Dabak N, ed. *Multidisipliner Yaklaşımla Kemik ve Yumuşak Doku Tümörleri*. Ankara, Bayt Yayıncılık; 2013:117–126.
- Yıldız Y. Osteosarkoma. In: Dabak N, ed. *Multidisipliner Yaklaşımla Kemik ve Yumuşak Doku Tümörleri*. Ankara, Bayt Yayıncılık; 2013:201–214.
- Heck RK. General Principles of Tumors. In: Canale ST, Beaty JH, eds. *Campbell's operative orthopaedics*. 11th ed. St Louis, MO: Mosby; 2008:775–854.