



Published in final edited form as:

*J Am Chem Soc.* 2018 December 05; 140(48): 16450–16454. doi:10.1021/jacs.8b11150.

## Enantioselective Coupling of Dienes and Phosphine Oxides

Shao-Zhen Nie<sup>†,§</sup>, Ryan T. Davison<sup>†</sup>, and Vy M. Dong<sup>\*,†</sup>


<sup>†</sup>Department of Chemistry, University of California, Irvine, California 92697, United States

<sup>§</sup>College of Pharmacy, Liaocheng University, Liaocheng, Shandong 252059, China

### Abstract

We report a Pd-catalyzed intermolecular hydrophosphinylation of 1,3-dienes to afford chiral allylic phosphine oxides. Commodity dienes and air stable phosphine oxides couple to generate organophosphorus building blocks with high enantio- and regiocontrol. This method constitutes the first asymmetric hydrophosphinylation of dienes.

### Graphical Abstract



Conjugated dienes are versatile motifs for constructing molecules that range from natural products to synthetic polymers.<sup>1,2</sup> In recent years, hydrofunctionalization has emerged as an attractive and atom-economical<sup>3</sup> method to transform dienes into valuable building blocks.<sup>4</sup> In comparison to other hydrofunctionalizations (e.g., hydroboration or hydroformylation), hydrophosphinylation remains in its infancy (Figure 1). Hirao first coupled isoprene and diethyl phosphonate to furnish an allylic phosphonate, albeit with low efficiency (10% yield) and at an elevated temperature (150 °C).<sup>5</sup> Tanaka later improved the hydrophosphorylation of 1,3-dienes by using a more reactive pinacol-derived phosphonate to synthesize allylphosphonates.<sup>6</sup> While promising, this strategy has been restricted to producing achiral regioisomers or racemic mixtures.<sup>7</sup> Given the potential for chiral phosphines in catalysis,<sup>8</sup> as well as the need for novel phosphine motifs in medicine<sup>9</sup> and agrochemical space,<sup>10</sup> we sought to develop an enantioselective hydrophosphinylation.<sup>11</sup> Herein, we report the transformation of several petroleum feedstocks and readily available dienes into chiral phosphine oxide building blocks, with high region- and enantiocontrol.

\*Corresponding Author dongv@uci.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and spectral data for all new compounds (PDF)

Crystallographic Data (CIF)

The authors declare no competing financial interests.

Given previously reported asymmetric hydroamination<sup>12</sup> and hydrothiolation<sup>13</sup> of 1,3-dienes, we chose to focus on a phosphorus nucleophile that would possess intermediate nucleophilicity compared to amines and thiols. As part of our reaction design, we imagined using phosphine oxides (**2**) as P-based nucleophiles because they are air stable, commercially available, and readily reduced to the corresponding phosphine.<sup>14</sup> In addition, the  $pK_a$  of **2** (ca. 25)<sup>15</sup> is between that of amines and thiols. Although the phosphine oxide reagent and its corresponding product could inhibit catalysis, hydrophosphinylation of alkenes<sup>16</sup> and alkynes<sup>17</sup> using transition-metal catalysis and photocatalysis has been reported. Encouraged by these examples, we set out to identify a catalyst that would overcome the established 1,4-addition pathway to furnish the desired chiral isomer.

We began our investigations with the coupling of 1-phenylbutadiene (**1a**) and commercially available **2a** (Table 1). We examined a range of achiral bisphosphine ligands, with both Rh and Pd precatalysts. While Rh showed no reactivity, Pd was promising for the hydrophosphinylation of **1a**. As highlighted in Table 1A, we observed that the ligand bite angle affected efficiency of the hydrophosphinylation.<sup>18</sup> Combining Pd<sub>2</sub>(dba)<sub>3</sub> and ferrocene-based dppf offered optimal results (90%, >20:1 *rr*). Catalytic amounts of acid provided an increase in the reaction rate; P(V)-based Brønsted acids proved to be the most effective for hydrophosphinylation (Table 1B). In the absence of an acid co-catalyst, we observe 16% of product **3aa** after 3 hours and 87% yield after 24 hours. Based on these results, we focused on the Josiphos ligand family with diphenylphosphinic acid as a co-catalyst.<sup>19</sup> As seen in Table 1C, with Pd(**L3**) we could lower the catalyst loading to 0.50 mol % and synthesize **3aa** on gram scale while retaining high reactivity (1.05 g, 91%) and selectivity (>20:1 *rr*, 95:5 *er*). The *er* in the presence of different acids shows little variation and ranges from 95:5 to 96:4.

With these conditions in hand, we investigated the hydrophosphinylation of various 1,3-dienes with phosphine oxide **2a** (Table 2). We found that a variety of 1-aryl substituted dienes could be transformed to chiral products **3ba-3ja** with moderate to high reactivity (36–88%) and selectivity (>20:1 *rr*, 88:12–96:4 *er*). Dienes containing aryl chlorides (**3ca**, **3ha**, **3ia**) offer higher reactivity than aryl bromides (**3da**), potentially due to the mitigation of side pathways initiated by oxidative addition into the C–X bond. The petroleum feedstocks butadiene (**1m**) and isoprene (**1n**) can be coupled with **2a** to furnish chiral building blocks **3ma** and **3na**, respectively. We observed product mixtures of **3ma** and **3na** that equally, or moderately, favor 3,4-addition over the established 1,4-addition previously reported for the hydrophosphorylation of butadiene<sup>6</sup> (**1m**) and isoprene<sup>5,6</sup> (**1n**). To examine if the allylic phosphine products (**3ma** and **3na**) could racemize by a sigmatropic rearrangement,<sup>20</sup> we resubjected **3ma** to the standard reaction conditions. After 12 hours, we observed no change in the enantiomeric excess. The 1,2-disubstituted diene (**1k**) and 1-alkyl substituted diene (**1l**) transform to products **3ka** and **3la**, respectively, in the presence of (*S*)-DTBM-SegPhos. This result suggests that the diene substitution pattern must be matched with the appropriate ligand family, an observation in agreement with our previous studies on Rh-catalyzed hydrothiolation of 1,3-dienes.<sup>13</sup>

Next, we investigated the hydrophosphinylation of **1a** with structurally and electronically different phosphine oxides (Table 3). We observed high reactivity (**3ab-3am**, 51–88%),

regioselectivity (>20:1 *rr*), and enantioselectivity (74:26–98:2 *er*). This coupling tolerates aryl (**3ab–3ai**), heterocyclic (**3aj**), and alkyl (**3ak**) phosphine oxides. Mono- (**2a–2g**), di- (**2h**), and trisubstituted (**2i**) aryl groups on the phosphine oxide partner can be coupled with **1a** to afford enantioenriched products (**3aa–3ai**). Fused ring motifs, which are the basis of a large class of ligand scaffolds, can also be incorporated in the phosphine oxide partner to generate products **3al** and **3am**.

Catalyst-controlled C–P bond formation would enable selective access to diastereomers. To test this idea, we prepared enantiopure phosphine oxide **2n** bearing a *tert*-butyl and phenyl group, a popular motif in chiral ligand design (Figure 2).<sup>21</sup> Depending on the enantiomer of the ligand **L3** used, the (*R,R*)-diastereomer **3an**<sup>22</sup> or (*R,S*)-diastereomer **3an'** can be obtained with high diastereocontrol (95:5 and 91:9 *dr*, respectively). This result represents a diastereodivergent strategy for making phosphine oxides.

Based on literature precedence and our own observations, we propose the mechanism depicted in Figure 3A. The Pd(0) precatalyst undergoes ligand substitution with the bisphosphine ligand to form a chiral monomeric species **I**, subsequent oxidative addition to diphenylphosphinic acid (HX) forms Pd–H species **II**. A related oxidative addition has been implicated as a key step in the hydrophosphinylation of terminal alkynes.<sup>17e</sup> In the absence of acid additives, we observe a significant induction period.<sup>23</sup> We reason that the addition of an acid co-catalyst (i.e., diphenylphosphinic acid) shortens the induction period and favors the Pd–H catalyst (e.g., **II**). At this point, two different modes of diene **1** coordination lead to the major product **3** (**path a**) and the minor product **4** (**path b**). In **path a**, species **III** undergoes hydropalladation to provide the key Pd– $\pi$ -allyl intermediate **IV**. Species **IV** then undergoes a ligand exchange with phosphine oxide **2** to form species **V**. Subsequent reductive elimination of **V** furnishes the allylic phosphine oxide **3** and regenerates the Pd-catalyst **I**.

To probe the mechanism, we conducted the following experiments (Figure 3B–D). First, a deuterium-labeled phosphine oxide **d-2a** was subjected to the standard reaction conditions. In this experiment, we see deuterium incorporation at the C1 (10% D) and C4 (64% D) positions of **d-3ea**. If hydropalladation were irreversible, we should observe about a 6:1 mixture of regioisomers. In contrast, we observe >20:1 *rr* and thus conclude that hydropalladation is reversible. Second, (*Z*)-1-phenylbutadiene (**Z-1a**) was subjected to the hydrophosphinylation. We observed only the (*E*)-product **3aa** (>95% *E* content) in similar yield (74%) and regioselectivity (>20:1 *rr*) compared to the model substrate (Table 1, **3aa**, 90% yield, >20:1 *rr*). This result suggests that isomerization occurs faster than C–P bond formation. Furthermore, excess diene **Z-1a** is recovered with *ca.* 25% *Z* content, which is consistent with a reversible hydropalladation and reversible diene coordination. By subjecting toluoyl phosphine oxide **2e** to product **3aa**, under otherwise standard conditions, we confirmed that the allylic phosphine oxide **3aa** cannot undergo further substitution to form **3ae**. Our proposal is in line with a study on alkyne hydrophosphinylation where Pd–P bond cleavage requires elevated temperatures and reductive elimination is the turnover-limiting step.<sup>17e</sup> We observed that alkyl-substituted dienes (**1l–1n**) form products (**3la–3na**)

with lower regioselectivity compared to the aryl-substituted dienes (**3ba-3ka**). Thus, reductive elimination to form the conjugated product appears to be favorable.

The direct construction of chiral phosphines and phosphine oxides has previously been achieved *via* additions to Michael acceptors or transition-metal catalyzed substitutions.<sup>24,25</sup> Herein, we reported a complementary way to access chiral phosphine oxides. This study features the first enantioselective hydrophosphinylation of dienes. Phosphine oxides and 1,3-dienes can be coupled to furnish chiral allylic products in high yields, regioselectivity, and enantioselectivity. Mechanistic studies suggest that the coupling proceeds through a reversible hypopalladation of the 1,3-diene partner followed by irreversible reductive elimination to afford the chiral phosphine oxide building blocks.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENT

Funding provided by UC Irvine, the National Institutes of Health (R35GM127071) and the National Science Foundation (CHE-1465263). Shao-Zhen Nie thanks Liaocheng University for a scholarship. We thank Dr. Joseph Ziller and Austin Ryan for X-ray crystallographic analysis and the Jarvo Lab for use of their SFC.

## REFERENCES

- (1). For selected reviews on 1,3-dienes as building blocks, see: Nicolaou KC; Snyder SA; Montagnon T; Vassilikogiannakis G *The Diels-Alder Reaction in Total Synthesis*. *Angew. Chem., Int. Ed* 2002, 41, 1668–1698. Reymond S; Cossy J *Copper-Catalyzed Diels-Alder Reactions*. *Chem. Rev* 2008, 108, 5359–5406. [PubMed: 18942879] Chen J-R; Hu X-Q; Lu L-Q; Xiao W-J *Formal [4+1] Annulation Reactions in the Synthesis of Carbocyclic and Heterocyclic Systems*. *Chem. Rev* 2015, 115, 5301–5365. [PubMed: 25992465] Büschleb M; Dorich S; Hanessian S; Tao D; Schenthal KB; Overman LE *Synthetic Strategies toward Natural Products Containing Contiguous Stereogenic Quaternary Carbon Atoms*. *Angew. Chem., Int. Ed* 2016, 55, 4156–4186.
- (2). For reviews on polymerization of 1,3-dienes, see: Monakov YB; Mullagaliev IR *Ionic and coordination diene polymerization and organic derivatives of main group metals*. *Russ. Chem. Bull.; Int. Ed* 2004, 53, 1–9. Friebe L; Nuyken O; Obrecht W *Neodymium Based Ziegler/Natta Catalysts and their Application in Diene Polymerization*. *Adv. Polym. Sci* 2006, 204, 1–154. Zhang Z; Cui D; Wang B; Liu B; Yang Y *Polymerization of 1,3-Conjugated Dienes with Rare-Earth Metal Precursors*. *Struct. Bonding (Berlin)* 2010, 137, 49–108. Valente A; Mortreux A; Visseaux M; Zinck P *Coordinative Chain Transfer Polymerization*. *Chem. Rev* 2013, 113, 3836–3857. [PubMed: 23391234] Takeuchi D *Stereoselective Polymerization of Conjugated Dienes* In *Encyclopedia of Polymer Science and Technology*; Wiley: New York, 2013; pp 1–25.
- (3). Trost BM *The atom economy—a search for synthetic efficiency*. *Science* 1991, 254, 1471–1477. [PubMed: 1962206]
- (4). For select hydrofunctionalizations of dienes, see: Löber O; Kawatsura M; Hartwig JF *Palladium-Catalyzed Hydroamination of 1,3-Dienes: A Colorimetric Assay and Enantioselective Additions*. *J. Am. Chem. Soc* 2001, 123, 4366–4367. [PubMed: 11457216] Page JP; RajanBabu TV *Asymmetric Hydrovinylation of 1-Vinylcycloalkenes. Reagent Control of Regio- and Stereoselectivity*. *J. Am. Chem. Soc* 2012, 134, 6556–6559. [PubMed: 22452442] Zbieg JR; Yamaguchi E; McInturff EL; Krische MJ *Enantioselective C–H Crotylation of Primary Alcohols via Hydrohydroxyalkylation of Butadiene*. *Science* 2012, 336, 324–327. [PubMed: 22442385] Park BY; Montgomery TP; Garza VJ; Krische MJ *Ruthenium Catalyzed Hydrohydroxyalkylation of Isoprene with Heteroaromatic Secondary Alcohols: Isolation and Reversible Formation of the Putative Metallocycle Intermediate*. *J. Am. Chem. Soc* 2013, 135, 16320–16323. [PubMed:

24156560] Chen Q-A; Kim DK; Dong VM Regioselective Hydroacylation of 1,3-Dienes by Cobalt Catalysis. *J. Am. Chem. Soc.* 2014, 136, 3772–3775. [PubMed: 24588202] Saini V; O'Dair M; Sigman MS Synthesis of Highly Functionalized Tri- and Tetrasubstituted Alkenes via Pd-Catalyzed 1,2-Hydrovinylation of Terminal 1,3-Dienes. *J. Am. Chem. Soc.* 2015, 137, 608–611. [PubMed: 25555197] Marcum JS; Roberts CC; Manan RS; Cervarich TN; Meek SJ Chiral Pincer Carbodicarbene Ligands for Enantioselective Rhodium-Catalyzed Hydroarylation of Terminal and Internal 1,3-Dienes with Indoles. *J. Am. Chem. Soc.* 2017, 139, 15580–15583. [PubMed: 29058881] Yang X-H; Lu A; Dong VM Intermolecular Hydroamination of 1,3-Dienes To Generate Homoallylic Amines. *J. Am. Chem. Soc.* 2017, 139, 14049–14052. [PubMed: 28953374] Gui Y-Y; Hu N; Chen X-W; Liao L-L; Ju T; Ye J-H; Zhang Z; Li J; Yu D-G Highly Regio- and Enantioselective Copper-Catalyzed Reductive Hydroxymethylation of Styrenes and 1,3-Dienes with CO<sub>2</sub>. *J. Am. Chem. Soc.* 2017, 139, 17011–17014. [PubMed: 29125759] Adamson NJ; Hull E; Malcolmson SJ Enantioselective Intermolecular Addition of Aliphatic Amines to Acyclic Dienes with a Pd-PHOX Catalyst. *J. Am. Chem. Soc.* 2017, 139, 7180–7183. [PubMed: 28453290] Adamson NJ; Wilbur KCE; Malcolmson SJ Enantioselective Intermolecular Pd-Catalyzed Hydroalkylation of Acyclic 1,3-Dienes with Activated Pronucleophiles. *J. Am. Chem. Soc.* 2018, 140, 2761–2764. [PubMed: 29446922] Schmidt VA; Kennedy CR; Bezdek MJ; Chirik PJ Selective [1,4]-Hydrovinylation of 1,3-Dienes with Unactivated Olefins Enabled by Iron Diimine Catalysts. *J. Am. Chem. Soc.* 2018, 140, 3443–3453. [PubMed: 29414238] For reviews, see: Hydrofunctionalization In Topics in Organometallic Chemistry; Ananikov VP, Tanaka M Eds.; Springer: Berlin, 2014; Vol. 343. McNeill E; Ritter T 1,4-Functionalization of 1,3-Dienes With Low-Valent Iron Catalysts. *Acc. Chem. Res.* 2015, 48, 2330–2343. [PubMed: 26214092] Bezenine-Lafollée S; Gil R; Prim D; Hannedouche J First-Row Late Transition Metals for Catalytic Alkene Hydrofunctionalisation: Recent Advances in C–N, C–O, and C–P Bond Formation. *Molecules* 2017, 22, 1901–1929.

- (5). Hirao T; Masunaga T; Yamada N; Ohshiro Y; Agawa T Palladium-catalyzed New Carbon-Phosphorous Bond Formation. *Bull. Chem. Soc. Jpn* 1982, 55, 909–913.
- (6). Mirzaei F; Han L-B; Tanaka M Palladium-catalyzed hydrophosphorylation of 1,3-dienes leading to allylphosphonates. *Tet. Lett* 2001, 42, 297–299.
- (7). Two additional examples have been reported for the coupling of hydrophosphorous acid (H<sub>3</sub>PO<sub>2</sub>) and a 1,3-diene that yield allylic phosphinic acids via Pd-catalyzed 1,4-addition, see: Bravo-Altamirano K; Abrunhosa-Thomas I; Montchamp J-L Palladium-Catalyzed Reactions of Hypophosphorous Compounds with Allenes, Dienes, and Allylic Electrophiles: Methodology for the Synthesis of *H*-Phosphinates. *J. Org. Chem* 2008, 73, 2292–2301. [PubMed: 18275217]
- (8). For select reviews of phosphines in catalysis, see: Tolman CA Steric effects of phosphorus ligands in organometallic chemistry and homogeneous catalysis. *Chem. Rev* 1977, 77, 313–348. van Leeuwen PWNM; Kamer PCJ; Reek JNH; Dierkes P Ligand Bite Angle Effects in Metal-catalyzed C–C Bond Formation. *Chem. Rev* 2000, 100, 2741–2770. [PubMed: 11749304] Noyori R; Ohkuma T Asymmetric Catalysis by Architectural and Functional Molecular Engineering: Practical Chemo- and Stereoselective Hydrogenation of Ketones. *Angew. Chem., Int. Ed* 2001, 40, 40–73. Xiao Y; Guo H; Kwon O Nucleophilic Chiral Phosphines: Powerful and Versatile Catalysts for Asymmetric Annulations. *Aldrichimica Acta.* 2016, 49, 3–13. [PubMed: 28077882] Ni H; Chan W-L; Lu Y Phosphine-Catalyzed Asymmetric Organic Reactions. *Chem. Rev* 2018, 118, 9344–9411. [PubMed: 30204423]
- (9). For reviews of phosphines in medicine, see: Tiekink ERT Phosphinegold(I) thiolates—pharmacological use and potential. *Bioinorg. Chem. Appl* 2003, 1, 53–67. Phillips AD; Gonsalvi L; Romerosa A; Vizza F; Peruzzini M Coordination chemistry of 1,3,5-triaza-7-phosphaadamantane (PTA): Transition metal complexes and related catalytic, medicinal, and photoluminescent applications. *Coord. Chem. Rev* 2004, 248, 955–993. Dominelli B; Correia JDG; Kühn FE Medicinal Applications of Gold(I/nI)-Based Complexes Bearing V-Heterocyclic Carbene and Phosphine Ligands. *J. Organomet. Chem* 2018, 866, 153–164.
- (10). For a select example of phosphines in agrochemicals, see: Schlipalius DI; Valmas N; Tuck AG; Jagadeesan R; Ma L; Kaur R; Goldinger A; Anderson C; Kuang J; Zuryn S; Mau YS; Cheng Q; Collins PJ; Nayak MK; Schirra HJ; Hilliard MA; Ebert PR A Core Metabolic Enzyme Mediates Resistance to Phosphine Gas. *Science* 2012, 338, 807–810. [PubMed: 23139334] For a review,




see: Nath NS; Bhattacharya I; Tuck AG; Schlipalius DI; Ebert PR Mechanisms of Phosphine Toxicity. *J. Toxicol* 2011, 2011, 1–9.

- (11). We are using the definition of hydrophosphinylation that refers to the addition of a phosphine oxide to a degree of unsaturation, see: Han L-B; Choi N; Tanaka M Oxidative Addition of  $\text{HP(O)Ph}_2$  to Platinum(0) and Palladium(0) Complexes and Palladium-Catalyzed Regio- and Stereoselective Hydrophosphinylation of Alkynes. *Organometallics* 1996, 15, 3259–3261.
- (12). Yang X-H; Dong VM Rhodium-Catalyzed Hydrofunctionalization: Enantioselective Coupling of Indolines and 1,3-Dienes. *J. Am. Chem. Soc* 2017, 139, 1774–1777. [PubMed: 28128936]
- (13). Yang X-H; Davison RT; Dong VM Catalytic Hydrothiolation: Regio- and Enantioselective Coupling of Thiols and Dienes. *J. Am. Chem. Soc* 2018, 140, 10443–10446.
- (14). For reviews on reducing phosphine oxides to phosphines, see: Héroult D; Nguyen DH; Nuel D; Buono G Reduction of secondary and tertiary phosphine oxides to phosphines. *Chem. Soc. Rev* 2015, 44, 2508–2528. [PubMed: 25714261] Kovács T; Keglevich G The Reduction of Tertiary Phosphine Oxides by Silanes. *Curr. Org. Res* 2017, 21, 569–585.
- (15). Grayson M; Farley CE; Streuli CA Secondary phosphine oxides: The effect of structure on acid strength and rates of cleavage of disulfides. *Tetrahedron* 1967, 23, 1065–1078.
- (16). For select hydrophosphinylations of alkenes, see: Kawaguchi S-I; Nomoto A; Sonoda M; Ogawa A Photoinduced hydrophosphinylation of alkenes with diphenylphosphine oxide. *Tet. Lett* 2009, 50, 624–626. Yoo W-J; Kobayashi S Hydrophosphinylation of unactivated alkenes with secondary phosphine oxides under visible-light photocatalysis. *Green. Chem* 2013, 15, 1844–1848. Li Z; Fan F; Zhang Z; Xiao Y; Liu D; Liu Z-Q A silver-initiated free-radical intermolecular hydrophosphinylation of unactivated alkenes. *RSC Adv.* 2015, 5, 27853–27856.
- (17). For select hydrophosphinylations of alkynes, see: Han L-B; Hua R; Tanaka M Phosphinic Acid Induced Reversal of Regioselectivity in Pd-Catalyzed Hydrophosphinylation of Alkynes with  $\text{Ph}_2\text{P(O)H}$ . *Angew. Chem., Int. Ed* 1998, 37, 94–96. Han L-B; Zhao C-Q; Tanaka M Rhodium-Catalyzed Regio- and Stereoselective Addition of Diphenylphosphine Oxide to Alkynes. *J. Org. Chem* 2001, 66, 5929–5932. [PubMed: 11511276] Han L-B; Zhang C; Yazawa H; Shimada S Efficient and Selective Nickel-Catalyzed Addition of H–P(O) and H–S Bonds to Alkynes. *J. Am. Chem. Soc* 2004, 126, 5080–5081. [PubMed: 15099086] Rooy SV; Cao C; Patrick BO; Lam A; Love JA Alkyne hydrophosphinylation catalyzed by rhodium pyrazolylborate complexes. *Inorg. Chim. Acta* 2006, 359, 2918–2923. Chen T; Zhao C-Q; Han L-B Hydrophosphorylation of Alkynes Catalyzed by Palladium: Generality and Mechanism. *J. Am. Chem. Soc* 2018, 140, 3139–3155. [PubMed: 29420024] For a review, see: Xu Q; Han L-B Metal-catalyzed additions of H–P(O) bonds to carbon–carbon unsaturated bonds. *J. Organomet. Chem* 2011, 696, 130–140.
- (18). For a review on ligand bite angle effects, see: ref 8b.
- (19). We also tried the PHOX ligand scaffold, which has demonstrated high reactivity and enantioselectivity in Pd-catalyzed diene hydroamination (see: ref 4j) and hydroalkylation (see: ref 4k), but observed no reactivity.
- (20). Herriott AW; Mislow K Rearrangement of allyl phosphinites and optical stability of allyl phosphine oxides. *Tet. Lett* 1968, 9, 3013–3016.
- (21). Lühr S; Holz J; Börner A The Synthesis of Chiral Phosphorus Ligands for use in Homogeneous Metal Catalysis. *Chem-CatChem* 2011, 3, 1708–1730.
- (22). X-ray crystallography data confirmed the absolute configuration of 3an, CCDC: 1868886. The absolute configuration of compounds 3aa–3am, 3an' and 3ba–3na were assigned by analogy.
- (23). See the Supporting Information for more details.
- (24). For select enantioselective additions to Michael acceptors, see: Carlone A; Bartoli G; Bosco M; Sambri L; Melchiorre P Organocatalytic Asymmetric Hydrophosphination of  $\alpha,\beta$ -Unsaturated Aldehydes. *Angew. Chem., Int. Ed* 2007, 46, 4504–4506. Ibrahim I; Rios R; Vesely J; Hammar P; Eriksson L; Himo F; Córdova A Enantioselective Organocatalytic Hydrophosphination of  $\alpha,\beta$ -Unsaturated Aldehydes. *Angew. Chem., Int. Ed* 2007, 46, 4507–4510. Feng J-J; Chen X-F; Shi M; Duan W-L Palladium-Catalyzed Asymmetric Addition of Diarylphosphines to Enones toward the Synthesis of Chiral Phosphines. *J. Am. Chem. Soc* 2010, 132, 5562–5563. [PubMed: 20359215] Chew RJ; Teo KY; Huang Y; Li B-B; Li Y; Pullarkat SA; Leung P-H Enantioselective phospho-Michael addition of diarylphosphines to  $\beta,\gamma$ -unsaturated  $\alpha$ -ketoesters and amides.


Chem. Commun 2014, 50, 8768–8770. For a review, see: Pullarkat SA Recent Progress in Palladium-Catalyzed Asymmetric Hydrophosphination. Synthesis 2016, 48, 493–503.

- (25). For select enantioselective transition-metal catalyzed sub-stitutions, see: Butti P; Rochat R; Sadow AD; Togni A Palladium-Catalyzed Enantioselective Allylic Phosphination. Angew. Chem., Int. Ed 2008, 47, 4878–4881. Zhang L; Liu W; Zhao X Carbon-Phosphorous Bond Formation by Enantioselective Palladium-Catalyzed Allylation of Diphenylphosphine Oxide. Eur. J. Org. Chem 2014, 6846–6849. Liu C; Wang Q Alkenylation of C(sp<sup>3</sup>)-H Bonds by Zincation/Copper-Catalyzed Cross-Coupling with Iodonium Salts. Angew. Chem., Int. Ed 2018, 57, 4727–4731.


A. First example of diene and phosphonate coupling (Hirao, 1982)



B. Improved hydrophosphorylation of dienes (Tanaka, 2001)




C. Proposed enantioselective hydrophosphinylation of dienes




**Figure 1.**  
Inspiration for asymmetric hydrophosphorylation of 1,3-dienes.






**Figure 2.**  
Diastereodivergent hydrophosphinylation of **1a**.


## A. Proposed catalytic cycle for the hydrophosphinylation of dienes




## B. Deuterium-labeling study



## C. Diene geometry study



## D. Cross-over study



**Figure 3.** Proposed mechanism and initial investigations of the Pd-catalyzed hydrophosphinylation of 1,3-dienes.

Table 1.

Ligand and Acid Effects on Asymmetric Hydrophosphinylation of **1a**<sup>a</sup>

**A. Ligand Bite Angle Effects<sup>b</sup>** Increasing Bite Angle  $\longrightarrow$

Ligand:	dppm	dppe	dppp	dppf	dppb	DPEphos	Xantphos
Yield <b>3aa</b> :	trace	trace	90%	90%	61%	34%	9%
Time:	16 h	16 h	16 h	3 h	16 h	16 h	16 h

---

**B. Acid Effects<sup>c</sup>** Increasing Acidity  $\longrightarrow$

Acid:	None	PhCOOH	(Ph) <sub>2</sub> P(O)OH	(PhO) <sub>2</sub> P(O)OH	MsOH
Yield <b>3aa</b> :	16%	51%	90%	87%	72%

---

**C. Chiral Ligand Effects**

**L1**  
88%, 73:27 *er*

**L2**  
87%, 77:23 *er*

**L3**  
91%<sup>d</sup>, 95:5 *er*

<sup>a</sup>Reaction conditions: **1a** (0.12 mmol), **2a** (0.10 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (2.5 mol%), ligand (5.0 mol%), acid (20 mol%), toluene (0.40 mL), 3 h (unless otherwise noted). Yield determined by GC-FID analysis of the reaction mixture, which was referenced to 1,3,5-trimethoxybenzene.


Regioselectivity ratio (*rr*) is the ratio of **3aa** to **4aa**, which is determined by <sup>31</sup>P NMR analysis of reaction mixture. Enantioselectivity ratio (*er*) determined by chiral SFC. See SI for full structure of abbreviations used. Unless otherwise noted, *rr* is >20:1.

<sup>b</sup>Standard conditions with (Ph)<sub>2</sub>P(O)OH as acid.

<sup>c</sup>Standard conditions with dppf as ligand.

<sup>d</sup>Isolated yield of **3aa**, 3.47 mmol scale, using Pd<sub>2</sub>(dba)<sub>3</sub> (0.50 mol%) and **L3** (1.0 mol%) with standard conditions, 18 h.

Table 2.


Hydrophosphinylation of Various 1,3-Dienes<sup>a</sup>

<sup>a</sup>Reaction conditions: **1** (0.12 mmol), **2a** (0.10 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (2.5 mol%), ligand (5.0 mol%), (Ph)<sub>2</sub>P(O)OH (20 mol%), toluene (0.40 mL), 6

h. Isolated yield of **3**. Regioselectivity ratio (*rr*) is the ratio of **3** to **4**, which is determined by <sup>31</sup>P NMR analysis of reaction mixture. Enantioselectivity determined by chiral SFC.

<sup>b</sup>(S)-DTBM-SegPhos (5.0 mol%) instead of **L3**, see SI for structure, 24 h.

Table 3.

Hydrophosphinylation of **1a** with Various Phosphine Oxides<sup>a</sup>

<sup>a</sup>Reaction conditions: **1a** (0.12 mmol), **2** (0.10 mmol),  $\text{Pd}_2(\text{dba})_3$  (2.5 mol%), ligand (5.0 mol%),  $(\text{Ph})_2\text{P}(\text{O})\text{OH}$  (20 mol%), toluene (0.40 mL), 6 h. Isolated yield of **3**. Regioselectivity ratio (*rr*) is the ratio of **3** to **4**, which is determined by  $^{31}\text{P}$  NMR analysis of reaction mixture. Enantioselectivity determined by chiral SFC. See SI for full structure of abbreviations used.

<sup>b</sup>Reaction time is 24 h.