

## Comparison of Kato-Katz Thick Smear, Mini-FLOTAC, and Flukefinder for the Detection and Quantification of *Fasciola hepatica* Eggs in Artificially Spiked Human Stool

Daniel A. Zárate-Rendón,<sup>1\*</sup> Johnny Vlamincq,<sup>2</sup> Bruno Levecke,<sup>2</sup> Andrea Briones-Montero,<sup>1</sup> and Peter Geldhof<sup>2</sup>

<sup>1</sup>Laboratorio de Parasitología, Facultad de Zootecnia, Universidad Nacional Agraria La Molina, Lima, Peru; <sup>2</sup>Laboratory of Parasitology, Faculty of Veterinary Medicine, University of Ghent, Merelbeke, Belgium

**Abstract.** We compared the diagnostic performance of the standard method (Kato-Katz) with two recently developed methods (Mini-FLOTAC and Flukefinder) for the detection and quantification of *Fasciola hepatica* eggs in human stool. Uninfected human stool samples were artificially spiked with *F. hepatica* eggs to reach final concentrations of 14, 28, 41, or 96 eggs per gram of stool (epg). Only Flukefinder showed 100% sensitivity in all but the samples with the lowest concentration of eggs (14 epg), in which it had a sensitivity of 60%. Each of the methods underestimated the true fecal egg counts (FECs), Flukefinder resulting in the most biased egg counts (egg counts 0.18 times lower than the expected FECs). Only the Flukefinder resulted in more precise results (coefficient of variance < 30%) from FECs of 96 epg onward. The outcome of this study indicates that the Flukefinder is a useful alternative diagnostic method for human fascioliasis in stool.

Fascioliasis is caused by the flatworms *Fasciola hepatica* and *F. gigantica* and poses a major problem to both animal and public health.<sup>1,2</sup> In humans, this parasitosis is an acknowledged (re-)emerging human disease in approximately 51 countries, its spread being closely linked to climatic conditions and affected by climatic change.<sup>3</sup> The more important regions for human disease in the world are located in South America,<sup>4</sup> with highly endemic areas in the Andean highlands.<sup>2,3</sup> In recent decades, a variety of studies reported a hyperendemic prevalence (> 10%) of human fascioliasis in these areas,<sup>5,6</sup> accounting for approximately 17 million infected people and 180 million at risk of infection.<sup>1</sup> Morbidity caused by fascioliasis is particularly pronounced in children,<sup>7–9</sup> contributing to anemia and poor development.<sup>10,11</sup> Despite the increased use of serological tests, the routine tests used to diagnose human fascioliasis in endemic rural areas are based on the microscopical demonstration and quantification of eggs in stool using coprological methods. One of the most widely used coprological methods is the Kato-Katz thick smear.<sup>9</sup> This method has been extensively used to determine both prevalence and intensity of infection (based on the number of eggs per gram of stool; epg) in highly endemic areas.<sup>12,13</sup> It is a cheap and simple method that has been standardized and recommended for a wider range of parasitic diseases in humans.<sup>5</sup> However, an important limitation of this method is that it fails to detect infection of low intensity.<sup>14</sup> Other coprological methods that hold promise as an alternative for Kato-Katz are the recently developed Mini-FLOTAC and Flukefinder test. Both methods have proven to have a high sensitivity for the diagnosis of fascioliasis in livestock,<sup>15–17</sup> but they have not yet been fully validated for the diagnosis of human fascioliasis. In this study, we compared the diagnostic performance of the standard method (Kato-Katz) with the recently developed methods Mini-FLOTAC and Flukefinder for the detection and quantification of *F. hepatica* eggs in human stool samples.

The study protocol was approved by the Ethics Committee of the Universidad Peruana Cayetano Heredia (Code SIDISI:

17010) and Ghent University (Belgium (Code OZ 2017/0869). Negative stool samples were collected from children attending Institutional Day Care of the Universidad Nacional Agraria la Molina in Lima (Peru). This day care lies in an area which is nonendemic for fascioliasis. The collected samples were repeatedly examined by a modified rapid sedimentation test<sup>5</sup> to confirm the absence of *F. hepatica* eggs. Confirmed *Fasciola*-negative stool samples were subsequently pooled and homogenized using wooden tongue depressors. The pooled sample was subsequently divided into four aliquots of 100 g each. This amount of stool allowed for repeated examination by all diagnostic methods. The aliquots were subsequently stored at 4°C for approximately 2 weeks until the experiments. Eggs of *F. hepatica* were obtained from bile and gall bladders collected from livers of naturally infected cattle that were slaughtered at an authorized abattoir in Lima. The eggs were purified and collected using the commercial Flukefinder method (Flukefinder) as described by the manufacturer's instructions. The purified eggs were suspended in distilled water, and their concentration was determined by counting the number of eggs present in 10 aliquots of 10 µL each and subsequently calculating the arithmetic mean of these egg counts. Different volumes of the egg suspension were subsequently pipetted to the four aliquots of the pooled stool sample to obtain a final concentration of approximately 14, 28, 41, and 96 epg. A homogeneous distribution of the eggs was obtained in the stool by vigorously mixing the pooled stool using a metal plaster spatula.

The Kato-Katz thick smear (SteriTech, Kent, WA),<sup>17</sup> Flukefinder (Flukefinder, Soda Springs, ID),<sup>16</sup> and Mini-FLOTAC (Zinc Chloride flotation solution with a specific gravity of 1.30; University of Naples, Naples, Italy)<sup>14</sup> methods were performed according to the manufacturer's instructions. The fecal egg counts (FECs; expressed in epg) were obtained by multiplying the egg counts by 24 for Kato-Katz (0.417 g of stool is examined), 10 for Mini-FLOTAC (0.1 g of stool is examined = 2 mL of stool suspension which contains 2 g in a total volume 40 mL), and 0.5 for Flukefinder (2 g of stool is examined). Each method was performed 10 times on each of the four spiked samples. To avoid any systematic error in the results (e.g., first examining all aliquots with one particular method), the sequence of stool analysis was randomized. The sensitivity

\* Address correspondence to Daniel A. Zárate-Rendón, Laboratorio de Parasitología, Facultad de Zootecnia, Universidad Nacional Agraria La Molina, Lima 12, Peru. E-mail: dazre@lamolina.edu.pe

(proportion of samples in which eggs were found) was calculated across the three diagnostic methods and the four egg concentrations. The accuracy and precision of the FECs measured by the methods were based on the ratio observed over the expected FECs and the coefficient of variance (CV) (= SD/mean), respectively.

Overall, Flukefinder was the most sensitive method (90.0% [95% CI: 80.0; 97.5]), followed by Mini-FLOTAC (67.5 [95% CI: 52.5; 80.0]) and Kato-Katz (32.5% [95% CI: 17.5; 47.5]). As shown in Table 1, the sensitivity increased as a function of the number of eggs spiked, although this increase varied across the methods. The Flukefinder already reached a sensitivity of 100% for FECs of at least 28 epg, whereas for Mini-FLOTAC, this was only the case for the highest FECs (96 epg). Kato-Katz never reached a sensitivity of 100%, the sensitivity being 70% at most.

The ratio of the observed epg over the expected FECs was < 1 for all the three methods, indicating that the methods

underestimated FECs. The least biased FECs were obtained by Kato-Katz and Mini-FLOTAC, both methods resulting in FECs that on average were 0.31 times lower than the expected FECs. Fecal egg counts obtained by Flukefinder were 0.19 times lower. Table 1 illustrates the accuracy across the different levels of seeded eggs for the different methods. For Mini-FLOTAC, the percentage of accuracy in FECs remained relatively unchanged, ranging from 24% to 35.7%. This was in contrast to the other methods for which a wider range was observed (Kato-Katz: 17.1–55%; Flukefinder: 4.3–33.6%). Flukefinder was the most precise method of all, resulting in the lowest CV values across the different levels of seeded eggs. A CV not higher than 30% was obtained for FECs of 96 epg. The two other methods did not achieve a CV < 30%, although the CV of Mini-FLOTAC was 35% for the highest FECs (98 epg).

To the best of our knowledge, this is the first work that compares Kato-Katz, Mini-FLOTAC, and Flukefinder egg detection and quantification of *F. hepatica* eggs in artificially

TABLE 1

The results of the spiking experiment using the Kato-Katz thick smear, Mini-FLOTAC, and Flukefinder on the samples artificially spiked with 14, 28, 41, or 96 *Fasciola hepatica* eggs per gram (epg)

Method	Replicate	Concentration of <i>F. hepatica</i> eggs per gram of stool (epg)			
		14	28	41	96
Kato-Katz	1	0	0	0	72
	2	0	24	0	120
	3	0	0	0	0
	4	0	48	0	0
	5	0	0	0	48
	6	0	24	48	168
	7	0	0	0	72
	8	24	24	0	24
	9	0	0	0	0
	10	0	0	0	24
Sensitivity (%)		10.0	40.0	10.0	70.0
Mean + SD		2.4 + 7.6	12 + 17.0	4.8 + 15.2	52.8 + 56.3
Accuracy (%)		17.1	42.9	11.7	55.0
CV (%)		44.3	39.6	129.7	102.4
Mini-FLOTAC	1	0	20	0	30
	2	10	20	0	30
	3	0	20	10	30
	4	20	10	0	20
	5	0	0	20	10
	6	10	0	20	30
	7	10	10	20	30
	8	0	10	10	10
	9	0	0	30	20
	10	0	0	30	20
Sensitivity (%)		40.0	60.0	70.0	100
Mean + SD		5 ± 7.1	9 ± 8.8	14 ± 11.7	23 ± 8.2
Accuracy (%)		35.7	32.1	34.1	24.9
CV (%)		19.8	27.2	34.4	34.3
Flukefinder	1	1	4.5	2.5	29.5
	2	0.5	9.5	1.5	28
	3	0	7	5	23
	4	1.5	6	5	32
	5	0	4	5	42.5
	6	1	11	2.5	30.5
	7	0.5	7.5	0.5	46
	8	0	6	3	24.5
	9	1.5	6.5	6	30
	10	0	17.5	6	37
Sensitivity (%)		60.0	100	100	100
Mean + SD		0.6 ± 0.6	7.95 ± 4.0	3.7 ± 1.9	32.3 ± 7.4
Accuracy (%)		4.3	28.4	9.0	33.6
CV (%)		14.3	14.0	21.6	22.1

CV = coefficient of variance; FEC = fecal egg count. Ten FEC replicates were performed for each method and the results are shown in epg. The accuracy equals the ratio observed over the expected FECs. CV = mean/SD.

spiked human stool samples. Our results indicate that Flukefinder holds promise as an alternative to Kato-Katz for the diagnosis of human fascioliasis in stool. In contrast to the other methods, it shows a high sensitivity at low FECs (up to FECs of 28 epg) and it results in highly precise egg counts (CV < 30%) at FECs of 96 epg. The method underestimates FECs, but this was observed by all methods. Moreover, the Flukefinder could potentially also be used for the diagnosis of other major trematodiasis, such as paragonimiasis or schistosomiasis. Further research is now required to assess the clinical sensitivity and the user-friendliness of the Flukefinder under field conditions.

Received December 14, 2018. Accepted for publication March 14, 2019.

Published online May 6, 2019.

Financial support: This work was supported by the Consortium of Flemish Universities of Belgium (VLIR) through the Project VLIR UOS-UNALM. Both J. V. and B. L. are financially supported by an FWO grant.

Authors' addresses: Daniel A. Zárate-Rendón and Andrea Briones-Montero, Facultad de Zootecnia, Laboratorio de Parasitología, Universidad Nacional Agraria La Molina, Lima, Peru, E-mails: dazre@lamolina.edu.pe and abriones.montero@gmail.com. Johnny Vlamincq, Bruno Levecke, and Peter Geldhof, Faculty of Veterinary Medicine, Laboratory of Parasitology, Ghent University, Ghent, Belgium, E-mails: johnny.vlaminck@ugent.be, bruno.levecke@ugent.be, and peter.geldhof@ugent.be.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## REFERENCES

- Cwiklinski K, O'Neill SM, Donnelly S, Dalton JP, 2016. A prospective view of animal and human fasciolosis. *Parasite Immunol* 38: 558–568.
- Mas-Coma S, Bargues MD, Valero MA, 2018. Human fascioliasis infection sources, their diversity, incidence factors, analytical methods and prevention measures. *Parasitology* 145: 1665–1699.
- Mas-Coma S, Bargues MD, Valero MA, 2005. Fascioliasis and other plant-borne trematode zoonoses. *Int J Parasitol* 35: 1255–1278.
- Adela Valero M, Perez-Crespo I, Khoubbane M, Artigas P, Panova M, 2012. *Fasciola hepatica* phenotypic characterization in Andean human endemic areas: valley versus altiplanic patterns analyzed in liver flukes from sheep from Cajamarca and Mantaro, Peru. *Infect Genet Evol* 12: 403–410.
- Marcos L, Maco V, Samalvides F, Terashima A, Espinoza JR, Gotuzzo E, 2006. Risk factors for *Fasciola hepatica* infection in children: a case-control study. *Trans R Soc Trop Med Hyg* 100: 158–166.
- Marcos LA, Terashima A, Leguia G, Canales M, Espinoza JR, Gotuzzo E, 2007. *Fasciola hepatica* infection in Peru: an emergent disease. *Rev Gastroenterol Peru* 27: 389–396.
- Cabada MM, Lopez M, Cruz M, Delgado JR, Hill V, White AC Jr., 2016. Treatment failure after multiple courses of triclabendazole among patients with fascioliasis in Cusco, Peru: a case series. *PLoS Negl Trop Dis* 10: e0004361.
- Espinoza JR, Terashima A, Herrera-Velit P, Marcos LA, 2010. Human and animal fascioliasis in Peru: impact in the economy of endemic zones. *Rev Peru Med Exp Salud Publica* 27: 604–612.
- Marcos L, Romani L, Florencio L, Terashima A, Canales M, Nestares J, Huayanay L, Gotuzzo E, 2007. Hyperendemic and mesoendemic zones of *Fasciola* infection surrounding urban Lima: an emerging disease?. *Rev Gastroenterol Peru* 27: 31–36.
- Marcos Raymundo LA, Maco Flores V, Terashima Iwashita A, Samalvides Cuba F, Gotuzzo Herencia E, 2002. Clinical characteristics of chronic infection by *Fasciola hepatica* in children. *Rev Gastroenterol Peru* 22: 228–233.
- Lopez M, White AC Jr., Cabada MM, 2012. Burden of *Fasciola hepatica* infection among children from Paucartambo in Cusco, Peru. *Am J Trop Med Hyg* 86: 481–485.
- Cabada MM, Goodrich MR, Graham B, Villanueva-Meyer PG, Lopez M, Arque E, White AC Jr., 2014. Fascioliasis and eosinophilia in the highlands of Cuzco, Peru and their association with water and socioeconomic factors. *Am J Trop Med Hyg* 91: 989–993.
- Gonzalez LC, Esteban JG, Bargues MD, Valero MA, Ortiz P, Naquira C, Mas-Coma S, 2011. Hyperendemic human fascioliasis in Andean valleys: an altitudinal transect analysis in children of Cajamarca province, Peru. *Acta Trop* 120: 119–129.
- Mas-Coma S, Bargues MD, Valero MA, 2014. Diagnosis of human fascioliasis by stool and blood techniques: update for the present global scenario. *Parasitology* 141: 1918–1946.
- Cringoli G, Maurelli MP, Levecke B, Bosco A, Vercruysse J, Utzinger J, Rinaldi L, 2017. The Mini-FLOTAC technique for the diagnosis of helminth and protozoan infections in humans and animals. *Nat Protoc* 12: 1723–1732.
- Rinaldi L, Gonzalez S, Guerrero J, Aguilera LC, Musella V, Genchi C, Cringoli G, 2012. A one-health integrated approach to control fascioliasis in the Cajamarca valley of Peru. *Geospat Health* 6: S67–S73.
- Faria RN, Cury MC, Lima WS, 2008. Concordância entre duas técnicas coproparasitológicas para diagnóstico de *Fasciola hepatica* em bovinos. *Arq Bras Med Vet Zootec* 60: 1023–1025.