

## MOLECULAR BIOLOGY

Single-base mapping of m<sup>6</sup>A by an antibody-independent methodZhang Zhang<sup>1</sup>, Li-Qian Chen<sup>1</sup>, Yu-Li Zhao<sup>1</sup>, Cai-Guang Yang<sup>2</sup>, Ian A. Roundtree<sup>3</sup>, Zijie Zhang<sup>3</sup>, Jian Ren<sup>1</sup>, Wei Xie<sup>1</sup>, Chuan He<sup>3</sup>, Guan-Zheng Luo<sup>1\*</sup>

*N*<sup>6</sup>-methyladenosine (m<sup>6</sup>A) is one of the most abundant messenger RNA modifications in eukaryotes involved in various pivotal processes of RNA metabolism. The most popular high-throughput m<sup>6</sup>A identification method depends on the anti-m<sup>6</sup>A antibody but suffers from poor reproducibility and limited resolution. Exact location information is of great value for understanding the dynamics, machinery, and functions of m<sup>6</sup>A. Here, we developed a precise and high-throughput antibody-independent m<sup>6</sup>A identification method based on the m<sup>6</sup>A-sensitive RNA endoribonuclease recognizing ACA motif (m<sup>6</sup>A-sensitive RNA-Endoribonuclease-Facilitated sequencing or m<sup>6</sup>A-REF-seq). Whole-transcriptomic, single-base m<sup>6</sup>A maps generated by m<sup>6</sup>A-REF-seq quantitatively displayed an explicit distribution pattern with enrichment near stop codons. We used independent methods to validate methylation status and abundance of individual m<sup>6</sup>A sites, confirming the high reliability and accuracy of m<sup>6</sup>A-REF-seq. We applied this method on five tissues from human, mouse, and rat, showing that m<sup>6</sup>A sites are conserved with single-nucleotide specificity and tend to cluster among species.

## INTRODUCTION

Among more than 100 RNA chemical modifications, *N*<sup>6</sup>-methyladenosine (m<sup>6</sup>A) is one of the most abundant forms in eukaryotic mRNA, accounting for 0.1 to 0.4% of all adenosine (1). The m<sup>6</sup>A modification influences various stages of mRNA metabolism and the biogenesis of long noncoding RNA and microRNA, with effects across diverse biological processes including neuronal development, cell fate transition, immune response, DNA damage response, and tumorigenesis (2–10). Most m<sup>6</sup>A sites were found in conserved motif DRACH (D=G/A/U, R=G/A, H=A/U/C) (1), with an estimated methylation ratio ranging from 6 to 80% (11). Previous whole-transcriptome m<sup>6</sup>A maps have suggested that m<sup>6</sup>A modifications are enriched around stop codon, implying the functional importance of distribution pattern for m<sup>6</sup>A (12–14).

The chemical properties of m<sup>6</sup>A are similar to adenosine, making it difficult to discriminate by chemical reactions (15). The recently developed high-sensitivity liquid chromatography–tandem mass spectrometry and blotting methods relying on antibodies were widely used to quantify the overall m<sup>6</sup>A level. To detect individual sites, methyl-sensitive ligase has been applied to confirm the methylation status of specific adenosines (16, 17), while the method called SCARLET (site-specific cleavage and radioactive-labeling followed by ligation-assisted extraction and thin-layer chromatography) can quantify the methylation level of individual m<sup>6</sup>A site (11). Other methods were developed to identify m<sup>6</sup>A in single-base resolution during reverse transcription, taking advantage of m<sup>6</sup>A-sensitive reverse transcriptase (18, 19), chemoenzymatic substitution of the *N*<sup>6</sup>-methyl group (20), or selective dTTP (deoxythymidine triphosphate) analog such as 4SedTTP (21). Along with the rapid progress of single-molecule se-

quencing technologies, Oxford Nanopore Technologies sequencing platform is able to detect modifications on model RNA (22), although the systematic error prevents the practical application in biological samples. Nevertheless, comprehensive interrogating of m<sup>6</sup>A at the transcriptome level is pivotal to reveal the biological importance of this mRNA modification. Methylated RNA immunoprecipitation and sequencing (MeRIP-seq or m<sup>6</sup>A-seq) has been widely used to profile m<sup>6</sup>A, identifying approximate region with m<sup>6</sup>A in ~100 nucleotide (nt) length, while the exact location of individual m<sup>6</sup>A site remains undetermined (12, 13). Many refined methods have been developed to improve resolution, such as PA-m<sup>6</sup>A-seq, miCLIP, and m<sup>6</sup>A-CLIP (23–26). However, all of these methods are dependent on m<sup>6</sup>A-specific antibodies, suffering from poor reproducibility and complicated process. In addition, affinity variation and batch effects of antibodies make it difficult to quantify the methylation level (27). Therefore, a convenient and single-base resolution method is still needed for whole-transcriptome m<sup>6</sup>A identification and quantification, advancing the comprehension of m<sup>6</sup>A for its dynamics and cellular functions in posttranscriptomic regulation.

Several DNA endonucleases belonging to the restriction-modification system have demonstrated sensitivity to DNA methylation. For instance, Dpn I specifically cleaves methylated sites, while Dpn II is blocked by the same modified sequence. This feature has been adopted in genome-wide studies to detect the DNA 6mA modification in single-base resolution (6mA-RE-seq/DA-6mA-seq) (28). For RNA endoribonucleases, a large group of sequence-specific enzymes belonging to the bacterial type II toxin-antitoxin system has been found with determined cleavage motifs. Recently, an *Escherichia coli* toxin and RNA endoribonuclease, MazF, was reported to be sensitive to m<sup>6</sup>A modification within ACA motif, specifically cleaving the unmethylated ACA motif, leaving methylated (m<sup>6</sup>A)CA motifs intact (29). By screening the endoribonuclease pool, we identified ChpBK as another m<sup>6</sup>A-sensitive endoribonuclease, which can discriminate m<sup>6</sup>A-modified motifs from unmodified sequences. Taking advantage of the m<sup>6</sup>A-sensitive endoribonucleases, we developed m<sup>6</sup>A-sensitive RNA-endoribonuclease-facilitated sequencing method or m<sup>6</sup>A-REF-seq, which can identify transcriptomic m<sup>6</sup>A sites at specific motifs

Copyright © 2019  
The Authors, some  
rights reserved;  
exclusive licensee  
American Association  
for the Advancement  
of Science. No claim to  
original U.S. Government  
Works. Distributed  
under a Creative  
Commons Attribution  
NonCommercial  
License 4.0 (CC BY-NC).

<sup>1</sup>MOE Key Laboratory of Gene Function and Regulation, State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou, Guangdong, China. <sup>2</sup>Laboratory of Chemical Biology, State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, China. <sup>3</sup>Department of Chemistry and Department of Biochemistry and Molecular Biology, Institute for Biophysical Dynamics, Howard Hughes Medical Institute, University of Chicago, Chicago, IL, USA.

\*Corresponding author. Email: luogzh5@mail.sysu.edu.cn

and quantify the methylation level in single-base resolution. To validate the m<sup>6</sup>A sites identified by this method, we used a ligation-based method testing individual sites and confirmed the high-reliability results. Further analysis revealed distribution pattern of m<sup>6</sup>A in single-nucleotide resolution. Last, we applied m<sup>6</sup>A-REF-seq to five tissues from human, mouse, and rat, revealing the conservation of m<sup>6</sup>A at both single-base and regional levels across diverse tissues and species.

## RESULTS

### Identification and quantification of m<sup>6</sup>A by endoribonuclease


The application of methylation-sensitive DNA endonuclease in genome-wide 6mA identification inspired us to inspect the possibility of finding similar tools for m<sup>6</sup>A determination (28). To discover endoribonuclease with m<sup>6</sup>A sensitivity, we expressed and screened the candidate enzymes by testing the cleavage capacity to synthetic RNA oligonucleotides (RNA oligos) with or without m<sup>6</sup>A. Two enzymes, MazF and ChpBK, both belonging to the bacterial toxin-antitoxin system, were able to distinguish m<sup>6</sup>A from unmethylated A. MazF recognized and cleaved the motif sequence ACA from the 5' side of first A, leaving the methylated (m<sup>6</sup>A)CA motif intact (Fig. 1A). The cleavage activity of ChpBK was also blocked by m<sup>6</sup>A at the recognition motif UAC (Fig. 1B). Synthetic RNA oligos with different m<sup>6</sup>A/A ratios were used to simulate the partially methylated sites on mRNA. The cleavage assay demonstrated that the fraction of digested

oligo versus intact part was proportional to the methylation ratio, implying that MazF has the potential to quantify the methylation level (Fig. 1, C and D, and fig. S1).


The high sensitivity and specificity of selected endoribonuclease that enzymatically distinguished m<sup>6</sup>A urged us to develop a new high-throughput method to identify single-base m<sup>6</sup>A at the transcriptome level. Since the MazF recognition motif ACA is more prevalent than that of ChpBK among known m<sup>6</sup>A sites, we chose MazF in the following studies. In general, mRNA from human embryonic kidney (HEK) 293T cells was digested into fragments by MazF. After end repair and purification, the mRNA fragments were ligated to next-generation sequencing (NGS) adaptors and amplified by polymerase chain reaction (PCR). To prevent potential false positives caused by unknown factors, we also included a parallel sample batch demethylated by fat mass and obesity-associated protein (FTO) in advance as the negative control (fig. S2). We call this method m<sup>6</sup>A-REF-seq, which stands for m<sup>6</sup>A-sensitive RNA-Endoribonuclease-Facilitated sequencing (Fig. 1E).

### Transcriptome-wide identification of m<sup>6</sup>A sites in single-base resolution

We developed a pipeline to analyze the high-throughput sequencing data generated by m<sup>6</sup>A-REF-seq (Fig. 2A). As we anticipated, most reads contained the ACA at the 5' terminus, in line with the effect of MazF treatment (fig. S3). The intact ACA motif sequence observed internally within an RNA fragment was supposed to be



**Fig. 1. m<sup>6</sup>A identification method based on m<sup>6</sup>A sensitive RNA endoribonuclease.** (A) Validation for the methylation sensitivity of endoribonuclease MazF by synthetic m<sup>6</sup>A-containing RNA oligonucleotide. (B) Validation for the methylation sensitivity of endoribonuclease ChpBK. (C) Various ratios of m<sup>6</sup>A-containing oligo mixed with unmethylated oligo digested by endoribonuclease. The m<sup>6</sup>A/A ratios of synthetic oligo are indicated to imitate practical RNAs with different m<sup>6</sup>A ratios. (D) Relative grayscale of proportionally mixed RNA oligo digested by endoribonuclease. (E) The schematic diagram of m<sup>6</sup>A-REF-seq.



**Fig. 2. Transcriptome-wide distribution of m<sup>6</sup>A revealed by m<sup>6</sup>A-REF-seq.** (A) The analysis pipeline of m<sup>6</sup>A-REF-seq data. (B) The snapshot of sequencing reads shows a known m<sup>6</sup>A site in 18S rRNA. (C) Overlap sites among three replicates after removing RNA secondary structure. (D) The overall shift of methylation ratio after FTO treatment. The value of methylation change indicates the methylation ratio of each m<sup>6</sup>A site in MazF minus that after FTO treatment. (E) Transcriptome-wide distribution of m<sup>6</sup>A. Pie chart shows the percentages of m<sup>6</sup>A sites located within CDS, 5'UTR, and 3'UTR. (F) Single-base m<sup>6</sup>A sites from m<sup>6</sup>A-REF-seq show a typical transcriptome-wide distribution pattern of m<sup>6</sup>A. (G) Overlap of m<sup>6</sup>A sites to the m<sup>6</sup>A peaks identified by antibody-based method. (H) The proportion of motifs containing m<sup>6</sup>A sites. The red square includes the RRACA motif and the green square includes the DRACA motif.

methylated. We verified this using the known m<sup>6</sup>A site with ACA motif on the 18S ribosomal RNA (rRNA) as an example. The unmethylated rRNA fragments were cleaved on the ACA motif, and the methylated fragments were retained (Fig. 2B). Next, we scanned each ACA motif within the transcriptome and interrogated the methylation sites by calculating the reads harboring intact internal ACA motifs. To further eliminate the potential false positive caused by RNA structure, we predicted the probability of each RNA fragment forming secondary structure according to minimum free energy (30) and removed the candidate sites tending to reside in double-stranded regions. We performed three biological replicates and retained the recurrent candidate sites detected in at least two replicates (Fig. 2C and table S1). For each site, the ratio of sequence reads with internal ACA motif versus reads split at the end represents the relative methylation ratio. We compared the methylation ratio of each candidate site in MazF treatment to the sample treated with

FTO in advance, which was considered as demethylated RNA. Most of m<sup>6</sup>A sites had marked reductions in methylation level with FTO treatment (Fig. 2D). Only candidate sites with a substantial decrease of methylation ratio in FTO treatment were considered as accurate m<sup>6</sup>A sites. In total, we identified 4260 high-confidence m<sup>6</sup>A sites for further studies.

About 65% of these sites were located in the coding sequence (CDS) region, and others were in untranslated regions (UTRs) (Fig. 2E). The transcriptome-wide distribution showed strong enrichment surrounding the stop codon (Fig. 2F), in line with the typical pattern found by antibody-dependent methods such as MeRIP-seq or m<sup>6</sup>A-seq. To compare the m<sup>6</sup>A location with MeRIP-seq in detail, we downloaded the profiling data deposited in Gene Expression Omnibus (GEO) database (GSE29714) (12) and calculated the cumulated reads distribution of input and IP (immunoprecipitation) samples within the range of -400 to 400 nt flanking to every m<sup>6</sup>A site identified

by m<sup>6</sup>A-REF-seq. The reads from IP samples were markedly concentrated in this region, with a peak summit at almost the exact position of individual m<sup>6</sup>A sites; whereas reads from input sample did not show any enrichment, implying that the m<sup>6</sup>A sites identified by m<sup>6</sup>A-REF-seq coincided well with the m<sup>6</sup>A peaks reported by MeRIP-seq (Fig. 2G). By expanding the context sequences flanking m<sup>6</sup>A sites, we found that GGACA was the most enriched consensus motif. More generally, the consensus motifs DRACA and RRACA were overrepresented, accounting for 56 and 45% of all identified m<sup>6</sup>A sites, respectively (Fig. 2H). These results were in concordance with previous studies, in which DRACH/RRACH was the most preferred m<sup>6</sup>A motif in native mRNA (12, 13, 24).

### High reliability of individual m<sup>6</sup>A sites identified by m<sup>6</sup>A-REF-seq

To show the reliability of m<sup>6</sup>A-REF-seq as a new method, we sought another independent principle to validate individual m<sup>6</sup>A sites. The T3 ligase was reported to be sensitive to m<sup>6</sup>A sites during the ligation reaction, which had been applied practically to determine m<sup>6</sup>A sites in mRNA (17). We adopted this method and designed the probe L (left) and probe R (right) based on the sequence flanking the exact site (table S2) and then used the T3 ligase to concatenate the two probes to an integrated template, which could be amplified by PCR. According to previous result, the ligation efficiency at m<sup>6</sup>A site was significantly suppressed comparing to that at unmethylated A sites (17), and PCR amplification magnified the difference of ligation efficiency (fig. S4). Thus, the amount of PCR products could be assessed on gel to represent the ligation efficiency, implying the methylation status of the interrogated site (Fig. 3A). We also applied the same protocol on FTO-demethylated mRNA as negative control. In total, we picked nine m<sup>6</sup>A sites for validation, including eight m<sup>6</sup>A sites and one unmethylated site. Six of eight m<sup>6</sup>A sites were validated by the ligase-based method, while the unmethylated site was also confirmed (Fig. 3B and fig. S5). To the best of our knowledge, this is the first batch of individual m<sup>6</sup>A sites confirmed by two independent methods. One validated m<sup>6</sup>A site (chr1: 29070177; Fig. 3B) is located at 343 nt upstream from the stop codon of m<sup>6</sup>A reader protein YTHDF2, suggesting a self-regulating feedback loop mediated by RNA modification.

Two other antibody-dependent methods, m<sup>6</sup>A-CLIP and miCLIP, had reported comprehensive m<sup>6</sup>A maps at transcriptome level. We checked whether the six individual m<sup>6</sup>A sites could be identified by these two methods. Peaks from MeRIP-seq data were also included as reference. Of the six sites, one of them was not reported in either previous method. The negative control site that supposed to be unmethylated was reported by miCLIP as an m<sup>6</sup>A site but not by other methods (Fig. 3B). We suspect that differences in samples, as well as the dynamic nature of m<sup>6</sup>A methylation, contribute to these inconsistencies and potential false positives. Nevertheless, these results underline the need of independent methods to confirm authentic m<sup>6</sup>A sites for downstream analysis, especially considering m<sup>6</sup>A as a highly dynamic RNA modification.

### Quantification of m<sup>6</sup>A abundance by m<sup>6</sup>A-REF-seq

m<sup>6</sup>A-REF-seq not only determines the methylation status but also can be used to quantify the methylation level of each m<sup>6</sup>A site by calculating the ratio of sequence reads with internal ACA versus reads split at the motif. To verify this, we designed a digestion-quantification assay that used reverse transcription quantitative PCR (RT-qPCR) to evaluate the methylation level of identified m<sup>6</sup>A sites. First, mRNA was treated by

MazF. The unmethylated sites located in ACA motif were cleaved, leaving intact methylated sites. After reverse transcription, complementary DNA (cDNA) spanning the intended m<sup>6</sup>A site was amplified by PCR to estimate the relative abundance. The same amount of mRNA without MazF digestion was amplified simultaneously as a control to compare with the digested sample (Fig. 3C). We picked six m<sup>6</sup>A sites including three sites with high methylation level and three sites with low methylation level determined by m<sup>6</sup>A-REF-seq (table S3). The  $\Delta\Delta C_t$  values calculated from RT-qPCR results were comparable to the average methylation ratio determined by m<sup>6</sup>A-REF-seq (Fig. 3D and fig. S6), demonstrating that m<sup>6</sup>A-REF-seq can principally quantify the methylation fraction of m<sup>6</sup>A sites.

### Characterizing m<sup>6</sup>A distribution pattern in single-nucleotide resolution


Previous studies demonstrated that one of the most remarkable features of m<sup>6</sup>A was the distribution enrichment surrounding stop codon (12, 13). However, limited by the resolution of antibody-based methods, it was hard to know the precise position relative to the exact stop codon site, which divides coding regions and 3'UTRs. The single-nucleotide m<sup>6</sup>A identification method largely increased the resolution of m<sup>6</sup>A map. Our results showed that this stop codon-concentrated pattern could be reproduced by using single-nucleotide m<sup>6</sup>A sites, without any preference to coding regions or 3'UTRs (Fig. 4A). Meanwhile, it was worth noting that consensus motifs DRACA and DRACH were also enriched near the stop codon, suggesting that sequence motif might be one factor to shape this pattern (Fig. 4A).

m<sup>6</sup>A had been proposed to be involved in various key regulatory processes of RNA metabolisms, including splicing (31). We calculated the relative distances of m<sup>6</sup>A sites to intron-exon boundaries in each exon and plotted in Fig. 4B. The result showed m<sup>6</sup>A distributed evenly along exon regions, with slight depletion toward the 3' end. By mapping single m<sup>6</sup>A sites to exons, we compared the lengths of exons containing m<sup>6</sup>A modifications versus all exons. The lengths of m<sup>6</sup>A-containing exons were significantly greater than all exons ( $P < 2.2 \times 10^{-16}$ , Wilcoxon test; Fig. 4C), in line with previous report that m<sup>6</sup>A modification usually occurred within long exons (13).

Some indirect evidence showed that m<sup>6</sup>A tended to locate in short regions, suggesting a specialized working model of m<sup>6</sup>A as clusters. However, antibody-based methods could barely divide multiple sites from one m<sup>6</sup>A peak to demonstrate clustering. Using the single-nucleotide m<sup>6</sup>A positions identified by m<sup>6</sup>A-REF-seq, we calculated the distances between each two adjacent m<sup>6</sup>A sites and compared to the control dataset, which contained the same number of sites randomly sampled from transcriptome 1000 times. Our results showed that two adjacent m<sup>6</sup>A sites were statistically prone to aggregate within 200-nt regions ( $P = 6.6 \times 10^{-5}$  and  $P = 0.0065$ , Fisher's exact test), supporting the hypothesis that m<sup>6</sup>A functions as clusters of modification (Fig. 4D).

### Conservation of m<sup>6</sup>A sites in mammals

We next applied m<sup>6</sup>A-REF-seq to identify m<sup>6</sup>A modifications in different tissues of human, mouse, and rat and identified more than 3500 individual m<sup>6</sup>A sites in each sample (tables S4 and S5). The transcriptomic distribution displayed similar patterns among tissues in three species (Fig. 5A and figs. S7 to S9). To examine the conservation of m<sup>6</sup>A at the individual gene level, we first collected 13,117 one-to-one orthologs in three species. By mapping the m<sup>6</sup>A sites to these ortholog genes in each type of tissues, we identified 2439




**Fig. 3. Single-base validation using ligation-amplification method and qPCR.** (A) Schematic diagram of ligation-amplification method for single-base m<sup>6</sup>A validation. (B) Validation results of six individual sites. Five of the sites are validated to be m<sup>6</sup>A sites, whereas the other one is confirmed to be not modified. Data for m<sup>6</sup>A-CLIP, miCLIP-CITS, miCLIP-CIMS, and MeRIP-seq are downloaded from published literatures. (C) Schematic diagram of qPCR to quantify the methylation level of a specific m<sup>6</sup>A site. The undigested mRNA sample is treated as control. (D) qPCR results and methylation ratios of six m<sup>6</sup>A sites. H1 to H3 represent the highly methylated sites (>0.75), while L1 to L3 represent the weakly methylated sites (<0.35). The left y axis represents the  $\Delta\Delta Ct$  values, and the right y axis represents the methylation ratio determined by m<sup>6</sup>A-REF-seq. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

human genes, 3596 mouse genes, and 1343 rat genes in the brain, among which 571 of them were shared in three species. The numbers of shared genes were 536 and 262 in the kidney and the liver, respectively (figs. S8 and S9). The ortholog genes modified by m<sup>6</sup>A in all three species were significantly more than expected even after adjusting for gene expression (Fig. 5B, fig. S10, and table S6), indicating a well-conserved distribution of m<sup>6</sup>A at the gene level.

Similarly, previous studies have reported that the count of shared m<sup>6</sup>A peaks between human and mouse were significantly higher than expected, proposing the conservation and important functions of m<sup>6</sup>A modification (13).

Since m<sup>6</sup>A-REF-seq identified high-confidence sites, we were able to investigate the conservation of m<sup>6</sup>A at single-nucleotide level. Previous work analyzed the evolutionary conservation of yeast and



**Fig. 4. Single-base method reveals high resolution features of m<sup>6</sup>A.** (A) The distance of individual m<sup>6</sup>A sites to the stop codons. DRACA and DRACH background motifs are extracted from the transcriptome. (B) The relative position of m<sup>6</sup>A sites within exons. (C) The length distribution of m<sup>6</sup>A-containing exons versus all exons in m<sup>6</sup>A-modified genes. (D) The distances between two m<sup>6</sup>A sites are significantly enriched within 200 bp regions compared to random sampling (Fisher's exact test, \*\* $P < 0.01$ ).

human m<sup>6</sup>A sites in protein-coding regions and concluded that m<sup>6</sup>A were overall not conserved in yeasts and only slightly conserved in mammals (32). However, the m<sup>6</sup>A sites used in this work were predicted based on the consensus motif DRACH, which could contain underestimated false positives. Therefore, we mapped the single-nucleotide m<sup>6</sup>A sites identified by m<sup>6</sup>A-REF-seq to orthologous genes in three species (table S6). By using similar method (32), we tested whether m<sup>6</sup>A sites were more conserved than unmethylated A. We randomly sampled the human unmethylated A sites as the same number of m<sup>6</sup>A sites in each orthologous gene and tested whether the corresponding sites in mouse and rat remaining to be base A (fig. S11). A total of 1854 to 3722 m<sup>6</sup>A sites were compared to the same number of unmethylated A sites from 1286 to 2306 genes across three tissues. The results indicated that m<sup>6</sup>A sites were more conserved than unmethylated A sites in brain, kidney, and liver tissues for all three codon positions ( $P = 0.002$ , 0.021, and 0.075, respectively; figs. S11 to S13), confirming the m<sup>6</sup>A conservation at single-nucleotide level.


Furthermore, we tested whether the nonpairwise m<sup>6</sup>A sites tended to cluster in narrow corresponding regions among species. The shortest distances between pairwise m<sup>6</sup>A sites in ortholog genes were averaged (Fig. 5C). Random sampling of the same number of ACA motifs were performed for 1000 times as control. The average distances between pairwise m<sup>6</sup>A sites were shorter than the control (Fig. 5D, and figs. S8 and S9), indicating that, although some m<sup>6</sup>A

sites were not conserved at single-nucleotide level, they tended to cluster within conserved short regions across multiple species.

More extensively, we computed the PhastCons scores representing the conservation of each nucleotide among multiple species (33). Identified m<sup>6</sup>A sites in the brain were attributed and classified into sites conserved in three species or sites residing in ortholog genes. As expected, conserved m<sup>6</sup>A sites at single-base level and sites conserved at gene level had highest conservation scores (medians 0.991 and 0.970, respectively). The scores of overall m<sup>6</sup>A sites were also distinctively higher than that of all A sites in ACA motifs with the medians 0.791 and 0.338, respectively ( $P < 9 \times 10^{-10}$ , Wilcoxon test; Fig. 5E). Similar results hold for other tissues in human, mouse, and rat (figs. S8, S9, and S14).

## DISCUSSION

Antibody-dependent pull-down methods are widely used to profile transcriptomic m<sup>6</sup>A, the most prevalent modification in mRNA. An antibody-independent, easy-to-use method is strongly desired to cross-validate the known m<sup>6</sup>A sites and study the dynamic functions in single-nucleotide resolution. We developed m<sup>6</sup>A-REF-seq, an antibody-independent, high-throughput, and single-base m<sup>6</sup>A detection method based on m<sup>6</sup>A-sensitive RNA endoribonuclease. High sensitivity and specificity of endoribonuclease that we found to discriminate m<sup>6</sup>A modification, as well as stringent data process



**Fig. 5. Conservation of m<sup>6</sup>A in mammals.** (A) Metagenome plots of m<sup>6</sup>A in the brain of human, mouse, and rat. (B) Shared m<sup>6</sup>A-modified genes among three species. (C) Diagram showing the m<sup>6</sup>A sites conserved in the corresponding short regions from different species. (D) Frequency of distances for pairwise m<sup>6</sup>A in brain. Randomly picked ACA motifs are assigned for the same analysis as control. (E) Conservation scores of all m<sup>6</sup>A sites, methylation sites in ortholog genes, and conserved m<sup>6</sup>A sites are compared to that for all A sites in ACA motifs ( $P < 9 \times 10^{-10}$ , Wilcoxon test).

by removing potential RNA secondary structure and the introducing of FTO demethylation control, ensured reliability of m<sup>6</sup>A sites identified by this method. Transcriptome-wide m<sup>6</sup>A maps in human HEK293T cell and mammalian tissues displayed similar distribution pattern to previous antibody-based pull-down methods. To validate the m<sup>6</sup>A sites identified by m<sup>6</sup>A-REF-seq, we introduced another independent method, which used T3 ligase to differentiate individual m<sup>6</sup>A or A at template RNA. The high consistency between different methods demonstrated the efficiency of m<sup>6</sup>A-REF-seq with high confidence. We also applied qPCR to show that m<sup>6</sup>A-REF-seq could quantify the methylation level of individual m<sup>6</sup>A site.

The RNA endoribonuclease MazF has high sensitivity and specificity for RNA substrates. It works very well on extremely limited substrate, as little as nanograms or even picograms. Rapid and simplified experiment design, without the need for antibody-enrichment step, largely reduces the requirement of starting RNA amount and sample preparation time. Therefore, this method could be applied to the studies for rare biological materials such as samples from pathological tissues or early embryos. In addition, m<sup>6</sup>A-REF-seq offers the potential to capture the subtle changes of m<sup>6</sup>A during metabolic processes, advancing the dynamic studies of this post-transcriptional RNA modification in different life stages.

By screening RNA endoribonucleases, we found two m<sup>6</sup>A-sensitive enzymes from the bacterial type II toxin-antitoxin system, which recognized specific motifs “ACA” and “UAC,” respectively. To cover

all the RRACH motifs in the transcriptome, we need new enzyme(s) capable of recognizing more universal sequence motifs while retaining the m<sup>6</sup>A sensitivity. On the basis of our preliminary screening, we speculated that the m<sup>6</sup>A sensitivity might be a general feature of endoribonucleases belonging to type II TA system. Further exploring in the endoribonucleases pool would be helpful to expand the toolbox for RNA modification studies. Another strategy is to transform the existing enzyme to adapt for new recognition sequences while retaining the methylation sensitivity. We studied the structure of MazF protein and mutated the 56th lysine to alanine. The MazF-K56A mutant showed irregular recognition sequences not meeting our anticipation, although the methyl sensitivity was retained (fig. S15). On the other hand, directed evolution would be a promising way to expand the recognition sequence scope of MazF or other related proteins, as long as dedicated screening system was created.

We applied m<sup>6</sup>A-REF-seq to five different tissues from human, mouse, and rat. The single-base m<sup>6</sup>A maps displayed a distribution pattern similar to previous findings that m<sup>6</sup>A tended to enrich surrounding stop codons. This unique pattern seemed to be conserved among different species. However, because of limited resolution, previous methods could not tell to what extent the modification is conserved. In this study, we mapped the m<sup>6</sup>A sites to orthologous genes in these three species and compared the conservation score at single-nucleotide level. We found that significantly more m<sup>6</sup>A-modified genes were shared among species than expected, and a large group

of m<sup>6</sup>A sites could be aligned to the syntenic regions, showing that the m<sup>6</sup>A modification was conserved at both the gene and individual base level. The well-conserved feature restated the biological importance of m<sup>6</sup>A, implying other uncovered roles of this RNA modification.

Together, m<sup>6</sup>A-REF-seq provides a new perspective for single-base m<sup>6</sup>A identification at the transcriptome level. The new principle eliminates the need for m<sup>6</sup>A-specific antibodies, making it unique and amenable for future applications in small and precious samples. The features of this efficient and convenient method will largely expand the scope of m<sup>6</sup>A studies.

## MATERIALS AND METHODS

### RNA materials

The RNA oligos synthesized from Takara Bio Inc. and Sangon Biotech were as follows: RNA-m<sup>6</sup>ACA (5'-UUGGUUUUUUUUGG(m<sup>6</sup>A)CAUGUAUAUAGU-3'), RNA-ACA (5'-UUGGUUUUUUUUG-GACAUGUAUAUAGU-3'), RB1-Um<sup>6</sup>AC (5'-GUUGUGUGAUU(m<sup>6</sup>A)-CAUAUGGUGUG-3'), and RB1-UAC (5' GUUGUGUGAUUACAUAUGGUGGUG-3').

HEK293T cell line was cultured with Dulbecco's modified Eagle's medium (DMEM; Corning) supplemented with 10% (v/v) fetal bovine serum (FBS; Gibco) at 37°C in the presence of 5% CO<sub>2</sub> and tested to be mycoplasma negative. Total RNA from HEK293T cells was extracted using TRIzol reagent (Thermo Fisher Scientific), and the mRNA was purified using the Dynabeads mRNA Purification Kit (Thermo Fisher Scientific, catalog no. 61006). The mRNA of human tissues (brain, liver, and kidney), mouse tissues (brain, liver, heart, testis, and kidney), and rat tissues (brain, liver, and kidney) were purchased from Takara Bio Inc.

### RNA endoribonuclease validation for MazF

RNA oligos (10 pmol) were incubated with 2.5 U of MazF (mRNA intereferase MazF, Takara Bio Inc., code no. 2415A) in the 20- $\mu$ l reaction mixture of MazF buffer [40 mM sodium phosphate (pH 7.5) and 0.01% Tween 20] at 37°C for 30 min. The RNA oligos were mixed with the m<sup>6</sup>A percentage of 0, 20, 40, 60, 80, and 100%, and the oligo mixtures were incubated with 2.5 U of MazF in the 20- $\mu$ l reaction mixtures. The samples were loaded into 15% urea polyacrylamide gel and electrophoresed in 0.5 $\times$  tris-boric acid-EDTA (TBE) buffer.

The FTO demethylation reaction was conducted in the reaction mixture that contained 10 pmol of m<sup>6</sup>A RNA oligo, 0 to 10  $\mu$ g of FTO demethylase, 283  $\mu$ M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O, 300  $\mu$ M  $\alpha$ -ketoglutarate ( $\alpha$ -KG), 2 mM L-ascorbic acid, 20 U of RNase inhibitor (Takara Bio Inc., code no. 2313A), and 50 mM tris-HCl buffer (pH 7.5). The reaction was stopped by 40 mM EDTA after 3 hours of incubation at room temperature. The MazF cleavage reaction was performed to test the efficiency of FTO demethylation.

### Dot-blot assay of FTO demethylation

m<sup>6</sup>A FTO demethylase with different concentrations (0, 0.25, and 2.5  $\mu$ g) was used in the dot-blot assay to test the demethylation of RNA. The total RNA was incubated with FTO, 283  $\mu$ M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O, 300  $\mu$ M  $\alpha$ -KG, and 2 mM L-ascorbic acid for 3 hours at room temperature. After purification by an RNA Clean & Concentrator-5 kit (Zymo Research), the concentration of total RNA was determined by the Qubit RNA HS Assay Kit (Thermo Fisher Scientific) and diluted to 10- $\mu$ l volume with 500 ng. Spotting

and thermal cross-linking were carried out on a metal base plate preheated to 70°C. After thermal cross-linking for 15 min, ultraviolet cross-linking was conducted twice with 2000 kJ. Then, 3% bovine serum albumin (BSA) was used to block at room temperature for 1 hour. The anti-m<sup>6</sup>A antibody (Synaptic Systems, catalog no. 202003) was diluted to 1:2000 and incubated with the nylon membrane for 2 hours. The nylon membrane was washed for three times with 1% tris-buffered saline with Tween 20 (TBST) and then incubated with the 1:5000 diluted anti-rabbit antibody for 1 hour. The membrane was washed for five times with 1% TBST. After draining off the water, the chemiluminescence detection signal was performed using the enhanced chemiluminescence (ECL) system. The membrane was washed twice with 1% TBST. After draining off the water, it was dyed with methylene blue solution for 8 min and then washed with 1% TBST. The signal was detected by a white light system.

### Expression and purification of recombinant ChpBK protein

To express recombinant ChpBK protein [National Center for Biotechnology Information (NCBI reference sequence: NP\_418646.1), *E. coli* BL21(DE3) strain carrying PET-28a-chpBK plasmid was grown in lysogeny broth (LB) medium (Sangon Biotech, A507002) supplemented with Kan (10  $\mu$ g/ml) at 37°C and 200 rpm until the optical density at 600 nm (OD<sub>600</sub>) reach to 0.5. Isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) was added to a final concentration of 0.4 mM, and cell growth resumed for 2 hours under the same conditions. Harvested cell pellets were lysed by sonication in 30 ml of buffer A [10 mM bis-tris (pH 7.0) and 30 mM imidazole] and centrifuged for 20 min at 16,000g and 4°C. The soluble fraction was incubated on ice for 30 min in the presence of DNase and RNase (10 g/ml; Thermo Fisher Scientific). The His-ChpBK fusion protein was purified from clarified lysates using Ni-NTA (nickel-nitrilotriacetic acid) beads (Invitrogen). Protein purity was determined by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) analysis, and samples were quantified using the Bio-Rad protein assay and stored at -80°C.

### Expression and purification of MazF-K56A in *E. coli*

The 56th lysine of MazF protein was mutated to alanine. The protein sequence was shown below, where the underlined bold "A" indicates the mutated amino acid: MazF-K56A (MVSRYVPDMGLIIVDFDPTK GSEQA GHRPA VVLS PFM YNNKTGMCLCVPCTTQSA **A**GYPFEVVLGQERDGVALADQVKSIAWRARGATKKGTVAPPEELQLIKAKINVLIG).

The plasmid DNA pCOLD-MazF was transformed into *E. coli* Rossetta strain, and a single colony was picked by sterilized loop and inoculated into a 50-ml culture (containing ampicillin) in sterile culture flask. *E. coli* was cultured at 37°C until OD<sub>600</sub> reached 0.6 to 0.8. IPTG (1 mM) was added to induce protein expression, and cells grew overnight at 18°C. Cells were harvested by centrifugation and suspended in 200 ml of lysis buffer [50 mM tris (pH 8.0), 350 mM NaCl, and 10 mM imidazole]. Cell pellets were lysed by sonication, and the supernatant was collected by centrifugation and applied to nickel affinity purification using HisTrap HP. The column was washed by lysis buffer to remove nonspecific binding protein, and target protein was eluted with linear gradient of elution buffer [50 mM tris (pH 8.0), 350 mM NaCl, and 500 mM imidazole]. We concentrated the protein and changed protein storage buffer to ion-exchange binding buffer [20 mM Hepes and 50 mM NaCl (pH 7.5)] by HiTrap Desalting column. Tobacco etch virus (TEV)



protease was used to remove the His-tag, and target protein was loaded to HiTrap HP Q column. Target protein was eluted by linear gradient of ion-exchange elution buffer [20 mM Hepes and 1 M NaCl (pH 7.5)]. The concentrated protein was analyzed by SDS-PAGE.

### RNA endoribonuclease validations for ChpBK and MazF-K56A

The digestion reaction of ChpBK was conducted in a 20- $\mu$ l reaction mixture with 10 pmol of RNA oligo (RB1), 1  $\mu$ g of ChpBK, 20 U of RNase inhibitor, and 10 mM tris-HCl buffer (pH 7.5). The mixture was incubated at 37°C for 30 min. The digestion reaction of ChpBK with different m<sup>6</sup>A percentage was conducted as described above. The MazF-K56A cleavage reaction was performed in a 20- $\mu$ l reaction mixture with 10 pmol of RNA oligo, 2  $\mu$ g of MazF-K56A, and 1 $\times$  MazF buffer [40 mM sodium phosphate (pH 7.5) and 0.01% Tween 20] at 37°C for 30 min. All samples were loaded on the 15% urea polyacrylamide gel and electrophoresed in 0.5 $\times$  TBE buffer.

### Library construction for NGS

mRNA was first heated at 85°C for 5 min and chilled on ice for 2 min to denature RNA secondary structure. The FTO demethylation reaction was conducted before MazF reaction in a 20- $\mu$ l reaction mixture containing 100 to 200 ng mRNA, 2.5  $\mu$ g of FTO demethylase, 283  $\mu$ M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O, 300  $\mu$ M  $\alpha$ -KG, 2 mM L-ascorbic acid, 20 U of RNase inhibitor, and 50 mM tris-HCl buffer (pH 7.5). After incubating at room temperature for 3 hours, the reaction was stopped by adding 40 mM EDTA. To prevent the influence of FTO buffer, the MazF treatment was supplemented with the same reaction mixture except FTO protein. The MazF and FTO-MazF samples were heated at 85°C for 5 min, chilled on ice for 2 min, and then incubated with 10 U of MazF in the MazF reaction buffer at 37°C for 30 min. The fragmented mRNA was purified by an RNA Clean & Concentrator-5 kit (Zymo Research) and eluted into 20  $\mu$ l of RNase-free water. The eluted RNA fragment was end repaired by the T4 polynucleotide kinase (T4PNK; Vazyme Biotech) in 50- $\mu$ l reaction mixture with 1 $\times$  T4PNK buffer and incubated at 37°C for 30 min. After purification by an RNA Clean & Concentrator-5 kit (Zymo Research), the concentration of fragmented mRNA was measured by the Qubit RNA HS Assay Kit (Thermo Fisher Scientific). The NGS library was constructed by using the VAHTS Small RNA Library Prep Kit (Vazyme Biotech, NR801) with 10 ng of fragmented mRNA and amplified by PCR for 15 cycles. Libraries were loaded on the 8% urea polyacrylamide gel and electrophoresed in 0.5 $\times$  TBE buffer, and purified libraries were sequenced on the Illumina HiSeq X10 platform. Three replicates have been conducted for mRNA of HEK293T cell line. The same protocol was also conducted for mammalian tissues.

### NGS data analyses

First, quality control was performed on raw sequencing reads by the open software FastQC ([www.bioinformatics.babraham.ac.uk/projects/fastqc/](http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)). Then, adaptors were filtered by cutadapt (34) with at least 15 nt remaining length of paired-end reads. The clean reads were mapped to the reference genome (hg19) using HISAT2 (35) with default parameters, except reporting only one result. Integrative Genomics Viewer was used for data visualization (36). We scanned the motif ACA (and the reverse complementary motif TGT) on the reference genome and mapping result file, respectively, counting the number of ACA/TGT at the internal or terminal of a sequence read. As short sequence reads with lengths less than 15 nt were filtered,

only ACA/TGT with the distance from the end of a read more than 15 nt was treated as “internal” of a read. Each ACA/TGT motif on the reference genome, which has more than 10 reads supporting the motif located internal of reads, was treated as a candidate m<sup>6</sup>A site. The ratio of sequence reads with internal ACA versus reads split at the end represents the relative methylation ratio of each site. Candidate sites with repeated or continuous ACA motifs were removed.

As the RNA secondary structure could affect the reaction efficiency, we predicted the probability of each RNA fragment forming secondary structure and removed the candidate sites tending to reside in double-stranded regions. The RNAfold program of ViennaRNA package (30) was used to predict intramolecular secondary structure of each read, which supported the candidate sites. The condensed representation of the pair probabilities of each nucleotide was parsed according to the tutorials of ViennaRNA package. The pairing probability value of NNACA motif was calculated, and candidate sites with high pairing probability were discarded. The FTO demethylation treatment was also treated as a negative control. Candidate sites with at least 10% methylation ratio decrease were remained as m<sup>6</sup>A sites. The metagene plot of m<sup>6</sup>A sites, which was described the relative location of each site on the 3'UTR, CDS, and 5'UTR of mRNA, was calculated and plotted as well. The raw reads of MeRIP-seq datasets were downloaded from Meyer *et al.* (12) (GSE29714) and mapped to human (hg19) reference genome using HISAT2. The reads within the range of -400 to 400 nt flanking to every m<sup>6</sup>A site were plotted. As the sequencing data were single end, we extended the reads to 200 nt to simulate the IP peak.

### Single-base validation using ligase-based method

Refined single-base T3 ligase-based method (17) was used to validate m<sup>6</sup>A sites identified by m<sup>6</sup>A-REF-seq. The FTO demethylation sample was used as negative control (i.e., the FTO<sup>+</sup> treatment in Fig. 3A). DNA probe L and probe R were designed to match the flanking sequences of intended site. Probe L was modified with a phosphate at 5' terminal. Probe R was modified with two ribonucleotides at 3' end. The ligation reaction mixture A consisted of 20 nM probe L, 20 nM probe R, 1 $\times$  T3 ligation buffer [New England Biolabs (NEB)], and 300 ng of total RNA with or without FTO treatment. The ligation reaction mixture B consisted of appropriate amount of T3 DNA ligase (NEB) with ligation buffer. Mixture A was heated at 85°C for 3 min and then incubated at 35°C for 10 min, and then the ligation reaction mixture B was added to the final volume of 10  $\mu$ l. This mixture was incubated at room temperature for ligation reaction for 8 to 10 min and chilled on ice immediately. The volume of 1  $\mu$ l of ligation product was used to PCR amplified for 24 cycles in the 2 $\times$  Taq mix (Vazyme Biotech). The samples were loaded into the 2 to 3% agarose gel and electrophoresed in 0.5 $\times$  tris-acetate-EDTA (TAE) buffer to detect the signal. The previous m<sup>6</sup>A-CLIP and miCLIP datasets were downloaded from Ke *et al.* (25) and Linder *et al.* (24), respectively.

### qPCR validation for m<sup>6</sup>A quantification

The MazF treatment was performed in 10- $\mu$ l volume with 500 ng of mRNA and 10 U of MazF at 37°C for 30 min. The digested mRNA and nondigested mRNA were subjected to reverse transcription using 5 $\times$  TransScript All-in-One First-Strand cDNA Synthesis SuperMix for qPCR (One-Step gDNA Removal, TransGen Biotech, AT341-02). TB Green Premix Ex Taq II (Tli RNaseH Plus, Takara Bio Inc., code no. RR820A) was used on BIO-CFX96 Touch.

## Characterizing m<sup>6</sup>A distribution pattern in single-base resolution

Consensus motifs DRACA and DRACH were scanned on the reference genome, and the distances of A sites in the two motifs to stop codons were calculated and plotted as well. The relative position in exon for m<sup>6</sup>A, regular A sites in DRACA, and DRACH was computed. To estimate the distance between two adjacent m<sup>6</sup>A sites, only genes with more than two m<sup>6</sup>A sites were analyzed. We randomly sampled the same number of ACA motifs as the number of observed m<sup>6</sup>A and repeated this step for 1000 times as control. The distances between adjacent A sites in control were calculated. Fisher's exact test was used to test the significance of first two bins.

## Conservation of m<sup>6</sup>A in the same mammal tissues

m<sup>6</sup>A-REF-seq protocol was used to identify m<sup>6</sup>A sites in the different tissues of human (brain, liver, and kidney), mouse (brain, liver, heart, testis, and kidney), and rat (brain, liver, and kidney). The one-to-one orthologous genes in pairwise species of human (GRCh37.p13), mouse (GRCm38.p6), and rat (Rnor\_6.0) were downloaded from Ensembl (Ensembl genes 94). A total of 13,117 one-to-one orthologous genes among three species were retained for downstream analyses. Protein sequences of 13,117 orthologs in human, mouse, and rat were aligned by MUSCLE (37) and then converted to corresponding coding sequence alignments using PAL2NAL (38). The shared m<sup>6</sup>A-modified genes and share m<sup>6</sup>A sites among species were calculated and tested for significances based on the well-aligned orthologous coding-region sequences as mentioned above. The average pairwise distances among closely m<sup>6</sup>A sites in three species were calculated and compared with random control sample. The control sample was randomly picked for the same number of ACA motifs as the number of observed m<sup>6</sup>A for 1000 times. Analyses of conservation score were conducted by using PhastCons scores of human (hg19), mouse (mm10), and rat (rn6) downloaded from University of California, Santa Cruz (UCSC).

## SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/5/7/eaax0250/DC1>

Fig. S1. Quantitative demonstration of various fractions of m<sup>6</sup>A-containing oligo mixed with unmethylated oligo digested by ChpBK.

Fig. S2. FTO demethylate assay.

Fig. S3. The base composition of mRNA products cleaved by MazF.

Fig. S4. The single-site validation for 185 rRNA control site and m<sup>6</sup>A site.

Fig. S5. The single-site validation of eight m<sup>6</sup>A sites and one unmethylated site.

Fig. S6. Quantitative PCR results of six m<sup>6</sup>A sites.

Fig. S7. Metagene plots of m<sup>6</sup>A in mouse heart and mouse testis.

Fig. S8. Conservation of m<sup>6</sup>A in mammalian kidney.

Fig. S9. Conservation of m<sup>6</sup>A in mammalian liver.

Fig. S10. Gene expression and the expected number of shared m<sup>6</sup>A-modified genes.

Fig. S11. Evolutionary conservation between methylated and unmethylated A sites in the same genes of brain tissue.

Fig. S12. Evolutionary conservation between methylated and unmethylated A sites in the same genes of kidney tissue.

Fig. S13. Evolutionary conservation between methylated and unmethylated A sites in the same genes of liver tissue.

Fig. S14. Conservation scores for brain.

Fig. S15. Validation for the methylation sensitivity of mutated MazF-K56A.

Table S1. Basic information of sequencing data for human HEK293T cell line.

Table S2. Designed probes and universal primers for T3 ligase-based validation.

Table S3. Designed primers for Quantitative PCR validation.

Table S4. Basic information of sequencing data for mammalian tissues.

Table S5. m<sup>6</sup>A sites identified by m<sup>6</sup>A-REF-seq for mammalian tissues.

Table S6. Shared m<sup>6</sup>A-modified genes and m<sup>6</sup>A sites between pairwise species and the significance.

## REFERENCES AND NOTES

- Y. Fu, D. Dominissini, G. Rechavi, C. He, Gene expression regulation mediated through reversible m<sup>6</sup>A RNA methylation. *Nat. Rev. Genet.* **15**, 293–306 (2014).
- I. Barbieri, K. Tzelepis, L. Pandolfini, J. Shi, G. Millán-Zambrano, S. C. Robson, D. Aspris, V. Migliori, A. J. Bannister, N. Han, E. De Braekeleer, H. Pongstingl, A. Hendrick, C. R. Vakoc, G. S. Vassiliou, T. Kouzarides, Promoter-bound METTL3 maintains myeloid leukaemia by m<sup>6</sup>A-dependent translation control. *Nature* **552**, 126–131 (2017).
- P. J. Batista, B. Molinier, J. Wang, K. Qu, J. Zhang, L. Li, D. M. Bouley, E. Lujan, B. Haddad, K. Daneshvar, A. C. Carter, R. A. Flynn, C. Zhou, K.-S. Lim, P. Dedon, M. Wernig, A. C. Mullen, Y. Xing, C. C. Giallourakis, H. Y. Chang, m<sup>6</sup>A RNA modification controls cell fate transition in mammalian embryonic stem cells. *Cell Stem Cell* **15**, 707–719 (2014).
- N. S. Gokhale, A. B. R. McIntyre, M. J. McFadden, A. E. Roder, E. M. Kennedy, J. A. Gandara, S. E. Hopcraft, K. M. Quicke, C. Vazquez, J. Willer, O. R. Ilkayeva, B. A. Law, C. L. Holley, M. A. Garcia-Blanco, M. J. Evans, M. S. Suthar, S. S. Bradrick, C. E. Mason, S. M. Horne, N<sup>6</sup>-Methyladenosine in *Flaviviridae* Viral RNA Genomes Regulates Infection. *Cell Host Microbe* **20**, 654–665 (2016).
- T. Lence, J. Akhtar, M. Bayer, K. Schmid, L. Spindler, C. H. Ho, N. Kreim, M. A. Andrade-Navarro, B. Poeck, M. Helm, J.-Y. Roignant, m<sup>6</sup>A modulates neuronal functions and sex determination in *Drosophila*. *Nature* **540**, 242–247 (2016).
- D. P. Patil, C.-K. Chen, B. F. Pickering, A. Chow, C. Jackson, M. Guttman, S. R. Jaffrey, m<sup>6</sup>A RNA methylation promotes XIST-mediated transcriptional repression. *Nature* **537**, 369–373 (2016).
- Y. Xiang, B. Laurent, C.-H. Hsu, S. Nachtergaele, Z. Lu, W. Sheng, C. Xu, H. Chen, J. Ouyang, S. Wang, D. Ling, P.-H. Hsu, L. Zou, A. Jambhekar, C. He, Y. Shi, RNA m<sup>6</sup>A methylation regulates the ultraviolet-induced DNA damage response. *Nature* **543**, 573–576 (2017).
- Y. Wang, Y. Li, M. Yue, J. Wang, S. Kumar, R. J. Wechsler-Reya, Z. Zhang, Y. Ogawa, M. Kellis, G. Duester, J. C. Zhao, N<sup>6</sup>-methyladenosine RNA modification regulates embryonic neural stem cell self-renewal through histone modifications. *Nat. Neurosci.* **21**, 195–206 (2018).
- R. Winkler, E. Gillis, L. Lasman, M. Safra, S. Geula, C. Soyris, A. Nachshon, J. Tai-Schmiedel, N. Friedman, V. T. K. Le-Trilling, M. Trilling, M. Mandelboim, J. H. Hanna, S. Schwartz, N. Stern-Ginossar, m<sup>6</sup>A modification controls the innate immune response to infection by targeting type I interferons. *Nat. Immunol.* **20**, 173–182 (2019).
- L. P. Vu, B. F. Pickering, Y. Cheng, S. Zaccara, D. Nguyen, G. Minuesa, T. Chou, A. Chow, Y. Saletore, M. MacKay, J. Schulman, C. Famulare, M. Patel, V. M. Klimek, F. E. Garrett-Bakelman, A. Melnick, M. Carroll, C. E. Mason, S. R. Jaffrey, M. G. Kharas, The N<sup>6</sup>-methyladenosine (m<sup>6</sup>A)-forming enzyme METTL3 controls myeloid differentiation of normal hematopoietic and leukemia cells. *Nat. Med.* **23**, 1369–1376 (2017).
- N. Liu, M. Parisien, Q. Dai, G. Zheng, C. He, T. Pan, Probing N<sup>6</sup>-methyladenosine RNA modification status at single nucleotide resolution in mRNA and long noncoding RNA. *RNA* **19**, 1848–1856 (2013).
- K. D. Meyer, Y. Saletore, P. Zumbo, O. Elemento, C. E. Mason, S. R. Jaffrey, Comprehensive analysis of mRNA methylation reveals enrichment in 3'UTRs and near stop codons. *Cell* **149**, 1635–1646 (2012).
- D. Dominissini, S. Moshitch-Moshkovitz, S. Schwartz, M. Salmon-Divon, L. Ungar, S. Osenberg, K. Cesarkas, J. Jacob-Hirsch, N. Amariglio, M. Kupiec, R. Sorek, G. Rechavi, Topology of the human and mouse m<sup>6</sup>A RNA methylomes revealed by m<sup>6</sup>A-seq. *Nature* **485**, 201–206 (2012).
- S. Schwartz, S. D. Agarwala, M. R. Mumbach, M. Jovanovic, P. Mertins, A. Shishkin, Y. Tabach, T. S. Mikkelsen, R. Satija, G. Ruvkun, S. A. Carr, E. S. Lander, G. R. Fink, A. Regev, High-Resolution mapping reveals a conserved, widespread, dynamic mRNA methylation program in yeast meiosis. *Cell* **155**, 1409–1421 (2013).
- Y. Yue, J. Liu, C. He, RNA N<sup>6</sup>-methyladenosine methylation in post-transcriptional gene expression regulation. *Genes Dev.* **29**, 1343–1355 (2015).
- Y. Xiao, Y. Wang, Q. Tang, L. Wei, X. Zhang, G. Jia, An elongation and ligation-based qPCR amplification method for the radiolabeling-free detection of locus-specific N<sup>6</sup>-methyladenosine modifications. *Angew. Chem.* **57**, 15995–16000 (2018).
- W. Liu, J. Yan, Z. Zhang, H. Pian, C. Liu, Z. Li, Identification of a selective DNA ligase for accurate recognition and ultrasensitive quantification of N<sup>6</sup>-methyladenosine in RNA at one-nucleotide resolution. *Chem. Sci.* **9**, 3354–3359 (2018).
- E. M. Harcourt, T. Ehrenschrwender, P. J. Batista, H. Y. Chang, E. T. Kool, Identification of a selective polymerase enables detection of N<sup>6</sup>-methyladenosine in RNA. *J. Am. Chem. Soc.* **135**, 19079–19082 (2013).
- S. Wang, J. Wang, X. Zhang, B. Fu, Y. Song, P. Ma, K. Gu, X. Zhou, X. Zhang, T. Tian, X. Zhou, N<sup>6</sup>-Methyladenine hinders RNA- and DNA-directed DNA synthesis: Application in human rRNA methylation analysis of clinical specimens. *Chem. Sci.* **7**, 1440–1446 (2016).
- X. Shu, Q. Dai, T. Wu, I. R. Bothwell, Y. Yue, Z. Zhang, J. Cao, Q. Fei, M. Luo, C. He, J. Liu, N<sup>6</sup>-Allyladenosine: A New Small Molecule for RNA Labeling Identified by Mutation Assay. *J. Am. Chem. Soc.* **139**, 17213–17216 (2017).

21. T. Hong, Y. Yuan, Z. Chen, K. Xi, T. Wang, Y. Xie, Z. He, H. Su, Y. Zhou, Z.-J. Tan, X. Weng, X. Zhou, Precise antibody-independent m6A identification via 4SeDTTP-involved and FTO-assisted strategy at single-nucleotide resolution. *J. Am. Chem. Soc.* **140**, 5886–5889 (2018).
22. D. R. Garalde, E. A. Snell, D. Jachimowicz, B. Sipos, J. H. Lloyd, M. Bruce, N. Pantic, T. Admassu, P. James, A. Warland, M. Jordan, J. Ciccone, S. Serra, J. Keenan, S. Martin, L. McNeill, E. J. Wallace, L. Jayasinghe, C. Wright, J. Blasco, S. Young, D. Brocklebank, S. Juul, J. Clarke, A. J. Heron, D. J. Turner, Highly parallel direct RNA sequencing on an array of nanopores. *Nat. Methods* **15**, 201–206 (2018).
23. K. Chen, Z. Lu, X. Wang, Y. Fu, G.-Z. Luo, N. Liu, D. Han, D. Dominissini, Q. Dai, T. Pan, C. He, High-resolution N<sup>6</sup>-methyladenosine (m<sup>6</sup>A) map using photo-crosslinking-assisted m6A sequencing. *Angew. Chem. Int. Ed.* **54**, 1587–1590 (2015).
24. B. Linder, A. V. Grozhik, A. O. Orlarin-George, C. Meydan, C. E. Mason, S. R. Jaffrey, Single-nucleotide-resolution mapping of m6A and m6Am throughout the transcriptome. *Nat. Methods* **12**, 767–772 (2015).
25. S. Ke, E. A. Alemu, C. Mertens, E. C. Gantman, J. J. Fak, A. Mele, B. Haripal, I. Zucker-Scharff, M. J. Moore, C. Y. Park, C. B. Vågbo, A. Kusnierczyk, A. Klungland, J. E. Darnell Jr., R. B. Darnell, A majority of m<sup>6</sup>A residues are in the last exons, allowing the potential for 3' UTR regulation. *Genes Dev.* **29**, 2037–2053 (2015).
26. X. Li, X. Xiong, C. Yi, Epitranscriptome sequencing technologies: Decoding RNA modifications. *Nat. Methods* **14**, 23–31 (2016).
27. B. Molinie, J. Wang, K. S. Lim, R. Hillebrand, Z.-x. Lu, N. Van Wittenberghe, B. D. Howard, K. Daneshvar, A. C. Mullen, P. Dedon, Y. Xing, C. C. Giallourakis, M<sup>6</sup>A-LAIC-seq reveals the census and complexity of the m6A epitranscriptome. *Nat. Methods* **13**, 692–698 (2016).
28. G.-Z. Luo, F. Wang, X. Weng, K. Chen, Z. Hao, M. Yu, X. Deng, J. Liu, C. He, Characterization of eukaryotic DNA N<sup>6</sup>-methyladenine by a highly sensitive restriction enzyme-assisted sequencing. *Nat. Commun.* **7**, 11301 (2016).
29. M. Imanishi, S. Tsuji, A. Suda, S. Futaki, Detection of N<sup>6</sup>-methyladenosine based on the methyl-sensitivity of MazF RNA endonuclease. *Chem. Commun.* **53**, 12930–12933 (2017).
30. R. Lorenz, S. H. Bernhart, C. H. zu Siederdissen, H. Tafer, C. Flamm, P. F. Stadler, I. L. Hofacker, ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 26 (2011).
31. W. Xiao, S. Adhikari, U. Dahal, Y.-S. Chen, Y.-J. Hao, B.-F. Sun, H.-Y. Sun, A. Li, X.-L. Ping, W.-Y. Lai, X. Wang, H.-L. Ma, C.-M. Huang, Y. Yang, N. Huang, G.-B. Jiang, H.-L. Wang, Q. Zhou, X.-J. Wang, Y.-L. Zhao, Y.-G. Yang, Nuclear m6A reader YTHDC1 regulates mRNA splicing. *Mol. Cell* **61**, 507–519 (2016).
32. Z. Liu, J. Zhang, Most m<sup>6</sup>A RNA modifications in protein-coding regions are evolutionary unconserved and likely nonfunctional. *Mol. Biol. Evol.* **35**, 666–675 (2018).
33. A. Siepel, G. Bejerano, J. S. Pedersen, A. S. Hinrichs, M. Hou, K. Rosenbloom, H. Clawson, J. Spieth, L. W. Hillier, S. Richards, G. M. Weinstock, R. K. Wilson, R. A. Gibbs, W. J. Kent, W. Miller, D. Haussler, Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res.* **15**, 1034–1050 (2005).
34. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* **17**, 10–12 (2013).
35. D. Kim, B. Langmead, S. L. Salzberg, HISAT: A fast spliced aligner with low memory requirements. *Nat. Methods* **12**, 357–360 (2015).
36. H. Thorvaldsdóttir, J. T. Robinson, J. P. Mesirov, Integrative Genomics Viewer (IGV): High-performance genomics data visualization and exploration. *Brief. Bioinform.* **14**, 178–192 (2013).
37. R. C. Edgar, MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
38. M. Suyama, D. Torrents, P. Bork, PAL2NAL: Robust conversion of protein sequence alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609–W612 (2006).

#### Acknowledgments

**Funding:** This work was supported by the Ministry of Science and Technology of China (National Science and Technology Major Project, grant no. 2018YFA0109100), National Science Foundation of China (grant nos. 91753129 and 31870808), and Natural Science Foundation of Guangdong Province (grant nos. 2018B030306044). **Author contributions:** Zhang Zhang, C.H., and G.Z.L. conceived the project. Zhang Zhang, L.-Q.C., and Y.-L.Z. conducted the experiments. Zhang Zhang analyzed the NGS data. Zijie Zhang, I.A.R., C.-G.Y., J.R., and W.X. provided the supporting materials and ideas. Zhang Zhang, L.-Q.C., C.H., and G.-Z.L. wrote the manuscript. All authors reviewed the results and approved the final version of the manuscript. **Competing interests:** C.H. is a scientific founder and a member of the scientific advisory board of Accent Therapeutics, Inc. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors. Data were deposited in NCBI's GEO under accession number GSE125240.

Submitted 15 February 2019

Accepted 30 May 2019

Published 3 July 2019

10.1126/sciadv.aax0250

**Citation:** Z. Zhang, L.-Q. Chen, Y.-L. Zhao, C.-G. Yang, I. A. Roundtree, Z. Zhang, J. Ren, W. Xie, C. He, G.-Z. Luo, Single-base mapping of m<sup>6</sup>A by an antibody-independent method. *Sci. Adv.* **5**, eaax0250 (2019).