

# Wellness and Work: Mixed Messages in Residency Training

Lisa M. Meeks, PhD, MA<sup>1</sup>, Jennifer Ramsey, MD, MS<sup>2</sup>, Maureen Lyons, MD<sup>3</sup>, Abby L. Spencer, MD, MS, FACP<sup>2</sup>, and Wei Wei Lee, MD, MPH, FACP<sup>4</sup>



<sup>1</sup>Department of Family Medicine, University of Michigan Medical School, Ann Arbor, MI, USA; <sup>2</sup>Cleveland Clinic Foundation, Cleveland, OH, USA; <sup>3</sup>SSM Health Saint Louis University Hospital, St. Louis, MO, USA; <sup>4</sup>Department of Medicine, University of Chicago, Chicago, IL, USA.

Resident physicians are at higher risk for depression, anxiety, and burnout when compared with same-age peers, resulting in substantive personal and professional consequences. Training programs across the country have acknowledged the gravity of this situation and many have implemented programs and curricula that address wellness and resilience, yet the benefits of such initiatives are still largely unknown. While the development of wellness programming is well intentioned, it is often incongruent with the residency training environment. The mixed messaging that occurs when wellness programs are implemented in environments that do not support self-care may unintentionally cause resident distress. Indeed, outside of the time dedicated to wellness curricula, residents are often rewarded for self-sacrifice. In this commentary, we describe how the complexities of the medical system and culture contribute to mixed messaging and we explore the potential impact on residents. We offer recommendations to strengthen wellness programs through efforts to promote structural change in the training environment.

**KEY WORDS:** workforce; burnout; medical education-curriculum development; self-care; postgraduate medical education; wellness; resilience.

J Gen Intern Med 34(7):1352–5  
DOI: 10.1007/s11606-019-04952-5  
© Society of General Internal Medicine 2019

## INTRODUCTION

Burnout among physicians is higher than in any other field in the USA and is characterized by exhaustion, cynicism, and reduced effectiveness.<sup>1, 2</sup> Multiple studies document an upward of 50% prevalence in burnout among residents,<sup>3–5</sup> coupled with symptoms of depression and anxiety.<sup>6–9</sup> The largest contributors to burnout include time pressure, chaotic environments, a lack of control over work conditions, and a lack of shared vision between providers and administrators.<sup>2</sup> These high levels of burnout negatively impact individual residents and their patients by contributing to medical errors, lapses in professionalism, and poor patient satisfaction.<sup>10</sup> In addition, burnout is also a key contributor to substance abuse, strained relationships, attrition, and physician suicide.<sup>11–14</sup> In a 2018 commentary, leaders from the National Academy of Medicine, The Association of American Medical Colleges, and The Accreditation Council for Graduate Medical

Education declared that clinician burnout, depression, and suicide had reached a “crisis level.”<sup>15</sup>

In response to the high prevalence of burnout, training programs around the country have developed wellness initiatives to improve resident well-being.<sup>16</sup> While the preliminary data on the impact of wellness programs on resident and faculty well-being is promising, academically rigorous research on the impact of resident-specific interventions is still needed.<sup>17–20</sup> Despite efforts to enhance well-being, residents identify that finding time to focus on their own medical and self-care is one of the biggest challenges in the current learning environment.<sup>21–25</sup>

Residents are often expected to balance learning, patient care, teaching, and documentation while working 80-hour weeks for months on end. When life events related to physical or mental health, relationships, or family occur, there is little flexibility in the system to allow residents to attend to their needs.<sup>21, 26, 27</sup> Instead, residents who require time off for health or personal reasons must navigate a complex coverage system, which could include the redistribution of work to their peers. Moreover, requesting time away from work often necessitates disclosure of a mental or physical health concern or disability to their superiors, who are often the same individuals responsible for writing their letters of recommendation and advocating for them to future training programs or employers.<sup>21</sup> These conflicts leave residents hesitant to disclose these issues or to request time off for fear of a lack of confidentiality, stigma, documentation on academic record, and downstream consequences for licensure and employment.<sup>28–32</sup>

In addition, many well-meaning wellness initiatives disproportionately focus on individual factors without adequate attention to systematic contributors to burnout, despite research that points to the learning environment as a major driver of resident burnout.<sup>33–35</sup>

## MIXED MESSAGES

As a profession, our response to the burnout crisis has been primarily educational. We develop curricula to teach trainees the symptoms of burnout and offer training that focuses on personal resilience (i.e., meditation, mindfulness, etc.).<sup>22, 36</sup> While the literature suggests that meditation, mindfulness, and yoga may be protective against burnout, the efficacy of these

programs lacks robust evidence-based support.<sup>22, 37-40</sup> The expectation to engage in wellness activities in addition to their other responsibilities may place an additional burden on trainees and suggests that burnout results from residents' personal failure to engage in wellness programming.

There is emerging evidence that organizationally directed wellness interventions may be more effective in combatting burnout,<sup>37</sup> which supports the notion that burnout is a systemic issue rather than an individual one. Additionally, residents report that personal resilience training, in a system that remains dysfunctional, is ineffective, and that effective interventions to reduce burnout must include improving workplace conditions.<sup>41</sup>

### RESPONSE FROM REGULATING BODIES AND LEADING PROFESSIONAL ASSOCIATIONS

Accrediting agencies, physician and medical education associations underscore the importance of well-being in the development of the competent, caring, and resilient physician. The new Accreditation Council for Graduate Medical Education (ACGME) standards call for a residency training culture focused on physician well-being and programs are required to provide 24/7 access to urgent and emergent care and confidential mental health assessments, counseling, and treatment.<sup>42</sup> The new standards also call on programs to create formalized systems for identifying burnout and depression in residents.<sup>42</sup> In addition, the Clinical Learning Environment Review (CLER) program requires residency programs to focus on workload and work conditions to address fatigue and burnout.<sup>43, 44</sup>

Others, including the National Academy of Medicine (NAM), also identify burnout as a critical issue requiring sustained attention and action at organizational, state, and national levels and in response developed the Action Collaborative on Clinician Wellbeing and Resilience.<sup>45</sup> The American Medical Association (AMA), as part of their STEPS Forward series, offers five educational modules on professional well-being, including a burnout prevention module for residents and fellows.<sup>46</sup> While the aforementioned initiatives call for programmatic integration of wellness, they do not address the barrier of protected time to engage with recommended self-care. Unfortunately, mandated policies and organizational guidance in the absence of systemic change makes translating these policies into practice challenging.

### INSTITUTIONAL POLICIES AND PRACTICE MISMATCH

Despite the increased spotlight on physician well-being and institutional efforts to promote wellness, the culture and practice of medicine continues to send mixed messages. Residents are told that self-care is critical to avoiding burnout, yet that same messaging is absent for their faculty leadership. Working while sick, sacrificing personal relationships for work, and

forgoing self-care are the norm, not the exception.<sup>47, 48</sup> In addition, even when residency programs provide protected time off to attend health appointments, trainees and faculty physicians remain reluctant to seek mental healthcare given the stigma and potential downstream consequences.<sup>17, 42, 49, 50</sup> Faculty who are not aware of the research on physician burnout, depression, anxiety, and suicide may label trainees who engage in self-care as soft, needy, or unprepared for the real world, thereby perpetuating expectations for residents to work through challenges that might otherwise be addressed.<sup>51</sup>

### FORWARD FOCUS

Wellness programs should include a combination of personal resilience training and initiatives to address organizational issues that contribute to burnout. More research is needed to identify the systemic solutions to improve physician well-being and address burnout and must rigorously explore the impact of wellness initiatives on trainee well-being and patient outcomes.<sup>34</sup> Critical commentaries call for fundamental changes in our training programs and recommend structural changes to promote resident autonomy, strong social relatedness, adequate sleep, and time away from work.<sup>52, 53</sup> A consistent recommendation is for training programs to demonstrate their commitment to resident well-being by creating schedules with built-in time off for health and well-being appointments.<sup>52, 54-56</sup>

In addition to individualized efforts at hospitals and residency programs, systemic change will involve regulators, payers, accreditation agencies, policymakers, and patients.<sup>57, 58</sup> The path forward must first focus on systems that measure and track physician well-being as an institutional goal and create opportunities and time for physicians to engage in program development.<sup>59</sup>

Another recommendation is that health systems adequately staff their hospitals with resident and non-resident personnel and reduce documentation burden on trainees for non-educational purposes. One example of a successful intervention to reduce burnout is the University of Colorado's Ambulatory Process Excellence (APEX) team-based care model for family medicine residents, which expanded the scope of practice for medical assistants to include documentation assistance in addition to other clinical and administrative support. Researchers found that the team-based care model allowed providers more time for patient engagement, improved quality of care, and decreased physician burnout rates by 50%.<sup>60</sup>

Finally, accrediting bodies need to hold training institutions accountable for implementing and monitoring the effectiveness of wellness initiatives and ensure adequate support for leave and healthcare appointments. When policies become realistic practices, the training community sends one clear message to our medical trainees: your well-being is important.

## CONCLUSION

The medical community's response to burnout through the creation of wellness programs may inadvertently send confusing and frustrating "mixed messages." As a profession, we must operate with one voice—and model and reward the behaviors we recommend. Our words, actions, systems, and curricula must align to support choices and behaviors which seek personal and professional balance in addition to clinical and educational excellence. Until such time as our actions match our messages, the intended goals of wellness training will go unrealized.

**Corresponding Author:** Wei Wei Lee, MD, MPH, FACP, Department of Medicine, University of Chicago, Chicago, IL, USA (e-mail: wlee6@uchicago.edu).

**Prior Presentations** This was previously presented as workshops at the 2017 Society of General Internal Medicine National Meeting and the 2016 Society of General Internal Medicine Midwest Regional Meeting.

### Compliance with Ethical Standards:

**Conflict of Interest:** The authors declare that they do not have a conflict of interest.

## REFERENCES

1. Shanafelt TD, Boone S, Tan L, et al. Burnout and satisfaction with work-life balance among US physicians relative to the general US population. *Arch Intern Med.* 2012; 172(18):1377–1385.
2. Linzer M, Manwell LB, Williams ES, et al. Working conditions in primary care: physician reactions and care quality. *Ann Intern Med.* 2009; 151(1):28–36, W6–W9.
3. Dyrbye LN, West CP, Satele D, et al. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med.* 2014;89(3):443–451.
4. Desai SV, Asch DA, Bellini LM, et al. Education outcomes in a duty-hour flexibility trial in internal medicine. *N Engl J Med.* 2018;378: 1494–1508.
5. Shanafelt TD, Bradley KA, Wipf JE, Back AL. Burnout and self-reported patient care in an internal medicine residency program. *Ann Intern Med.* 2002; 136:358–367.
6. Dyrbye LN, Shanafelt T. A narrative review on burnout experienced by medical students and residents. *Med Educ.* 2016;50:132–49.
7. Mata DA, Ramos MA, Bansal N, et al. Prevalence of depression and depressive symptoms among resident physicians: a systematic review and meta-analysis. *JAMA* 2015;314: 2373–83.
8. Pereira-Lima K, Loureiro SR. Burnout, anxiety, depression, and social skills in medical residents. *Psychol Health Med.* 2015;20(3):353–62.
9. Goebert D, Thompson D, Takeshita J, et al. Depressive symptoms in medical students and residents: a multi-school study. *Acad Med.* 2009;84(2):236–241.
10. Panagioti M, Geraghty K, Johnson J, et al. Association between physician burnout and patient safety, professionalism, and patient satisfaction: a systematic review and meta-analysis. *JAMA Intern Med.* 2018;178(10):1317–1330.
11. Dyrbye LN, Massie FS Jr, Eacker A, et al. Relationship between burnout and professional conduct and attitudes among US medical students. *JAMA.* 2010;304:1173–1180.
12. Dyrbye LN, Thomas MR, Massie FS, et al. Burnout and suicidal ideation among U.S. medical students. *Ann Intern Med.* 2008;149:334–341.
13. Dyrbye LN, Thomas MR, Power DV, et al. Burnout and serious thoughts of dropping out of medical school: A multi-institutional study. *Acad Med.* 2010;85:94–102.
14. Shanafelt TD, Sloan JA, Habermann TM. The well-being of physicians. *Am J Med.* 2003;114:513–519.
15. Dzau VJ, Kirch DG, Nasca TJ. To care is human—collectively confronting the clinician-burnout crisis. *N Engl J Med.* 2018;378(4):312–4.
16. Raj KS. Well-being in residency: a systematic review. *J Grad Med Educ.* 2016;8(5):674–84.
17. Kay M, Mitchell G, Clavarino A, Doust J. Doctors as patients: a systematic review of doctors' health access and the barriers they experience. *Br J Gen Pract.* 2008;58(552):501–508.
18. Rosen IM, Christie JD, Bellini LM, Asch DA. Health and health care among housestaff in four U.S. internal medicine residency programs. *J Gen Intern Med.* 2000;15(2):116–121.
19. Campbell S, Delva D. Physician do not heal thyself. Survey of personal health practices among medical residents. *Can Fam Physician.* 2003;49:1121–1127.
20. Shadbolt NE. Attitudes to healthcare and self-care among junior medical officers: a preliminary report. *Med J Aust.* 2002;177(suppl):S19–20.
21. Meeks LM, Jain NR. Accessibility, Inclusion, and Action in Medical Education: Lived Experiences of Learners and Physicians with Disabilities. Washington, DC: Association of American Medical Colleges. March 13, 2018.
22. West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *Lancet.* 2016;388:2272–81.
23. Shiralkar MT, Harris TB, Eddins-Folensbee FF, Coverdale JH. A systematic review of stress-management programs for medical students. *Academic Psychiatry.* 2013;37(3):158–64.
24. Williams D, Tricomi G, Gupta J, Janise A. Efficacy of burnout interventions in the medical education pipeline. *Academic Psychiatry.* 2015;39(1):47–54.
25. Busireddy KR, Miller JA, Ellison K, Ren V, Gayyum R, Panda M. Efficacy of interventions to reduce resident physician burnout: a systematic review. *J Grad Med Educ.* 2017;9(3):294–301.
26. Jagsi R, Tarbell NJ, Weinstein DF. Becoming a doctor, starting a family—leaves of absence from graduate medical education. *N Engl J Med.* 2007;357(19):1889–91.
27. Grimm LJ, Nagler A, Maxfield CM. Survey of the incidence and effect of major life events on graduate medical education trainees. *Med Educ Online.* 2015;20(1):27597.
28. Gold KJ, Andrew LB, Goldman EB, Schwenk TL. "I would never want to have a mental health diagnosis on my record": a survey of female physicians on mental health diagnosis, treatment, and reporting. *Gen Hosp Psychiatry.* 2016;43:51–57.
29. Aaronson AL, Backes K, Agarwal G, Goldstein JL, Anzia J. Mental health during residency training: assessing the barriers to seeking care. *Acad Psychiatry.* 2018.
30. Givens JL, Tjia J. Depressed medical students' use of mental health services and barriers to use. *Acad Med.* 2002;77:918–21.
31. Dyrbye LN, Eacker A, Durning SJ, et al. The impact of stigma and personal experiences on the help-seeking behaviors of medical students with burnout. *Acad Med.* 2015; 90(7): 961–969.
32. Schwenk TL, Davis L, Wimsatt LA. Depression, stigma and suicidal ideation in medical students. *JAMA.* 2010; 304(11):1181–90.
33. van Vendeloo SN, Prins DJ, Verheyen CC, et al. The learning environment and resident burnout: a national study. *Perspect Med Educ.* 2018;7(2):120–125.
34. Shanafelt TD, Noseworthy JH. Executive leadership and physician well-being: nine organizational strategies to promote engagement and reduce burnout. *Mayo Clin Proc.* 2017;92(1):129–146.
35. Krug MF, Golob AL, Wander PL, Wipf JE. Changes in resident well-being at one institution across a decade of progressive work hours limitations. *Acad Med.* 2017;92:1480–4.
36. Wright AA, Katz IT. Beyond burnout—re-designing care to restore meaning and sanity for physicians. *N Engl J Med.* 2018; 378(4):309–311.
37. Panagioti M, Panagopoulou E, Bower P, et al. Controlled interventions to reduce burnout in physicians: A systematic review and meta-analysis. *JAMA Intern Med.* 2017;177:195–205.
38. Card AJ. Physician burnout: resilience training is only part of the solution. *Ann Fam Med.* 2018;16(3):267–270.
39. Krasner MS, Epstein RM, Beckman H. Association of an educational program in mindful communication with burnout, empathy, and attitudes among primary care physicians. *JAMA.* 2009;302(12):1284–1293.
40. Shapiro SL, Astin JA, Bishop ST, et al. Mindfulness-based stress reduction for health care professionals: results from a randomized trial. *Int J Stress Manag.* 2005; 12(2):164–176.
41. Abedini NC, Stack SW, Goodman JL, Steinberg KP. "It's Not Just Time Off": a Framework for Understanding Factors Promoting Recovery From

- Burnout among internal medicine residents. *J Grad Med Educ.* 2018;10(1):26–32.
42. Accreditation Council for Graduate Medical Education. ACGME Common Program Requirements Section VI with Background and Intent. [http://www.acgme.org/Portals/0/PFAssets/ProgramRequirements/CPRs\\_Section%20VI\\_with-Background-and-Intent\\_2017-01.pdf](http://www.acgme.org/Portals/0/PFAssets/ProgramRequirements/CPRs_Section%20VI_with-Background-and-Intent_2017-01.pdf) Published February 2017. Accessed December 15, 2018.
  43. **Weiss KB, Wagner R, Nasca TJ.** Development, testing, and implementation of the ACGME clinical learning environment review (CLER) program. *J Grad Med Educ.* 2012;4:396–8.
  44. **Weiss KB, Baglan JP.** Evaluation Committee CLER. Challenges and opportunities in the six focus areas: CLER National Report of Findings 2016. *J Grad Med Educ.* 2016;8(2 Suppl 1):25–34.
  45. National Academy of Medicine. (2018). Clinician Resilience and Well-being - National Academy of Medicine. [online] Available at: <https://nam.edu/initiatives/clinician-resilience-and-well-being/>. Accessed December 15, 2018.
  46. **Okanlawon T.** Physician Wellness: Preventing Resident and Fellow Burnout - STEPS Forward. 2018. [online] [STEPSforward.org](https://www.stepsforward.org). Available at: <https://www.stepsforward.org/modules/physician-wellness>. Accessed December 15, 2018.
  47. **Szymczak JE, Smathers S, Hoegg C, Klieger S, Coffin SE, Sammons JS.** Reasons why physicians and advanced practice clinicians work while sick. A mixed-methods analysis. *JAMA Pediatr.* 2015;169(9):815–821.
  48. **Wallace JE, Kemaire JV, Ghali WA.** Physician wellness: a missing quality indicator. *Lancet.* 2009;374:1714–21
  49. LCME. Functions and Structure of a Medical School: Standards for Accreditation of Medical Education Programs Leading to the MD Degree. Washington, DC, and Chicago, IL: LCME; 2016. Effective July 1, 2017. <http://lcme.org/publications/> Accessed December 15, 2018.
  50. **Dyrbye LN, West CP, Sinsky CA, Goeders LE, Satele DV, Shanafelt TD.** Medical Licensure Questions and Physician Reluctance to Seek Care for Mental Health Conditions. *Mayo Clin Proc.* 2017;92(10):1486–1493.
  51. **Jena AB, Baldwin DC Jr, Daugherty SR, Meltzer DO, Arora VM.** Presenteeism among resident physicians. *JAMA.* 2010;304(11):1166–1168
  52. **Hashimoto DA, Von Hofe JB, McDougall MA.** The trainee perspective: what can residency programs do to promote learner well-being? *Acad Med.* 2017;92(1):12.
  53. **Raj KS.** Well-being in residency: a systematic review. *J Grad Med Educ.* 2016;8(5):674–684.
  54. **Schwenk TL.** Resident depression: the tip of a graduate medical education iceberg. *JAMA.* 2015;314(22):2357–58.
  55. **Sklar DP.** Fostering student, resident, and faculty wellness to produce healthy doctors and a healthy population. *Acad Med.* 2016;91:1185–88.
  56. **Dyrbye LN, West CP.** Enhancing resident well-being: Illuminating the path forward. *J Gen Intern Med.* 2018;33(4):400–402.
  57. **Jennings ML, Slavin SJ.** Resident wellness matters: optimizing resident education and wellness through the learning environment. *Acad Med.* 2015;90(9):1246–50.
  58. **Ejnes YD.** Empowering trainees to aim for physician wellness. *J Grad Med Educ.* 2016;8(5):775–776.
  59. **Bodenheimer T, Sinsky C.** From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med.* 2014;12(6):573–6.
  60. **Lyon C, English AF, Smith PC.** A team-based care model that improves job satisfaction. *Fam Pract Manag.* 2018;25(2):6–11.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.