


A Ball-Milling-Enabled Reformatsky Reaction

Qun Cao, Roderick T. Stark, Ian A. Fallis, and Duncan L. Browne*^[a]


An operationally simple one-jar one-step mechanochemical Reformatsky reaction using *in situ* generated organozinc intermediates under neat grinding conditions has been developed. Notable features of this reaction protocol are that it requires no solvent, no inert gases, and no pre-activation of the bulk zinc source. The developed process is demonstrated to have good substrate scope (39–82% yield) and is effective irrespective of the initial morphology of the zinc source.

Metal-mediated C–C bond formation is an essential tool in modern organic synthesis. Numerous reactions consisting of metal-mediated nucleophilic addition to electrophiles have been developed for the synthesis of complex organic molecules.^[1] However, the generally high basicity and/or nucleophilicity of some organometallic reagents restricts their use in late-stage modification, where sensitive functional groups may already exist in the chemical structure. Conversely, organozinc species represent a class of “mild” organometallic compounds that demonstrate excellent functional group compatibility.^[2] Nevertheless, the preparation of organozinc species often requires initial access to more reactive organometallics, which are then transmetalated to give the desired organozinc reactant by metathesis with Zn^{II} salts. Alternatively, activated Zn⁰ can be used for the oxidative addition into carbon–halogen bonds (Scheme 1A). In general the formation and manipulation of organometallic compounds is not particularly clean or green when considering that solvents often have to be distilled and dried prior to use; inert gases are commonly required, and, in the case of organozinc reagents, the form of the bulk metal can play an important role and chemical additives are typically required to generate the activated zinc species.^[3] Recently, we identified mechanochemistry and ball-milling as a tool for the straightforward generation of organozinc species without the requirement for carefully prepared solvents or inert gases. Under these conditions the input of mechanical energy is enough to break down the resilient metal oxide surface and, in the presence of the alkyl/aryl halide, generate the corresponding organozinc species (Scheme 1B).


A) overview - formation and use of organozinc reagents



B) previous work - mechanochemical Negishi reaction



C) this work - mechanochemical one-pot Reformatsky reaction



Scheme 1. A) Formation and use of organozinc reagents. B) Previous work: one-jar two-step Negishi coupling.^[4] C) This work: one-jar one-step mechanochemical Reformatsky reaction.

These organozinc species may then be intercepted by opening the grinding jar, adding both catalyst and coupling partner before then running a telescoped Negishi coupling reaction; such a process is applicable to both sp³-sp³ and sp³-sp² coupling reactions. We have exploited this concept to carry out the one-jar, one-step preparation and use of organozinc species in a robust mechanochemical Reformatsky reaction (Scheme 1C). The Reformatsky reaction offers excellent potential for the formation of C–C bonds through (1) predictable C–C formation, (2) neutral reaction conditions (in comparison to obtaining the same products through aldol condensation), (3) broad functional group tolerance, and (4) the ability to impart a high degree of stereocontrol.^[5]

The classical Reformatsky reaction, in which β-hydroxy esters are formed by the reaction of aldehydes/ketones with α-halo esters in the presence of metallic zinc, was reported in 1887.^[6] Since this seminal work, a variety of latent nucleophiles and electrophiles have been studied and applied in this reaction and it has been routinely used in the synthesis of complex natural products.^[7] However, to carry out the Reformatsky reaction, organozinc reagents must be prepared at the point of use; the majority of organozinc reagents are not commercially available.^[8] This process can be problematic, owing to the formation of a layer of passivating zinc oxide on the surface of zinc powder, which hampers the formation of organozinc species and requires removal through treatment with chemical additives. Such additives include aqueous acid,^[9] iodine,^[10] 1,2-di-

[a] Dr. Q. Cao, R. T. Stark, Dr. I. A. Fallis, Dr. D. L. Browne

School of Chemistry, Cardiff University
Main Building, Park Place, Cardiff, CF10 3AT (UK)
E-mail: dlbbrowne@cardiff.ac.uk

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/cssc.201900886>.

© 2019 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

bromoethane,^[11] or chlorotrimethylsilane.^[12] Highly reactive Rieke zinc is an alternative, but its preparation is nontrivial and requires the reduction of ZnCl₂ with alkali metals (Li, Na, K) and naphthalene.^[13] Furthermore, the air-sensitivity of this method renders the green credentials of the Rieke approach poor, and the success of the outcome is highly dependent on the physical form of zinc used. Multiple different forms of zinc are commercially available (see the Supporting Information; Figure S1). Therefore, the focus of this work is on the mechanical activation of zinc, which not only renders the process more operationally simple and cost effective, but also delivers significant improvements to some of the green metrics typically associated with this reaction.^[14]

Mechanochemistry has been widely used among the crystal engineering and metal–organic framework communities.^[15] Recently, ball milling and other mechanochemical techniques have been explored as methods to complement the synthetic toolkit.^[16] Running reactions under mechanochemical conditions not only offers a more sustainable way to carry out solvent-minimized/free reactions but can also lead to decreased reaction times, increased selectivity, or different reaction outcomes when compared to results obtained from solution-based reactions.^[17] Herein we describe a green method for the Reformatsky reaction by using the ball mill mechanical activation of elemental zinc in air.

Studies commenced by treating model substrates benzaldehyde (**1**, 1 mmol) and ethyl 2-bromoacetate (**2**, 1.2 mmol) with 1.6 equivalents of zinc (20–30 mesh zinc granular) at 30 Hz in a 10 mL grinding jar with a single ball of mass 4 g (Table 1). After 2 hours of grinding, ¹H NMR spectroscopy of the crude mixture (mesitylene as internal standard) confirmed that a 70% yield of ethyl 3-hydroxy-3-phenylpropanoate (**3**) was produced during the milling process (Table 1, entry 1).

Increasing the amount of zinc to 2 equivalents resulted in an 81% yield of the desired hydroxy ester product **3** (Table 1, entry 2). Further increasing the amount of zinc led to no significant increase in yield (Table 1, entries 3 and 4). Rather than increasing the equivalents of zinc, increasing the amount of ethyl 2-bromoacetate (**2**) from 1.2 to 2.0 equivalents also led to no real difference in the observed yield (Table 1, entries 5 and 6). With the optimized ratio of reagents in hand, a reaction time study assessed four individual reaction times of 0.5, 1, 1.5, and 2 h (Table 1, entries 2 and 7–9), which indicated that the reaction needs 2 h to afford complete conversion. A control experiment, whereby zinc was omitted from the reaction, returned none of the desired product.

We then applied the optimized conditions to a further 11 commercially available forms of zinc (Figure S1). Although, the zinc forms had various particle sizes, which may lead to differences in the ratio between zinc oxide layer and zinc metal, a fixed mass (2 mmol, 0.130 g) of each sample was employed. Pleasingly, we found that in all cases the mechanochemical Reformatsky reaction was successful irrespective of form of zinc

Table 1. Optimization of one-jar one-step Reformatsky reaction using a ball mill.

Entry	Zn [equiv]	2 [equiv]	t [h]	Conv. [%] ^[a]	Yield [%] ^[a]
1	1.6	1.2	2	96	70
2	2	1.2	2	97	81 (72)
3	3	1.2	2	98	81
4	5	1.2	2	100	82
5	2	1.5	2	99	83
6	2	2.0	2	99	81
7	2	1.2	0.5	58	33
8	2	1.2	1	70	58
9	2	1.2	1.5	96	76
10	0	1.2	2	0	0

Reaction conditions: benzaldehyde (1 mmol), ethyl 2-bromoacetate (as specified), 20–30 mesh zinc granular (as specified), mixer mill, 30 Hz. ^[a] Conversion and yield were determined by ¹H NMR using mesitylene as internal standard; value in parentheses refers to yield of isolated product.


Table 2. Mechanochemical Reformatsky reaction using different zinc forms.

Entry	Zn form	Supplier	Conv. [%] ^[a]	Yield [%] ^[a]
1	zinc granular, 20–30 mesh	Sigma–Aldrich	97	81
2	zinc granular, 20 mesh	Sigma–Aldrich	99	72
3	zinc foil, 0.25 mm thick, 99.9%	Sigma–Aldrich	99	77
4	zinc dust, < 10 mm	Sigma–Aldrich	98	80
5	zinc puriss, ACS reagent, > 99.9%	Sigma–Aldrich	91	66
6	zinc shot, 10 mm dia., 2 mm thick, 99.99 %	Sigma–Aldrich	87	53
7	zinc flake, ca. 325 mesh, 99.9%	Alfa Aesar	99	87
8	zinc wire, 0.04 in dia., 99.95 %	Alfa Aesar	95	75
9	zinc powder, 6–9 µm, 97.5 %	Alfa Aesar	99	83
10	zinc metal powder	Fisher Scientific	98	68
11	zinc, ≥ 99%, mossy	Acros	96	73
12	zinc foil, 0.38 mm	BDH chemicals	99	73

Reaction conditions: benzaldehyde (1 mmol), ethyl 2-bromoacetate (1.2 mmol), zinc form as specified (2 equiv), mixer mill, 30 Hz, 2 h. ^[a] Conversion and yield were determined by ¹H NMR spectroscopy with mesitylene as an internal standard.

that was used (Table 2). Notably, there appears to be a general trend that the forms with a higher surface area/volume ratio performed better for the Reformatsky reaction under neat ball-milling conditions, this is perhaps contrary to prediction as these metal forms should also contain a higher proportion of zinc oxide.


With the optimized conditions established, application to a small range of 11 carbonyl compounds was investigated for reactivity with ethyl 2-bromoacetate and zinc flakes (ca. 325 mesh) under mechanical grinding (Scheme 2). We found that the mechanochemical Reformatsky reaction demonstrates good functional group tolerance. The highly sterically hindered substrate 2,4,6-trimethylbenzaldehyde afforded the



corresponding hydroxy ester, **5**, in good yield (66%). Halogen-containing aromatic aldehydes were also effective substrates, leading to good yields of the isolated halogen-containing products (Scheme 2, **6–8**, 77–82%). Hydrodehalogenation was not observed under these one-pot reaction conditions. Acetophenone was also a competent electrophile under these conditions, affording the isolated tertiary benzylic alcohol product **10** in 69% yield. Aliphatic aldehydes offered mixed results. Cyclohexylcarboxaldehyde gave the product in 39% yield, 2-phenylethyl aldehyde afforded 73% yield of the corresponding product, and cinnamaldehyde resulted in 61% yield (Scheme 2).


Eight different latent nucleophiles were also examined in this process with benzaldehyde as the model electrophile. Pleasingly, the corresponding organozinc intermediates could be generated under mechanochemical conditions (Scheme 3) and both methyl and *tert*-butyl α-halo esters could be used to effectively form the β-hydroxy esters **14** and **15** in good to excellent yields. Reaction with ethyl 2-bromopropionate provided 67% yield of **16** as a mixture of diastereoisomers (*syn/anti*=58:42). Reaction with more stereocongested ethyl α-bromoisobutyrate maintained a good yield of 68% (**17**, Scheme 3) and α,α-difluoro-β-hydroxyester **18** could also be prepared by this method. The ball-milling-enabled Reformatsky reaction with ethyl 4-bromobut-2-enoate resulted in 52% yield of the α-substituted product **19** with moderate diastereoselectivity (*syn/anti*=62:38). 2-Bromoacetonitrile also participated in the *in situ* generation of an organozinc reagent and formed the corresponding Reformatsky product 3-hydroxy-3-phenylpropanenitrile (**20**). Notably, in all cases explored, no reductive aldehyde coupling (pinacol reaction) was observed.

The applicability of imines as electrophiles was also briefly explored under the developed ball-milling conditions. Besides



aldehydes and ketones, nonclassical Reformatsky electrophiles such as azomethines, nitriles, lactones, anhydrides, Y-thioalactams, and amides can also be used for the Reformatsky reaction.^[18] Although not fully optimized, *N*-benzylideneaniline (**22**) underwent a mechanochemical Reformatsky reaction (Scheme 4A) to afford the β-amino ester **23** in 48% yield, alongside a small amount of the corresponding β-lactam **24** (7% yield).

The convenience of the method is well demonstrated by comparing a solution-based reaction with that under ball-mill-



Scheme 4. A) Example of mechanochemical Reformatsky-type reaction with imine electrophile. B) Reformatsky reaction with unactivated zinc source: ball-milling enabled reaction vs. solution-based reaction. [a] Yield determined by ¹H NMR spectroscopy with mesitylene as internal standard. [b] Yield determined by GC with mesitylene as internal standard.

ing conditions using identical reagents (Scheme 4B). The solution-based reaction in dry THF and under nitrogen atmosphere at 50 °C with either zinc flakes or granular zinc (no additive used) resulted in low yields of 7 and 4%, respectively (Scheme 4B), whereas under ball-milling conditions without any solvent, inert gas, or additive (Table 2), the reaction proceeded smoothly in 2 h and all forms of zinc explored were effective for this transformation.

In conclusion, a reliable, operationally simple one-jar one-step mechanochemical Reformatsky reaction has been developed. By utilizing the organozinc generated *in situ* upon milling, this method avoids the requirement for dry solvents, inert gases, and chemical additives and thus furnishes a process where the green metrics are improved.

Acknowledgements

D.L.B. is grateful to the EPSRC for a First Grant (EP/P002951/1). I.A.F. and D.L.B. are grateful to the European Union, Tenovus Cancer Charity, and Cardiff University for a Knowledge Economy Skills Scholarship (KESS) to R.S. We thank the EPSRC UK National Mass Spectrometry Facility at Swansea University for mass spectrometry measurements

Conflict of interest

The authors declare no conflict of interest.

Keywords: ball milling • mechanochemistry • organozinc formation • reformatsky reaction

- [1] A. Boudier, L. O. Bromm, M. Lotz, P. Knochel, *Angew. Chem. Int. Ed.* **2000**, *39*, 4414–4435; *Angew. Chem.* **2000**, *112*, 4584–4606.
- [2] a) P. Knochel, J. Perea, P. Jones, *Tetrahedron* **1998**, *54*, 8275–8319; b) J. H. Kim, Y. O. Ko, J. Bouffard, S.-G. Lee, *Chem. Soc. Rev.* **2015**, *44*, 2489–2507.
- [3] a) P. Knochel, R. D. Singer, *Chem. Rev.* **1993**, *93*, 2117–2188; b) G. T. Achonduh, N. Hadei, C. Valente, S. Avola, C. J. O'Brien, M. G. Organ, *Chem. Commun.* **2010**, *46*, 4109–4111; c) H. N. Hunter, N. Hadei, V. Blagojevic, P. Patschinski, G. T. Achonduh, S. Avola, D. K. Bohme, M. G. Organ, *Chem. Eur. J.* **2011**, *17*, 7845–7851; d) L. C. McCann, H. N. Hunter, J. A. C. Clyburne, M. G. Organ, *Angew. Chem. Int. Ed.* **2012**, *51*, 7024–7027; *Angew. Chem.* **2012**, *124*, 7130–7133; e) L. C. McCann, M. G. Organ, *Angew. Chem. Int. Ed.* **2014**, *53*, 4386–4389; *Angew. Chem.* **2014**, *126*, 4475–4478; f) Y. H. Chen, M. Ellwart, V. Malakhov, P. Knochel, *Synthesis* **2017**, *49*, 3215–3223; g) Y. H. Chen, M. Ellwart, G. Toupalas, Y. Ebe, P. Knochel, *Angew. Chem. Int. Ed.* **2017**, *56*, 4612–4616; *Angew. Chem.* **2017**, *129*, 4683–4687; h) S. M. Manolikakes, M. Ellwart, C. I. Stathakis, P. Knochel, *Chem. Eur. J.* **2014**, *20*, 12289–12297.
- [4] Q. Cao, J. L. Howard, E. Wheatley, D. L. Browne, *Angew. Chem. Int. Ed.* **2018**, *57*, 11339–11343; *Angew. Chem.* **2018**, *130*, 11509–11513.
- [5] R. Ocampo, W. R. Dolbier, Jr., *Tetrahedron* **2004**, *60*, 9325–9374.
- [6] S. Reformatsky, *Ber. Dtsch. Chem. Ges.* **1887**, *20*, 1210–1211.
- [7] a) A. Fürstner, *Synthesis* **1989**, *571*–*590*; b) S. Witkowski, Y. K. Rao, R. H. Premchandran, P. V. Halushka, J. Fried, *J. Am. Chem. Soc.* **1992**, *114*, 8464–8472; c) T. Mukaiyama, I. Shiina, H. Iwadare, M. Saitoh, T. Nishimura, N. Ohkawa, H. Sakoh, K. Nishimura, Y.-I. Tani, M. Hasegawa, K. Yamada, K. Saitoh, *Chem. Eur. J.* **1999**, *5*, 121–161; d) J. I. DeGraw, P. H. Christie, R. L. Kisliuk, Y. Gaumont, F. M. Sirotnak, *J. Med. Chem.* **1990**, *33*, 212–215.
- [8] A. Herath, V. Molteni, S. Pan, J. Loren, *Org. Lett.* **2018**, *20*, 7429–7432.
- [9] M. Berton, L. Huck, J. Alcázar, *Nat. Protoc.* **2018**, *13*, 324–334.
- [10] a) P. Knochel, M. C. P. Yeh, S. C. Berk, J. Talbert, *J. Org. Chem.* **1988**, *53*, 2390–2392; b) M. S. Newman, *J. Am. Chem. Soc.* **1942**, *64*, 2131–2133; c) A. K. Bose, K. Gupta, M. S. Manhas, *J. Chem. Soc. Chem. Commun.* **1984**, 86–87; d) S. Huo, *Org. Lett.* **2003**, *5*, 423–425.
- [11] a) S. C. Berk, P. Knochel, M. C. P. Yeh, *J. Org. Chem.* **1988**, *53*, 5789–5791; b) P. Knochel, J. F. Normant, *Tetrahedron Lett.* **1984**, *25*, 1475–1478.
- [12] a) J. K. Gawroński, *Tetrahedron Lett.* **1984**, *25*, 2605–2608; b) G. Picotin, P. Miginiac, *J. Org. Chem.* **1987**, *52*, 4796–4798; c) R. Ikegami, A. Koresawa, T. Shibata, K. Takagi, *J. Org. Chem.* **2003**, *68*, 2195–2199.
- [13] a) R. D. Rieke, S. J. Uhm, P. M. Hudnall, *J. Chem. Soc. Chem. Commun.* **1973**, 269b–270; b) R. D. Rieke, P. T.-J. Li, T. P. Burns, S. T. Uhm, *J. Org. Chem.* **1981**, *46*, 4323–4324; c) L. Zhu, R. M. Wehmeyer, R. D. Rieke, *J. Org. Chem.* **1991**, *56*, 1445–1453; d) A. Guijarro, D. M. Rosenberg, R. D. Rieke, *J. Am. Chem. Soc.* **1999**, *121*, 4155–4167; e) M. V. Hanson, R. D. Rieke, *J. Am. Chem. Soc.* **1995**, *117*, 10775–10776; f) R. D. Rieke, M. V. Hanson, J. D. Brown, Q. J. Niu, *J. Org. Chem.* **1996**, *61*, 2726–2730.
- [14] We are aware of a previous report of the use of a mortar and pestle to perform the Reformatsky reaction: a) K. Tanaka, S. Kishigami, F. Toda, *J. Org. Chem.* **1991**, *56*, 4333–4334. The use of a ball mill helps to standardize this process amongst different operators, reduce the energy required from the operator, and improve the safety of the reaction process. There exists a single unoptimized example of this process: b) C. Falenczyk, B. Pölloth, P. Hilgers, B. König, *Synth. Commun.* **2015**, *45*, 348–354. For examples of zinc mediated modification of [60]Fullerene under ball milling conditions see: c) G.-W. Wang, Y. Murata, K. Komatsu, T. S. M. Wan, *Chem. Commun.* **1996**, 2059–2060; d) H.-W. Liu, H. Xu, G. Shao, G.-W. Wang, *Org. Lett.* **2019**, *21*, 2625–2628.
- [15] a) A. M. Belenguer, G. I. Lampert, A. J. Cruz-Cabeza, C. A. Hunter, J. K. M. Sanders, *Chem. Sci.* **2016**, *7*, 6617–6627; b) D. Hasa, E. Miniussi, W. Jones, *Cryst. Growth Des.* **2016**, *16*, 4582–4588; c) A. V. Trask, N. Shan, W. D. S. Motherwell, W. Jones, S. Feng, R. B. H. Tan, K. J. Carpenter, *Chem. Commun.* **2005**, 880–882; d) Y. Zhou, F. Guo, C. E. Hughes, D. L. Browne, T. R. Peskett, K. D. M. Harris, *Cryst. Growth Des.* **2015**, *15*, 2901–2907; e) D. Crawford, J. Casabán, R. Haydon, N. Giri, T. McNally, S. L. James, *Chem. Sci.* **2015**, *6*, 1645–1649; f) T. Friščić, *Chem. Soc. Rev.* **2012**, *41*, 3493–3510; g) T. Friščić, *J. Mater. Chem.* **2010**, *20*, 7599–7605; h) P. J. Beldon, L. Fábián, R. S. Stein, A. Thirumurugan, A. K. Cheetham, T. Friščić, *Angew. Chem. Int. Ed.* **2010**, *49*, 9640–9643; *Angew. Chem.* **2010**, *122*, 9834–9837; i) K. Užarević, T. C. Wang, S. Y. Moon, A. M. Fidelli, J. T. Hupp, O. K. Farha, T. Friščić, *Chem. Commun.* **2016**, *52*, 2133–2136; j) M. Klimakow, P. Klobes, A. F. Thünemann, K. Rademann, F. Emmerling, *Chem. Mater.* **2010**, *22*, 5216–5221.
- [16] a) G.-W. Wang, *Chem. Soc. Rev.* **2013**, *42*, 7668–7700; b) S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Friščić, F. Grepioni, K. D. M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin, W. C. Shearouse, J. W. Steed, D. C. Waddell, *Chem. Soc. Rev.* **2012**, *41*, 413–447; c) J.-L. Do, T. Friščić, *ACS Cent. Sci.* **2017**, *3*, 13–19; d) D. Tan, T. Friščić, *Eur. J. Org. Chem.* **2018**, 18–33; e) O. Eguaogie, J. S. Vyle, P. F. Conlon, M. A. Gilea, Y. Liang, *Beilstein J. Org. Chem.* **2018**, *14*, 955–970; f) J. Andersen, J. Mack, *Green Chem.* **2018**, *20*, 1435–1443.
- [17] a) J. L. Howard, Q. Cao, D. L. Browne, *Chem. Sci.* **2018**, *9*, 3080–3094; b) J. G. Hernández, C. Bolm, *J. Org. Chem.* **2017**, *82*, 4007–4019.
- [18] a) S. M. Hannick, Y. Kishi, *J. Org. Chem.* **1983**, *48*, 3833–3835; b) K. Mohri, Y. Yoshida, S. Mataga, Y. Hisatsune, K. Isobe, Y. Tsuda, *Heterocycles* **1997**, *45*, 2405–2412; c) M.-X. Wang, Y. Liu, H.-Y. Gao, Y. Zhang, C.-Y. Yu, Z.-T. Huang, G. W. J. Fleet, *J. Org. Chem.* **2003**, *68*, 3281–3286; d) K. H. Schwarz, K. Kleiner, R. Ludwig, H. Schick, *J. Org. Chem.* **1992**, *57*, 4013–4015; e) J. P. Michael, C. B. de Koning, T. V. Stanbury, *Tetrahedron Lett.* **1996**, *37*, 9403–9406; f) C. Kashima, X. C. Huang, Y. Harada, A. Hosomi, *J. Org. Chem.* **1993**, *58*, 793–794; g) T. Honda, H. Wakabayashi, K. Kanai, *Chem. Pharm. Bull.* **2002**, *50*, 307–308; h) C. Baldoli, P. Del Buttero, E. Licandro, A. Papagni, T. Pilati, *Tetrahedron* **1996**, *52*, 4849–4856; i) J. C. Adrian, J. L. Barkin, L. Hassib, *Tetrahedron Lett.* **1999**, *40*, 2457–2460.

Manuscript received: April 1, 2019

Revised manuscript received: April 24, 2019

Accepted manuscript online: April 29, 2019

Version of record online: June 5, 2019