



Published in final edited form as:

Harv Rev Psychiatry. 2018 ; 26(4): 216–227. doi:10.1097/HRP.000000000000176.

Conceptualization, Assessment, and Treatment of Traumatic Stress in First Responders: A Review of Critical Issues

Nina F. Lewis-Schroeder, PhD, Kathryn Kieran, NP, Beth L. Murphy, MD, PhD, Jonathan D. Wolff, BS, Matthew A. Robinson, PhD, and Milissa L. Kaufman, MD, PhD

Harvard Medical School (Drs. Lewis-Schroeder, Murphy, Robinson, and Kaufman) and McLean Hospital, Belmont, MA (all).

Abstract

First responders are regularly confronted with exposure to traumatic events, including potentially life-threatening situations as well as the grave injuries and deaths of colleagues and civilians. Evidence indicates that the prevalence of posttraumatic stress disorder (PTSD) is substantially higher among first responders than the general population. This article provides information about the outpatient trauma services at McLean Hospital's LEADER (Law Enforcement, Active Duty, Emergency Responder) program to assist clinicians who encounter these first responders in their practices or who are specifically interested in working with this patient population. We begin by synthesizing the literature on the prevalence of PTSD in first responders following work-related exposure to traumatic stress, and by addressing the occupation-specific risk factors and the third-variable risk factors that may contribute to potentiated risk. We then discuss assessment strategies and treatment options used in our program, which is tailored for individuals who are dealing with mental health issues stemming from occupation-specific traumatic-stress exposure. We also address the unique challenges of treating traumatized first responders with more complex issues such as traumatic stress exposure across the lifespan and safety issues, including acute suicidality. We conclude by discussing notable gaps in the literature, including the need to investigate why and how women present with different PTSD symptoms than men and how these differences need to be taken into account in determining appropriate treatment for women.

Keywords

assessment; first responders; posttraumatic stress disorder; trauma; treatment

The Boston Marathon bombing and subsequent suspect-apprehension efforts the week of 15 April 2013 brought the unique mental health needs of first responders to the attention of McLean Hospital. To meet these needs, local representatives from police, government, and the hospital came together to begin shaping a treatment program. Out of these meetings, the LEADER (Law Enforcement, Active Duty, Emergency Responder) program at McLean Hospital was initiated. It is now a robust program offering services across multiple

Correspondence: Nina F. Lewis-Schroeder, PhD, McLean Hospital, 115 Mill St., Belmont, MA02478. nlewis3@partners.org.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the article.

diagnostic areas and levels of care, including inpatient, residential, partial hospital, and outpatient services. Working with first responders is a privilege afforded to those who participate in the LEADER program. This article seeks to help other clinicians to provide first responders with culturally sensitive and diagnostically sound assessment and treatment.

For individuals with posttraumatic stress disorder (PTSD), the LEADER program was developed within the context of well-established, hospital-wide services for assessing and treating traumatic stress, with a focus on childhood and interpersonal trauma. Currently, our trauma programs offer a comprehensive, phase-oriented approach to PTSD treatment, including multimodal interventions for both male and female first responders.¹ As expertise regarding this unique population has grown, the unmet needs of the broader community have become more apparent. This article provides the background and experiences of our program to assist clinicians who will encounter these first responders in their practices. We focus here on law enforcement officers (LEOs), ambulance personnel (EMT/paramedics), and firefighters and other fire personnel (FFs), although we well recognize the important, high-risk work of all emergency-response personnel.

PTSD PREVALENCE RATES

First responders are exposed to potentially traumatic events repeatedly while on the job. For example, LEOs, EMT/ paramedics, and FFs are exposed to death, serious injury, and violence at significantly higher rates than most civilian professionals.²⁻⁴ Given the high frequency and severity of traumatic exposures, it is not surprising that first responders are at an elevated risk for developing PTSD.

Over the past two decades, a growing body of research has yielded prevalence estimates for the development of PTSD in first responders following work-related exposure to traumatic events. These data were derived mostly from small-scale, retrospective studies utilizing self-report measures, rather than from diagnostic clinical interviews. As such, results must be interpreted judiciously. Nevertheless, cumulative range estimates can serve as potentially reliable indicators of PTSD prevalence. Studies show that LEOs develop PTSD at rates ranging from 6% to 32%,⁴⁻⁹ EMT/paramedics at rates ranging from 9% to 22%,^{2,5,7,10,11} and FFs at rates ranging from 17% to 32%.¹²⁻²⁴ By contrast, approximately 7% to 12% of adults in the United States will develop PTSD at some point in their lifetimes.^{25,26}

These wide ranges indicate that third-variable risk factors influencing PTSD prevalence exist both within groups (at the level of individual) and between groups (at the level of occupation type). For example, the lower ranges of PTSD prevalence rates found in some studies of LEOs may be explained by underreporting of symptoms due to fears of being considered unfit for duty and the requirements attendant to carrying a firearm.⁷ However, using a conservative estimation calculation,^{27,28} prevalence data suggest that more than 87,000 LEOs, 21,000 EMT/paramedics, and 804,000 FFs suffer from PTSD in the United States. Likely, many more suffer from subthreshold PTSD symptoms, causing significant occupational and social impairment.^{6,9,18} As such, LEOs, EMT/ paramedics, and FFs should be considered as special populations at increased relative risk.

RISK FACTORS FOR DEVELOPING PTSD

Several non-occupational and occupation-specific risk factors may increase the likelihood of developing PTSD subsequent to traumatic stress exposure in LEOs, EMT/paramedics, and FFs. Non-occupational risk factors can be organized into three categories: historical, peritraumatic, and posttraumatic.^{29–34} Historical risk factors include family history of psychiatric disorders, intelligence, education, early conduct problems, childhood adversity, and childhood abuse.^{35,36}

Peritraumatic risk factors include severity of the traumatic event, perception of the trauma as life threatening, actual physical injury or assault, dissociation during the event, and magnitude of the dissociative response.^{30–33,37} Posttraumatic risk factors include the absence of social support, poor access to healthy coping skills, limited access to mental health resources, and other life stressors.^{9,32,34}

Occupation-specific risk factors include the cumulative nature of the traumatic events encountered on the job, the types of traumatic events encountered, routine occupational stress, the perception of inadequate workplace social support, and the concurrent experience of gender or ethnic discrimination or stigmatization.^{2,5,24,29,35,38–41} In addition, hostile occupational environments that include exposure to extreme heat, fire, smoke, risk for repeated physical injury, and erratic sleep patterns may alter inflammatory and physiological stress responses, and compromise resilience in the face of PTSD risk factors.^{33,42–44} Given that first responders are repeatedly exposed to high-magnitude stressors, occupational risk may be compounded throughout their careers, placing even the most resilient at increased risk for problematic posttraumatic responding.^{20,30}

Individuals confronted with high-impact and high-frequency stressors are at an increased risk of experiencing an acute stress response.^{45,46} While such responses are normal, some of these individuals subsequently develop more serious impairments that require assessment and treatment—for example, acute stress disorder (ASD) or PTSD.^{45,46} An acute stress response at the time of, or shortly after, exposure to traumatic stress involves well-documented biological and psychological sequelae, which may include transient experiences of hyperarousal, anger/irritability, sadness, numbing, nightmares, and intrusive thoughts.^{31,32} Although rates of treatment seeking vary, the culture and selfimage of first responders may discourage them from seeking formal mental health interventions that are seen as stigmatizing.^{47–49}

Some of the individuals acutely affected by traumatic events subsequently develop ASD, which can be associated with increased risk for developing PTSD.^{45,46,50,51} ASD can be identified in individuals starting three days after an event and includes multiple symptoms within five diagnostic categories: intrusions, negative mood, dissociation, avoidance, and arousal.^{33,34} This disorder is differentiated from a more typical, transient response to acute stress exposure by the persistent severity of symptoms beyond the time frame of an acute stress reaction.^{33,52} ASD was formally introduced in the fourth edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) in an attempt to identify a PTSD prodrome.⁵³ The predictive value of the criteria, however, has been questioned.^{33,52,54,55}

While a diagnosis of ASD is associated with high risk for subsequently developing PTSD, the majority of those with PTSD will not have met criteria for ASD.^{54,55} In addition, the requirement of dissociation in the diagnosis of ASD may identify a specific subgroup at risk for PTSD.⁵⁶⁻⁵⁹ Dissociation experienced prior to or during a traumatic event appears to confer added risk for PTSD in first re-responders,⁴² though not consistently in the general population.³³

DIAGNOSIS AND ASSESSMENT OF PTSD

Diagnostic assessment is the cornerstone of the clinician's ability to develop the best treatment plan, and constitutes the first phase of treatment for each first responder. A thorough diagnostic evaluation clarifies which problems and symptoms have priority in treatment, and will guide decision making about pacing into later phases of treatment. Further, while careful assessment and symptom stabilization for PTSD can be provided safely to any traumatized first responder, the important decision about when to proceed to the trauma-focused processing phase of treatment is dependent upon a thorough understanding of the first responder's past history, the presence of co-occurring psychiatric and medical conditions, and functional status. A number of both standardized clinical interviews and self-report measures for assessing traumatic event exposure^{43,44} and PTSD symptoms^{43,44} in adults have been developed. Some of these assessment tools are available only for purchase, whereas others can be found in the articles cited^{43,44} or through the National Center for PTSD (<http://www.ptsd.va.gov>). Comprehensive reviews comparing the psychometric and cost-utility analyses of these tools are readily available to clinicians and researchers.^{43,44}

We will focus on PTSD assessment measures that have been updated to reflect the most current diagnostic criteria for PTSD, as presented in DSM-5.³⁴ These include the Clinician Administered PTSD Scale for DSM-5 (CAPS-5)^{60,61} and the PTSD Checklist for DSM-5 (PCL-5).⁶² Diagnostic assessment is geared toward the generation of a full psychiatric differential with an added emphasis on trauma-spectrum disorders. All assessments should include a comprehensive clinical interview as well as the administration of a validated measure for assessing PTSD. The CAPS-5 is a structured clinical interview considered to be the gold standard for diagnostic assessment of PTSD. The PCL-5 is a brief, self-report measure of current PTSD symptomatology that can be used to monitor PTSD treatment response longitudinally. Both of these measures are licensed by the National Center for PTSD, Behavioral Science Division.

Researchers at the National Center for PTSD in Boston initially developed the CAPS in 1990.⁶¹ Recently, this same group updated the CAPS to reflect changes in DSM-5.⁶⁰ The CAPS-5 is a structured clinical interview containing 30 items assessing for the full spectrum of DSM-5 PTSD criteria, including exposure to a traumatic event (Criterion A) and the core symptom clusters of intrusion (Criterion B), avoidance (Cluster C), negative alterations in cognition and mood (Criterion D), and alterations in arousal and reactivity (Criterion E). It is a diagnostic tool capable of assessment for both current and lifetime PTSD diagnoses. It includes items assessing duration of symptoms, severity of symptoms, response validity, and extent of social and occupational impairment. The CAPS-5 also includes assessment for a

dissociative subtype of PTSD^{63,64} using two items probing for experiences of depersonalization and derealization. As reviewed in detail by Weathers and colleagues (2001),⁶⁵ psychometric properties of the CAPS have been evaluated extensively across a wide range of traumatized sample types and, overall, are excellent. The CAPS has been found to show consistency over repeated administration times (test-retest reliability), over different interviewers or raters (interrater reliability), and across its various test items (internal consistency). In addition, CAPS scores correlate strongly with other measures of PTSD (convergent validity) and weakly with measures of different diagnostic constructs (discriminant validity). The CAPS also has been found to perform well in analyses designed to examine capacity to predict PTSD diagnostic status. In a personal communication with Dr. Terry Keane at the National Center for PTSD, he stated studies are currently under way to evaluate the psychometric properties of the recent version update.

The PCL was developed in 1993⁶⁵ at the National Center for PTSD. It has become, over the years, one of the most widely used self-report measures of PTSD symptomatology⁶⁶ and has consistently demonstrated excellent psychometric properties across many different settings.⁶⁷ The PCL recently was revised to reflect DSM-5 changes to the PTSD criteria.⁶² The PCL-5 is a 20-item self-report measure employing a rating scale from 0 to 4 for each symptom within the clusters of intrusion (Criterion B), avoidance (Cluster C), negative alterations in cognition and mood (Criterion D), and alterations in arousal and reactivity (Criterion E). The PCL-5 does not assess for symptoms of dissociation, duration of symptoms, or occupational and social impairment. The PCL-5 can be administered at the time of assessment or on a regular basis to assess for changes in symptoms. Recent findings suggest that, since its modification to adapt to the DSM-5, the PCL-5 is a psychometrically sound measure of PTSD. For example, the PCL-5 consistently demonstrated strong internal consistency, test-retest reliability, and both convergent and discriminative validity across traumatic event-exposed samples, including active military service members,⁶⁸ military veterans,⁶⁹ and college students.⁷⁰

First responders may be reluctant to discuss their PTSD symptoms or to acknowledge the presence of traumatic events in their lives, because of the stigma associated with diagnosis and treatment of psychiatric disorders.¹ As such, the therapeutic benefit they may derive from disclosing traumatic experiences and distress with a respectful, understanding professional should not be underestimated. Clinicians should take care to provide skillful pacing and containment during discussion to counteract potential flooding during the assessment. The clinician can also provide feedback about assessment results and diagnosis in a clinically sensitive manner, building the first therapeutic step in what will become an ongoing, psychoeducational process regarding the management of PTSD symptoms. As previously noted, it is rare for individuals exposed to traumatic events to present solely with symptoms of PTSD, which highlights the important need for comprehensive diagnostic evaluation for first responders.¹

Many first responders report that they are expected to minimize the impact of traumatic exposures in their professional and personal lives.¹ Therefore, first responders often engage in avoidance and may employ substance use or high-risk behavior to that end.⁷¹ Avoidance may present as absenteeism from work and result in early retirement.^{1,24} First responders

may also describe extreme irritability or an intense anger response that they may not perceive as a posttraumatic response.⁷² Further, they are likely to endorse constant hypervigilance and may experience sleep disruption due to nightmares or the challenges of sleeping while working on shifts.^{1,73,74} Many first responders note that their symptoms create tension with significant others and children.⁷⁵ Others may report conflict in the workplace, noting that they engage in greater discord with colleagues and supervisors than in the past.^{11,74,76} These factors indicate the importance of comprehensive assessment and psychoeducation regarding symptoms. Psychoeducation also plays a significant role in reducing subcultural boundaries that may prohibit a first responder from seeking and receiving appropriate treatment.¹

TREATMENT CONSIDERATIONS: A PHASE-ORIENTED APPROACH

Since the initial inclusion of PTSD in 1980 as a diagnostic entity in DSM-III,⁷⁷ the development of psychosocial and psychopharmacologic PTSD treatments has progressed at an impressive rate. Various forms of treatment have been evaluated by outcome studies using rigorous methodology.^{45,46,78,79} Currently, definitive professional practice guidelines exist, providing well-documented treatment recommendations for PTSD caused by a range of traumatic events experienced by a variety of different patient populations.^{50,80}

Trauma-focused processing may be contraindicated for those with actively concurrent suicidal or homicidal ideation, recent (past two months) nonsuicidal self-harm, active substance dependence, or psychosis.⁵¹ Furthermore, trauma-focused processing may be contraindicated in those without adequate social support, safety, and daily structure.^{81–83} Historically, trauma-focused processing treatment has been contraindicated prior to a period of stabilization and skills training in those with complex PTSD.^{81,82} However, recent studies have shown that cognitive-behavioral approaches combined with brief exposure elements can be used effectively in this population.^{51,84,85} This treatment advance is especially important for treating first responders with job-related PTSD complicated by a childhood abuse history and significant dissociation or mood dysregulation.

The approach to PTSD treatment in first responders is phase oriented through the outpatient trauma services at McLean Hospital's LEADER program. It has the following structure:

- Phase 1: Diagnostic assessment (as discussed above)
- Phase 2: Symptom stabilization and skills training
- Phase 3: Trauma-focused processing
- Phase 4: Consolidation and aftercare

Treatment Phase 2: Symptom Stabilization and Skills Training

The phase-oriented approach to treatment recognizes that some individuals with PTSD and other co-occurring symptoms may require an initial period of stabilization or skills training prior to undertaking trauma-focused processing.^{81–83} Immediate stabilization is warranted to address issues involving physical safety (e.g., suicidal or homicidal intention/plan) or active self-harm and neglect. In addition, rapid stabilization may be warranted in cases of severe

co-occurring depression, medically urgent co-occurring substance dependence, or debilitating reexperiencing symptoms (nightmares, flashbacks, intrusions) and accompanying hyperarousal symptoms (agitation and severely disrupted sleep). Such acute issues typically require inpatient level of care. In such cases, focus is on providing safety, medication evaluation, rapid symptom containment, and crisis management. During the hospitalization, fundamental coping skills, such as safety planning, grounding, self-care, and sleep hygiene, are taught and actively reinforced in a collaborative manner.

Our program has found that skills-acquisition training can be an appropriate next step for first responders with PTSD who are newly stabilized or for those with PTSD who are without acute safety/medical issues or severe debilitation but are nevertheless dealing with symptoms that intrude significantly upon daily functioning.⁴⁷ This treatment includes the following: psychoeducation; containment of PTSD-specific symptoms; management of co-occurring psychiatric health conditions; skills training in distress tolerance, emotion regulation, and impulse control; maintenance of supportive relationships with others; maintenance of healthy life style; and resilience training. Skills-acquisition training may not be required as a separate stage of treatment, however, for individuals requiring trauma-focused treatment.⁸⁶

Treatment modalities such as the dialectical behavior therapy-prolonged exposure (DBT PE) protocol integrate treatment approaches that increase skills in regulating emotion and tolerating distress while also addressing symptoms of PTSD.^{75,87,88} This growing body of research indicates significant improvements in PTSD symptoms, suicidal behaviors and urges, dissociation, depression, anxiety, and trauma-related guilt and shame in patients presenting with low-to-moderate suicidality behaviors and self-injurious behaviors.^{51,75,87-89} This research indicates that co-occurring symptoms may improve alongside symptoms of PTSD without completing skills-acquisition treatment in advance of trauma-focused treatment;⁸⁹ future studies are needed, however, to replicate these intriguing but early findings that, to date, have been conducted primarily by a single research group working with adults.

Treatment Phase 3: Trauma-Focused Processing

Well-documented, empirical support exists for several trauma-focused treatment modalities, including cognitive processing therapy (CPT), prolonged-exposure therapy (PE), and eye-movement desensitization and reprocessing. All have received “ A ” ratings, established on the strength of their respective published evidence bases, in practice guidelines issued by the International Society for Traumatic Stress Studies⁸⁰ and other national oversight agencies.^{50,90-92} Substantial literature supports several trauma-focused treatment modalities;^{85,93-96} for the purpose of this article, however, we focus on CPT, PE, and EMDR. Although no single treatment modality has been identified as most effective for first responders,²⁸ one study of LEOs with varying trauma histories found that 90% of participants chose PE or CPT as their first or second most preferred treatment.⁹⁷

COGNITIVE PROCESSING THERAPY CPT is a type of cognitive-behavioral therapy developed specifically for PTSD.⁹⁸ Among many skills, patients learn to challenge distorted negative self-cognitions resulting from traumatic experiences.^{99,100} This 12-session treatment has

successful, long-lasting recovery outcomes as measured by randomized, clinical trials and has a high rate of success in patients with co-occurring conditions (e.g., depression, substance abuse, anxiety disorders, and personality disorders).^{100–104}

Given CPT's unique focus on distorted negative self-cognitions, it may be especially helpful for individuals who have a history of chronic, childhood traumatic-event exposure—for example, individuals who have experienced early childhood abuse and neglect—who may have less favorable outcomes with therapies focused primarily on exposure-based interventions.^{84,105,106} For individuals with complex PTSD, CPT also can be an effective treatment without the exposure component or may be used as an actively planned, paced intervention to control dissociative symptoms while constructing the trauma-relevant narrative.⁹⁸

There is a long-standing, well-studied history of using CPT to treat PTSD among veterans, ^{103,107–111} including those with co-occurring alcohol use disorder.¹¹² No treatment-outcome studies have been published to date, however, with a specific focus on LEOs, EMT/paramedics, or FFs. In one study of 36 LEOs diagnosed with PTSD, 39% selected CPT as their first choice of treatment⁹⁷ when presented with multiple empirically based PTSD treatment descriptions. Research on the efficacy of CPT, coupled with indicated first-responder interest, suggests that CPT may be a treatment of choice for this unique population.

PROLONGED EXPOSURE PE is an internationally utilized cognitive-behavioral approach and treatment of choice for PTSD.¹¹³ It has a significant evidence base supporting its effectiveness in treating PTSD, with lasting results both posttreatment and at follow-up.^{114,115} PE incorporates components of PTSD psychoeducation, in vivo exposure to safe but feared stimuli related to trauma, imaginal exposure to traumatic memories, self-assessment of anxiety using subjective units of distress, and processing of trauma memories.^{51,116} Treatment consists of 8 to 15 90-minute individual sessions that engage in some form of exposure, with the goal of achieving physiological activation and habituation within and across exposure sessions.¹¹⁷

PE treats both PTSD-related distress and more general trauma-related distress,¹¹⁸ its focus on fear reduction also indicates great effectiveness in treating avoidance symptoms.^{119,120} It has been found effective in reducing negative trauma-related cognitions and depression symptoms.^{121,122} It was studied with first responders following the 11 September 2001 terrorist attacks and found to be highly effective for this group in reducing symptoms of PTSD.¹²³

EYE MOVEMENT DESENSITIZATION AND REPROCESSING EMDR is a treatment approach in which the patient's attention is directed to an external stimulus while concentrating on an emotionally disturbing experience, such as a traumatic event.¹²⁴ Saccadic eye movements, hand taps, and auditory tones are utilized while the patient concurrently engages in sequential exposure, desensitization, cognitive restructuring, and rehearsal.¹²⁴ This form of treatment has been found effective in reducing symptoms of PTSD as measured by psychometrically sound self-report measures.^{95,125–130} It remains unclear, however, whether an

external stimulus is required to achieve symptom improvement or whether the primary benefit of EMDR is derived from the emotional-processing component of treatment.^{120,125}

A number of studies have highlighted the effectiveness of EMDR in mass trauma situations, including natural disasters^{131,132} and terrorist attacks.¹³³ EMDR has been studied as a first-line PTSD treatment for first responders to good effect, suggesting its utility with the first-responder population.^{1,134–136} EMDR may also serve to help first responders reintegrate to work through a reduction in avoidance symptoms, leading to increased social and occupational productivity and a faster return to work following occupation-specific traumatic events.¹³⁷

CONTRAINDICATIONS TO TRAUMA-PROCESSING TREATMENT ALTHOUGH trauma-processing treatment may be indicated for individuals with co-occurring diagnoses, there are several definitive contraindications to the use of trauma-processing treatments. Individuals with PTSD who are involved in an ongoing violent relationship (for example, victims of domestic or other familial violence) should not undergo trauma-processing therapy until safety has been established and maintained. In addition, these treatment approaches are frequently contraindicated for those with suicidality, homicidality, current dependence on substances, and severe dissociative symptoms. Despite these contraindications, however, CPT has been used to successfully treat individuals with histories of childhood abuse and complex PTSD presentations, once problematic dissociation has been controlled. Additionally, exposure therapy may benefit individuals diagnosed with the dissociative subtype of PTSD^{51,138}—despite beliefs about the negative impact of dissociation on this form of therapy.

Treatment Phase 4: Consolidation and Aftercare

Continued treatment following trauma-focused processing is recommended for first responders seeking treatment through our program. This phase of treatment may incorporate self-assessment, cognitive restructuring, and behavioral strategies, including exposure and distress tolerance for ongoing maintenance of therapeutic gains. By this stage, first responders will have received psychoeducation regarding the clinical characteristics, etiology, course, and treatment of trauma-related diagnoses, and have developed self-assessment skills to gain insight into how trauma-related symptoms are affecting daily functioning.¹ The goal of this fourth stage of treatment is to help them to identify resources and build skills to help manage recurring or new stressors, to integrate new skill acquisition into daily routines, and to reintegrate to social and occupational roles, including engagement in peer support, while utilizing new skill sets to avoid the recurrence of behaviors targeted in treatment, such as avoidance.¹

SPECIAL TREATMENT CONSIDERATIONS: SUICIDALITY AND OTHER SAFETY ISSUES

Research findings have been mixed on whether suicide rates among police officers and firefighters are higher than in the general population or equivalent to them.^{139–144} An association between the presence of PTSD symptoms and elevated suicidality in the first-responder population has been consistently demonstrated,^{9,27,139,140} though this effect may

be driven or exacerbated by co-occurring depression.^{27,141} Given that negative alterations in mood and cognition are part and parcel of PTSD, determining whether the first responders' depressive symptomatology is related to past traumatic events has important clinical ramifications.

Other important risk factors for suicidality have been documented. When LEOs believe that their traumatic experiences are not manageable, their risk of suicidality increases.¹³⁹ Alcohol consumption with severe PTSD also increases the risk of suicidality in LEOs as much as tenfold when compared to LEOs with lesser PTSD symptomatology.¹⁴² In FFs, one study noted that numbing and reexperiencing symptoms were particularly associated with suicidal ideation.²⁷ In addition, job loss or demotion, and attendant loss of status, as well as public shame, may contribute to suicidality.¹³⁹ Personal and family adversity, including inability to provide for family or a sense of being a burden to family, are also associated with significant suicidality.¹⁴³

Shift work has not been specifically implicated in increased suicidality among LEOs and FFs. Working evening and overnight shifts, longer hours, and sleep deprivation are associated, however, with a greater frequency of exposure to stressful events.^{145–147} Some studies have found that LEOs and FFs with more experience on the job have a decreased risk for depression and suicidality.^{142,144,145} First responders with more than 20 years of experience, however, may be self-selecting for retirement versus continued work. Studies of whether race or ethnicity affect suicidality in firefighters have reported conflicting results.^{140,145}

It is critical to keep in mind that LEOs who are depressed or repeatedly flooded with PTSD symptoms—who may potentially become suicidal—have access to high-lethality weapons. Recent research also indicates that other types of safety concerns are important to monitor when working with first responders diagnosed with PTSD. For example, corrections officers who dissociate are at greater risk of assault by inmates, and paramedics with insomnia or nightmares may be fatigued and more prone to make medical errors.

First-responder groups have been trained in a particular subculture that may affect their perceptions of mental health and treatments.¹ Culturally sensitive care includes maintaining an ongoing awareness and curiosity about occupational norms and their implications for clinical management.¹⁴⁶ As with all individuals entering into treatment, an important initial step is to establish an alliance. This is particularly true for first responders, as many have concerns about disclosure (especially if they have been referred by their units). Fears about whether treaters will prioritize their well-being, behave with commitment, and treat them with respect can be especially problematic for police, sheriff, and corrections officers, who have high levels of interaction with hostile members of society or impersonal bureaucratic systems. Once an alliance is established, ongoing assessment of safety is important. Frank discussion about a range of concerns is critical, including those concerns that do not merit disclosure or hospitalization. In addition, education about medication use that will not affect alertness, cognitive processing, or reaction time is an important consideration when working with active-duty first responders.

Given the unique responsibilities of first responders, other concerns include potential public safety concerns and occupational interference. Many clinicians are familiar with suicidal behaviors but less so with behaviors posing a risk to others. While mandated reporting requirements do not vary by patient population, it is important to know that a restraining order or child-protection reporting can result in occupational suspension for police officers. Also concerning in relation to safe job performance are the intrusion of flashback symptoms into current activities and the chronically heightened assessment of threat. Limited duty may not exist for some occupational categories. Where concerns about fitness for duty or public safety exist, a specialty consultation is recommended.

SPECIAL POPULATION CONSIDERATIONS: FEMALE FIRST RESPONDERS

Special considerations for female first responders are of great importance because of both the scarcity of research and the heterogeneity of this population. In the existing literature on female first responders, LEOs are better represented than either EMT/paramedics or those who work as volunteer or paid FFs (Bureau of Labor Statistics). In some ways the samples of female LEOs “break all the rules” of the conventional wisdom on PTSD. Women generally are considered at higher risk than men of experiencing traumatic stress exposure within relationships, particularly sexual violence,^{41,147} and studies have found that females are at higher risk of developing PTSD.⁴¹ Nevertheless, some studies have found that female LEOs have lower rates of PTSD than civilian samples.^{39,57,148}

The interpretation of traumatic events is highly important to the development of PTSD, and one’s identity obviously and notably affects this attribution and interpretation. Studies have found differences in the types of incidents that precipitate PTSD symptoms in female (child abuse cases) versus male (shooting incidents) police officers.¹⁴⁹ Others have identified differences in how personality characteristics, as well as coping or attribution styles, may differ between male and female LEOs.¹⁵⁰ Emotional distress has been seen as a key factor in developing PTSD, but questions remain regarding the impact of first-responder culture and experiences in mediating increased emotional reactivity and cumulative PTSD symptoms among female first responders.^{29,39,151}

Women who work in guarding or “protective” professions may acculturate to the traditionally masculine gender roles associated with LEOs or other subcultures. This acculturation may explain why female LEOs may experience less peritraumatic emotional distress than civilian females.¹⁵¹ Female LEOs also appear to be at increased risk for somatization following traumatic exposure—a characteristic often found in male LEOs.³⁹ Recent alcohol use and PTSD have not always been connected in policing samples,⁵⁷ which is a stark difference from civilian data.^{152,153} Recent findings show, however, that both male and female LEOs increase alcohol use as a means of coping with PTSD.^{3,39}

Female first responders with both interpersonal and occupational exposure to traumatic stress may also experience the complication of institutional trauma—trauma that originates from, or is reinforced by, a previously trusted authority; this complication is especially common when the interpersonal traumas occur in the workplace.^{152,153} Such situations present a challenge in identifying the best treatment options for female first responders.

More evidence regarding the potential benefits of the common practice of treating males and females separately across occupational subpopulations, versus cohort therapy by occupation, would be invaluable in determining the best practice for treating female first responders. The benefits of sex-separated group therapy for individuals who have suffered work-related sexual harassment have been clinically observed, but evidence for a standard of care regarding first responders would be highly welcome.¹⁵⁴ In addition, confirmation, or an elaboration, of the differences in treating civilian trauma versus occupation-specific trauma, with recommendations for the latter, would be beneficial for clinicians in pharmacotherapy and other therapeutic modalities. This question is of particular importance as female first responders may respond differently than men to treatments for PTSD. For example, prazosin treatment for nightmares has been an important tool arising out of Veterans Administration research,^{155,156} but many women with interpersonal trauma find less daytime benefit and cannot tolerate the high dosing patterns used in the classic VA studies. It might be that dissociative-subtype PTSD needs to be separated out in research samples, for better understanding of the treatment nuances associated with that disorder. It is important to note that the only medications with an Food and Drug Administration approval for treating PTSD are sertraline and paroxetine. All other medications mentioned in this article are being used “off label” based on research and empirical support only.

In treating female first responders with PTSD, numerous factors need to be kept in mind, including their potentially different attachment styles from male first responders and also the likelihood of their having relational versus individualistic coping styles. Successful treatment is best achieved through person-centered care and careful attunement to both the stated and inferred individual goals of the patient. Consistently adopting a clarifying and learning stance toward the content of a patient’s history will also increase clinical attunement and improve the clinician’s odds of making beneficial treatment recommendations resulting in reduced treatment frustration and dropout. Practical and logistical difficulties are also common among female first responders presenting for treatment. Female first responders are frequently responsible for child and elder care, which creates challenges in finding time for to be available for treatment.¹⁵⁷ Access to flexible day treatment and outpatient options is valuable for this cohort. Additionally, standards of care developed for male first responders may need to be adapted in order to provide women with individualized trauma-informed care.

CONCLUSION

As referenced throughout, the assessment and treatment of PTSD and trauma spectrum disorders in the heterogeneous category of first responders requires consideration of their unique subcultural concerns and risk factors. The risk of retraumatization is great among first responders, requiring additional sensitivity in clinical work. Further, additional research specifically focused on assessing and treating PTSD among first responders is greatly needed. In a systematic review of 845 randomized, control trials of PTSD treatment outcomes, only 2 (0.2%) focused on first responders,²⁸ and even less is known about female first responders.

In addition to the dearth of treatment studies, treatment guidelines are lacking for first-responder populations. Therefore, clinicians are encouraged to use international guidelines for PTSD treatment and to focus on the importance of assessing PTSD and ASD in this population. With regard to assessment, clinicians are encouraged to inquire about occupational and childhood experiences, which may be useful in providing psychoeducation to patients regarding the interconnections between their present symptoms, recent traumas, and occupational and childhood history.

Following a thorough assessment, preferably using a standardized measure, a stage-based approach to treatment is recommended. In stage 1, it is critical to identify acute symptoms such as suicidality or co-occurring medical or psychiatric illnesses that require additional interventions. Stage 2 places an emphasis on achieving safety and acute-symptom stabilization through skills management. For example, medication and behavioral interventions to restore sleep hygiene may be necessary before concentration and energy are adequate for intensive work. Initial work also will be hampered if the patient does not have the skills to manage acute exacerbations with grounding and soothing techniques (often along with additional safety planning). Stage 3 of trauma-focused treatment involves intensive evidence-based trauma processing with a prerequisite of a general skills base to effectively manage acute distress without undue safety risks. Finally, stage 4 involves transitioning individuals to greater self-management, including the ability to identify the need for additional assistance. For individuals with a disruption in work or other independent living activities, this phase will focus on integration of skills to return to social and occupational routines.

First responders undergoing treatment for traumatic stress may require ongoing safety assessments and should return to an earlier phase of treatment as needed. It is important to educate all individuals beginning this treatment about the possibility of returning to an earlier stage when confronted with increased unsafe or treatment-interfering behaviors. Consultation with peers or specialists in traumatic-stress treatment, safety assessment, or first-responder subculture is necessary when issues of public safety may exist. Additionally, special considerations for treatment settings or treatment approaches developed primarily for men may need to be adapted for female first responders. Finally, the first-responder population and clinicians working with first responders will benefit from increased awareness of their special needs and of the best practices in assessing and treating PTSD.

Acknowledgments

We would like to acknowledge the many dedicated members of McLean Hospital leadership in providing support for the development of the LEADER Adult Outpatient Trauma Track. We would also like to thank the many first responders with whom we have had the privilege to work.

Supported by the McLean Hospital Trauma Scholar Fund, Anonymous Women's Mental Health Fund, and O'Keefe Family Foundation Fund (to Dr. Kaufman).

REFERENCES

1. Stergiopoulos E, Cimo A, Cheng C, Bonato S, Dewa CS. Interventions to improve work outcomes in work-related PTSD: a systematic review. *BMC Public Health* 2011;11:838. [PubMed: 22040066]

2. Alexander DA, Klein S. Ambulance personnel and critical incidents: impact of accident and emergency work on mental health and emotional well-being. *Br J Psychiatry* 2001;178: 76–81. [PubMed: 11136215]
3. Marmar CR, McCaslin SE, Metzler TJ, et al. Predictors of posttraumatic stress in police and other first responders. *Ann N Y Acad Sci* 2006;1071:1–18.
4. Marchand A, Nadeau C, Beaulieu-Prévost D, Boyer R, Martin M. Predictors of posttraumatic stress disorder among police officers: a prospective study. *Psychol Trauma* 2015;7: 212–21. [PubMed: 25793514]
5. Berger W, Coutinho ES, Figueira I, et al. Rescuers at risk: a systematic review and meta-regression analysis of the worldwide current prevalence and correlates of PTSD in rescue workers. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:1001–11. [PubMed: 21681455]
6. Carlier IV, Lamberts RD, Gersons BP. Risk factors for posttraumatic stress symptomatology in police officers: a prospective analysis. *J Nerv Ment Dis* 1997;185:498–506. [PubMed: 9284863]
7. Perrin MA, DiGrande L, Wheeler K, Thorpe L, Farfel M, Brackbill R. Differences in PTSD prevalence and associated risk factors among World Trade Center disaster rescue and recovery workers. *Am J Psychiatry* 2007;164:1385–94. [PubMed: 17728424]
8. Robinson HM, Sigman MR, Wilson JP. Duty-related stressors and PTSD symptoms in suburban police officers. *Psychol Rep* 1997;81:835–45. [PubMed: 9400075]
9. Maia DB, Marmar CR, Metzler T, et al. Post-traumatic stress symptoms in an elite unit of Brazilian police officers: prevalence and impact on psychosocial functioning and on physical and mental health. *J Affect Disord* 2007;97:241–5. [PubMed: 16859752]
10. Bennett P, Williams Y, Page N, Hood K, Woollard M. Levels of mental health problems among UK emergency ambulance workers. *Emerg Med J* 2004;21:235–6. [PubMed: 14988360]
11. van der Ploeg E, Kleber RJ. Acute and chronic job stressors among ambulance personnel: predictors of health symptoms. *Occup Environ Med* 2003;60 suppl 1:i40–6.
12. Arbona C, Fan W, Noor N. Factor structure and external correlates of posttraumatic stress disorder symptoms among African American firefighters. *Psychol Res Behav Manag* 2016;9:201–9. [PubMed: 27563263]
13. Armstrong D, Shakespeare-Finch J, Shochet I. Predicting post-traumatic growth and post-traumatic stress in firefighters. *Aust J Psychol* 2014;66:38–46.
14. Berninger A, Webber MP, Cohen HW, et al. Trends of elevated PTSD risk in firefighters exposed to the World Trade Center disaster: 2001–2005. *Public Health Rep* 2010;125:556–66. [PubMed: 20597456]
15. Corneil W, Beaton R, Murphy S, Johnson C, Pike K. Exposure to traumatic incidents and prevalence of posttraumatic stress symptomatology in urban firefighters in two countries. *J Occup Health Psychol* 1999;4:131–41. [PubMed: 10212865]
16. Haslam C, Mallon K. A preliminary investigation of posttraumatic stress symptoms among firefighters. *Work Stress* 2003;17:277–85.
17. North CS, Kawasaki A, Spitznagel EL, Hong BA. The course of PTSD, major depression, substance abuse, and somatization after a natural disaster. *J Nerv Ment Dis* 2004;192:823–9. [PubMed: 15583503]
18. Heinrichs M, Wagner D, Schoch W, Soravia LM, Hellhammer DH, Ehlert U. Predicting posttraumatic stress symptoms from pretraumatic risk factors: a 2-year prospective follow-up study in firefighters. *Am J Psychiatry* 2005;162: 2276–86. [PubMed: 16330591]
19. Lee JS, Ahn YS, Jeong KS, Chae JH, Choi KS. Resilience buffers the impact of traumatic events on the development of PTSD symptoms in firefighters. *J Affect Disord* 2014;162: 128–33. [PubMed: 24767017]
20. Pinto RJ, Henriques SP, Jongenelen I, Carvalho C, Maia AC. The strongest correlates of PTSD for firefighters: number, recency, frequency, or perceived threat of traumatic events? *J Trauma Stress* 2015;28:434–40. [PubMed: 26389531]
21. Skeffington PM, Rees CS, Mazzucchelli TG, Kane RT. The primary prevention of PTSD in firefighters: preliminary results of an RCT with 12-month follow-up. *PLoS One* 2016;11:e0155873.

22. Yuan C, Wang Z, Inslicht SS, et al. Protective factors for posttraumatic stress disorder symptoms in a prospective study of police officers. *Psychiatry Res* 2011;188:45–50. [PubMed: 21095622]
23. Bryant RA, Harvey AG. Posttraumatic stress in volunteer firefighters: predictors of distress. *J Nerv Ment Dis* 1995;183: 267–71. [PubMed: 7714516]
24. Wagner D, Heinrichs M, Ehler U. Prevalence of symptoms of posttraumatic stress disorder in German professional fire-fighters. *Am J Psychiatry* 1998;155:1727–32. [PubMed: 9842783]
25. Kessler RC. Posttraumatic stress disorder: the burden to the individual and to society. *J Clin Psychiatry* 2000;61:4–14.
26. Kolkow TT, Spira JL, Morse JS, Grieger TA. Post-traumatic stress disorder and depression in health care providers returning from deployment to Iraq and Afghanistan. *Mil Med* 2007;172: 451–5. [PubMed: 17521088]
27. Boffa JW, Stanley IH, Hom MA, Norr AM, Joiner TE, Schmidt NB. PTSD symptoms and suicidal thoughts and behaviors among firefighters. *J Psychiatr Res* 2017;84:277–83. [PubMed: 27810667]
28. Haugen PT, Evces M, Weiss DS. Treating posttraumatic stress disorder in first responders: a systematic review. *Clin Psychol Rev* 2012;32:370–80. [PubMed: 22561967]
29. Breslau N, Chilcoat HD, Kessler RC, Davis GC. Previous exposure to trauma and PTSD effects of subsequent trauma: results from the Detroit Area Survey of Trauma. *Am J Psychiatry* 1999;156:902–7. [PubMed: 10360130]
30. Walker A, McKune A, Ferguson S, Pyne DB, Rattray B. Chronic occupational exposures can influence the rate of PTSD and depressive disorders in first responders and military personnel. *Extrem Physiol Med* 2016;5:8. [PubMed: 27429749]
31. U.S. Department of Veterans Affairs. Common reactions after trauma. 2015 <http://www.ptsd.va.gov/public/problems/common-reactions-after-trauma.asp>
32. American Psychological Association. Recovering emotionally from disaster. 2018 <http://www.apa.org/helpcenter/recovering-disasters.aspx>
33. Bryant RA, Friedman MJ, Spiegel D, Ursano R, Strain J. A review of acute stress disorder in DSM-5. *Depress Anxiety* 2011; 28:802–17. [PubMed: 21910186]
34. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed Arlington, VA: American Psychiatric Press, 2013.
35. Maguen S, Metzler TJ, McCaslin SE, et al. Routine work environment stress and PTSD symptoms in police officers. *J Nerv Ment Dis* 2009;197:754–60. [PubMed: 19829204]
36. McFarlane AC. Posttraumatic stress disorder: a model of the longitudinal course and the role of risk factors. *J Clin Psychiatry* 2000;61:15–23.
37. Sayed S, Iacoviello BM, Charney DS. Risk factors for the development of psychopathology following trauma. *Curr Psychiatry Rep* 2015;17:612. [PubMed: 26206108]
38. Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychol Bull* 2003;129:52–73. [PubMed: 12555794]
39. Lilly MM, Pole N, Best SR, Metzler T, Marmar CR. Gender and PTSD: what can we learn from female police officers? *J Anxiety Disord* 2009;23:767–74. [PubMed: 19345556]
40. Thompson BM, Kirk A, Brown D. Sources of stress in policewomen: a three-factor model. *Int J Stress Manag* 2006;13: 309–28.
41. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995;52:1048–60. [PubMed: 7492257]
42. Fullerton CS, Ursano RJ, Wang L. Acute stress disorder, posttraumatic stress disorder, and depression in disaster or rescue workers. *Am J Psychiatry* 2004;161:1370–6. [PubMed: 15285961]
43. Norris FH, Hamblen JL. Standardized self-report measures of civilian trauma and PTSD In: Wilson JP, Keane TM, eds. *Assessing psychological trauma and PTSD*. New York: Guilford, 2004:63–102.
44. Weiss DS. Structured clinical interview techniques for PTSD In: Wilson JP, Keane TM, eds. *Assessing psychological trauma and PTSD*. New York: Guilford, 2004:103–21.
45. Ehlers A, Grey N, Wild J, et al. Implementation of cognitive therapy for PTSD in routine clinical care: effectiveness and moderators of outcome in a consecutive sample. *Behav Res Ther* 2013;51:742–52. [PubMed: 24076408]

46. Morina N, Wicherts JM, Lobbrecht J, Priebe S. Remission from post-traumatic stress disorder in adults: a systematic review and meta-analysis of long term outcome studies. *Clin Psychol Rev* 2014;34:249–55. [PubMed: 24681171]
47. Waters JA, Ussery W. Police stress: history, contributing factors, symptoms, and interventions. *Policing* 2007;30:169–88.
48. Hom MA, Stanley IH, Ringer FB, Joiner TE. Mental health service use among firefighters with suicidal thoughts and behaviors. *Psychiatr Serv* 2016;67:688–91. [PubMed: 26927574]
49. Royle L, Keenan P, Farrell D. Issues of stigma for first responders accessing support for post traumatic stress. *Int J Emerg Ment Health* 2009;11:79–85. [PubMed: 19927494]
50. National Institute for Health and Care Excellence. Posttraumatic stress disorder (PTSD): the management of PTSD in adults and children in primary and secondary care (clinical guideline 26) 2005 <https://www.nice.org.uk/guidance/CG26/uptake>
51. van Minnen A, Harned MS, Zoellner L, Mills K. Examining potential contraindications for prolonged exposure therapy for PTSD. *Eur J Psychotraumatol* 2012;3.
52. Isserlin L, Zerach G, Solomon Z. Acute stress responses: a review and synthesis of ASD, ASR, and CSR. *Am J Orthopsychiatry* 2008;78:423–9. [PubMed: 19123763]
53. Bryant RA. Acute stress disorder as a predictor of posttraumatic stress disorder: a systematic review. *J Clin Psychiatry* 2011;72: 233–9. [PubMed: 21208593]
54. Bryant RA, Creamer M, O'Donnell ML, Silove D, McFarlane AC. A multisite study of the capacity of acute stress disorder diagnosis to predict posttraumatic stress disorder. *J Clin Psychiatry* 2008;69:923–9. [PubMed: 18422396]
55. Brewin CR, Andrews B, Rose S, Kirk M. Acute stress disorder and posttraumatic stress disorder in victims of violent crime. *Am J Psychiatry* 1999;156:360–6. [PubMed: 10080549]
56. Bryant RA, Harvey AG. Relationship between acute stress disorder and posttraumatic stress disorder following mild traumatic brain injury. *Am J Psychiatry* 1998;155:625–9. [PubMed: 9585713]
57. Ballenger JF, Best SR, Metzler TJ, et al. Patterns and predictors of alcohol use in male and female urban police officers. *Am J Addict* 2011;20:21–9. [PubMed: 21175917]
58. Kaufman ML, Kimble MO, Kaloupek DG, et al. Peritraumatic dissociation and physiological response to trauma-relevant stimuli in Vietnam combat veterans with posttraumatic stress disorder. *J Nerv Ment Dis* 2002;190:167–74. [PubMed: 11923651]
59. Keane T, Kaufman M, Kimble M. Peritraumatic dissociative symptoms, acute stress disorder, and the development of post-traumatic stress disorder: causation, correlation or epiphenom-ena In: Sanchez-Planell L, Diez-Quevedo C, eds. *Dissociative states*. Barcelona: Springer Verlag, 2001:21–43.
60. U.S. Department of Veterans Affairs. Clinician-administered PTSD scale for DSM-5 (CAPS-5). 2017 <http://www.ptsd.va.gov/professional/assessment/adult-int/caps.asp>
61. Blake DD, Weathers FW, Nagy LM, et al. The development of a clinician-administered PTSD scale. *J Trauma Stress* 1995;8: 75–90. [PubMed: 7712061]
62. U.S. Department of Veterans Affairs. PTSD Checklist for DSM-5 (PCL-5). 2017 <http://www.ptsd.va.gov/professional/assessment/adult-sr/ptsd-checklist.asp>
63. Wolf EJ, Miller MW, Reardon AF, Ryabchenko KA, Castillo D, Freund R. A latent class analysis of dissociation and posttraumatic stress disorder: evidence for a dissociative subtype. *Arch Gen Psychiatry* 2012;69:698–705. [PubMed: 22752235]
64. Lanius RA, Brand B, Vermetten E, Frewen PA, Spiegel D. The dissociative subtype of posttraumatic stress disorder: rationale, clinical and neurobiological evidence, and implications. *Depress Anxiety* 2012;29:701–8. [PubMed: 22431063]
65. Weathers FW, Keane TM, Davidson JR. Clinician-administered PTSD scale: a review of the first ten years of research. *Depress Anxiety* 2001;13:132–56. [PubMed: 11387733]
66. Weathers F Posttraumatic stress disorder checklist In: Reyes G, Elhai JD, Ford JD, eds. *Encyclopedia of psychological trauma*. Hoboken, NJ: Wiley, 2008:491–4.
67. Keen SM, Kutter CJ, Niles BL, Krinsley KE. Psychometric properties of PTSD checklist in sample of male veterans. *J Rehab Res Dev* 2008;45:465–74.

68. Wortmann JH, Jordan AH, Weathers FW, et al. Psychometric analysis of the PTSD Checklist-5 (PCL-5) among treatment-seeking military service members. *Psychol Assess* 2016;28:1392–403. [PubMed: 26751087]
69. Bovin M, Marx B, Weathers F, et al. Psychometric properties of the PTSD Checklist for Diagnostic and Statistical Manual of Mental Disorders-Fifth Edition (PCL-5) in veterans. *Psychol Asses* 2016;28:1379–91.
70. Blevins CA, Weathers FW, Davis MT, Witte TK, Domino JL. The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): development and initial psychometric evaluation. *J Trauma Stress* 2015;28:489–98. [PubMed: 26606250]
71. Haisch DC, Meyers LS. MMPI-2 assessed post-traumatic stress disorder related to job stress, coping, and personality in police agencies. *Stress and Health* 2004;20:223–9.
72. Meffert SM, Metzler TJ, Henn-Haase C, et al. A prospective study of trait anger and PTSD symptoms in police. *J Trauma Stress* 2008;21:410–6. [PubMed: 18720397]
73. Liberman AM, Best SR, Metzler TJ, Fagan JA, Weiss DS, Marmar CR. Routine occupational stress and psychological distress in police. *Policing* 2002;25:421–41.
74. Beaton R, Murphy S, Pike K. Work and nonwork stressors, negative affective states, and pain complaints among fire-fighters and paramedics. *Int J Stress Manag* 1996;3:223–37.
75. Harned MS, Korslund KE, Linehan MM. A pilot randomized controlled trial of dialectical behavior therapy with and without the dialectical behavior therapy prolonged exposure protocol for suicidal and self-injuring women with borderline personality disorder and PTSD. *Behav Res Ther* 2014;55:7–17. [PubMed: 24562087]
76. Young KM, Cooper CL. Occupational stress in the ambulance service: a diagnostic study. *J Managerial Psychol* 1995;10:29–36.
77. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 3rd ed Washington, DC: American Psychiatric Press, 1980.
78. Steenkamp MM, Litz BT. Psychotherapy for military-related posttraumatic stress disorder: review of the evidence. *Clin Psychol Rev* 2013;33:45–53. [PubMed: 23123570]
79. van Minnen A, Foa EB. The effect of imaginal exposure length on outcome of treatment for PTSD. *J Trauma Stress* 2006;19: 427–38. [PubMed: 16929519]
80. Foa EB, Keane TM, Friedman MJ, Cohen JA. Effective treatments for PTSD: practice guidelines from the international society for traumatic stress studies. 2nd ed New York: Guilford, 2008.
81. Chu JA. Rebuilding shattered lives: treating complex PTSD and dissociative disorders. Hoboken, NJ: Wiley, 2011.
82. Courtois CA, Ford JD. Treating complex traumatic stress disorders: an evidence-based guide. New York: Guilford, 2009.
83. Herman JL. Trauma and recovery: the aftermath of violence from domestic violence to political terrorism. New York: Guilford, 1992.
84. Resick PA, Nishith P, Griffin MG. How well does cognitive-behavioral therapy treat symptoms of complex PTSD? An examination of child sexual abuse survivors within a clinical trial. *CNS Spectr* 2003;8:340–55. [PubMed: 12766690]
85. Cloitre M, Koenen KC, Cohen LR, Han H. Skills training in affective and interpersonal regulation followed by exposure: a phase-based treatment for PTSD related to childhood abuse. *J Consult Clin Psychol* 2002;70:1067–74. [PubMed: 12362957]
86. Hamblen JL, Schnurr PP, Rosenberg A, Eftekhari A. A guide to the literature on psychotherapy for PTSD. *Psychiatr Ann* 2009; 39:348–54.
87. Harned MS, Korslund KE, Foa EB, Linehan MM. Treating PTSD in suicidal and self-injuring women with borderline personality disorder: development and preliminary evaluation of a Dialectical Behavior Therapy Prolonged Exposure Protocol. *Behav Res Ther* 2012;50:381–6. [PubMed: 22503959]
88. Harned MS, Linehan MM. Integrating dialectical behavior therapy and prolonged exposure to treat co-occurring border-line personality disorder and PTSD: two case studies. *Cogn Behav Pract* 2008;15:263–76.
89. van Minnen A, Zoellner LA, Harned MS, Mills K. Changes in comorbid conditions after prolonged exposure for PTSD: a literature review. *Curr Psychiatry Rep* 2015;17:549. [PubMed: 25736701]

90. Nemeroff CB, Bremner JD, Foa EB, Mayberg HS, North CS, Stein MB. Posttraumatic stress disorder: a state-of-the-science review. *J Psychiatr Res* 2006;40:1–21. [PubMed: 16242154]
91. Institute of Medicine. Treatment of PTSD: an assessment of the evidence. Washington, DC: National Academies, 2008.
92. Substance Abuse and Mental Health Services Administration. Trauma-informed approach and trauma-specific interventions. 2015 <https://www.samhsa.gov/nctic/trauma-interventions>
93. Bradley R, Greene J, Russ E, Dutra L, Westen D. A multidimensional meta-analysis of psychotherapy for PTSD. *Am J Psychiatry* 2005;162:214–27. [PubMed: 15677582]
94. Cloitre M, Courtois CA, Charuvastra A, Carapezza R, Stolbach BC, Green BL. Treatment of complex PTSD: results of the ISTSS expert clinician survey on best practices. *J Trauma Stress* 2011;24:615–27. [PubMed: 22147449]
95. Van Etten ML, Taylor S. Comparative efficacy of treatments for post-traumatic stress disorder: a meta-analysis. *Clin Psychol Psychother* 1998;5:126–44.
96. Bisson JI, Roberts NP, Andrew M, Cooper R, Lewis C. Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *Cochrane Database Syst Rev* 2013;(12): CD003388.
97. Becker CB, Meyer G, Price JS, et al. Law enforcement preferences for PTSD treatment and crisis management alternatives. *Behav Res Ther* 2009;47:245–53. [PubMed: 19200945]
98. Resick PA, Monson CM, Chard KM. Cognitive processing therapy: veteran/military version: therapist's manual. Washington, DC: Department of Veterans Affairs, 2014.
99. Resick PA. Cognitive therapy for posttraumatic stress disorder. *J Cogn Psychother* 2001;15:321–9.
100. Chard KM, Schuster PA, Resick P. Empirically supported psychological treatments: cognitive processing therapy In: Beck JG, Sloan DM, eds. *The Oxford handbook of traumatic stress disorders*. New York: Oxford University Press, 2012: 439–48.
101. Sobel AA, Resick PA, Rabalais AE. The effect of cognitive processing therapy on cognitions: impact statement coding. *J Trauma Stress* 2009;22:205–11. [PubMed: 19479979]
102. Schumm JA, Dickstein BD, Walter KH, Owens GP, Chard KM. Changes in posttraumatic cognitions predict changes in post-traumatic stress disorder symptoms during cognitive processing therapy. *J Consult Clin Psychol* 2015;83:1161–6. [PubMed: 26214540]
103. Monson CM, Schnurr PP, Resick PA, Friedman MJ, Young-Xu Y, Stevens SP. Cognitive processing therapy for veterans with military-related posttraumatic stress disorder. *J Consult Clin Psychol* 2006;74:898–907. [PubMed: 17032094]
104. Iverson KM, King MW, Cunningham KC, Resick PA. Rape survivors' trauma-related beliefs before and after cognitive processing therapy: associations with PTSD and depression symptoms. *Behav Res Ther* 2015;66:49–55. [PubMed: 25698164]
105. Chard KM. An evaluation of cognitive processing therapy for the treatment of posttraumatic stress disorder related to child-hood sexual abuse. *J Consult Clin Psychol* 2005;73:965–71. [PubMed: 16287396]
106. Resick PA, Williams LF, Suvak MK, Monson CM, Gradus JL. Long-term outcomes of cognitive-behavioral treatments for posttraumatic stress disorder among female rape survivors. *J Consult Clin Psychol* 2012;80:201–10. [PubMed: 22182261]
107. Alvarez J, McLean C, Harris AH, Rosen CS, Ruzek JI, Kimerling R. The comparative effectiveness of cognitive processing therapy for male veterans treated in a VHA posttraumatic stress disorder residential rehabilitation program. *J Consult Clin Psychol* 2011;79:590–9. [PubMed: 21744946]
108. Chard KM, Schumm JA, McIlvain SM, Bailey GW, Parkinson RB. Exploring the efficacy of a residential treatment program incorporating cognitive processing therapy-cognitive for veterans with PTSD and traumatic brain injury. *J Trauma Stress* 2011;24:347–51. [PubMed: 21626573]
109. Chard KM, Schumm JA, Owens GP, Cottingham SM. A comparison of OEF and OIF veterans and Vietnam veterans receiving cognitive processing therapy. *J Trauma Stress* 2010; 23:25–32. [PubMed: 20146255]
110. Macdonald A, Monson CM, Doron-Lamarca S, Resick PA, Palfai TP. Identifying patterns of symptom change during a randomized controlled trial of cognitive processing therapy for military-related posttraumatic stress disorder. *J Trauma Stress* 2011;24:268–76. [PubMed: 21626572]

111. Morland LA, Mackintosh M-A, Greene CJ, et al. Cognitive processing therapy for posttraumatic stress disorder delivered to rural veterans via telemental health: a randomized noninferiority clinical trial. *J Clin Psychiatry* 2014;75:470–6. [PubMed: 24922484]
112. Kaysen D, Schumm J, Pedersen ER, Seim RW, Bedard-Gilligan M, Chard K. Cognitive processing therapy for veterans with comorbid PTSD and alcohol use disorders. *Addict Behav* 2014;39:420–7. [PubMed: 24035644]
113. Foa EB, Rothbaum BO, Riggs DS, Murdock TB. Treatment of posttraumatic stress disorder in rape victims: a comparison between cognitive-behavioral procedures and counseling. *J Consult Clin Psychol* 1991;59:715–23. [PubMed: 1955605]
114. Foa EB, Hembree EA, Cahill SP, et al. Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: outcome at academic and community clinics. *J Consult Clin Psychol* 2005;73:953–64. [PubMed: 16287395]
115. Foa EB, Steketee G, Rothbaum BO. Behavioral/cognitive conceptualizations of post-traumatic stress disorder. *Behav Ther* 1989;20:155–76.
116. Foa EB, Hembree EA, Rothbaum BO. Prolonged exposure therapy for PTSD: emotional processing of traumatic experiences therapist guide. New York: Oxford University Press, 2007.
117. Foa EB, Kozak MJ. Emotional processing of fear: exposure to corrective information. *Psychol Bull* 1986;99:20–35. [PubMed: 2871574]
118. Powers MB, Halpern JM, Ferenschak MP, Gillihan SJ, Foa EB. A meta-analytic review of prolonged exposure for posttraumatic stress disorder. *Clin Psychol Rev* 2010;30:635–41. [PubMed: 20546985]
119. Foa EB, Hearst-Ikeda D, Perry KJ. Evaluation of a brief cognitive-behavioral program for the prevention of chronic PTSD in recent assault victims. *J Consult Clin Psychol* 1995; 63:948–55. [PubMed: 8543717]
120. Pitman RK, Orr SP, Altman B, Longpre RE, Poiré RE, Macklin ML. Emotional processing during eye movement desensitization and reprocessing therapy of Vietnam veterans with chronic posttraumatic stress disorder. *Compr Psychiatry* 1996;37:419–29. [PubMed: 8932966]
121. Rauch SAM, Defever E, Favorite T, et al. Prolonged exposure for PTSD in a Veterans Health Administration PTSD clinic. *J Trauma Stress* 2009;22:60–4. [PubMed: 19145643]
122. Zalta AK, Gillihan SJ, Fisher AJ, et al. Change in negative cognitions associated with PTSD predicts symptom reduction in prolonged exposure. *J Consult Clin Psychol* 2014;82:171–5. [PubMed: 24188512]
123. Schneier FR, Neria Y, Pavlicova M, et al. Combined prolonged exposure therapy and paroxetine for PTSD related to the World Trade Center attack: a randomized controlled trial. *Am J Psychiatry* 2012;169:80–8. [PubMed: 21908494]
124. Shapiro F Eye movement desensitization and reprocessing (EMDR): evaluation of controlled PTSD research. *J Behav Ther Exp Psychiatry* 1996;27:209–18. [PubMed: 8959422]
125. Cahill SP, Carrigan MH, Frueh BC. Does EMDR work? And if so, why?: a critical review of controlled outcome and dismantling research. *J Anxiety Disord* 1999;13:5–33. [PubMed: 10225499]
126. Maxfield L, Hyer L. The relationship between efficacy and methodology in studies investigating EMDR treatment of PTSD. *J Clin Psychol* 2002;58:23–41. [PubMed: 11748595]
127. Rothbaum BO. A controlled study of eye movement desensitization and reprocessing in the treatment of posttraumatic stress disorder sexual assault victims. *Bull Menninger Clin* 1997;61:317–33. [PubMed: 9260344]
128. Boudewyns PA, Hyer L. Physiological response to combat memories and preliminary treatment outcome in Vietnam veteran PTSD patients treated with direct therapeutic exposure. *Behav Ther* 1991;21:63–87.
129. Carlson JG, Chemtob CM, Rusnak K, Hedlund NL, Muraoka MY. Eye movement desensitization and reprocessing (EDMR) treatment for combat-related posttraumatic stress disorder. *J Trauma Stress* 1998;11:3–24. [PubMed: 9479673]
130. Davidson PR, Parker KC. Eye movement desensitization and reprocessing (EMDR): a meta-analysis. *J Consult Clin Psychol* 2001;69:305–16. [PubMed: 11393607]

131. Grainger RD, Levin C, Allen-Byrd L, Doctor RM, Lee H. An empirical evaluation of eye movement desensitization and reprocessing (EMDR) with survivors of a natural disaster. *J Trauma Stress* 1997;10:665–71. [PubMed: 9391949]
132. Chemtob CM, Nakashima J, Carlson JG. Brief treatment for elementary school children with disaster-related posttraumatic stress disorder: a field study. *J Clin Psychol* 2002;58:99–112. [PubMed: 11748599]
133. Silver SM, Rogers S, Knipe J, Colelli G. EMDR therapy following the 9/11 terrorist attacks: a community-based intervention project in New York City. *Int J Stress Manag* 2005;12:29–42.
134. Keenan P, Royle L. Vicarious trauma and first responders: a case study utilizing eye movement desensitization and reprocessing (EMDR) as the primary treatment modality. *Int J Emerg Ment Health* 2008;9:291–8.
135. Wilson SA, Tinker RH, Becker LA, Logan CR. Stress management with law enforcement personnel: a controlled outcome study of EMDR versus a traditional stress management program. *Int J Stress Manag* 2001;8:179–200.
136. Jarero I, Amaya C, Givaudan M, Miranda A. EMDR individual protocol for paraprofessional use: a randomized controlled trial with first responders. *J EMDR Pract Res* 2013;7:55–64.
137. Summerfield D Metropolitan Police blues: protracted sickness absence, ill health retirement, and the occupational psychiatrist. *BMJ* 2011;342:d2127.
138. Hagenaaers MA, van Minnen A, Hoogduin KAL. The impact of dissociation and depression on the efficacy of prolonged ex-posure treatment for PTSD. *Behav Res Ther* 2010;48:19–27. [PubMed: 19766987]
139. Rothmann S, Strijdom G. Suicide ideation in the South African police services in the North West Province. *SA J Ind Psychol* 2002;28:44–8.
140. Martin CE, Tran JK, Buser SJ. Correlates of suicidality in firefighter/EMS personnel. *J Affect Disord* 2017;208:177–83. [PubMed: 27788381]
141. Chopko BA, Palmieri PA, Facemire VC. Prevalence and predictors of suicidal ideation among US law enforcement officers. *J Police Crim Psychol* 2014;29:1–9.
142. Violanti JM. Predictors of police suicide ideation. *Suicide Life Threat Behav* 2004;34:277–83. [PubMed: 15385182]
143. Berg AM, Hem E, Lau B, Loeb M, Ekeberg O. Suicidal ideation and attempts in Norwegian police. *Suicide Life Threat Behav* 2003;33:302–12. [PubMed: 14582840]
144. Darensburg T, Andrew ME, Hartley TA, Burchfiel CM, Fekedulegn D, Violanti JM. Gender and age differences in posttraumatic stress disorder and depression among Buffalo police officers. *Traumatology* 2006;12:220–8.
145. Stanley IH, Hom MA, Hagan CR, Joiner TE. Career prevalence and correlates of suicidal thoughts and behaviors among firefighters. *J Affect Disord* 2015;187:163–71. [PubMed: 26339926]
146. Kirschman E, Kamena M, Fay J. *Counseling cops: what clinicians need to know*. New York: Guilford, 2014.
147. Barth SK, Kimerling RE, Pavao J, et al. Military sexual trauma among recent veterans: correlates of sexual assault and sexual harassment. *Am J Prev Med* 2016;50:77–86. [PubMed: 26232906]
148. Pole N, Best SR, Weiss DS, et al. Effects of gender and ethnicity on duty-related posttraumatic stress symptoms among urban police officers. *J Nerv Ment Dis* 2001;189:442–8. [PubMed: 11504321]
149. Gehrke A, Violanti JM. Gender differences and posttraumatic stress disorder: the role of trauma type and frequency of exposure. *Traumatology* 2006;12:229–35.
150. Andrew ME, Mnatsakanova A, Howsare JL, et al. Associations between protective factors and psychological distress vary by gender: the Buffalo cardio-metabolic occupational police stress study. *Int J Emerg Ment Health* 2013;15:277–88. [PubMed: 24707590]
151. Fullerton CS, Ursano RJ, Epstein RS, et al. Gender differences in posttraumatic stress disorder after motor vehicle accidents. *Am J Psychiatry* 2001;158:1486–91. [PubMed: 11532736]
152. Najavits L, Weiss RD, Shaw SR. The link between substance abuse and posttraumatic stress disorder in women. *Am J Addict* 1997;6:273–83.

153. Greenfield SF, Back SE, Lawson K, Brady KT. Substance abuse in women. *Psychiatr Clin North Am* 2010;33:339–55. [PubMed: 20385341]
154. Muskett C Trauma-informed care in inpatient mental health settings: a review of the literature. *Int J Ment Health Nurs* 2014;23:51–9. [PubMed: 23363398]
155. Raskind MA, Peskind ER, Kanter ED, et al. Reduction of nightmares and other PTSD symptoms in combat veterans by prazosin: a placebo-controlled study. *Am J Psychiatry* 2003;160:371–3. [PubMed: 12562588]
156. Dierks MR, Jordan JK, Sheehan AH. Prazosin treatment of nightmares related to posttraumatic stress disorder. *Ann Pharmacother* 2007;41:1013–7. [PubMed: 17504838]
157. Schulz R, Sherwood PR. Physical and mental health effects of family caregiving. *Am J Nurs* 2008;108:23–7.