



Published in final edited form as:

*Methods Enzymol.* 2018 ; 604: 367–388. doi:10.1016/bs.mie.2018.02.012.

## Preparation, Assay and Application of Chlorinase SalL for the Chemoenzymatic Synthesis of S-Adenosyl-L-Methionine and Analogs

TD Davis<sup>\*</sup>, S Kunakom<sup>§</sup>, MD Burkart<sup>\*,1</sup>, and AS Eustáquio<sup>§,1</sup>

<sup>\*</sup>University of California San Diego, Department of Chemistry and Biochemistry, San Diego, CA, United States

<sup>§</sup>University of Illinois at Chicago, College of Pharmacy, Department of Medicinal Chemistry and Pharmacognosy, and Center for Biomolecular Sciences, Chicago, IL, United States

### Abstract

*S*-Adenosyl-L-methionine (SAM) is universal in biology, serving as the second most common cofactor in a variety of enzymatic reactions. One of the main roles of SAM is the methylation of nucleic acids, proteins and metabolites. Methylation often imparts regulatory control to DNA and proteins, and leads to an increase in the activity of specialized metabolites such as those developed as pharmaceuticals. There has been increased interest in using SAM analogs in methyltransferase-catalyzed modification of biomolecules. However, SAM and its analogs are expensive and unstable, degrading rapidly under physiological conditions. Thus, the availability of methods to prepare SAM *in situ* are desirable. In addition, synthetic methods to generate SAM analogs suffer from low yields and poor diastereoselectivity. The chlorinase SalL from the marine bacterium *Salinispora tropica* catalyzes the reversible, nucleophilic attack of chloride at the C5' ribosyl carbon of SAM leading to the formation of 5'-chloro-5'-deoxyadenosine (CIDA) with concomitant displacement of L-methionine. It has been demonstrated that the *in vitro* equilibrium of the SalL-catalyzed reaction favors the synthesis of SAM. In this chapter, we describe methods for the preparation of SalL, and the chemoenzymatic synthesis of SAM and SAM analogs from CIDA and L-methionine congeners using SalL. In addition, we describe procedures for the *in situ* chemoenzymatic synthesis of SAM coupled to DNA, peptide and metabolite methylation, and to the incorporation of isotopes into alkylated products.

### Keywords

*S*-adenosylmethionine (SAM); chlorodeoxyadenosine synthase; chlorinase; methylation; isotope incorporation; chemoenzymatic

## 1. INTRODUCTION

*S*-Adenosyl-L-methionine (SAM or AdoMet) is ubiquitous in nature and the second most utilized enzyme substrate after ATP. SAM is not only a source of methyl groups to methylate

<sup>1</sup>Corresponding authors: mburkart@ucsd.edu and ase@uic.edu.

a variety of substrates, including nucleic acid, proteins and small molecules (Cantoni, 1975; Thomsen, Vogensen, Buchardt, Burkart & Clausen, 2013), but also a source of methylene, amino, ribosyl and aminopropyl groups, in addition to 5'-deoxyadenosyl radicals (Fontecave, Atta & Mulliez, 2004). Remarkably, SAM has also recently been shown to serve as cofactor in coordinating pericyclic reactions via electrostatic catalysis (Ohashi, Liu, Hai, Chen, Tang, Yang, et al., 2017). Another role for SAM is to serve as a substrate in reactions catalyzed by SAM-dependent halogenases of the 5'-halo-5'-deoxyadenosine synthase class (Dong, Huang, Deng, Schaffrath, Spencer, O'Hagan & Naismith, 2004; Eustáquio, Pojer, Noel & Moore, 2008; Agarwal, Miles, Winter, Eustáquio, El Gamal & Moore, 2017).

Halogenated compounds are pervasive across all domains of life. Over 5,000 organohalogens are known. Organochlorines and organobromines are the most abundant with >2,000 known examples of each, whereas organoiodines are more scarce (<200) and organofluorines are very rare with only five rigorously identified compounds (Gribble, 2012; Chan & O'Hagan, 2012, and Wang, Zhou, Fredimoses, Liao & Liu, 2014). Halogenating enzymes employ various mechanistic strategies depending on the properties of the halogen atom and the reactivity of the organic substrate. Most halogenases activate chloride, bromide and iodide via oxidation by utilizing cofactors such as heme-iron, vanadium, flavin and Fe(II)/ $\alpha$ -ketoglutarate (Agarwal et al., 2017). For fluoride, on the other hand, which evades oxidation due to its extreme electronegativity, an orthogonal strategy has been described that uses SAM as co-substrate in an  $S_N2$ -type, nucleophilic substitution reaction (Figure 1). The only fluorinase described thus far (FIA) catalyzes the first committed step in the biosynthesis of fluoroacetate and fluorotheonine in the soil bacterium *Streptomyces cattleya* in which fluoride attacks the C5' electrophilic carbon of SAM, displacing L-methionine and generating 5'-fluoro-5'-deoxyadenosine (Dong et al., 2004).

Four years after the structural characterization of the fluorinase, the chlorinase SalL was reported from the marine bacterium *Salinispora tropica* (Eustáquio et al., 2008). SalL is analogous to fluorinase and the only other characterized SAM-dependent halogenase of the 5'-halo-5'-deoxyadenosine (XDA) synthase class (Figure 1). SalL catalyzes the first committed step in the biosynthesis of proteasome inhibitor salinosporamide A (Eustáquio et al., 2008 and 2009; Feling, Buchanan, Mincer, Kauffman, Jensen & Fenical, 2003). Both FIA and SalL can also catalyze the reverse reaction, i.e. synthesis of SAM from XDA. For FIA, the *in vitro* equilibrium lies in favor of FDA by a factor of three (Dong et al., 2004). That the forward reaction is favored by FIA is unsurprising given that the C-F bond is the strongest bond in organic chemistry. For chlorinase SalL, on the other hand, synthesis of SAM from CIDA and L-met is favored, with the relative enzyme efficiency ( $k_{cat}/K_m$ ) for the reverse reaction (SAM synthesis) being several orders of magnitude higher than the forward halogenation reaction (Eustáquio et al., 2008). The fluorinase was also reported to favor the reverse reaction *in vitro* when using chloride as the halide (Deng, Cobb, McEwan, McGlinchey, Naismith, O'Hagan et al., 2006).

As a versatile methyl donor to biomolecules, there has been considerable interest in exploring the synthesis of SAM analogs of chemical and biological utility. For instance, non-natural SAM analogs could be used to modify cellular targets such as DNA, RNA, proteins and metabolites. Moreover, SAM itself is an expensive and unstable cofactor,

degrading rapidly under physiological or alkaline conditions. Thus, the availability of methods to prepare SAM *in situ* are desirable.

Traditionally, SAM analogs have been chemically synthesized from *S*-adenosyl-L-homocysteine. However, this synthesis suffers from low yields and lack of diastereoselectivity (Dalhoff, Lukinavicius, Klimasauskas & Weinhold, 2006). Furthermore, SAM synthesis is challenging due to its inherent lability and tendency to racemize (Hoffman, 1986). Chemoenzymatic synthesis using chlorinase SalL and CIDA offers an alternative to chemical synthesis. In addition to being stable and commercially available, CIDA can also be synthesized economically from adenosine and SOCl<sub>2</sub> (Sinhababu, Bartel, Pochopin & Borchardt 1985). Chemoenzymatic synthesis of SAM and analogs *in situ* could also be coupled with downstream applications such as methylation or alkylation by methyltransferases. This strategy can also be used for the facile incorporation of isotopes into alkylated products.

We have demonstrated the use of SalL for the diastereoselective synthesis of SAM and SAM congeners from CIDA and L-met analogs (See section 3). We have also coupled the SalL-catalyzed production of SAM with DNA and with metabolite methylation (Lipson, Thomsen, Moore, Clausen, La Clair & Burkart 2013). Importantly, we have demonstrated that *in situ* generated SAM analogs were recognized by rat protein arginine methyltransferase 1 (rPRMT1) which alkylated its peptide substrate (Thomsen, Vogensen, Buchardt, Burkart, & Clausen 2013).

In this chapter, we describe procedures for the preparation and assay of SalL, the SalL-mediated chemoenzymatic synthesis of SAM analogs from CIDA and L-met congeners, and the *in situ* chemoenzymatic synthesis of SAM coupled to DNA, peptide and metabolite methylation, and to the incorporation of isotopes into alkylated products.

## 2. Preparation and Assay of Chlorinase SalL

### 2.1 Purification of Recombinant SalL from *Escherichia coli*

Recombinant SalL containing a N-terminal His<sub>8</sub> tag was obtained by gene expression in and protein purification from *E. coli* BL21(DE3) (Eustáquio et al., 2008).

**2.1.1 Cloning of the salL Gene into an Expression Vector**—The *salL* gene can be amplified by PCR using total genomic DNA isolated from *S. tropica* CNB-440 as a template and appropriate primers. The following are suggested primers that were used to clone *salL* into pHis8 (Jez, Ferrer, Bowman, Dixon & Noel, 2000) by restriction/ligation, yielding plasmid pAEM7 (Eustáquio et al., 2008):

for: 5'-CGTGGTTCCCATGG**CATG**CAGCACAATCTCATTGC-3' (NcoI site underlined)

rev: 5'-GCTCGAATTCAAGCTTGTCAGCTACCCGAGCACCG-3' (HindIII site underlined)

The primers were designed to ensure that the ATG (Met start codon, bold) was in-frame with the His<sub>8</sub> tag. The priming sites are in italics. Bases upstream of the restriction sites are

random sequences to facilitate cutting of the PCR product with the respective restriction enzyme. *salL* was amplified by PCR using the Expand High Fidelity PCR system (Roche) following the manufacturer's instructions and using 5% DMSO (v/v) and annealing temperature of 58 °C. Alternative, proofreading DNA polymerases include Q5® High Fidelity polymerase and Phusion® High Fidelity polymerase (New England Biolabs). The expected size of the PCR product when using primers above is 888 bp. Alternatively, synthetic DNA can be ordered from GenScript. The accession code for the *salL* gene (Strop\_1026) of *S. tropica* CNB-440 is NC\_009380.1 (position 1,162,700 to 1,163,551, complement). If using the primers above, digest the PCR product with NcoI and HindIII and ligate it into the same sites of pHIS8 to yield the pHIS8-*salL* expression vector (pAEM7). While the PCR product can be digested and cloned directly into pHIS8, we subcloned the PCR product into the pGEM-T Easy vector (Promega) by following the manufacturer's instructions. The insert of the obtained plasmid (pAEM5) was sequenced and confirmed to contain no mistakes. Following digestion of pAEM5 with HindIII, NcoI and DraI (DraI cuts only the vector backbone), and isopropanol precipitation, the HindIII-NcoI *salL* fragment (868 bp) was cloned into the same sites of pHIS8. pHIS8 contains an N-terminal, 8xHis tag that will facilitate protein purification. Note that the pAEM7 *salL* expression plasmid used here contains native DNA from *S. tropica* rather than DNA which is codon-optimized for expression in *E. coli*.

## 2.1.2 Expression of *salL* in *E. coli* BL21(DE3) and Purification of Recombinant *SalL* Enzyme

### 2.1.2.1 Materials

- Competent *E. coli* BL-21 (DE3) cells (e.g. Novagen)
- *salL* expression vector (pAEM7)
- Luria-Bertani (LB) agar (Becton Dickinson)
- LB broth (Becton Dickinson)
- Terrific Broth (TB) medium (Becton Dickinson)
- Ni<sup>2+</sup>-NTA resin (Qiagen)
- Lysis buffer: 50 mM sodium phosphate buffer (pH 8.0), 500 mM NaCl, 20 mM imidazole (pH 8.0), 20 mM β-mercaptoethanol, 10% (v/v) glycerol, and 1% (v/v) Tween-20, 1 mg/mL lysozyme
- Wash buffer: 50 mM sodium phosphate buffer (pH 8.0), 500 mM NaCl, 20 mM imidazole (pH 8.0), 20 mM β-mercaptoethanol, and 10% (v/v) glycerol
- Elution buffer: 50 mM sodium phosphate buffer (pH 8.0), 500 mM NaCl, 250 mM imidazole (pH 8.0), 20 mM β-mercaptoethanol, and 10% (v/v) glycerol
- Storage buffer: 50 mM sodium phosphate buffer (pH 7.9), 20% glycerol
- Disposable polypropylene column (e.g. from Qiagen)
- Disposable PD-10 desalting columns (Sephadex G-25, GE Healthcare)

- SDS-PAGE (e.g. 10% Bis-Tris NuPAGE gel with MOPS running buffer, ThermoFisher)

### 2.1.2.2 Procedure

1. Competent *E. coli* BL-21 (DE3) cells are commercially available (e.g. Novagen, cat. No. 69450–3). Follow the manufacturer’s instructions to introduce the *sall*-containing expression vector into *E. coli* BL-21 (DE3) by chemical transformation. Plate the transformation reaction on LB agar containing 50 mg/L kanamycin (pHIS8 contains a kanamycin-resistance gene) and incubate the plates overnight at 37 °C.
2. Inoculate a single colony onto 50 mL LB containing 50 mg/L kanamycin (use a 250-mL flask) and incubate the flask overnight at 37 °C, 200 rpm.
3. Prepare cryo stocks for long-term storage by mixing the *E. coli* BL21(DE3) culture containing the *sall* plasmid 1:1 with sterile glycerol 40%, giving a final glycerol concentration of 20%. Store cryo stocks at –80 °C.
4. Use the overnight LB culture to inoculate 1 L of TB media containing 50 mg/L kanamycin so that the starting  $A_{600} = 0.1$  (10 to 20 mL of the overnight *E. coli* seed culture are expected to be necessary). Incubate the flask at 25 °C, 225 rpm until  $A_{600} = 0.8$ . Once an  $A_{600}$  of 0.8 is reached, induce cells with 0.25 mM IPTG (final concentration) and incubate overnight at 20 °C, 225 rpm. A 2.8-L Fernbach flask is recommended for cultivation of 1-L cultures.
5. Harvest cells by centrifugation (e.g. 15 min at  $5,000 \times g$ ) and resuspend the cell pellet in ~20 mL of cold lysis buffer. Incubate on ice for 30 min.
6. Note: take care to keep the preparation cold by cooling centrifuges, buffers, and keeping the sample on ice.
7. Lyse cells by sonication on ice (e.g. 2 sec on, 2 sec off for 3 min total). Alternatively, a French press can be used for cell lysis.
8. Centrifuge the lysate at high speed ( $> 10,000 \times g$ ) for 30 min at 4 °C.
9. The clear lysate is then purified using  $Ni^{2+}$ -NTA resin (Qiagen) and a modified batch purification method as briefly described below (refer to the manufacturer’s instructions “The QIAexpressionist” for more details).
10. Transfer the clear lysate to a clean, autoclaved beaker on ice, and add 5 mL  $Ni^{2+}$ -NTA to the clear lysate. Mix gently using a magnetic stirrer for 1 hour. Alternatively, use a rotary shaker for mixing the sample contained in a conical tube.
11. If using a beaker for mixing, transfer the lysate /  $Ni$ -NTA suspension to a conical tube and centrifuge at  $3,000 \times g$  for 5 min at 4 °C to collect the  $Ni$ -NTA resin.
12. Discard the supernatant and wash the resin twice with 25 mL wash buffer each as following. Add 25 mL wash buffer to the  $Ni$ -NTA pellet, resuspend gently, and centrifuge at  $3,000 \times g$  for 5 min at 4 °C. Discard supernatant and repeat.

13. After the last wash, resuspend the Ni-NTA pellet in ~5 mL wash buffer and transfer the resin to an empty polypropylene column with the bottom outlet capped. Remove bottom cap and discard flow-through. Wash once with 25 mL cold wash buffer. Discard flow through.
14. Elute the His8-tagged SaLL with 2× 5 mL elution buffer.
15. Assess the quality of the protein preparation using SDS-PAGE (e.g. 10% Bis-Tris NuPAGE gel with MOPS running buffer, ThermoFisher). The expected size of the His<sub>8</sub>-tagged protein is 32.4 kDa (Figure 2).
16. The expected yield is > 60 mg recombinant, soluble SaLL per liter of culture. Protein yield can be calculated using the Bradford assay (Bradford 1976).
17. Desalting can be performed using PD-10 columns (Sephadex G-25): cut the bottom tip off the column and let the storage liquid flow (discard); equilibrate the column with 25 mL storage buffer; load sample in a total volume of 2.5 mL per PD-10 column; discard the flow through; elute with 3.5 mL storage buffer. Alternatively, dialysis can be used for desalting.

## 2.2 SaLL Activity Assay

The reaction catalyzed by chlorinase SaLL is reversible. Chloride, bromide and iodide are accepted as the halide co-substrates in the reaction with SAM (forward reaction) to yield 5'-halo-5'-deoxyadenosine (XDA). Kinetic studies demonstrated that the reverse reaction (formation of SAM from XDA and L-met) is favored *in vitro*, with the relative enzyme efficiency ( $k_{cat}/K_m$ ) of the reverse reaction being orders of magnitude higher than the forward reaction (Eustáquio et al., 2008). The activity of recombinant SaLL can be investigated by a HPLC-based method as described below.

### 2.2.1 Reagents

- S-adenosyl-L-methionine (Sigma-Aldrich)
- KCl (analytical grade)
- L-methionine (Sigma-Aldrich)
- 5'-Chloro-5'-deoxyadenosine (Sigma-Aldrich)
- 50 mM sodium phosphate buffer pH 7.9
- Recombinant SaLL (section 2.1)

### 2.2.2 Instrumentation and Solvents

- Agilent HPLC 1200
- C18 column (250 × 4.60 mm, 5 μm particle size)
- Mobile phase A: 95% 50 mM KH<sub>2</sub>PO<sub>4</sub>; 5% acetonitrile
- Mobile phase B: 80% 50 mM KH<sub>2</sub>PO<sub>4</sub>; 20% acetonitrile
- Note: 50 mM KH<sub>2</sub>PO<sub>4</sub> gives pH ~ 4.5; there is no need to adjust the pH.

### 2.2.3 Procedure

1. In order to test if your SaL preparation is active, set up two different reactions by incubating SaL (400 nM) with
  - i. SAM (0.5 mM) and KCl (400 mM)
  - ii. L-methionine (10 mM) and CIDA (0.5 mM)in 50 mM sodium phosphate buffer (pH 7.9) at 37°C for 1 hour. In addition, set up following controls: reactions with inactivated enzyme (the enzyme is first boiled for 5 min and then, the reagents are added) and boiled enzyme only (the enzyme is boiled for 5 min in buffer).
2. Boil the reaction mixture for 2 min at 95 °C and centrifuge for 30 min, 10,000 × *g* to remove precipitated protein. 10 µl of clear supernatant is then analyzed by HPLC using a C18 column (250 × 4.60 mm, 5 µm particle size) and 1 mL/min flow rate. After equilibrating the system for at least 15 min using 100 % mobile phase A, run following gradient: 100% A for 1 min, 0 to 100% B for 29 min, 100% to 0% B over 5 min, 100% A for 5 min. Inject also SAM and CIDA standards for comparison. The retention times obtained under the conditions described were CIDA (16.4 min) and SAM (18.1 min).
3. Notes:
  - SAM is unstable and SAM solutions should be freshly prepared
  - Recombinant SaL purified from *E. coli* co-purifies with adenosine and CIDA. Thus, the boiled enzyme negative control may have a small peak for CIDA.

## 3. CHEMOENZYMATIC SYNTHESIS OF SAM ANALOGS

### 3.1 Overview of Chemoenzymatic Methods to Prepare SAM and SAM Analogs

SAM is biosynthesized by methionine adenosyltransferases (MAT) from methionine and adenosine triphosphate (ATP) (Cantoni, 1975; Figure 3). SAM is an essential cofactor for SAM-dependent methyltransferases to methylate various substrates including nucleic acids, proteins, lipids, and small molecules (Martin & McMillan 2002; Lin, 2011; Struck, Thompson, Wong & Micklefield, 2012; Bauerle, Schwalm, & Booker, 2015). While transmethylation produces small and modest chemical changes, these drastically affect molecular recognition of substrates by cellular targets to control processes such as epigenetic regulation of protein expression, neurotransmission, and natural products biosynthesis. The prevalence of SAM in cellular processes has inspired the development of analogs as chemical reporters for nucleic acid (Dalhoff Lukinavicius, Klimasauskas & Weinhold, 2006a; Dalhoff et al., 2006b; Lukinavicius, Lapiene, Stasevskij, Dalhoff, Weinhold & Klimasauskas 2007; Motorin, Burhenne, Teimer, Koynov, Willnow, Weinhold, et al., 2011; Vranken, Fin, Tufar, Hofkens, Burkart & Tor, 2016), protein (Peters, Willnow, Duisken, Kleine, Macherey, Duncan et al., 2010; Wang, Zheng, Yu, Deng & Luo, 2011; Islam, Zheng, Yu, Deng & Luo, 2011; Islam, Bothwell, Chen, Sengelaub, Wang, Deng & Luo, 2011; Wang, Islam, Liu, Zheng, Tang, Lailler et al., 2013; Wang & Luo, 2013; Islam, Chen, Wu,



Bothwell, Blum, Zeng et al., 2013; Guo, Wang, Zheng, Chen, Blum, Deng et al., 2014), small molecule (Zhang, Weller, Thorson & Rajska, 2006; Stecher, Teng, Ueberbacher, Remler, Schwab, Griengl et al., 2009; Lee, Sun, Zang, Kim, Alfaro & Zhou, 2010; Thomsen et al., 2013; Singh, Zhang, Huber, Sunkara, Hurley, Goff et al., 2014), and natural products methyltransferases (Zhang et al., 2006; Winter, Chiou, Bothwell, Xu, Garg, Luo et al., 2013).

The preparation of SAM and its analogs is especially challenging due to inherent lability, tendency to racemize under experimental conditions, and rapid degradation at physiological pH (Hoffman, 1986). Chemical synthesis of SAM and SAM derivatives involves nucleophilic substitution between *S*-adenosyl-L-homocysteine (SAH or AdoCy) and an alkyl halide to produce a mixture of two sulfonium diastereomers, usually in variable yields (de la Haba, Jamieson, Mudd & Richards, 1959; Dalhoff et al., 2006; Stecher et al., 2009). The *S,S* diastereomer serves as the methyl donor, while the *S,R* diastereomer inhibits the methyltransferase (Borchardt & Wu, 1976; Khani-Oskouee, Jones, & Woodard, 1984). Additionally, analogs of SAM where sulfur is replaced with selenium are more reactive methyl donors due to the longer, weaker Se-C bonds and do not racemize (Iwig & Booker, 2004). Notably, select *Se*-adenosyl-L-selenomethionines (SeAM), i.e. propargyl SeAM, are more stable than their SAM analogs (Bothwell, Islam, Chen, Zheng, Blum, Deng et al., 2012; Bothwell & Luo, 2014). Overall, these early synthetic efforts ignited investigations in which non-natural SAM analogs could be used by endogenous methyltransferases to modify and profile various cellular targets.

To complement chemical synthesis of SAM analogs, we pioneered chemoenzymatic methods to expand the range of SAM analogs available for labeling methyltransferase substrates (Lipson et al., 2013; Thomsen et al., 2013; Vranken et al., 2016). We employed two homologous halogenases, SalL and FIA, to produce SAM and SAM analogs. As discussed earlier, SalL catalyzes the reaction of SAM with chloride to produce 5'-chloro-5'-deoxyadenosine (CIDA) and L-methionine via nucleophilic attack in *Salinispora tropica* (Figure 1). Similarly, *Streptomyces cattleya* uses fluorinase FIA to react fluoride with SAM to afford 5'-fluoro-5'-deoxyadenosine (FDA) and L-methionine. Based on Le Chatelier's Principle, we hypothesized that under low salt concentrations and excess L-methionine, these halogenases could catalyze the biologically reversed reaction to produce SAM *in situ* (Figure 3) from easily prepared CIDA. Indeed, we observed SAM production by both SalL and FIA in various buffers, including Tris, HEPES, and phosphate buffers at pH 7–8, although SalL produced SAM in substantially higher yields than FIA. Furthermore, we generated a W190A point mutant of SalL that could accommodate larger alkyl substrates (Thomsen et al., 2013). Both wild type (WT) and mutant SalL generated alkyl, allylic, and benzylic SAM analogs, although the WT was most effective at generating small SAM analogs (methyl, ethyl), whereas both the WT and W190A mutant were equally effective at producing larger SAM analogs (propyl, butyl, allyl, and benzyl). Importantly, we demonstrated that our *in situ* generated SAM congeners were recognized by nucleic acid, peptide, and small molecule methyltransferases, and alkylated their respective substrates (Thomsen et al., 2013).



## 3.2 SaIL-Mediated Chemoenzymatic Synthesis of SAM and SAM Analogs

### 3.2.1 Reagents

- DMSO (Sigma-Aldrich)
- HPLC grade acetonitrile (Fisher Scientific)
- 5'-chloro-5'-deoxyadenosine (CIDA) (Sigma-Aldrich)
- L-methionine (Sigma-Aldrich)
- *S*-ethyl-L-homocysteine (Sigma-Aldrich)
- *S*-propyl-L-homocysteine (prepared as described by Thomsen et al., 2013)
- *S*-butyl-L-homocysteine (Toronto Research Chemicals, Inc.)
- *S*-allyl-L-homocysteine (prepared as described by Thomsen et al., 2013)
- *S*-propargyl-L-homocysteine (prepared as described by Thomsen et al., 2013)
- *S*-benzyl-L-homocysteine (prepared as described by Thomsen et al., 2013)
- *S*-phenylethyl-L-homocysteine (prepared as described by Thomsen et al., 2013)
- *S*-(2-amino-2-carboxylethyl)-L-homocysteine (Sigma-Aldrich)
- *S*-(2-amino-2-oxoethyl)-L-homocysteine (prepared as described by Thomsen et al., 2013)
- *S*-adenosyl-L-methionine (New England Biolabs)

### 3.2.2 Enzyme

- SaIL (expressed and purified as described above in section 2.1.2)

### 3.2.3 Stock Solutions and Buffers

- 300 mM CIDA in DMSO
- 150 mM L-methionine in 250 mM NaOH
- 15–150 mM L-methionine analog in 250 mM NaOH
- 32 mM SAM in 5 mM H<sub>2</sub>SO<sub>4</sub> and 10% ethanol
- 50 mM sodium phosphate buffer, pH 6.8
- 400 mM sodium formate, pH 3.0
- 50 mM HEPES, pH 7.3

### 3.2.4 Analytical Instrumentation

- Agilent 6230 ESI-TOF LC-MS
- Agilent HPLC 1200
- Agilent Poroshell 300SB (5 μm, 2.1×75 mm) column

### 3.2.5 Procedure

1. Equilibrate 200  $\mu$ M CIDA and 3  $\mu$ M SalL in 50 mM phosphate buffer, pH 6.8 at 37  $^{\circ}$ C.
2. Add 1.5–15 mM L-methionine or analog from stock. Due to the poor solubility of the benzyl and phenylethyl analogs of L-methionine, these were added at a final concentration of 1.5 mM from a 15 mM stock solution in 250 mM NaOH. L-Methionine and all other analogs were added at a final concentration of 15 mM from a 150 mM stock solution in 250 mM NaOH. In all assays, the addition of L-methionine or analog was 10% of the total reaction volume, resulting in a final pH of 7.4.
3. Quench reactions at various time points (0–5 hours) by the addition of equal volumes of 400 mM sodium formate. Store quenched aliquots at 4  $^{\circ}$ C until HPLC analysis.
4. Analyze aliquots using analytical reversed phase HPLC using a Synergi<sup>TM</sup> Hydro RP column (4  $\mu$ m, 4.6 $\times$ 150 mm, Phenomenex) using the following gradient:  
Buffer A: 10 mM  $\text{NH}_4\text{HCO}_2$ , pH 3.2  
Buffer B: 90%  $\text{CH}_3\text{CN}$ , 10 mM  $\text{NH}_4\text{HCO}_2$ , pH 3.2  
0 min: 100% buffer A, 1 mL/min  
5 min: 98% buffer A, 1.5 mL/min  
10 min: 70% buffer A, 2 mL/min  
12 min: 0% buffer A, 2 mL/min  
Note: Complete conversion of CIDA (retention time = 11 min) into SAM (retention time = 4 min) was observed within 30 min.
5. Verify the identity of CIDA and SAM by collecting 0.1 mL fractions from the HPLC and analyze by LC-MS. Table 1 provides a summary of LC-MS data for select SAM analogs. See Thomsen, et al. 2013 for representative LC-MS traces and chromatograms.

### 3.3 *In situ* Chemoenzymatic Synthesis of SAM Coupled to DNA Methylation by HhaI

HhaI is a DNA methyltransferase that methylates cytosine at C5 (Sankpal & Rao, 2002). In this assay, HhaI is coupled with *in situ* SAM production with methylation of lambda DNA.

#### 3.3.1 Reagents

- N6-methyladenine free lambda DNA (New England Biolabs)

#### 3.3.2 Enzymes

- SalL (expressed and purified as described above in section 2.1.2)
- HhaI methyltransferase (New England Biolabs)
- 0.2 U/ $\mu$ L HinPII restriction enzyme (New England Biolabs)

### 3.3.3 Stock Solutions and Buffers

- 300 mM CIDA in DMSO
- 150 mM L-methionine in 250 mM NaOH
- HhaI buffer (5 mM 2-mercaptoethanol, 10 mM EDTA, 50 mM Tris•HCl, pH 7.5)
- NEB4 buffer (10 mM KOAc, 4 mM Tris•HOAc, 2 mM MgOAc<sub>2</sub>, 0.2 mM DTT, pH 7.9)

### 3.3.4 Procedure

1. Equilibrate 50 ng/μL N<sup>6</sup>-methyladenine free lambda DNA, 4 U/μL HhaI, 200 μM CIDA, and 15 mM L-methionine in HhaI buffer at 37 °C.

Note: The buffers used for DNA methyltransferase HhaI and SalL activity assays were incompatible. HhaI was inactive in 50 mM sodium phosphate buffer, pH 7.9, however, the activity of SalL was reduced by 30% in HhaI buffer. Thus, coupled assays were conducted in HhaI buffer.

2. Add SalL to a final concentration of 900 nM.
3. After 1 hour, quench reaction by heating at 80 °C for 5 min.
4. Dilute reaction in an equal volume of HinP1I in 2x NEB4 buffer to afford a final concentration of 0.1 U/μL HinP1I.
5. Incubate 1 hour at 37 °C.
6. Analyze lambda DNA cleavage by gel electrophoresis on 2% agarose gel.

## 3.4 *In situ* Chemoenzymatic Synthesis of SAM Coupled to Teicoplanin Methylation by MtfA

MtfA is a small molecule methyltransferase from the chloroeremomycin biosynthetic pathway in *Amycolatopsis orientalis* that catalyzes *N*-methylation of vancomycin-like glycopeptides (Shi et al., 2009). In this assay, MtfA is used to couple *in situ* SAM production with methylation of teicoplanin.

### 3.4.1 Reagents

- Teicoplanin (TargetMol)

### 3.4.2 Enzymes

- SalL (expressed and purified as described above in section 2.1.2)
- MtfA methyltransferase (expressed and purified as described below)

### 3.4.3 Stock Solutions and Buffers

- 300 mM CIDA in DMSO
- 150 mM L-methionine (or [*methyl*-<sup>13</sup>C]-L-methionine) in 250 mM NaOH
- 50 mM HEPES, pH 7.3

**3.4.4 Expression and Purification of MtfA Methyltransferase**—One liter of 2x YT medium (Sigma-Aldrich) supplemented with 50 µg/mL kanamycin was inoculated with *E. coli* BL21 (DE3) cells transformed with *mtfA*/pET28a plasmid (obtained from Prof. Gerard D. Wright, McMaster University). The cells were cultured at 37 °C to an OD<sub>600</sub> of 0.6. The cells were cooled to 0 °C on ice, induced with 100 µM isopropyl β-D-1-thiogalactopyranoside (IPTG), and grown for 20 h at 16 °C. Cells were harvested via centrifugation at 2,000 × g, 30 min, 4 °C and the pellet was resuspended in lysis buffer (50 mM HEPES, 500 mM NaCl, 20 mM imidazole, and 1 mM dithiothreitol (DTT), pH 7.5) and lysed by passage through a French pressure cell. The lysate was clarified by centrifugation at 4150 × g, 1.5 h, 4 °C. His<sub>6</sub>-MtfA was purified using 2 mL Ni-NTA resin (GE Healthcare) that was equilibrated with lysis buffer. The column was washed with 40 mL of buffer A (50 mM HEPES, 1 M NaCl, 20 mM imidazole, 1 mM DTT, pH 7.5), followed by 40 mL of buffer B (50 mM HEPES, 400 mM NaCl, 30 mM imidazole, 1 mM DTT, pH 7.5). The protein was eluted with buffer C (50 mM HEPES, 400 mM NaCl, 200 mM imidazole, 1 mM DTT, pH 7.5). We obtained around 50 mg of MtfA per L of cell culture. Eluted proteins were visualized with 12% SDS-PAGE with Coomassie Brilliant Blue staining. Imidazole was removed using spin concentration into 20 mM HEPES, 20 mM NaCl, and 5 mM DTT, pH 7.5. Pure protein was stored at −20 °C in 10 mM HEPES, 10 mM NaCl, 2.5 mM DTT, pH 7.5 containing 50% glycerol. The protein is stable as a glycerol stock at −20 °C.

#### 3.4.5 Procedure

1. Equilibrate a 50 µL mixture of 20 mM CIDA, 50 mM L-methionine, and 1.4 µM SalL in 32 mM phosphate buffer, pH 7.9 at 37 °C.
2. After 1–3 h, add 250 µL of a solution containing: 1 mM teicoplanin, 1 mg/mL BSA, and 54 µM MtfA in 50 mM HEPES, pH 7.3.
3. Monitor the methylation of teicoplanin by removing aliquots at various times and analyzing by LC-MS using a C18 column (3 µm, 2.1×150 mm, Atlantis) and an isocratic mixture of 3:1 H<sub>2</sub>O/CH<sub>3</sub>CN (v/v) with 0.1% formic acid. Under these conditions, the retention times of teicoplanin and *N*-methyl-teicoplanin were 9 min and 9.5 min, respectively.

### 3.5 *In situ* Chemoenzymatic Synthesis of <sup>13</sup>C-labeled SAM Coupled with Isotope Incorporation into Teicoplanin via Methylation by MtfA

We envisioned SalL-mediated *in situ* chemoenzymatic synthesis of SAM as a compelling and facile way to incorporate isotopes into biomolecules. Here we describe the use of this technology to incorporate <sup>13</sup>C into teicoplanin for sensitive NMR identification (Lipson et al., 2013).

#### 3.5.1 Reagents

- d<sub>6</sub>-DMSO (Cambridge Isotope Laboratories, Item No. DLM-10–10)
- [*Methyl*-<sup>13</sup>C]-L-methionine (Cambridge Isotope Laboratories, Item No. CLM-206-PK)
- Teicoplanin (TargetMol)

### 3.5.2 Enzymes

- SalL (expressed and purified as described above in section 2.1.2)
- MtfA methyltransferase (expressed and purified as described in section 3.4.4)

### 3.5.3 Stock Solutions and Buffers

- 300 mM CIDA in d<sub>6</sub>-DMSO
- 300 mM [*methyl*-<sup>13</sup>C]-L-methionine in d<sub>6</sub>-DMSO
- 10 mM phosphate buffer, pH 7.5

### 3.5.4 Analytical Instrumentation

- Varian VX500 NMR spectrometer equipped with an X-Sens cold probe

### 3.5.5 Procedure

1. Prepare 800 μL of a mixture of 350 nM SalL and 350 nM MtfA in 10 mM phosphate buffer, pH 7.5.
2. Prepare 200 μL of a mixture of 7 mM teicoplanin, 142 mM [*methyl*-<sup>13</sup>C]-L-methionine, and 14.2 mM CIDA in 10 mM phosphate buffer, pH 7.5 containing 50% d<sub>6</sub>-DMSO.
3. Transfer the mixture of teicoplanin, [*methyl*-<sup>13</sup>C]-L-methionine, and CIDA mixture to the mixture of SalL and MtfA.
4. Transfer the entire solution to a clean 5 mm NMR tube.
5. Collect <sup>13</sup>C-NMR spectra on a Varian VX500 NMR spectrometer at various time points. For each time point, we acquired data for 64 scans with a delay time (d1) of 5 seconds. In our hands, we observed the appearance of [*methyl*-<sup>13</sup>C]-*N*-teicoplanin within 3–9 hours, as evidenced by a <sup>13</sup>C-NMR signal at δ = 33.7 ppm.

## 3.6 *In situ* Chemoenzymatic Synthesis of SAM Coupled to RGG Peptide Methylation by rPRMT1

Protein arginine methyltransferase 1 (PRMT1) is responsible for the majority of arginine methylation in mammalian cells. In the assay described below, rPRMT1, a rat homolog that dimethylates arginine residues, is used to couple *in situ* SAM production with methylation of arginine residues in the RGG peptide. Transfer of ethyl, allyl, and benzyl congeners of L-methionine has also been observed (Thomsen et al., 2013). Overall, sterics likely influences transfer since we observed RGG allylation, but did not observe RGG propylation, although the corresponding SAM analogs have similar sizes and are formed *in situ* at similar rates. RGG allylation most likely occurs via an S<sub>N</sub>2' mechanism.

### 3.6.1 Reagents

- RGG peptide (expressed and purified as described by Thomsen et al., 2013)

### 3.6.2 Enzymes

1. SalL (expressed and purified as described above in section 2.1.2)
2. Rat protein arginine methyltransferase 1 (rPRMT1, expressed and purified as described by Thomsen et al., 2013)

### 3.6.3 Stock Solutions and Buffers

- 300 mM ClDA in DMSO
- 150 mM L-methionine in 250 mM NaOH
- 400 mM sodium formate, pH 3.0
- 50 mM phosphate buffer, pH 7.4

### 3.6.4. Procedure Peptide Methylation using rPRMT1

1. Equilibrate a mixture containing 200  $\mu$ M ClDA, 15 mM L-methionine, 1  $\mu$ M SalL, 10  $\mu$ M RGG peptide, and 2  $\mu$ M rPRMT1 in 50 mM phosphate buffer, pH 7.4 at 37 °C.
2. After 12 hours, quench reaction by the addition of an equal volume of 400 mM sodium formate.
3. Assess the formation of methylated RGG peptide using LC-MS.

## 4. SUMMARY AND CONCLUSIONS

We described procedures for the chemoenzymatic synthesis of SAM and analogs that employ chlorinase SalL discovered from a marine actinomycete bacterium. Recombinant SalL can be purified from *E. coli* in relatively high yields, i.e. over 60 mg soluble enzyme per liter culture. The SalL-catalyzed synthesis of SAM analogs from L-methionine congeners and the stable and commercially available substrate 5'-chloro-5'-deoxyadenosine is diastereoselective. Moreover, we described methods for *in situ* synthesis of SAM coupled with DNA, protein and metabolite methylation, and with the incorporation of isotopes into biomolecules.

In addition to SAM-dependent halogenases, numerous groups have used methionine adenosyltransferase wild type and mutants that accommodate structural changes in L-methionine and ATP to generate stable SAM analogs that alkylate a variety of acceptor substrates (Wang et al., 2013; Singh et al., 2014; Huber, Wang, Singh, Johnson, Zhang, Sunkara et al., 2016; Huber, Johnson, Zhang, & Thorson, 2016). Overall, the development of chemoenzymatic methods and protein engineering efforts has expanded the structural scope of SAM analogs that can be generated for biological studies.

## ACKNOWLEDGEMENTS

Financial support for this work was provided by the National Center for Advancing Translational Sciences, National Institutes of Health (NIH), under grant KL2TR002002 (to A.S.E.), by NIH GM095970 (to M.D.B) and by startup funds from the Department of Medicinal Chemistry and Pharmacognosy and the Center for Biomolecular Sciences, University of Illinois at Chicago (to A.S.E.). T.D.D. was supported by the NIH/NIGMS IRACDA K12




GM068524 award. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Plasmid pAEM7 for the expression of *salL* was constructed in Prof. Bradley S. Moore's laboratory (Scripps Institution of Oceanography). The authors also graciously thank Prof. Joseph P. Noel (Salk Institute for Biological Studies) for vector pHIS8, Prof. Gerard D. Wright (McMaster University) for the *mtfA* gene construct, Dr. Yongxuan Su (UC San Diego) for mass spectral analyses, and Drs. Anthony Mrse and Xuemei Huang (UC San Diego) for assistance with acquiring NMR spectral data.

## REFERENCES

- Agarwal V, Miles ZD, Winter JM, Eustáquio AS, El Gamal AA, & Moore BS (2017). Enzymatic halogenation and dehalogenation reactions: pervasive and mechanistically diverse. *Chemical Reviews*, 117, 5619–5674. [PubMed: 28106994]
- Bauerle MR, Schwalm EL, & Booker SJ (2015). Mechanistic diversity of radical S-adenosylmethionine (SAM)-dependent methylation. *Journal of Biological Chemistry*, 290, 3995–4002. [PubMed: 25477520]
- Borchardt RT & Wu YS (1976). Potential inhibitors of S-adenosylmethionine-dependent methyltransferases V: role of the asymmetric sulfonium pole in the enzymatic binding of S-adenosyl-L-methionine. *Journal of Medicinal Chemistry*, 19, 1099–1103. [PubMed: 978674]
- Bothwell IR, Islam K, Chen Y, Zheng W, Blum G, Deng H, & Luo M (2012). Se-adeonsyl-L-selenomethionine cofactor analogue as a reporter of protein methylation. *Journal of the American Chemical Society*, 134, 14905–14912. [PubMed: 22917021]
- Bothwell IR & Luo M (2014). Large-scale, protection free synthesis of Se-adenosyl-L-selenomethionine analogues and their application as cofactor surrogates of methyltransferases. *Organic Letters*, 16, 3056–3059. [PubMed: 24852128]
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72, 248–254. [PubMed: 942051]
- Cantoni GL (1975). Biological methylation: selected aspects. *Annual Review of Biochemistry*, 44, 435–451.
- Chan KK, & O'Hagan D (2012). The rare fluorinated natural products and biotechnological prospects for fluorine enzymology. *Methods in Enzymology*, 516, 219–235. [PubMed: 23034231]
- Dalhoff C, Lukinavicius G, Klimasauskas S, & Weinhold E (2006). Direct transfer of extended groups from synthetic cofactors by DNA methyltransferases. *Nature Chemical Biology*, 2, 31–32. [PubMed: 16408089]
- Dalhoff C, Lukinavicius G, Klimasauskas S, & Weinhold E (2006). Synthesis of S-adenosyl-L-methionine analogs and their use for sequence-specific transalkylation of DNA by methyltransferases. *Nature Protocols*, 1, 1879–1886. [PubMed: 17487172]
- Deng H, Cobb SL, McEwan AR, McGlinchey RP, Naismith JH, O'Hagan D, Robinson DA, & Spencer JB (2006). The fluorinase from *Streptomyces cattleya* is also a chlorinase. *Angewandte Chemie*, 45, 759–762. [PubMed: 16370017]
- Dong C, Huang F, Deng H, Schaffrath C, Spencer JB, O'Hagan D, & Naismith JH (2004). Crystal structure and mechanism of a bacterial fluorinating enzyme. *Nature*, 427, 561–565. [PubMed: 14765200]
- Eustáquio AS, Pojer F, Noel JP & Moore BS (2008). Discovery and characterization of a marine bacterial SAM-dependent chlorinase. *Nature Chemical Biology*, 4, 66–74.
- Eustáquio AS, McGlinchey RP, Liu Y, Hazzard C, Beer LL, Florova G, Alhamadsheh MM, Lechner A, Kale AJ, Kobayashi Y, Reynolds KA & Moore BS (2009). Biosynthesis of the salinosporamide A polyketide synthase substrate chloroethylmalonyl-coenzyme A from S-adenosyl-L-methionine. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 12295–12300. [PubMed: 19590008]
- Feling RH, Buchanan GO, Mincer TJ, Kauffman CA, Jensen PR & Fenical W (2003). Salinosporamide A: a highly cytotoxic proteasome inhibitor from a novel microbial source, a marine bacterium of the new genus *Salinospira*. *Angewandte Chemie*, 42, 355–357. [PubMed: 12548698]


- Fontecave M, Atta M, & Mulliez E (2004). S-adenosylmethionine: nothing goes to waste. *Trends in Biochemical Sciences*, 29, 243–249. [PubMed: 15130560]
- Gribble GW (2012). Recently discovered naturally occurring heterocyclic organohalogen compounds. *Heterocycles*, 84, 157–207.
- Guo H, Wang R, Zheng W, Chen Y, Blum G, Deng H, & Luo M (2014). Profiling substrates of protein arginine N-methyltransferase 3 with S-adenosyl-l-methionine analogues. *ACS Chemical Biology*, 9, 476–484. [PubMed: 24320160]
- de la Haba G, Jamieson GA, Mudd SH, & Richards HH (1959). S-adenosylmethionine: the relation of configuration at the sulfonium center to enzymatic reactivity. *Journal of the American Chemical Society*, 81, 3975–3980.
- Hoffman JL (1986). Chromatographic analysis of the chiral and covalent instability of S-adenosyl-l-methionine. *Biochemistry*, 25, 4444–4449. [PubMed: 3530324]
- Huber TD, Johnson BR, Zhang J, & Thorson JS (2016). AdoMet analog synthesis and utilization: current state of the art. *Current Opinion in Biotechnology*, 42, 189–197. [PubMed: 27506965]
- Huber TD, Wang F, Singh S, Johnson BR, Zhang J, Sunkara M, Van Lanen SG, Morris AJ, Philips GN Jr., & Thorson JS (2016). Functional AdoMet isosteres resistant to classical AdoMet degradation pathways. *ACS Chemical Biology*, 11, 2484–2491. [PubMed: 27351335]
- Islam K, Zheng W, Yu H, Deng H, & Luo M (2011). Expanding cofactor repertoire of protein lysine methyltransferase for substrate labeling. *ACS Chemical Biology*, 6, 679–684.
- Islam K, Bothwell I, Chen Y, Sengelaub C, Wang R, Deng H, & Luo M (2012). Bioorthogonal profiling of protein methylation using azido derivative of S-adenosyl-l-methionine. *Journal of the American Chemical Society*, 134, 5909–5915. [PubMed: 22404544]
- Islam K, Chen Y, Wu H, Bothwell IR, Blum GJ, Zeng H, Dong A, Zheng W, Min J, Deng H, & Luo M (2013). Defining efficient enzyme-cofactor pairs for bioorthogonal profiling of protein methylation. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 16778–16783. [PubMed: 24082136]
- Iwig DF, & Booker SJ (2004). Insight into the polar reactivity of the onium chalcogen analogues of S-adenosyl-l-methionine. *Biochemistry*, 43, 13496–13509. [PubMed: 15491157]
- Jez JM, Ferrer JL, Bowman ME, Dixon RA & Noel JP (2000). Dissection of malonyl-coenzyme A decarboxylation from polyketide formation in the reaction mechanism of a plate polyketide synthase. *Biochemistry*, 39, 890–902. [PubMed: 10653632]
- Khani-Oskouee S, Jones JP, & Woodard RW (1984). Stereochemical course of the biosynthesis of l-aminocyclopropane-1-carboxylic acid. I. Role of the asymmetric sulfonium pole and the  $\alpha$ -amino acid center. *Biochemical and Biophysical Research Communications*, 121, 181–187. [PubMed: 6732801]
- Lee BW, Sun HG, Zang T, Kim BJ, Alfaro JF, & Zhou ZS (2010). Enzyme-catalyzed transfer of a ketone group from an S-adenosylmethionine analogue: a tool for the functional analysis of methyltransferases. *Journal of the American Chemical Society*, 132, 3642–3643. [PubMed: 20196537]
- Lin H (2011). S-adenosylmethionine-dependent alkylation reactions: when are radical reactions used? *Bioorganic Chemistry*, 39, 161–170. [PubMed: 21762947]
- Lipson JM, Thomsen M, Moore BS, Clausen RP, La Clair JJ, & Burkart MD (2013). A tandem chemoenzymatic methylation of S-adenosyl-l-methionine. *ChemBioChem*, 14, 950–953. [PubMed: 23650044]
- Lukinavicius G, Lapiene V, Stasevskij Z, Dalhoff C, Weinhold E, & Klimasauskas S (2007). Targeted labeling of DNA by methyltransferase-directed transfer of activated groups (mTAG). *Journal of the American Chemical Society*, 129, 2758–2759. [PubMed: 17309265]
- Martin JL, & McMillan FM (2002). SAM (dependent) I AM: the S-adenosylmethionine-dependent methyltransferase fold. *Current Opinion in Structural Biology*, 12, 783–793. [PubMed: 12504684]
- Motorin Y, Burhenne J, Teimer R, Koynov K, Willnow S, Weinhold E, & Helm M (2011). Expanding the chemical scope of RNA: methyltransferases to site-specific alkynylation of RNA for click labeling. *Nucleic Acids Research*, 39, 1943–1952. [PubMed: 21037259]

- Ohashi M, Liu F, Hai Y, Chen M, Tang MC, Yang Z, Sato M, Watanabe K, Houk KN, Tang Y (2017). SAM-dependent enzyme-catalysed pericyclic reactions in natural product biosynthesis. *Nature*, 549, 502–506. [PubMed: 28902839]
- Peters W, Willnow S, Duisken M, Kleine H, Macherey T, Duncan KE, Litchfield DW, Luscher B, & Weinhold E (2010). Enzymatic site-specific functionalization of protein methyltransferase substrates with alkynes for click labeling. *Angewandte Chemie International Edition England*, 49, 5170–5173.
- Sankpal UT & Rao DN (2002). Structure, function, and mechanism of HhaI DNA methyltransferases. *Critical Reviews in Biochemistry and Molecular Biology*, 37, 167–197. [PubMed: 12139442]
- Shi R, Lamb SS, Zakeri B, Proteau A, Cui Q, Sulean T, Matte A, Wright GD, & Cygler M (2009). Structure and function of the glycopeptide N-methyltransferase MtfA, a tool for the biosynthesis of modified glycopeptide antibiotics. *Chemistry & Biology*, 16, 401–410. [PubMed: 19389626]
- Singh S, Zhang J, Huber TD, Sunkara M, Hurley K, Goff RD, Wang G, Zhang W, Liu C, Rohr J, Van Lanen SG, Morris AJ, & Thorson JS (2014). Facile chemoenzymatic strategies for the synthesis and utilization of S-adenosyl-L-methionine analogues. *Angewandte Chemie International Edition*, 53, 3965–3969. [PubMed: 24616228]
- Sinhababu AK, Bartel RL, Pochopin N, & Borchardt RT (1985). Mechanism of action of S-adenosyl-L-homocysteine hydrolase. Measurement of kinetic isotope effects using adenosine-3'-d and S-adenosyl-L-homocysteine-3'-d as substrates. *Journal of American Chemical Society*, 107, 7628–7632.
- Stecher H, Teng M, Ueberbacher BJ, Remler P, Schwab H, Griengl H, & Gruber-Khadjawi M (2009). Biocatalytic Friedel-Crafts alkylation using non-natural cofactors. *Angewandte Chemie International Edition England*, 48, 9546–9548.
- Struck AW, Thompson ML, Wong LS, & Micklefield J (2012). S-adenosyl-methionine-dependent methyltransferases: highly versatile enzymes in biocatalysis, biosynthesis and other biotechnological applications. *ChemBioChem*, 13, 2642–2655. [PubMed: 23180741]
- Thomsen M, Vogensen SB, Buchardt J, Burkart MD, & Clausen RP (2013). Chemoenzymatic synthesis and in situ application of S-adenosyl-L-methionine analogs. *Organic and Biomolecular Chemistry*, 11, 7606–7610. [PubMed: 24100405]
- Vranken C, Fin A, Tufar P, Hofkens J, Burkart MD, & Tor Y (2016). Chemoenzymatic synthesis and utilization of a SAM analog with an isomorphous nucleobase. *Organic and Biomolecular Chemistry*, 14, 6189–6192. [PubMed: 27270873]
- Wang R, Zheng W, Yu H, Deng H, & Luo M (2011). Labeling substrates of protein arginine methyltransferase with engineered enzymes and matched S-adenosyl-L-methionine analogues. *Journal of the American Chemical Society*, 133, 7648–7651. [PubMed: 21539310]
- Wang R, Islam K, Liu Y, Zheng W, Tang H, Lailier N, Blum G, Deng H, & Luo M (2013). Profiling genome-wide chromatin methylation with engineered posttranslation apparatus within living cells. *Journal of the American Chemical Society*, 135, 1048–1056. [PubMed: 23244065]
- Wang R, & Luo M (2013). A journey toward bioorthogonal profiling of protein methylation inside living cells. *Current Opinion in Chemical Biology*, 17, 729–737. [PubMed: 24035694]
- Wang L, Zhou X, Fredimoses M, Liao S & Liu Y (2014). Naturally occurring organoiodines. *RSC Advances*, 4, 57350–57376.
- Winter JM, Chiou G, Bothwell IR, Xu W, Garg NK, Luo M, & Tang Y (2013). Expanding the structural diversity of polyketides by exploring the cofactor tolerance of an inline methyltransferase domain. *Organic Letters*, 15, 3774–3777. [PubMed: 23837609]
- Zhang C, Weller RL, Thorson JS, & Rajski SR (2006). Natural product diversification using a non-natural cofactor analogue of S-adenosyl-L-methionine. *Journal of the American Chemical Society*, 128, 2760–2761. [PubMed: 16506729]




**Figure 1. Reaction catalyzed by SAM-dependent halogenases.**

Fluorinase FIA from soil bacterium *Streptomyces cattleya* (Dong et al., 2004) and chlorinase SaL from marine bacterium *Salinispora tropica* (Eustáquio et al., 2008) catalyze reversible S<sub>N</sub>2-type, nucleophilic substitution reactions in which the halide attacks the C5' ribosyl carbon of SAM, generating 5'-halo-5'-deoxyadenosine (XDA) and releasing L-methionine. FIA and SaL catalyze the first committed step in the biosynthesis of fluorometabolites and salinosporamide A, respectively.



**Figure 2. SDS-PAGE analysis of purified, recombinant SalL.**

10% Bis-Tris NuPAGE gel with MOPS running buffer (ThermoFisher). L, Fisherbrand EZ-Run protein marker; 1, soluble protein before induction; 2, soluble protein after induction; 3, his<sub>8</sub>-SalL purified by Ni-NTA affinity chromatography.



**Figure 3. Biosynthesis, utilization, and chemoenzymatic synthesis of *S*-adenosyl-L-methionine (SAM) analogs.**

(A) Biosynthesis of SAM *in vivo* is catalyzed by L-methionine-adenosyltransferases (MAT). Subsequently, nucleic acid, protein, or small molecule substrates are methylated by SAM-dependent methyltransferases (MT). SAH, *S*-adenosyl-L-homocysteine. (B) Chemoenzymatic synthesis of SAM analogs *in vitro* using chlorinase SalL from *Salinispora tropica*. CIDA, 5'-chloro-5'-deoxyadenosine.



**Table 1**

LC-MS Retention Times and Chromatographic Analyses of Select SAM Analogs

<b>Analog</b>	<b>Retention Time</b>	<b>Observed Monoisotopic Mass</b>
SAM	4 min.	399.1555
Ethyl-SAM	5 min.	413.1671
Propyl-SAM	8.5, 8.9 min. <sup>a</sup>	427.1784
Butyl-SAM	9.5 min.	441.2366
Allyl-SAM	7.2, 7.9 min. <sup>a</sup>	425.1862
Benzyl-SAM	10.4 min.	475.2242

<sup>a</sup>Mixture of diastereomers.