

ORIGINAL RESEARCH

Pre-Hospital Delay and Its Contributing Factors in Patients with ST-Elevation Myocardial Infarction; a Cross sectional Study

Hamidreza Poorhosseini¹, Mohammad Saadat², Mojtaba Salarifar¹, Seyedeh Hamideh Mortazavi², Babak Geraiely^{1*}

1. Interventional Cardiology Department, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran.

2. Cardiology Department, Tehran Heart Center, Tehran University of Medical Sciences, Tehran, Iran.

Received: March 2019; Accepted: April 2019; Published online: 29 May 2019

Abstract: **Introduction:** The outcome of ST-elevation myocardial infarction (STEMI) is significantly influenced by the total tissue ischemic time. In spite of efforts for reducing the in-hospital delay by full-time provision of primary percutaneous coronary intervention (P-PCI) in the 24/7 program, pre-hospital delay still persists. As a first report in Iran, we aimed to assess the duration of pre-hospital delay and its contributing factors in STEMI patients in the P-PCI era. **Methods:** The present cross-sectional study evaluated 2103 STEMI patients who underwent primary PCI from 2016 to 2018. Demographic, personal and socioeconomic factors, index event characteristics, past medical history, pain onset and door times of patients were recorded and independent factors of pre-hospital delay were calculated. **Results:** Median (IQR) of pain to door (P2D) time was 279 (120-630) minutes. In multivariate analysis, female gender [Beta=0.064 (95%CI: 0.003-0.125); p=0.038], being uneducated [Beta=0.213 (95%CI: 0.115-0.311); p<0.001], the onset of chest pain between 00:00 to 6:00 [Beta=0.130 (95%CI: 0.058-0.202); p<0.001] or 7:00 to 12:00 [Beta=0.119 (95%CI: 0.049-0.190); p=0.001], self-transportation [Beta=0.098 (95%CI: 0.015-0.181); p=0.020] or referral from another hospital [Beta=0.253 (95%CI: 0.117-0.389); p<0.001], atypical chest pain [Beta=0.170 (95%CI: 0.048-0.293); p=0.006], history of hypertension [Beta=0.052 (95%CI: 0.002-0.102); p=0.041], and opium abuse [Beta=0.076 (95%CI: 0.007-0.146); p=0.031] were associated with significantly higher log(P2D), while history of CABG was associated with shorter P2D. **Conclusion:** Our study showed that P2D is still very high in Iran and revealed the high-risk groups associated with longer P2D. Effective actions should be implemented to increase the public awareness about the symptoms of STEMI, and the importance of immediate appropriate help-seeking.

Keywords: ST-elevation myocardial infarction; myocardial infarction, STEMI; time-to-treatment

Cite this article as: Poorhosseini H, Saadat M, Salarifar M, Mortazavi S H, Geraiely B. Pre-Hospital Delay and Its Contributing Factors in Patients with ST-Elevation Myocardial Infarction; a Cross sectional Study. Arch Acad Emerg Med. 2019; 7(1): e29.

1. Introduction

Ischemic heart disease is still the most common cause of death worldwide (1-3). Several studies have shown that the morbidity and mortality of patients with ST-elevation myocardial infarction (STEMI) is significantly influenced by the total tissue ischemic time, which consists of pre-hospital and/or in-hospital delays (4-7). High expenditure strategies like primary percutaneous coronary intervention (P-PCI) for

STEMI and early invasive strategy for Non-STEMI are developed to reduce the in-hospital component of ischemic time; while a huge amount of golden time is lost in the pre-hospital phase. Efforts have been made in different countries to reduce the total ischemic time. While in-hospital delay has been reduced in many countries, even developing ones (8, 9), only developed countries have been able to reduce the pre-hospital delay by focusing on total ischemic time through increasing the general population's awareness via public educational programs in social media (10, 11).

Due to the implementation of full-time (24/7) provision of P-PCI services in our country by the ministry of health and medical education, the in-hospital delay has been reduced in

*Corresponding Author: Babak Geraiely; Tehran Heart Center, North Kargar Street, Tehran-Iran, P.O: 1411713138, Tel & Fax: +98 21 88029600, Email: bgeraiely@sina.tums.ac.ir



recent years (12). However, as long as the pre-hospital delay remains too long, the benefits of 24/7 P-PCI will be limited. There is no large-scale study evaluating the accurate duration of pre-hospital delay in STEMI patients in our country. Given this lack of information, we aimed to assess the duration of prehospital delay and its contributing factors in STEMI patients undergoing P-PCI.

2. Methods

2.1. Study design and setting

In the present cross-sectional study, we enrolled 2407 consecutive STEMI patients who underwent P-PCI between January 2016 and December 2018 at a tertiary cardiac center (Tehran Heart center)(13), Tehran, Iran. The hospital's local review board and Ethics Committee approved the study protocol (Ethics number: IR.TUMS.MEDICINE.REC.1397.954).

2.2. Participants

The study population consisted of all STEMI patients who were referred to the mentioned hospital during the study period and underwent P-PCI. Patients were excluded if the STEMI had occurred in the hospital (n=23). In addition, patients with missed data on pain or door times were excluded from the analysis (n=281). Finally, 2103 STEMI patients were included.

2.3. Data gathering

Data on the patients' demographic information, personal and socioeconomic factors, marital status, educational level, ethnicity, place of longest stay, insurance type, physical activity level, mode of transfer to hospital, pain characteristics, pain onset time, door times, cardiovascular risk factors, and patients' past medical history, as well as the infarct related artery were extracted from ischemic heart disease, angiography and angioplasty registries of the hospital, which have been described in details before (14).

Physical activity level was defined as high in professional athletes, intermediate in those who do usual daily activities and low in patients with the least or lack of physical activity.

2.4. Statistical Analysis

Continuous variables were presented as mean \pm standard deviations (SDs) if they assumed normal distributions and as medians (25th – 75th interquartile ranges: IQR) if they failed to assume normal distributions. Discrete variables were presented as numbers (percentages). Pain to door (P2D) was compared between groups using Mann–Whitney test and Kruskal–Wallis test as appropriate. The predictors exhibiting a borderline statistical relationship with pain to door time in the univariate analysis ($P \leq 0.15$) were taken for a multivariate logistic regression analysis to investigate their indepen-

dence. Backward elimination regression analysis was used to remove insignificant variables and log(P2D) was considered as a dependent variable. A $P \leq 0.05$ was considered statistically significant. All the statistical analyses were conducted using IBM SPSS Statistics for Windows, version 24.0 (IBM Corp, Armonk, NY).

3. Results

3.1. Baseline characteristics of participants

2407 consecutive patients were studied out of which 23 cases were excluded due to occurrence of STEMI within the hospital and 281 were excluded because of missed data on pain or door times. Finally, 2103 STEMI patients with the mean age of 59.49 ± 11.79 years were enrolled for analysis (76.4% male). Table 1 and 2 summarize the baseline characteristics of studied patients. 94.3% of the patients were married, 79.0% had a diploma or university level education, 77.1% were of Fars ethnicity. Self-transport was the most common form of transfer (86%). Table 3 shows the index event's characteristics. Median (IQR) of P2D time of patients was 279 (120-630) minutes.

3.2. Contributing factors of P2D delay (univariate analysis)

The results of univariate analysis are presented in table 1-3. Based on these analyses, female gender was associated with longer median of P2D time ($p < 0.001$) and higher educational level was associated with shorter P2D time ($p < 0.001$). Age had a significant relationship ($r = 0.036$, $p = 0.095$) with log(P2D), while the association was insignificant for BMI ($r = 0.004$, $p = 0.865$).

The P2D time was significantly shorter in those who were transferred to the hospital by EMS ($p < 0.001$). Despite the presence of some meaningful patterns, statistical significance was not observed regarding marital status ($p = 0.137$) and physical activity status ($p = 0.507$). Description of symptoms as atypical or typical chest pain ($p = 0.005$) and also epigastric pain ($p = 0.007$) was significantly associated with longer P2D. In addition, the history of diabetes ($p = 0.029$) and hypertension ($p = 0.004$) were associated with longer P2D. Although P2D was not different among those with and without the history of coronary stenting ($p = 0.924$), the history of coronary artery bypass graft (CABG) was associated with shorter P2D with borderline significance [191.0 (97.50-425.50) vs. 280.0 (120.0-630.0), $p = 0.085$].

3.3. Contributing factors of P2D delay (multivariate analysis)

After nine steps of the backward elimination method, eight variables remained in the final model (table 4). Female gender (Beta-Coefficient: 0.064, 95%CI: 0.003 - 0.125, $p = 0.038$), being uneducated (Beta: 0.213, 95%CI: 0.115 - 0.311,



Table 1: Demographic and socioeconomic characteristics

Variables	Number (%)	Prehospital Delay (minute)		P
		Median	IQR 25% - 75%	
Gender				
Male	1607(76.4)	255.00	110.00 - 595.00	<0.001
Female	496 (23.6)	337.00	160.00 - 790.00	
Marital status				
Married	1983 (94.3)	270.50	117.00 - 619.75	0.173
Single	17 (0.8)	314.00	122.50 - 625.50	
Divorced	18 (0.9)	227.50	83.00 - 1020.25	
Widowed	85 (4.0)	349.00	172.50 - 858.00	
Education				
University	279 (13.4)	206.00	90.00 - 465.00	<0.001
High school diploma	1362 (65.6)	265.00	115.00 - 606.75	
Elementary education	175 (8.4)	310.00	136.00 - 662.00	
Uneducated	260 (12.5)	400.50	184.25 - 1014.00	
Ethnicity				
Fars	1622 (77.1)	270.00	117.00 - 603.75	0.267
Turk	299 (14.2)	303.00	130.00 - 730.00	
Other	182 (8.7)	318.00	120.00 - 806.00	
Longest linger				
Tehran	1507 (71.7)	260.00	110.00 - 603.25	0.011
Main Cities	162 (7.7)	355.00	141.00 - 732.50	
Small Cities	161 (7.7)	312.00	140.25 - 689.75	
Village	273 (13.0)	300.00	138.00 - 659.50	
Insurance				
Social security	843 (40.1)	273.00	120.00 - 648.50	0.609
Health facilities	704 (33.5)	285.00	120.25 - 625.75	
Other companies	390 (18.5)	289.00	120.00 - 600.00	
No insurance	166 (7.9)	239.00	90.75 - 601.50	
Physical activity level				
High	49 (2.3)	206.00	128.75 - 574.00	0.507
Intermediate	1986 (94.4)	279.00	118.50 - 624.50	
Low	68 (3.2)	310.00	129.00 - 790.00	

$p < 0.001$), the onset of chest pain in 00:00 to 6:00 (Beta: 0.130, 95%CI: 0.058 - 0.202, $p < 0.001$) or 7:00 to 12:00 (Beta: 0.119, 95%CI: 0.049 - 0.190, $p = 0.001$), self-transportation (Beta: 0.098, 95%CI: 0.015 - 0.181, $p = 0.020$) or referral from another hospital (Beta: 0.253, 95%CI: 0.117 - 0.389, $p < 0.001$), description of symptoms as atypical chest pain (Beta: 0.170, 95%CI: 0.048 - 0.293, $p = 0.006$), history of hypertension (Beta: 0.052, 95%CI: 0.002 - 0.102, $p = 0.041$), and opium abuse (Beta: 0.076, 95%CI: 0.007 - 0.146, $p = 0.031$) were associated with longer P2D and the history of CABG (Beta: -0.124, 95%CI: -0.252 - -0.004, $p = 0.048$) was associated with shorter P2D time.

4. Discussion

Based on the findings of the present study, female gender, being uneducated, the onset of chest pain in 00:00 to 6:00 or 7:00 to 12:00, self-transportation or referral from another hospital, description of symptoms as atypical chest pain, history of hypertension and opium abuse were associated with longer P2D while history of CABG was associated with shorter P2D time.

Several studies have been performed in different countries to estimate the interval between pain onset and hospital arrival time. Table 5 demonstrates the median of prehospital delay in STEMI patients in various countries. As is evident grossly, developed countries have succeeded in reducing P2D to around 2 hours, while India as a developing country hasn't shown any obvious progress during these years. Limited studies with small sample sizes have been conducted regarding prehospital delay in Iran (Table 6). As is evident, all of them were performed before implementation of 24/7 program. Except for one study, all of them have small sample sizes and their results are greatly discordant. To the best of our knowledge, this is the first study to evaluate predictors of prehospital delay in Iran in a large population of STEMI patients undergoing P-PCI. In the current study, using multivariate analysis, pain to door time was found to be significantly higher in female gender, uneducated patients, those with onset of chest pain between 00:00 to 6:00 or 7:00 to 12:00, self-transported patients or individuals who were referred from other hospitals, patients with atypical chest pain



Table 2: Past medical history of the patients

Variables	Number (%)	Prehospital Delay (minute)		P
		Median	IQR 25% - 75%	
Diabetes mellitus				
Yes	657 (31.2)	309.50	123.25 - 629.25	0.029
No	1446 (68.8)	264.00	115.25 - 628.00	
Hypertension				
Yes	852 (40.5)	315.50	130.75 - 670.00	0.004
No	1251 (59.5)	250.00	110.00 - 596.25	
Hyperlipidemia				
Yes	880 (41.8)	266.50	116.25 - 612.00	0.383
No	1223 (58.2)	285.50	120.00 - 640.25	
Smoking				
Yes	739 (35.1)	252.00	111.50 - 612.00	0.235
No	1364 (64.9)	290.00	120.00 - 649.00	
Opium abuse				
Yes	303 (14.4)	340.00	133.00 - 663.00	0.081
No	1800 (85.6)	270.00	117.00 - 618.00	
Family history of CAD				
Yes	354 (16.8)	253.00	102.00 - 607.50	0.066
No	1749 (83.2)	285.00	120.00 - 642.00	
Cerebrovascular event				
Yes	78 (3.7)	266.00	111.00 - 547.50	0.852
No	2025 (96.3)	279.00	120.00 - 630.00	
Chronic kidney disease				
Yes	46 (2.2)	350.50	157.00 - 679.50	0.324
No	2057 (97.8)	274.00	119.75 - 627.75	
History of CABG				
Yes	77 (3.7)	191.00	97.50 - 425.50	0.085
No	2026 (96.3)	280.00	120.00 - 630.00	
History of myocardial infarction				
Yes	153 (7.3)	330.00	125.00 - 890.00	0.053
No	1950 (92.7)	270.00	119.00 - 610.00	
History of coronary stenting				
Yes	122 (5.8)	266.00	110.00 - 733.75	0.924
No	1981 (94.2)	279.50	120.00 - 621.25	
Infarct related artery				
LAD	1144 (54.4)	270.00	120.00 - 642.50	0.621
LCX	297 (14.1)	310.00	119.25 - 583.25	
RCA	612 (29.1)	287.00	120.00 - 645.75	
SVG	50 (2.4)	162.00	98.75 - 898.00	

Hx, history; CAD, coronary artery disease; CABG, coronary artery bypass graft; LAD, left anterior descending; LCX, left circumflex; RCA, right coronary artery; SVG, saphenous vein graft.

and history of hypertension and opium abuse; while history of CABG was associated with shorter pain to door time. In a study by Noorani et al. (15), prehospital delay has been shown to be associated with long distance from hospital, lower socioeconomic status and using ambulance. In a study by Moser et al. (11) several factors have been mentioned to be associated with prehospital delay including female gender, older age, worse socioeconomic status, history of angina, having cardiovascular risk factors and poor knowledge of the individual. In the current study we found that patients with chest pain between 00:00 to 6:00 or 7:00 to 12:00 had higher prehospital delays. On the contrary, patients

transferred by EMS and educated individuals had lower pain to door time. Infarct related artery had no significant effect in pain to door time in our study population. Our findings are in line with those of Peng et al. (16) who assessed 1088 STEMI patients. They demonstrated that prehospital delay was negatively correlated with high educational level, previous history of MI, transportation by ambulance, onset of pain during the daytime (6:00-18:00) and anterior and posterior MI. In our study, the level of education was negatively correlated with pain to door time. Similar to our work, the study of Heo et al. (17), reported a pain to door time of 144, 76 and 68 minutes in STEMI patients with low, moderate and high educa-

Table 3: Index event's characteristics

Index	Number	Prehospital Delay (minute)		P
		Median	IQR 25% - 75%	
Mode of transfer				
Ambulance	196 (9.3)	209.00	91.25 - 458.00	<0.001
Self-transport	1808 (86.8)	280.00	120.00 - 650.00	
Referral	99 (4.7)	364.00	208.50 - 684.00	
Pain onset time				
0 to 6	554 (26.4)	345.00	112.00 - 872.00	<0.001
7 to 12	617 (29.4)	324.00	135.00 - 677.50	
13 to 18	519 (24.7)	264.00	120.00 - 462.00	
19 to 24	412 (19.6)	205.00	106.25 - 549.50	
Pain description				
Typical chest pain	1984 (94.3)	270.00	117.00 - 613.00	0.005
Atypical chest pain	84 (4.0)	488.00	176.00 - 895.75	
No chest pain	35 (1.7)	307.00	99.00 - 705.00	
Pain duration				
>30 min	1183 (56.3)	300.00	120.00 - 663.00	0.022
11-30 min	801 (38.1)	248.00	112.50 - 581.50	
1-10 min	84 (4.0)	324.00	142.00 - 847.00	
No chest pain	35 (1.7)	307.00	99.00 - 705.00	
Back pain				
Yes	26 (1.2)	377.50	149.25 - 904.50	0.146
No	2077 (98.8)	275.50	119.00 - 620.25	
Epigastric pain				
Yes	277 (13.2)	335.50	144.25 - 701.75	0.007
No	1826 (86.8)	270.00	115.00 - 615.00	
Jaw pain				
Yes	14 (0.7)	221.50	131.25 - 553.75	0.640
No	2089 (99.3)	278.50	120.00 - 630.00	
Left precordial pain				
Yes	1022 (48.6)	265.00	113.50 - 612.00	0.162
No	1081 (51.4)	289.00	120.00 - 645.00	
Retro-sternal pain				
Yes	1400 (66.6)	270.00	115.00 - 630.00	0.223
No	703 (33.4)	285.00	127.50 - 616.00	
Right precordial pain				
Yes	11 (0.5)	345.00	205.00 - 610.00	0.495
No	2092 (99.5)	276.00	119.50 - 630.00	
Arm & shoulder pain				
Yes	71 (3.4)	227.50	140.25 - 574.00	0.815
No	2032 (96.6)	279.50	117.25 - 630.75	

tional levels, respectively. In MEDEA Study (18) on 486 acute MI patients, prehospital delay was higher in patients with low MI-knowledge. They also found that patients with atypical symptoms had higher prehospital delays, which corresponds to our findings.

Our study showed that P2D is still very high in Iran and revealed the high-risk groups associated with longer P2D. We assume that effective actions should be implemented to increase the general population's knowledge about the presentations of acute MI in order to decrease the time to seek treatment.

5. Limitations and Strengths

Being single-centered and retrospective design of the current work can be considered as our study limitations. We had missing values in pain or door times in 281 patients and thus we excluded them from the final analysis. We had no information regarding the patients' place of living and could not retrieve the data on their distance from the hospital. Meanwhile, the present study is the largest study that has been done to evaluate P2D in Iranian patients and the first study that has published after starting the 24/7 program. Unlike IPACE2 study, only STEMI patients, for whom P2D is applicable and can be defined, were included in our study.



Table 4: Multivariate analysis for prediction of log (p2d)

Variable	Beta Coefficient	95% Confidence Interval		P
		Lower	Upper	
Gender				
Male	-	-	-	-
Female	0.064	0.003	0.125	0.038
Education				
University	-	-	-	-
High school diploma	0.070	-0.002	0.143	0.058
Elementary education	0.082	-0.025	0.189	0.135
Uneducated	0.213	0.115	0.311	<0.001
Pain Onset Time				
19 to 24	-	-	-	-
13 to 18	0.043	-0.030	0.116	0.243
7 to 12	0.119	0.049	0.190	0.001
0 to 6	0.130	0.058	0.202	<0.001
Mode of Transfer				
Ambulance	-	-	-	-
Self-transfer	0.098	0.015	0.181	0.020
Referral	0.253	0.117	0.389	<0.001
Pain Description				
Typical chest pain	-	-	-	-
Atypical chest pain	0.170	0.048	0.293	0.006
No chest pain	0.066	-0.121	0.253	0.491
Hypertension				
No	-	-	-	-
Yes	0.052	0.002	0.102	0.041
Opium				
No	-	-	-	-
Yes	0.076	0.007	0.146	0.031
CABG				
No	-	-	-	-
Yes	-0.124	-0.252	-0.004	0.048

CABG: coronary artery bypass graft.

Table 5: Median of prehospital delay in ST-elevation myocardial infarction patients in various countries according to published reports

Country	Prehospital delay (minute)	Year
	290	1990
United States (10, 19)	84 in males	2002
	121 in females	
	59 in males	2006
	81 in females	
Denmark (20)	125	1998
Australia and New Zealand (21)	145	2008
South Korea (22)	130	2012
India (23, 24)	310	2003
	290	2016

6. Conclusion

In the present study female gender, transfer via vehicles other than ambulance, atypical chest pain, low level of education, late night and morning onset of pain, history of hypertension and opium abuse were associated with higher prehospital delay while history of CABG was associated with shortened P2D.

7. Appendix

7.1. Acknowledgements

The present work was supported by Tehran Heart Center, Tehran University of Medical Sciences.



Table 6: Iranian studies on prehospital delay

Study	Year	City	Number*	ACS forms	P2D
Momeni (25)	2011	Rasht	162	STEMI	120
Khosravi (26)	2011	Isfahan	103	STEMI	255
Farshidi (27)	2012	Hormozgan	227	STEMI & NSTEMI	N/A
IPACE2 (28)	2012	Tehran, Mashhad, Isfahan, Shiraz, Tabriz	1997	UA & NSTEMI & STEMI	265
Taghadosi (29)	2013	Kashan	117	STEMI & NSTEMI	129

*: number of patients, P2D: pain to door time (minutes), ACS: acute coronary syndrome.

7.2. Author contribution

All the authors met the criteria of authorship based on the recommendations of the international committee of medical journal editors.

Authors ORCIDs

Hamidreza Poorhosseini: 0000-0002-5733-9588
 Mohammad Saadat: 0000-0003-4950-8334
 Mojtaba Salarifar: 0000-0001-9062-3495
 Seyedeh Hamideh Mortazavi: 0000-0002-8167-2241
 Babak Geraiely: 0000-0001-6695-7751

7.3. Funding/Support

None.

7.4. Conflict of interest

None.

References

- Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. 2013;380(9859):2197-223.
- Finegold JA, Asaria P, Francis DP. Mortality from ischaemic heart disease by country, region, and age: statistics from World Health Organisation and United Nations. *Int J Cardiol*. 2013;168(2):934-45.
- Mendis S. Global progress in prevention of cardiovascular disease. *Cardiovasc Diagn Ther*. 2017;7(Suppl 1):S32-s8.
- Avorn J, Knight E, Ganz DA, Schneeweiss S. Therapeutic delay and reduced functional status six months after thrombolysis for acute myocardial infarction. *The American journal of cardiology*. 2004;94(4):415-20.
- Cullen L, Greenslade JH, Menzies L, et al. Time to presentation and 12-month health outcomes in patients presenting to the emergency department with symptoms of possible acute coronary syndrome. *Emergency Medicine Journal*. 2016;33(6):390-5.
- Nilsson G, Moee T, Soderstrom L, Samuelsson E. Pre-hospital delay in patients with first time myocardial infarction: an observational study in a northern Swedish population. *BMC cardiovascular disorders*. 2016;16(1):93.
- De Luca G, Suryapranata H, Ottervanger JP, Antman EM. Time delay to treatment and mortality in primary angioplasty for acute myocardial infarction: every minute of delay counts. *Circulation*. 2004;109(10):1223-5.
- Akimbaeva Z, Ismailov Z, Akanov AA, Radisauskas R, Padaiga Z. Assessment of coronary care management and hospital mortality from ST-segment elevation myocardial infarction in the Kazakhstan population: Data from 2012 to 2015. *Medicina (Kaunas)*. 2017;53(1):58-65.
- Beig JR, Trambo NA, Kumar K, et al. Components and determinants of therapeutic delay in patients with acute ST-elevation myocardial infarction: A tertiary care hospital-based study. *J Saudi Heart Assoc*. 2017;29(1):7-14.
- Kaul P, Armstrong PW, Sookram S, Leung BK, Brass N, Welsh RC. Temporal trends in patient and treatment delay among men and women presenting with ST-elevation myocardial infarction. *Am Heart J*. 2011;161(1):91-7.
- Moser DK, Kimble LP, Alberts MJ, et al. Reducing delay in seeking treatment by patients with acute coronary syndrome and stroke: a scientific statement from the American Heart Association Council on cardiovascular nursing and stroke council. *Circulation*. 2006;114(2):168-82.
- Salarifar M, Askari J, Saadat M, et al. Strategies to Reduce the Door-to-Device time in ST-Elevation Myocardial Infarction Patients. *The Journal of Tehran University Heart Center*. 2019;14(1):18-27.
- Poorhosseini H, Abbasi SH. The Tehran Heart Center. *Eur Heart J*. 2018;39(29):2695-6.
- Salarifar M, Mousavi MR, Saroukhani S, et al. Percutaneous coronary intervention to treat chronic total occlusion: predictors of technical success and one-year clinical outcome. *Tex Heart Inst J*. 2014;41(1):40-7.
- Noorani F, Runge M, Tripathi S, et al. Pre-Hospital Delays in Care for STEMI Patients in Mumbai: Challenges and Opportunities. *Am Heart Assoc*; 2016.
- Peng YG, Feng JJ, Guo LF, et al. Factors associated with



- prehospital delay in patients with ST-segment elevation acute myocardial infarction in China. *Am J Emerg Med.* 2014;32(4):349-55.
17. Heo JY, Hong KJ, Shin SD, Song KJ, Ro YS. Association of educational level with delay of prehospital care before reperfusion in STEMI. *Am J Emerg Med.* 2015;33(12):1760-9.
 18. Albarqouni L, Smenes K, Meinertz T, et al. Patients' knowledge about symptoms and adequate behaviour during acute myocardial infarction and its impact on delay time: Findings from the multicentre MEDEA Study. *Patient Educ Couns.* 2016;99(11):1845-51.
 19. Yarzebski J, Goldberg RJ, Gore JM, Alpert JS. Temporal trends and factors associated with extent of delay to hospital arrival in patients with acute myocardial infarction: the Worcester Heart Attack Study. *American heart journal.* 1994;128(2):255-63.
 20. Rasmussen CH, Munck A, Kragstrup J, Haghfelt T. Patient delay from onset of chest pain suggesting acute coronary syndrome to hospital admission. *Scand Cardiovasc J.* 2003;37(4):183-6.
 21. McKinley S, Aitken LM, Marshall AP, et al. Delays in presentation with acute coronary syndrome in people with coronary artery disease in Australia and New Zealand. *Emerg Med Australas.* 2011;23(2):153-61.
 22. Lee MR, Yun KH, Kim DH, et al. Factors Related to Pre-hospital Delay in Korean Patients with ST-segment Elevation Myocardial Infarction: A Data from the Province of Jeonbuk Regional Cardiovascular Center. *Journal of Lipid and Atherosclerosis.* 2016;5(1):21-6.
 23. Malhotra S, Gupta M, Chandra KK, Grover A, Pandhi P. Prehospital delay in patients hospitalized with acute myocardial infarction in the emergency unit of a North Indian tertiary care hospital. *Indian Heart J.* 2003;55(4):349-53.
 24. George L, Ramamoorthy L, Satheesh S, Saya RP, Subrahmanyam DK. Prehospital delay and time to reperfusion therapy in ST elevation myocardial infarction. *J Emerg Trauma Shock.* 2017;10(2):64-9.
 25. Momeni M, Salari A, Shafighnia S, Ghanbari A, Mirbolouk F. Factors influencing pre-hospital delay among patients with acute myocardial infarction in Iran. *Chin Med J.* 2012;125(19):3404-9.
 26. Khosravi AR, Hoseinabadi M, Pourmoghaddas M, et al. Primary percutaneous coronary intervention in the Isfahan province, Iran; A situation analysis and needs assessment. *ARYA Atheroscler.* 2013;9(1):38-44.
 27. Taghaddosi M, Dianati M, Fath Gharib Bidgoli J, Bahonaran J. Delay and its related factors in seeking treatment in patients with acute myocardial infarction. *ARYA Atheroscler.* 2010;6(1):35-41.
 28. Kassaian SE, Masoudkabar F, Sezavar H, et al. Clinical characteristics, management and 1-year outcomes of patients with acute coronary syndrome in Iran: the Iranian Project for Assessment of Coronary Events 2 (IPACE2). *BMJ open.* 2015;5(12):e007786.
 29. Farshidi H, Rahimi S, Abdi A, Salehi S, Madani A. Factors Associated With Pre-hospital Delay in Patients With Acute Myocardial Infarction. *Iran Red Crescent Med J.* 2013;15(4):312-6.

