

A dimeric PR-1-type pathogenesis-related protein interacts with ToxA and potentially mediates ToxA-induced necrosis in sensitive wheat

SHUNWEN LU^{1,*}, JUSTIN D. FARIS¹, ROBERT SHERWOOD², TIMOTHY L. FRIESEN¹ AND MICHAEL C. EDWARDS¹

¹Cereal Crops Research Unit, USDA-ARS, Fargo, ND 58102, USA

²Cornell University Biotechnology Resource Center, Ithaca, NY 14853, USA

SUMMARY

A dimeric PR-1-type pathogenesis-related protein (PR-1-5), recently identified in wheat, was found to interact with *Stagonospora nodorum* ToxA in both yeast two-hybrid and co-immunoprecipitation assays. Site-specific mutational analyses revealed that the RGD motif of ToxA is not targeted by PR-1-5, whereas two surface-exposed asparagine residues are essential for the interaction: the N102 residue of the turning loop between β 2 and β 3 in ToxA and the N141 residue of the turning loop between β C and β D in PR-1-5. Recombinant PR-1-5 and ToxA mutant proteins carrying alanine substitutions at the interacting sites were expressed in *Pichia pastoris*, together with the wild-type proteins. Native polyacrylamide gel electrophoresis (PAGE) confirmed that the PR-1-5-N141A mutant retains the ability to form dimers. Plant assays indicated that the ToxA-N102A mutant fails to induce necrosis, whereas the PR-1-5-N141A mutant is impaired in the 'necrosis-promoting' activity shown by the wild-type PR-1-5 when co-infiltrated with ToxA in sensitive wheat. Reverse transcriptase-polymerase chain reaction and Western blot analyses revealed that the native PR-1-5 protein is differentially expressed between ToxA-sensitive and ToxA-insensitive wheat lines in response to ToxA treatment. These results suggest that PR-1-5 is a potential target of ToxA and the site-specific interaction between PR-1-5 and ToxA may mediate ToxA-induced necrosis in sensitive wheat.

Keywords: host-selective toxin, inducible defence-related protein, protein dimerization, protein–protein interaction.

INTRODUCTION

PR-1-type pathogenesis-related (PR) proteins are best known among the plant PR families for being hallmarks of the activation

*Correspondence: Email: shunwen.lu@ars.usda.gov

Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture (USDA). USDA is an equal opportunity provider and employer.

of hypersensitive response/defence pathways and the development of salicylic acid-dependent systemic acquired resistance (Buchel and Linthorst, 1999; van Loon *et al.*, 2006; van Loon and van Strien, 1999). The accumulation of PR-1 proteins in a wide range of plant species has been associated with both incompatible and compatible pathogen–host interactions culminating in cell death (Gruner *et al.*, 2003; Hong *et al.*, 2005; Lazniewska and Macioszek, 2010; Leon-Reyes *et al.*, 2009; Lu *et al.*, 2011; Santamaria *et al.*, 2001). However, the biochemical function and biological activity of PR-1 proteins are still obscure, although antifungal activities have been described for a few proteins purified from dicotyledonous plants (Kiba *et al.*, 2007; Niderman *et al.*, 1995; Rauscher *et al.*, 1999), and enhanced disease resistance has been observed in transgenic tobacco plants overexpressing certain PR-1 genes (Alexander *et al.*, 1993; Li *et al.*, 2011). Direct roles in the mediation of plant cell death associated with host–pathogen interactions have not been demonstrated for any members of the plant PR-1 family.

Homologues of PR-1 proteins are widely spread among other kingdoms, including Animalia, Fungi and Bacteria (Cantacessi *et al.*, 2012; Gibbs *et al.*, 2008; Henriksen *et al.*, 2001; Schuren *et al.*, 1993). Among these PR-1-like proteins, the best known is the human glioma pathogenesis-related (GliPR) protein implicated in cancer development in brain and prostate cells (Bonafe *et al.*, 2010; Szyperski *et al.*, 1998). Early studies on the tomato PR-1 protein P14a (Fernandez *et al.*, 1997) and GliPR (Szyperski *et al.*, 1998) revealed a conserved 'PR-1 fold' (also called SCP-like extracellular protein domain, pfam00188) that features four α -helices and one four-strand β -sheet, with histidine (His)-72, glutamic acid (Glu)-77, Glu-98 and His-117 (in the P14a nomenclature) identified to be potential active sites. Homology-based modelling for the cone snail (*Conus textile*) Tex31 (Milne *et al.*, 2003) and human GAPR-1 (Serrano *et al.*, 2004) proteins suggested a 'catalytic triad' consisting of serine (Ser)-73 with Glu-98/His-117 (or Glu-77/His-72) within the largest cavity in the PR-1 fold, which is reminiscent of that of serine proteases, but none of these PR-1-like proteins has been confirmed to possess protease activity. Recent studies in human/animal systems have suggested 'novel' functions for PR-1-like proteins based on several newly identified properties, including coordination with divalent metal ions, such

as Zn^{2+} (Asojo *et al.*, 2011; Wang *et al.*, 2010), and binding to negatively charged membrane-associated lipids (van Galen *et al.*, 2012) or certain fatty acid-derived lipid signalling molecules, such as leukotrienes (Xu *et al.*, 2012). The importance of these unusual features in specific physiological processes has not been well established.

ToxA is the first discovered fungal proteinaceous host-selective toxin originally identified from the tan spot fungus *Pyrenophora tritici-repentis* (Ptr) (Ballance *et al.*, 1989, 1996; Ciuffetti *et al.*, 1997; Tomas *et al.*, 1990) and, more recently, from the leaf/glume blotch fungus *Stagonospora nodorum* (Sn) (Friesen *et al.*, 2006). SnToxA is nearly identical (>99% similarity) to PtrToxA and is thought to have been acquired by Ptr from Sn through a recent horizontal gene transfer event (Friesen *et al.*, 2006). How ToxA induces cell death in sensitive wheat is still under investigation. A current model suggests that ToxA binds to a membrane-located receptor through its RGD motif and is internalized through a yet-to-be-identified mechanism; within the plant cells, ToxA targets ToxABP1 (Manning *et al.*, 2007), a chloroplastic protein homologous to the *Arabidopsis* plastid membrane protein THYLAKOID FORMATION 1 (THF1) (Wang *et al.*, 2004), essential for photosynthesis, thus leading to cell death (reviewed by Ciuffetti *et al.*, 2010). It has also been reported that ToxA interacts with a second chloroplast-associated protein, plastocyanin (Tai *et al.*, 2007). Sensitivity to ToxA and susceptibility to the ToxA-producing fungus are controlled by a single dominant gene *Tsn1* (Faris *et al.*, 1996), which encodes a protein belonging to the nucleotide-binding site-leucine-rich repeat (NBS-LRR) family of plant disease resistance proteins, but featuring an unusual N-terminal fusion with a serine/threonine protein kinase (S/TPK) domain (Faris *et al.*, 2010). *Tsn1* is unlikely to be the long-assumed ToxA receptor as it lacks a transmembrane domain and does not interact directly (at least in yeast) with ToxA (Faris *et al.*, 2010). ToxA may act as both an elicitor and a virulence factor because a large number of defence-related genes, including those for PR proteins, are up-regulated in ToxA-treated sensitive wheat, like those observed in classical gene-for-gene interactions (Adhikari *et al.*, 2009; Pandelova *et al.*, 2009, 2012). It has been proposed that ToxA, and perhaps other fungal effectors, may exploit plant defence mechanisms to induce cell death for survival of the necrotrophic pathogen (Ciuffetti *et al.*, 2010; Friesen *et al.*, 2008; Mengiste, 2012; Oliver *et al.*, 2012).

Previously, we have obtained two partial cDNA clones that encode PR-1-like proteins potentially interacting with SnToxA from a yeast two-hybrid (Y2H) library screening experiment (Lu *et al.*, 2009). Subsequent genome-wide analysis of the wheat *PR-1* family identified the corresponding full-length genes, named *TaPr-1-1* and *TaPr-1-5* (Lu *et al.*, 2011). Recent characterization of the recombinant PR-1-1 and PR-1-5 proteins revealed that the former exists as monomers, whereas the latter forms dimers contributing to protease resistance (Lu *et al.*, 2013). Here, we report that the

dimeric PR-1-5 physically interacts with SnToxA in a sequence-specific manner, and that the interacting sites on both proteins are important for their respective biological activities. The data suggest that PR-1-5 is a potential target of ToxA and that the site-specific interaction between PR-1-5 and ToxA may mediate ToxA-induced necrosis governed by the cognate *Tsn1* gene in sensitive wheat.

RESULTS

PR-1-5 physically interacts with ToxA

Our initial Y2H library screening (Lu *et al.*, 2009) identified two partial cDNA clones encoding PR-1-like proteins (named PR-1-1 and PR-1-5) potentially interacting with SnToxA (hereafter ToxA unless otherwise noted). To confirm the PR-1–ToxA interaction, we performed additional Y2H assays using the full-length cDNA sequences encoding the mature protein of ToxA and PR-1-1 or PR-1-5. The ToxA protein was N-terminally fused to the GAL4-DNA-binding domain (BD) and the two PR-1 proteins were fused to the GAL4-activation domain (AD). ‘Positive’ interactions were determined by blue coloration derived from α -galactosidase activity on selective agar plates, and the strength of the interaction was simultaneously ‘quantified’ using serially diluted cell suspensions of the transformant co-expressing the bait/prey proteins. In all tests, the yeast transformant co-expressing the full-length mature ToxA and PR-1-5 proteins produced blue coloration, with the intensity observed in undiluted cells comparable with that of 10^{-2} -diluted cells in a standard positive control (Fig. 1A, compare rows 1 and 5). In contrast, the transformant co-expressing the full-length mature ToxA and PR-1-1 proteins did not produce any blue coloration (row 2). Replacement of either the PR-1-5 prey or the ToxA bait with the yeast proteins RecT or p53 caused a loss of blue coloration (rows 3 and 4). A bait–prey swap assay was pursued, but was found not to be feasible because the GAL4-AD–ToxA fusion appeared to be ‘toxic’ when expressed in yeast (data not shown). Nevertheless, the Y2H assay results were sufficient to show that the full-length mature PR-1-5 protein, but not PR-1-1 protein, interacted with ToxA, although the interaction appeared to be weak (about two orders of magnitude lower than the standard control).

The interaction was further validated by co-immunoprecipitation assays in which PR-1-5 and ToxA were N-terminally labelled with c-Myc and haemagglutinin (HA) epitope tags, respectively, and translated *in vitro*, and the protein complex was detected using anti-c-Myc and anti-HA antibodies. As shown in Fig. 1B, the 17.5-kDa c-Myc-tagged PR-1-5 protein (15.1-kDa PR-1-5 plus 2.4-kDa c-Myc) was detected using anti-c-Myc antibody, whereas the 15.6-kDa HA-tagged ToxA protein (13.2-kDa ToxA plus 2.4-kDa HA) was detected in the same protein sample using anti-HA antibody (lane 1). In the negative control,




Fig. 1 Characterization of the interaction between PR-1-5 and ToxA. (A) Yeast two-hybrid assay. Serially diluted cell suspensions from a single colony of the transformant co-expressing the bait/prey proteins were inoculated onto selective agar plates and incubated at 30 °C for 36 h. p53/RecT and p53/Lam are standard 'positive' and 'negative' controls, respectively. Blue coloration on the X- α -gal plates reflects the strength of protein–protein interactions. (B) Co-immunoprecipitation assay. Epitope-tagged bait and prey proteins were separated by sodium dodecylsulphate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto poly(vinylidene difluoride) (PVDF) membrane; the blot was probed first with anti-c-Myc antibody (top panel) and then with anti-haemagglutinin (HA) antibody (bottom panel) after stripping. Lane 3 was loaded with a 'blank' control. Arrows indicate the respective bait and prey proteins (molecular masses in kDa). The faint band found in the bottom panel (lane 2) was caused by non-specific binding of the HA tag to the protein G beads. Numbers on the left indicate protein size markers.

the 38.1-kDa c-Myc-tagged p53 protein (35.7-kDa p53 plus 2.4-kDa c-Myc) was detected with a signal intensity equal to the 17.5-kDa c-Myc-tagged PR-1-5 protein, and a faint signal was seen in the vicinity of the 15.6-kDa HA-tagged ToxA protein (lane 2). The latter was a result of the non-specific binding of the HA tag to protein G beads, as confirmed in a separate experiment (data not shown). Nothing was detected in the background control (lane 3), which only contained the lysate used for *in vitro* translation. These experiments demonstrate that PR-1-5 physically interacts

with ToxA both *in vivo* (at least in yeast) and *in vitro*. Attempts to pull down the native PR-1-5 protein in leaf tissue extracts isolated from ToxA-treated sensitive wheat using the 15.6-kDa HA-tagged ToxA protein were unsuccessful. This may have been a result of the fact that the concentration of the native PR-1-5 protein in the wheat leaf extracts was relatively lower than that of the *in vitro*-translated c-Myc-tagged PR-1-5 protein. However, it was more likely a result of competitive binding from other ToxA-interacting proteins, e.g. ToxABP1 (Manning *et al.*, 2007) and plastocyanin (Tai *et al.*, 2007), both of which are constitutively expressed at high levels (see the final section of the Results).

Two surface-exposed asparagine residues are essential for the interaction between PR-1-5 and ToxA

A series of deletion constructs for both PR-1-5 and ToxA proteins was tested by Y2H assays to locate regions containing the interacting sites. In PR-1-5, the deletion at the N-terminal region [up to 62 amino acids (aa)] did not affect the interaction (Fig. 2A, row 2, right), whereas the two deletions (13 and 36 aa, respectively) at the C-terminal region greatly reduced or abolished the interaction (rows 3 and 4). These results delineate the ToxA-interacting region to positions 128–151, which include the two β strands (β C and β D) and the turning loop in the PR-1 fold. In ToxA, the 39-aa deletion at the C-terminus, which includes three β strands (β 6, β 7 and β 8) and the entire RGD motif, did not affect the interaction (Fig. 2B, row 2), whereas the two deletions (31 and 44 aa) at the N-terminal region apparently reduced or abolished the interaction (rows 3 and 4). These results indicate that the ToxA RGD motif, which is essential for ToxA activity (Manning *et al.*, 2004, 2008; Sarma *et al.*, 2005; Tuori *et al.*, 2000), is not involved in the PR-1-5–ToxA interaction. Instead, the two β strands (β 2 and β 3) and the turning loop (positions 92–105) contain the interacting sites.

In order to identify critical residues in the targeted interacting regions, we performed site-specific mutagenesis, with a focus on the residues within the turning loops, because mutations in these amino acids are less likely to be structurally disruptive. In PR-1-5, the turning loop between β C and β D contains six residues (Fig. 2C, left). We mutated all of these residues individually, except for cysteine (Cys)-140, which is predicted to be part of a disulphide bridge. The N141A mutation affected the interaction most significantly, as indicated by the complete loss of blue coloration in the 10^{-1} -diluted cells and the faint coloration in the undiluted cells (Fig. 2C, left, row 2). The N142A mutation reduced the interaction slightly, as indicated by the weakened coloration in the 10^{-1} -diluted cells (Fig. 2C, left, row 3). The N143A mutation also caused the loss of blue coloration in the 10^{-1} -diluted cells, but only reduced the coloration slightly in the undiluted cells (Fig. 2C, left, row 4). In contrast, the L144R and G145A mutations did not change the blue coloration patterns (Fig. 2C, left, rows 5 and 6). These results suggest that L144 and G145 residues are not




Fig. 2 Identification of residues essential for the PR-1-5-ToxA interaction. (A, B) Deletion analyses. Open boxes indicate the mature proteins with secondary structures drawn at the top following P14a (Fernandez *et al.*, 1997) and Ptr ToxA (Sarma *et al.*, 2005) for PR-1-5 and SnToxA, respectively. Broken and full bars indicate the positions of the five putative active sites in the PR-1 domain and the ToxA RGD motif, respectively. Red bars indicate the positions of the asparagine-rich loops. Residue numbering is based on the full-length PR-1-5 protein (GenBank accession number HQ541965) and SnToxA (DQ423483) proteins with 'Δ' indicating the deleted residues. (C) Site-specific mutagenesis. Open and filled circles indicate residues consisting of the loops and the connecting β -strands, respectively. Asparagine residues are highlighted in red. Asterisks indicate residues conserved in P14a. (D) Ribbon diagrams showing the predicted site-specific interaction between PR-1-5 and ToxA. α helices, β strands and loops are in light blue, green and orange, respectively. The side-chains of the two interacting asparagine residues are shown in a 'stick-ball' scheme with red, grey, blue and white balls for oxygen, carbon, nitrogen and hydrogen atoms, respectively. Broken lines indicate potential hydrogen bonds.

essential for interaction with ToxA. They also indicate that a single-residue mutation within the turning loop is less likely to affect the expression level/stability of the PR-1-5 protein. In ToxA, the turning loop between $\beta 2$ and $\beta 3$ contains two asparagine residues only (Fig. 2C, right). We mutated both asparagine residues individually, and also substituted the adjacent phenylalanine (F103) residue with leucine (L) to generate a site-related control with a conservative substitution (F and L are both hydrophobic). The N101A and F103L mutations did not affect the interaction (Fig. 2C, right, rows 2 and 4). In contrast, the N102A mutation greatly reduced the interaction, as indicated by the complete loss of blue coloration in the 10^{-1} -diluted cells and the weak coloration

in the undiluted cells (Fig. 2C, right, row 3). Thus, the N141 residue in PR-1-5 and the N102 residue in ToxA were identified as the primary interacting sites. Interestingly, mutations at the five putative active sites in the PR-1-5 protein did not affect the interaction significantly (data not shown). The N141A mutation did not affect dimerization (see next section) or the protease resistance characteristics of the wild-type PR-1-5 protein (Lu *et al.*, 2013).

To determine the surface accessibility of the two asparagine residues, we examined the structural data published for P14a [Protein Data Bank (PDB) ID: 1CFE; Fernandez *et al.*, 1997] and PtrToxA (PDB ID: 1ZLE; Sarma *et al.*, 2005) using the Jmol (Hanson, 2010) and Ribbon (Carson, 1997) programs. The 'PR-1 fold' is

highly conserved among PR-1-like proteins, as demonstrated for the human GliPR protein, which has only 35% identity to P14a, but nevertheless adopts the same three-dimensional structure (Szyperski *et al.*, 1998). The sequences that comprise the β C, β D strands and the turning loop between P14a (*RLGCGRARCNNGWWFISCNYD*) and PR-1-5 (*SIGCARVVCNNNLGVFITCNYE*) (italics indicate similar residues) have 67% similarity and two conserved asparagine residues at the targeted sites (N137/N138 in P14a aligned to N141/N142 in PR-1-5). PtrToxA and SnToxA are nearly identical (Friesen *et al.*, 2006; Sarma *et al.*, 2005) and have 100% similarity within the β 2, β 3 strands and the turning loop. Both Jmol and Ribbon programs placed the side chains of the N102 residue in ToxA and the N137 residue in P14a at a position protruding from the surface of the folded protein when the corresponding PDB data were examined. We also confirmed that the asparagine residues at the PR-1-5–ToxA interacting sites were not likely to be glycosylated, as predicted by several web servers specializing in the prediction of N-linked glycosylation sites in

eukaryotic proteins (e.g. GlycoEP; Chauhan *et al.*, 2013). Thus, we predicted that the interaction between PR-1-5 and ToxA is probably established through hydrogen bonding between the side chains of the N102 and N141 residues (Fig. 2D).

The recombinant ToxA-N102A mutant protein fails to induce necrosis in ToxA-sensitive wheat

For the functional characterization of the PR-1-interacting site in ToxA, we expressed the wild-type ToxA and the N102A mutant proteins in *Pichia pastoris*. Very low yields were obtained in initial trials with the constructs for the 118-aa mature ToxA proteins [not detectable by sodium dodecylsulphate-polyacrylamide gel electrophoresis (SDS-PAGE)]. We then attempted to express the 17-kDa proToxA with the N-domain (Tuori *et al.*, 2000). To our surprise, a single major species of 13 kDa was detected for both proteins (Fig. 3A, lanes 1 and 2). Matrix-assisted laser desorption ionization-time of flight/time of flight (MALDI-TOF/TOF) analysis




Fig. 3 Characterization of the recombinant ToxA proteins. (A) Sodium dodecylsulphate-polyacrylamide gel electrophoresis (SDS-PAGE) gel showing ToxA and the ToxA-N102A mutant proteins expressed in *Pichia pastoris*. Each lane was loaded with 20 μ L of cell-free culture supernatant. The control sample was from a 'mock' strain transformed with the expression vector only. Numbers on the left indicate the molecular masses (in kDa) of protein markers. (B) Matrix-assisted laser desorption ionization-time of flight/time of flight mass spectrometry (MALDI-TOF/TOF MS) data showing the 11-amino-acid fragments containing the wild-type sequence (top, peak = 1274.71) and the alanine substitution (highlighted in red) in the mutant protein (bottom, peak = 1231.75). (C) Plant assays showing the activity of the recombinant ToxA proteins. ToxA-sensitive wheat (Grandin) plants were infiltrated with ToxA at the estimated final concentrations (ng/ μ L) indicated on the right. Control plants were treated with the protein sample of the 'mock' strain shown in (A) (lane 3). Marks delimit the infiltrated area. Photographs were taken at 72 h post-infiltration.

confirmed the sequence identity of the 13-kDa bands in which the N-domain was apparently removed, suggesting that ToxA was 'correctly' processed in *Pichia*, probably through a mechanism common in ascomycete fungi. No significant background proteins were detected in the mock strain transformed with the expression vector only (lane 3). The amount of recombinant ToxA proteins was estimated at ~5 ng/μL. Examination of the tandem mass spectrometry (MS/MS) data confirmed the presence of the 11-aa protein fragments containing the interacting site (LNMFITIGLNR, positions 100–110 in the full-length ToxA protein) in the wild-type ToxA and the alanine substitution (LNAFITIGLNR) in the mutant protein (Fig. 3B, MS data are shown for convenience; detailed MS/MS data are available on request).

The ToxA-sensitive wheats Grandin (hexaploid) and Langdon (tetraploid) and ToxA-insensitive wheat BR34 (hexaploid) were infiltrated with varying amounts of recombinant ToxA proteins (ranging from 0.1 to 1 ng/μL) to determine their activity and specificity. Typical necrosis symptoms were observed in Grandin plants treated with 0.25–1 ng/μL of ToxA at 72 h post-infiltration (Fig. 3C, top); this activity was comparable with that of purified PtrToxA, which induces the same necrosis symptoms when applied at 0.1–0.5 μM (roughly 1–5 ng/μL) (Manning *et al.*, 2008). Necrosis symptoms were also observed in ToxA-treated plants of Langdon, which appeared to be less sensitive than Grandin to ToxA, as indicated by the weaker necrosis at lower concentrations (e.g. 0.25 ng/μL) (Fig. S1, see Supporting Information). The ToxA-insensitive BR34 plants did not show any necrosis in the same treatments (data not shown). These observations indicate that the recombinant ToxA protein is active with the same specificity as the native toxin. Under the same conditions, the ToxA-N102A mutant failed to induce necrosis in sensitive wheat; some weak chlorosis-like symptoms were observed only in plants treated with the highest concentration (1 ng/μL) (Fig. 3C, middle). These results confirm that the surface-exposed, PR-1-5-interacting N102 residue is essential for ToxA activity. The control plants treated with the cell-free culture supernatant from a yeast strain transformed with the 'empty' expression vector did not show any symptoms (Fig. 3C, bottom), indicating that the ToxA activity was solely responsible for the observed necrosis symptoms in sensitive wheat.

The recombinant PR-1-5-N141A mutant protein shows impaired necrosis-promoting activity compared with the wild-type

For the functional characterization of the ToxA-interacting site in PR-1-5, we expressed the PR-1-5-N141A mutant protein in *P. pastoris* and obtained its purified form, together with the previously expressed wild-type PR-1-5 and the dimerization-defective E102A mutant proteins (Lu *et al.*, 2013). The three purified PR-1-5 proteins were all found to have the expected molecular mass of

15 kDa as determined by SDS-PAGE analysis (Fig. 4A). Low-temperature (LT) SDS-PAGE analysis indicated that the PR-1-5-N141A mutant protein retained the ability to form dimers, in contrast with the E102A mutant, which was found only as a monomer (Fig. 4B). Native PAGE analysis confirmed that the PR-1-5 dimers show the expected molecular masses of ~30 kDa as compared with the native proteinase K (EC 3.4.21.14; molecular mass, 28.9 kDa; isoelectric point, 8.9), which migrated to a position very close to the PR-1-5 dimers, but distant to the E102A monomers (Fig. 4C). The monomeric bands detected in LT SDS-PAGE for the two dimeric PR-1-5 proteins were absent in native PAGE (compare arrows with 'm' in B and C), suggesting that the PR-1-5 protein may exist exclusively as a dimer in its native configuration. Proteolytic digestion tests indicated that the PR-1-5-N141A mutant protein remained resistant to both serine and cysteine proteases, as did the wild-type (data not shown). To ensure sequence identity, all three purified PR-1-5 proteins were re-validated by size exclusion chromatography and/or MALDI-TOF/TOF analysis, and the MS/MS data confirmed the presence of the 18-aa protein fragments containing the mutated interacting site (VVCANNLGVFITCNYEPR) in the N141A mutant protein (Fig. S2, see Supporting Information).

The biological activity of the PR-1-5 proteins was tested by co-infiltration assays in different wheat lines. In attempts to mimic a natural situation, we adjusted the ToxA concentration to 0.05 ng/μL at which ToxA alone does not induce apparent necrosis at 48 h (or even longer) post-infiltration; high concentrations (>0.1 ng/μL) of ToxA were not suitable for these assays because they cause rapid cell death, making it difficult to detect the activity of the recombinant PR-1-5 proteins. The final concentrations of the PR-1-5 proteins were adjusted to 0.25–0.5 ng/μL, which is close to the concentration of the native PR-1-5 protein in ToxA-treated sensitive wheat (based on a comparison with serially diluted recombinant PR-1-5 protein in Western blots, data not shown). The best results were obtained when the co-infiltration assays were performed using tetraploid Langdon wheat (which is less sensitive to ToxA, as mentioned above) with an extended incubation time (i.e. 5 days) as used in previous studies (e.g. Manning *et al.*, 2008). At the defined concentrations, the Langdon plants treated with ToxA alone did not show apparent necrosis symptoms at 5 days post-infiltration (Fig. 4D, leaf 1). In contrast, those co-infiltrated with ToxA and the wild-type PR-1-5 protein all developed necrotic symptoms with tissue collapse seen across the infiltration area (Fig. 4D, leaves 2–4). Under the same conditions, plants co-infiltrated with the PR-1-5-N141A mutant protein all looked similar to those treated with ToxA alone, although some minor necrosis was seen near the infiltration site (Fig. 4D, leaves 5–7), indicating that the mutant protein is impaired in its ability to promote the development of necrosis. Infiltration with PR-1-5 or the PR-1-5-N141A mutant protein alone did not induce necrosis (Fig. 4D, leaf 8), and the same was true for co-infiltration with the




Fig. 4 Characterization of the recombinant PR-1-5 proteins. (A) Sodium dodecylsulphate-polyacrylamide gel electrophoresis (SDS-PAGE) gel showing the purified PR-1-5 proteins (arrow). E102A is a dimerization-defective mutant (Lu *et al.*, 2013) included as a control. (B) Low-temperature SDS-PAGE gel showing the dimers (d) and monomers (m) of the PR-1-5 proteins under non-reducing conditions. (C) Native PAGE gel showing the ~30-kDa PR-1-5 dimers (d) in comparison with the 15-kDa E102A monomers (m) (loaded twice in lanes 3 and 4). Lane 5 was loaded with proteinase K (pro-K, MW = 28.9 kDa; isoelectric point, 8.9). Numbers on the left in (A) and (B) indicate the molecular masses (in kDa) of the protein markers. (D) Plant assays showing the necrosis-promoting activity of the PR-1-5 protein. ToxA-sensitive wheat (Langdon) plants were infiltrated with the recombinant ToxA and/or PR-1-5 proteins as indicated on the right. Photographs were taken at 5 days post-infiltration (one or three representative leaves of each treatment are shown).

ToxA-N102A mutant and the PR-1-5 protein (Fig. 4D, leaf 9). When tested with hexaploid wheat lines, PR-1-5-associated necrosis-promoting activity was observed in ToxA-sensitive Grandin, but not in ToxA-insensitive BR34, wheat plants (Fig. S3, see Supporting Information). No necrosis was observed when several ToxA-insensitive *tsn1* mutant wheat lines (Faris *et al.*, 2010) were tested under the same co-infiltration conditions (S. Lu and J. D. Faris, unpublished data). These results demonstrate that the expression of the PR-1-5-related necrosis-promoting activity depends on the co-existence of ToxA and its cognate sensitivity gene *Tsn1* (Faris *et al.*, 2010), and such 'death-promoting' activity requires a physical interaction between PR-1-5 and ToxA.

In addition, preliminary experiments suggest that the monomeric E102A mutant protein is less active in promoting necrosis, but the results are not conclusive because the tested protein sample contains a second minor species detectable under non-reducing conditions (Fig. 4B, lane 3). We will try to further purify this mutant protein, together with the monomeric PR-1-1 protein (Lu *et al.*, 2013), to address whether the dimerization of PR-1-5 is essential for the observed 'death-promoting' activity.

The native PR-1-5 protein is differentially expressed between ToxA-sensitive and ToxA-insensitive wheat lines

The *PR-1-5* gene is present in the genomes of both ToxA-sensitive and ToxA-insensitive wheat lines, but has been shown to be differentially expressed in response to infection by a ToxA-producing isolate of Sn (Lu *et al.*, 2009, 2011). To determine whether the native PR-1-5 protein is expressed in response to ToxA in the absence of the pathogen, we analysed mRNA and protein samples extracted from ToxA-treated plants of different wheat lines. Reverse transcriptase-polymerase chain reaction (RT-PCR) revealed that *PR-1-5* transcripts were absent in mock-infiltrated ToxA-insensitive wheat BR34 (Fig. 5, lanes 1–4) and detectable only at low levels (24 and 48 h post-infiltration) in mock-infiltrated ToxA-sensitive wheat Grandin (Fig. 5, lanes 11 and 12). When BR34 plants were treated with ToxA, only low levels of *PR-1-5* transcripts became detectable at 48 h (Fig. 5, lane 8). In contrast, the transcripts were up-regulated sharply in Grandin, as indicated by the strong band seen as early as 12 h, when necrosis




Fig. 5 Differential expression of the native PR-1-5 protein between ToxA-insensitive (BR34) and ToxA-sensitive (Grandin) wheat lines in response to ToxA treatment. Plant assays (top panel): second leaves of 2-week-old plants were 'mock' infiltrated (–) or infiltrated with ToxA protein (+) at a final concentration of 1.0 ng/μL. Photographs were taken at four time points post-infiltration as indicated at the top. A representative leaf is shown for each treatment. Reverse transcriptase-polymerase chain reaction (RT-PCR) (second and third panels): PCR products were separated on ethidium bromide-stained 1% agarose gels. Numbers on the right indicate size (in base pairs) of the amplicons. Transcripts up-regulated relative to those of controls are indicated by red asterisks. *ACT-1*, wheat actin gene (internal control). The oligonucleotide sequences of the primers used for RT-PCR are given in Table S1. Western blot analysis (bottom two panels): protein samples isolated from the plants shown at the top were separated by sodium dodecylsulphate-polyacrylamide gel electrophoresis (SDS-PAGE), transferred onto poly(vinylidene difluoride) (PVDF) membrane and probed with the anti-PR-1-5 antibody (α -PR-1-5) (Lu *et al.*, 2013). Signals were detected by exposure to X-ray films for 30 s. Membranes were stained with Coomassie blue after signal detection (bottom). Numbers on the right indicate the positions of size markers (molecular masses in kDa).

had just become visible (Fig. 5, lane 14), and by the greatly intensified band seen at 48 h, at which time the entire infiltrated area had collapsed (Fig. 5, lane 16). In Western blot analysis, the anti-PR-1-5 antibody (Lu *et al.*, 2013) hybridized to a 15-kDa band in ToxA-treated Grandin plants at 12, 24 and 48 h, with the signal intensity correlating with the abundance of the *PR-1-5* transcripts shown in the RT-PCR results (Fig. 5, lanes 14–16, compare white and red asterisks). These data confirm that the native PR-1-5 protein is expressed and accumulates coincidentally with necrosis development in ToxA-sensitive wheat. In contrast, under the same conditions, the transcripts encoding three other proteins were found to be either up-regulated in ToxA-treated plants (as for the PR-1-1 protein) or constitutively expressed at high levels in all plants [as for the two ToxA-interacting chloroplast proteins ToxABP1 (Manning *et al.*, 2007) and plastocyanin (Tai *et al.*, 2007)] of both sensitive and insensitive wheat lines (Fig. S4, see Supporting Information). The up-regulation of *PR-1-1* in both wheat lines indicates that ToxA may induce the expression of multiple *PR-1* genes through *Tsn-1*-independent pathways not necessarily associated with the development of necrosis. Functional tests of the recombinant PR-1-1 protein, as mentioned

above, would help to determine whether the native PR-1-1 protein also contributes to ToxA-induced necrosis in sensitive wheat.

DISCUSSION

Our initial Y2H library screening experiments identified PR-1 proteins as potential targets of ToxA (Lu *et al.*, 2009). In the same experiments, we also recovered several cDNA clones encoding plastocyanin identical to that reported by Tai *et al.* (2007), but were unable to identify ToxABP1 (Manning *et al.*, 2007). We noticed that the ToxA bait used by Manning *et al.* (2007) included residues 23–178, with the first 22 amino acids removed based on the original signal peptide analysis (Ciuffetti *et al.*, 1997). In our experiments, we used residues 17–178 with the first 16 amino acids removed based on the prediction from the SignalP server (Petersen *et al.*, 2011). In addition, we used mRNA for cDNA synthesis instead of total RNA as used by Manning *et al.* (2007). These technical differences may have affected the chance to pull out ToxABP1 from a particular cDNA library, and may also explain why no PR-1-like clones were recovered from previous studies

using the same Y2H library screening systems (Manning *et al.*, 2007; Tai *et al.*, 2007).

In this study, we confirmed the physical interaction between PR-1-5 and ToxA through Y2H and *in vitro* co-immunoprecipitation assays, and determined that the interaction is mediated by two surface-exposed asparagine residues. The polar side-chain of asparagine can act as both a hydrogen bond donor and an acceptor, thus playing an important role in intermolecular interactions (Carpena *et al.*, 2009; Shimoni and Glusker, 1995). The binding of PR-1-5 to ToxA may rely on hydrogen bonds formed between the oxygen (or hydrogen) atom in the side-chain of the N141 residue in PR-1-5 and the hydrogen (or oxygen) atom in the side-chain of the N102 residue in ToxA (Fig. 2D). Amino acid sequence alignments revealed that the N141 residue in the PR-1-5 protein is actually a part of a unique motif-like sequence (NNNL) found only in PR-1-5 and PR-1-4 proteins in the wheat PR-1 family; extensive database searches failed to identify a homologue having this 'NNNL' motif in plant PR-1 proteins (Fig. S5, see Supporting Information). However, the N102 residue in ToxA is located within a region whose counterpart is completely missing in the mammalian FnIII proteins topologically most closely related to ToxA (Sarma *et al.*, 2005). Thus, the specificity of the interaction between PR-1-5 and ToxA may have stemmed from a beneficial 'matching' of two unusual structures, each acquired by an unrelated protein during natural selection. Like the RGD motif, the N101–N102 loop is conserved; no substitutions were found at either asparagine residue in the natural ToxA isoforms investigated in a previous study (Tan *et al.*, 2012). Further mutational analysis is needed to determine the functional importance of the two asparagine-rich motifs in the PR-1-5 and ToxA interaction. It remains to be determined whether PR-1-5 interacts directly with ToxA *in planta*. The HA-tagged ToxA protein failed to pull down the native PR-1-5 protein from the plant extracts used for co-immunoprecipitation assays, probably as a result of potential competitive binding from ToxABP1 (Manning *et al.*, 2007) and plastocyanin (Tai *et al.*, 2007), as mentioned in the previous section; however, the negative results may also suggest that PR-1-5 does not interact directly with ToxA *in planta* or the interaction is too weak to be detected using the pull-down method. Bimolecular fluorescence complementation assays (reviewed by Kodama and Hu, 2012) will be pursued using an *Agrobacterium*-based transient expression system to further evaluate the PR-1-5–ToxA interaction *in planta*.

Previous studies on ToxA have been largely focused on the RGD motif (Manning *et al.*, 2004, 2008; Meinhardt *et al.*, 2002; Sarma *et al.*, 2005; Tuori *et al.*, 1995, 2000; Zhang *et al.*, 1997) because of its similarity to the classical mammalian RGD-containing domains that bind to membrane-associated integrin receptors. More than 20 single-residue mutations have been made to ToxA (Manning *et al.*, 2004, 2008; Meinhardt *et al.*, 2002); none of them targeted the N102 residue or immediately adjacent amino acids. Despite the demonstrated importance of the RGD motif, the

fact that plants lack any bona fide integrins makes it difficult to draw parallels with the mammalian systems. Some integrin-like proteins have been identified in Arabidopsis (Gouget *et al.*, 2006; Knepper *et al.*, 2011), but their homologues in wheat have not been reported. The RGD-independent interaction between PR-1-5 and ToxA identified in this study opens a door to explore alternative receptors directly linked to plant pathways. The successful expression of ToxA in *Pichia* will facilitate functional studies. Unlike the *Escherichia coli*-expressed PtrToxA (Manning *et al.*, 2008) and SnToxA (Tan *et al.*, 2012), which are His-tagged and less active, the *Pichia*-expressed SnToxA obtained in this study is tag free and appears to be as active as the native PtrToxA, although further purification is needed to precisely quantify the activity.

By using LT SDS-PAGE and native PAGE analyses, we confirmed that PR-1-5 represents the first dimeric member in the plant PR-1 family, and may exist exclusively as a dimer, at least under the defined native PAGE conditions (Fig. 4). The fact that the PR-1-5-N141A mutant protein retains the ability to form dimers (Fig. 4) indicates that the ToxA-binding site is not essential for dimerization in PR-1-5. This is in contrast with the E102A mutant, which fails to form dimers because of the mutation at one of the active sites known to be involved in dimerization (Lu *et al.*, 2013; Serrano *et al.*, 2004). In ToxA, the RGD motif is not targeted by PR-1-5 (Fig. 2). Together, these results suggest that the binding of ToxA to PR-1-5 (or vice versa) would not disturb the active sites in either protein; thus, the resulting protein complex would be dually functional in downstream operations. Recent studies have shown that the dimerization of the GPR-1 protein is required for its function involving membrane-binding activity (Eberle *et al.*, 2002; Serrano *et al.*, 2004; van Galen *et al.*, 2012). It will be important to determine whether the PR-1-5 dimer is the functional subunit in the PR-1-5–ToxA protein complex and how dimerization affects the necrosis-promoting activity.

In planta characterization of PR-1-5 function is a challenging task because of the genomic redundancy of the PR-1 family. There are an estimated total of >60 PR-1-like genes in the hexaploid wheat genome, with several having >80% nucleotide identity to PR-1-5 (Lu *et al.*, 2011). Attempts to specifically knock down the PR-1-5 gene by virus-induced gene silencing (Holzberg *et al.*, 2002) have been unsuccessful in our laboratory. The co-infiltration assays presented in this study provide an alternative way to test the biological activity of the PR-1 proteins in their native host. The concentration of ToxA is important for co-infiltration assays. It is known that ToxA produced by Ptr isolates in liquid culture under laboratory conditions can reach a final concentration of 1–2 ng/μL (Tomas *et al.*, 1990). Purified native or recombinant ToxA proteins are routinely used for infiltration assays with a final concentration of 1–25 ng/μL; the minimal concentration required for the induction of necrosis has been reported to be 0.7–1.3 ng/μL (Manning *et al.*, 2004; Meinhardt *et al.*, 2002; Tomas *et al.*, 1990). Realistic

cally, the apoplastic concentrations of ToxA secreted by invading mycelia of the fungus in a susceptible host may never reach such minimal levels. Thus, we believe that the use of subminimal concentrations (below 0.7 ng/ μ L) for co-infiltration assays is a legitimate choice to mimic a natural situation. Similarly, the amount of the PR-1 protein must be taken into account. It is essential to use a concentration close to its native status, because the impact of forcefully delivered excess 'exogenous' proteins on biological processes associated with plant apoplastic spaces, cell walls or cell wall–plasma membrane interfaces cannot be predicted.

How the PR-1-5–ToxA interaction mediates ToxA-induced necrosis in wheat remains to be investigated. One possibility is that the PR-1-5–ToxA interaction may be involved in the activation of cell death pathway(s) in sensitive wheat. The native PR-1-5 protein is also expressed in mock-treated sensitive wheat and is easily detectable at 24 and 48 h post-infiltration (Fig. 5, lanes 11 and 12). We speculate that, in sensitive wheat, the PR-1-5 protein can be induced by certain stress conditions, such as cell wall damage associated with artificial infiltration or fungal penetration in a disease-favourable natural environment. ToxA can target these 'stress-induced' PR-1-5 proteins (although with low abundance) to establish an initial interaction, which is sufficient for the activation of *Tsn1*-controlled cell death pathway(s). Once the pathway(s) is initiated and local cell death starts, there could be a feedback control loop that up-regulates the PR-1-5 protein whilst necrosis advances (Fig. 5, lanes 14–16). We noticed that the sensitive wheat plants treated with ToxA alone in the co-infiltration assays (Figs 4D and 53A, leaf 1) eventually developed necrosis symptoms, similar to plants co-infiltrated with ToxA and PR-1-5, if incubated for a longer period (4–5 days instead of 48 h post-infiltration in the case of Grandin). Thus, the exogenous application of PR-1-5 through co-infiltration might mimic an 'ahead of time' expression that allows necrosis to develop earlier than in the case of infiltration with ToxA alone, in which the expression of necrosis may depend on the progressive up-regulation of the native PR-1-5 protein.

It is known that, in the presence of specific death stimuli, plant programmed cell death pathways can be activated by extracellular enzymes, such as serine protease-related saspases/phytaspases having caspase-like activities essential for apoptosis in human/animal systems (Bonneau *et al.*, 2008; Chichkova *et al.*, 2010; Vartapetian *et al.*, 2011). Although lacking similarity to saspases/phytaspases, PR-1-5 shares certain commonalities with caspases, including resistance to serine/cysteine proteases, formation of dimers and conserved motif-like sequences (Lu *et al.*, 2013). The necrosis-promoting activity observed in this study further supports the notion that PR-1-5 may have caspase-like activities. It is possible that the binding of ToxA to PR-1-5 causes a specific conformational change in the PR-1-5 dimer, leading to the activation of its substrate-specific protease activity, which would turn on (probably through interaction with other death signalling-related

enzymes/proteins) *Tsn1*-controlled defence-related pathways, such as those in classical gene-for-gene interactions, eventually resulting in cell death. The identification of the potential substrates of PR-1-5 activity through protease profiling may provide useful information with regard to this hypothesis. Interestingly, a recent study (Zhang *et al.*, 2012) has reported that the powdery mildew effector CSEP0055 targets three barley PR proteins, including PR17c, PR1a and PR1b, as revealed by Y2H library screening, and the CSEP0055–PR17c interaction may contribute to disease resistance, although the functional significance of the CSEP0055–PR1a/PR1b interaction has not been tested. Nevertheless, these findings suggest further that plant PR-1 proteins can be potential targets of both necrotrophic and biotrophic effectors.

In summary, the PR-1-5–ToxA interaction identified in this study suggests that specific recognition between a host-selective toxin and its target (or cofactor) may be achieved through affinity binding mediated by unique surface-exposed sites on the two interacting partners. The ToxA-binding asparagine-rich loop identified in the dimeric PR-1-5 protein may represent an effector-binding domain in the PR-1 family. Thus, PR-1, and perhaps also other extracellular defence proteins, may be directly targeted by specific proteinaceous toxins produced by necrotrophic pathogens, such as Ptr and Sn. Further characterization of the PR-1-5–ToxA interaction will help us to understand how necrotrophic pathogens exploit plant defence mechanisms to induce diseases in the host plants.

EXPERIMENTAL PROCEDURES

Y2H assays

The targeted coding regions in the mature proteins of SnToxA (bait) and PR-1-1/PR-1-5 (prey) were PCR amplified from laboratory cDNA clones of *SnToxA* (GenBank accession number DQ423483; Friesen *et al.*, 2006) and *PR-1-1* (HQ541961)/*PR-1-5* (HQ541965) (Lu *et al.*, 2011), respectively. Mutations were generated by PCR fusion using the primers listed in Table S1 (see Supporting Information). PCR products were subcloned into the pGBK-T7 (bait) or the pGAD-T7 (prey) vector and transformed into yeast strain AH109, as described previously (Lu, 2012). For α -galactosidase activity assays, yeast cells were picked up from a single colony of the transformant grown on agar plates containing synthetic defined medium lacking leucine and tryptophan (SD/–LT, selecting for the bait and prey proteins) and suspended in 50 μ L of sterile water, followed by three 1 : 10 dilutions in series. Five microlitres of each cell suspension in the dilution sets were inoculated onto agar plates of SD/–LT and SD/–LTHA + X- α -gal (lacking leucine, tryptophan, histidine and adenine, and supplemented with the chromogenic substrate 5-bromo-4-chloro-3-indolyl- α -D-galactopyranoside) at the same time to select for reporter genes. Plates were incubated at 30 °C for 36–48 h until blue coloration became visible in the 10^{–3}-diluted cells of the standard positive control (p53/RecT) before photographing. All α -galactosidase activity assays were duplicated and repeated at least once.

Co-immunoprecipitation assays

The cMyc-PR-1-5 (bait) and HA-SnToxA (prey) epitope-tagged proteins (constructed in the pGBK-T7/pGAD-T7 plasmids with the T7 promoter) were generated by *in vitro* translation using the TNT® Coupled Reticulocyte Lysate System (Promega, Madison, WI, USA), and co-immunoprecipitation assays were performed using the Dynabeads Protein G package (Life Technologies, Grand Island, NY, USA) following the manufacturer's instructions. The protein complex was finally eluted from the magnetic beads in 20 µL of elution buffer with 10 µL of reducing agent provided by the manufacturer. The co-precipitated bait and prey proteins were detected by Western blot analysis using horseradish peroxidase (HRP)-conjugated anti-c-Myc (Sigma-Aldrich, St. Louis, MO, USA) and HRP-conjugated anti-HA (Roche Diagnostics, Indianapolis, IN, USA) antibodies.

Expression of the recombinant SnToxA and PR-1-5 proteins in yeast

The coding region corresponding to the 17-kDa proToxA (Tuori *et al.*, 2000) was PCR amplified from the *SnToxA* cDNA clone. ToxA-N102A and PR-1-5-N141A mutant constructs were generated by PCR fusion. PCR primers are listed in Table S2 (see Supporting Information). Yeast transformation and protein expression/isolation were performed as described previously (Lu *et al.*, 2013). The concentration of the target protein was estimated by SDS-PAGE in comparison with a series of dilutions of bovine serum albumin (BSA) (New England Biolabs, Ipswich, MA, USA) included in the same gel. Large-scale expression (from the same yeast strain) and subsequent purification of the PR-1-5-N141A mutant protein were performed through a commercial service (NeoBioLab, Cambridge, MA, USA), together with the wild-type PR-1-5 and the E102A mutant proteins (Lu *et al.*, 2013). The purified PR-1-5 proteins were finally provided in phosphate-buffered saline with a concentration of 0.2–0.27 mg/mL. The sequence identity of the recombinant SnToxA and PR-1-5 proteins was confirmed by MALDI-TOF/TOF analysis.

Protein gel electrophoresis and MS analysis

SDS-PAGE was performed following standard protocols (Green and Sambrook, 2012). Low-temperature SDS-PAGE was performed as described by Lu *et al.* (2013). Size exclusion chromatography and MS analyses were performed at the Proteomics and Mass Spectrometry Facility, Cornell University Biotechnology Resource Center, Ithaca, NY, USA. Native PAGE was performed using procedures adapted from a Bio-Rad (Hercules, CA, USA) protocol. Purified proteins (1 µg per lane) were separated on SDS-free 8% straight polyacrylamide gels without stacking gel in SDS-free Tris-glycine running buffer at 4 °C, followed by Coomassie blue staining. The native proteinase K (EC 3.4.21.14) (New England Biolabs, Ipswich, MA, USA) was included as a reference protein because it has a molecular mass of 28.9 kDa, close to that of the PR-1-5 dimers (~30 kDa), and is a basic protein [isoelectric point (pI), 8.9], like the PR-1-5 protein (pI, 8.3).

Plant assays

ToxA-sensitive hexaploid (cultivar Grandin) and tetraploid (cultivar Langdon) wheat and ToxA-insensitive hexaploid wheat (cultivar BR34)

were used for infiltration/co-infiltration assays. Plants were grown in 6-in (15.24-cm) pots (10–16 plants per pot) containing SB100 professional growing mix (Sungrow Horticulture, Delleve, WA, USA) in a growth chamber at an average temperature of 21 °C with a 16-h photoperiod for 2 weeks. For ToxA activity assays, the second leaf of each plant was infiltrated with ~20 µL of cell-free culture supernatant (diluted in series with water) containing the recombinant ToxA protein (Fig. 3A) at the desired concentrations using a 1-mL syringe without the needle (Delasco, Council Bluffs, IA, USA). Control plants were infiltrated with the culture supernatant from a yeast strain transformed with the expression vector only (Fig. 3A). Treated plants were kept in a growth chamber and symptoms were examined 2–5 days after infiltration. Each treatment included at least five plants. For co-infiltration assays, ToxA and PR-1-5 proteins were diluted with water and mixed together, with the final concentrations adjusted to 0.05 ng/µL for ToxA and 0.25 ng/µL for PR-1-5. The protein solution was incubated at room temperature for 10–60 min before use. Plants were treated in the same way as mentioned above, except that both the first and second leaves of the plants were infiltrated. Each treatment included 5–10 plants and was repeated at least twice.

RT-PCR and Western blot analysis

mRNA samples were isolated from leaf tissues of 2-week-old ToxA-treated and control plants of cultivars BR34 and Grandin that were collected at different time points (4, 12, 24 and 48 h post-infiltration). RT-PCR was performed as described previously (Lu *et al.*, 2011) using the following primers: 5'-AATACGCCGAGGACTACGTT-3' and 5'-AAGACTGCCATA GAGAAGCTCA-3' for PR-1-5 and 5'-ATGGCTGACGGTGAGGACAT-3' and 5'-GATCAGAAGCACTTCCTGTGGA-3' for ACT-1 (internal control) (Lu *et al.*, 2011). Protein extraction and Western blot analysis using the HRP-conjugated anti-PR-1-5 polyclonal antibody were performed as described by Lu *et al.* (2013).

Sequence analysis

Amino acid sequences of plant PR-1 proteins from wheat and other species were retrieved from GenBank databases available online from the National Center for Biotechnology Information. Alignments were generated using the MegAlign programs from Lasergene 8.1 software (DNASTAR Inc., Madison, WI, USA). Structural data for P14a (Fernandez *et al.*, 1997) and PtrToxA (Sarma *et al.*, 2005) were retrieved from PDB available online from the European Bioinformatics Institute; the corresponding PDB files were examined using the Jmol (Hanson, 2010) and Ribbon (Carson, 1997) programs.

ACKNOWLEDGEMENTS

We thank Kelsey Dunnell and Tyler Lewandowski for technical assistance. This study was supported by USDA-ARS CRIS project 5442-21000-037-00D.

REFERENCES

- Adhikari, T.B., Bai, J., Meinhardt, S.W., Gurung, S., Myrfield, M., Patel, J., Ali, S., Gudmestad, N.C. and Rasmussen, J.B. (2009) Tsn1-mediated host responses to ToxA from *Pyrenophora tritici-repentis*. *Mol Plant Microbe Interact.* **22**, 1056–1068.

- Alexander, D., Goodman, R.M., Gut-Rella, M., Glascock, C., Weymann, K., Friedrich, L., Maddox, D., Ahl-Goy, P., Luntz, T. and Ward, E. (1993) Increased tolerance to two oomycete pathogens in transgenic tobacco expressing pathogenesis-related protein 1a. *Proc. Natl. Acad. Sci. USA*, **90**, 7327–7331.
- Asojo, O.A., Koski, R.A. and Bonafe, N. (2011) Structural studies of human glioma pathogenesis-related protein 1. *Acta Crystallogr. D: Biol. Crystallogr.* **67**, 847–855.
- Ballance, G.M., Lamari, L. and Bernier, C.C. (1989) Purification and characterization of a host-selective necrosis toxin from *Pyrenophora tritici-repentis*. *Physiol. Mol. Plant Pathol.* **35**, 203–213.
- Ballance, G.M., Lamari, L., Kowatsch, R. and Bernier, C.C. (1996) Cloning, expression and occurrence of a gene encoding the Ptr necrosis toxin from *Pyrenophora tritici-repentis*. *Mol. Plant Pathol. On-Line* (<http://www.bspp.org.uk/mppl/1996/1209ballance>).
- Bonafe, N., Zhan, B., Bottazzi, M.E., Perez, O.A., Koski, R.A. and Asojo, O.A. (2010) Expression, purification, crystallization and preliminary X-ray analysis of a truncated soluble domain of human glioma pathogenesis-related protein 1. *Acta Crystallogr. Sect. F: Struct. Biol. Cryst. Commun.* **66**, 1487–1489.
- Bonneau, L., Ge, Y., Drury, G.E. and Gallois, P. (2008) What happened to plant caspases? *J. Exp. Bot.* **59**, 491–499.
- Buchel, A.S. and Linthorst, H.J.M. (1999) PR-1: a group of plant proteins induced upon pathogen infection. In: *Pathogenesis-Related Proteins in Plants* (Datta, S.K. and Muthukrishnan, S., eds), pp. 21–47. Boca Raton, FL: CRC Press.
- Cantacessi, C., Hofmann, A., Young, N.D., Broder, U., Hall, R.S., Loukas, A. et al. (2012) Insights into SCP/TAPS proteins of liver flukes based on large-scale bioinformatic analyses of sequence datasets. *PLoS ONE*, **7**, e31164.
- Carpene, X., Vidossich, P., Schroettner, K., Calisto, B.M., Banerjee, S., Stamper, J. et al. (2009) Essential role of proximal histidine–asparagine interaction in mammalian peroxidases. *J. Biol. Chem.* **284**, 25 929–25 937.
- Carson, M. (1997) Ribbons. *Methods Enzymol.* **277**, 493–505.
- Chauhan, J.S., Rao, A. and Raghava, G.P. (2013) *In silico* platform for prediction of N-, O- and C-glycosites in eukaryotic protein sequences. *PLoS ONE*, **8**, e67008.
- Chichkova, N.V., Shaw, J., Galiullina, R.A., Drury, G.E., Tuzhikov, A.I., Kim, S.H. et al. (2010) Phytaspase, a relocatable cell death promoting plant protease with caspase specificity. *EMBO J.* **29**, 1149–1161.
- Ciuffetti, L.M., Tuori, R.P. and Gaventa, J.M. (1997) A single gene encodes a selective toxin causal to the development of tan spot of wheat. *Plant Cell*, **9**, 135–144.
- Ciuffetti, L.M., Manning, V.A., Pandelova, I., Betts, M.F. and Martinez, J.P. (2010) Host selective toxins, Ptr ToxA and Ptr ToxB, as necrotrophic effectors in the *Pyrenophora tritici-repentis*–wheat interaction. *New Phytol.* **187**, 911–919.
- Eberle, H.B., Serrano, R.L., Fullekrug, J., Schlosser, A., Lehmann, W.D., Lottspeich, F., Kaloyanova, D., Wieland, F.T. and Helms, J.B. (2002) Identification and characterization of a novel human plant pathogenesis related protein that localizes to lipid-enriched microdomains in the Golgi complex. *J. Cell Sci.* **115**, 827–838.
- Faris, J.D., Anderson, J.A., Francl, L.J. and Jordahl, J.G. (1996) Chromosomal location of a gene conditioning insensitivity in wheat to a necrosis-inducing culture filtrate from *Pyrenophora tritici-repentis*. *Phytopathology*, **86**, 459–463.
- Faris, J.D., Zhang, Z., Lu, H., Lu, S., Reddy, L., Cloutier, S., Fellers, J.P., Meinhardt, S.W., Rasmussen, J.B., Xu, S.S., Oliver, R.P., Simons, K.J. and Friesen, T.L. (2010) A unique wheat disease resistance-like gene governs effector-triggered susceptibility to necrotrophic pathogens. *Proc. Natl. Acad. Sci. USA*, **107**, 13 544–13 549.
- Fernandez, C., Szyperski, T., Bruyere, T., Ramage, P., Mosinger, E. and Wuthrich, K. (1997) NMR solution structure of the pathogenesis-related protein P14a. *J. Mol. Biol.* **266**, 576–593.
- Friesen, T.L., Stukenbrock, E.H., Liu, Z., Meinhardt, S., Ling, H., Faris, J.D., Rasmussen, J.B., Solomon, P.S., McDonald, B.A. and Oliver, R.P. (2006) Emergence of a new disease as a result of interspecific virulence gene transfer. *Nat. Genet.* **38**, 953–956.
- Friesen, T.L., Faris, J.D., Solomon, P.S. and Oliver, R.P. (2008) Host-specific toxins: effectors of necrotrophic pathogenicity. *Cell. Microbiol.* **10**, 1421–1428.
- van Galen, J., Orlachs, N.K., Schouten, A., Serrano, R.L., Nolte-t Hoen, E.N., Eerland, R. et al. (2012) Interaction of GAPR-1 with lipid bilayers is regulated by alternative homodimerization. *Biochim. Biophys. Acta*, **1818**, 2175–2183.
- Gibbs, G.M., Roelants, K. and O'Bryan, M.K. (2008) The CAP superfamily: cysteine rich secretory proteins, antigen 5, and pathogenesis-related 1 proteins – roles in reproduction, cancer, and immune defense. *Endocr. Rev.* **29**, 865–897.
- Gougout, A., Senchou, V., Govers, F., Sanson, A., Barre, A., Rouge, P., Pont-Lezica, R. and Canut, H. (2006) Lectin receptor kinases participate in protein–protein interactions to mediate plasma membrane cell wall adhesions in Arabidopsis. *Plant Physiol.* **140**, 81–90.
- Green, M.R. and Sambrook, J. (2012) *Molecular Cloning: A Laboratory Manual*, 4th edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Gruner, R., Strompen, G., Pfitzner, A.J. and Pfitzner, U.M. (2003) Salicylic acid and the hypersensitive response initiate distinct signal transduction pathways in tobacco that converge on the AS-1-like element of the PR-1 promoter. *Eur. J. Biochem.* **270**, 4876–4886.
- Hanson, R.M. (2010) Jmol – a paradigm shift in crystallographic visualization. *J. Appl. Crystallogr.* **43**, 1250–1260.
- Henriksen, A., King, T.P., Mirza, O., Monsalve, R.I., Meno, K., Ipsen, H., Larsen, J.N., Gajhede, M. and Spangfort, M.D. (2001) Major venom allergen of yellow jackets, Ves v 5: structural characterization of a pathogenesis-related protein superfamily. *Proteins*, **45**, 438–448.
- Holzberg, S., Brosio, P., Gross, C. and Pogue, G.P. (2002) Barley stripe mosaic virus induced gene silencing in a monocot plant. *Plant J.* **30**, 315–327.
- Hong, J.K., Lee, S.C. and Hwang, B.K. (2005) Activation of pepper basic PR-1 gene promoter during defense signaling to pathogen, abiotic and environmental stresses. *Gene*, **356**, 169–180.
- Kiba, A., Nishihara, M., Nakatsuka, T. and Yamamura, S. (2007) Pathogenesis related protein 1 homologue is an antifungal protein in *Wasabia japonica* leaves and confers resistance to *Botrytis cinerea* in transgenic tobacco. *Plant Biotechnol.* **24**, 247–253.
- Knepper, C., Savory, E.A. and Day, B. (2011) Arabidopsis NDR1 is an integrin-like protein with a role in fluid loss and plasma membrane-cell wall adhesion. *Plant Physiol.* **156**, 286–300.
- Kodama, Y. and Hu, C.D. (2012) Bimolecular fluorescence complementation (BiFC): a 5-year update and future perspectives. *Biotechniques*, **53**, 285–298.
- Lazniewska, J. and Macioszek, V.K. (2010) Fight to the death: *Arabidopsis thaliana* defense response to fungal necrotrophic pathogens. *Acta Physiol. Plant.* **32**, 1–10.
- Leon-Reyes, A., Spoel, S.H., De Lange, E.S., Abe, H., Kobayashi, M., Tsuda, S. et al. (2009) Ethylene modulates the role of NONEXPRESSOR OF PATHOGENESIS RELATED GENES1 in cross talk between salicylate and jasmonate signaling. *Plant Physiol.* **149**, 1797–1809.
- Li, Z.T., Dhekney, S.A. and Gray, D.J. (2011) PR-1 gene family of grapevine: a uniquely duplicated PR-1 gene from a Vitis interspecific hybrid confers high level resistance to bacterial disease in transgenic tobacco. *Plant Cell Rep.* **30**, 1–11.
- van Loon, L.C. and van Strien, E.A. (1999) The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiol. Mol. Plant Pathol.* **55**, 85–97.
- van Loon, L.C., Rep, M. and Pieterse, C.M.J. (2006) Significance of inducible defense related proteins in infected plants. *Annu. Rev. Phytopathol.* **44**, 135–162.
- Lu, S. (2012) Use of the yeast two-hybrid system to identify targets of fungal effectors. *Methods Mol. Biol.* **835**, 165–189.
- Lu, S., Friesen, T.L. and Faris, J.D. (2009) Identification of ToxA-interacting proteins suggests a possible role for pathogenesis-related protein 1 (PR-1) in mediating *Stagonospora nodorum*–wheat interactions. In: *XIV International Congress on Molecular Plant–Microbe Interactions, Quebec City, Canada, IS-MPMI 2009 XIV Congress 2009*.
- Lu, S., Friesen, T.L. and Faris, J.D. (2011) Molecular characterization and genomic mapping of the pathogenesis-related protein 1 (PR-1) gene family in hexaploid wheat (*Triticum aestivum* L.). *Mol. Genet. Genomics*, **285**, 485–503.
- Lu, S., Faris, J.D., Sherwood, R. and Edwards, M.C. (2013) Dimerization and protease resistance: new insight into the function of PR-1. *J. Plant Physiol.* **170**, 105–110.
- Manning, V.A., Andrie, R.M., Trippe, A.F. and Ciuffetti, L.M. (2004) Ptr ToxA requires multiple motifs for complete activity. *Mol Plant Microbe Interact.* **17**, 491–501.
- Manning, V.A., Hardison, L.K. and Ciuffetti, L.M. (2007) Ptr ToxA interacts with a chloroplast-localized protein. *Mol. Plant–Microbe Interact.* **20**, 168–177.
- Manning, V.A., Hamilton, S.M., Karplus, P.A. and Ciuffetti, L.M. (2008) The Arg-Gly-Asp-containing, solvent-exposed loop of Ptr ToxA is required for internalization. *Mol. Plant–Microbe Interact.* **21**, 315–325.
- Meinhardt, S.W., Cheng, W., Kwon, C.Y., Donohue, C.M. and Rasmussen, J.B. (2002) Role of the arginyl-glycyl-aspartic motif in the action of Ptr ToxA produced by *Pyrenophora tritici-repentis*. *Plant Physiol.* **130**, 1545–1551.
- Mengiste, T. (2012) Plant immunity to necrotrophs. *Annu. Rev. Phytopathol.* **50**, 267–294.
- Milne, T.J., Abbenante, G., Tyndall, J.D., Halliday, J. and Lewis, R.J. (2003) Isolation and characterization of a cone snail protease with homology to CRISP proteins of the pathogenesis-related protein superfamily. *J. Biol. Chem.* **278**, 31 105–31 110.
- Niderman, T., Genetet, I., Bruyere, T., Gees, R., Stintzi, A., Legrand, M., Fritig, B. and Mosinger, E. (1995) Pathogenesis-related PR-1 proteins are antifungal. Isola-

- tion and characterization of three 14-kilodalton proteins of tomato and of a basic PR-1 of tobacco with inhibitory activity against *Phytophthora infestans*. *Plant Physiol.* **108**, 17–27.
- Oliver, R.P., Friesen, T.L., Faris, J.D. and Solomon, P.S. (2012) *Stagonospora nodorum*: from pathology to genomics and host resistance. *Annu. Rev. Phytopathol.* **50**, 23–43.
- Pandelova, I., Betts, M.F., Manning, V.A., Wilhelm, L.J., Mockler, T.C. and Ciuffetti, L.M. (2009) Analysis of transcriptome changes induced by Ptr ToxA in wheat provides insights into the mechanisms of plant susceptibility. *Mol. Plant*, **2**, 1067–1083.
- Pandelova, I., Figueroa, M., Wilhelm, L.J., Manning, V.A., Mankaney, A.N., Mockler, T.C. and Ciuffetti, L.M. (2012) Host-selective toxins of *Pyrenophora tritici-repentis* induce common responses associated with host susceptibility. *PLoS ONE*, **7**, e40240.
- Petersen, T.N., Brunak, S., von Heijne, G. and Nielsen, H. (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods*, **8**, 785–786.
- Rauscher, M., Adam, A.L., Wirtz, S., Guggenheim, R., Mendgen, K. and Deising, H.B. (1999) PR-1 protein inhibits the differentiation of rust infection hyphae in leaves of acquired resistant broad bean. *Plant J.* **19**, 625–633.
- Santamaria, M., Thomson, C.J., Read, N.D. and Loake, G.J. (2001) The promoter of a basic PR1-like gene, *AtPRB1*, from *Arabidopsis* establishes an organ-specific expression pattern and responsiveness to ethylene and methyl jasmonate. *Plant Mol. Biol.* **47**, 641–652.
- Sarma, G.N., Manning, V.A., Ciuffetti, L.M. and Karplus, P.A. (2005) Structure of Ptr ToxA: an RGD-containing host-selective toxin from *Pyrenophora tritici-repentis*. *Plant Cell*, **17**, 3190–3202.
- Schuren, F.H., Asgeirsdottir, S.A., Kothe, E.M., Scheer, J.M. and Wessels, J.G. (1993) The *Sc7/Sc14* gene family of *Schizophyllum commune* codes for extracellular proteins specifically expressed during fruit-body formation. *J. Gen. Microbiol.* **139**, 2083–2090.
- Serrano, R.L., Kuhn, A., Hendricks, A., Helms, J.B., Sinning, I. and Groves, M.R. (2004) Structural analysis of the human Golgi-associated plant pathogenesis related protein GAPP-1 implicates dimerization as a regulatory mechanism. *J. Mol. Biol.* **339**, 173–183.
- Shimoni, L. and Glusker, J.P. (1995) Hydrogen bonding motifs of protein side chains: descriptions of binding of arginine and amide groups. *Protein Sci.* **4**, 65–74.
- Szyperki, T., Fernandez, C., Mumenthaler, C. and Wuthrich, K. (1998) Structure comparison of human glioma pathogenesis-related protein GliPR and the plant pathogenesis-related protein P14a indicates a functional link between the human immune system and a plant defense system. *Proc. Natl. Acad. Sci. USA*, **95**, 2262–2266.
- Tai, Y.-S., Bragg, J. and Meinhardt, S.W. (2007) Functional characterization of ToxA and molecular identification of its intracellular targeting protein in wheat. *Am. J. Plant Physiol.* **2**, 76–89.
- Tan, K.C., Ferguson-Hunt, M., Rybak, K., Waters, O.D., Stanley, W.A., Bond, C.S. *et al.* (2012) Quantitative variation in effector activity of ToxA isoforms from *Stagonospora nodorum* and *Pyrenophora tritici-repentis*. *Mol. Plant-Microbe Interact.* **25**, 515–522.
- Tomas, A., Feng, G.H., Reeck, G.R., Bockus, W.W. and Leach, J.E. (1990) Purification of a cultivar-specific toxin from *Pyrenophora tritici-repentis*, causal agent of tan spot of wheat. *Mol. Plant-Microbe Interact.* **3**, 221–224.
- Tuori, R.P., Wolpert, T.J. and Ciuffetti, L.M. (1995) Purification and immunological characterization of toxic components from cultures of *Pyrenophora tritici-repentis*. *Mol Plant-Microbe Interact.* **8**, 41–48.
- Tuori, R.P., Wolpert, T.J. and Ciuffetti, L.M. (2000) Heterologous expression of functional Ptr ToxA. *Mol. Plant-Microbe Interact.* **13**, 456–464.
- Vartapetian, A.B., Tuzhikov, A.I., Chichkova, N.V., Taliensky, M. and Wolpert, T.J. (2011) A plant alternative to animal caspases: subtilisin-like proteases. *Cell Death Differ.* **18**, 1289–1297.
- Wang, Q., Sullivan, R.W., Kight, A., Henry, R.L., Huang, J., Jones, A.M. and Korth, K.L. (2004) Deletion of the chloroplast-localized thylakoid formation1 gene product in *Arabidopsis* leads to deficient thylakoid formation and variegated leaves. *Plant Physiol.* **136**, 3594–3604.
- Wang, Y.L., Kuo, J.H., Lee, S.C., Liu, J.S., Hsieh, Y.C., Shih, Y.T., Chen, C.J., Chiu, J.J. and Wu, W.G. (2010) Cobra CRISP functions as an inflammatory modulator via a novel Zn²⁺- and heparan sulfate dependent transcriptional regulation of endothelial cell adhesion molecules. *J. Biol. Chem.* **285**, 37 872–37 883.
- Xu, X., Francischetti, I.M., Lai, R., Ribeiro, J.M. and Andersen, J.F. (2012) Structure of protein having inhibitory disintegrin and leukotriene scavenging functions contained in single domain. *J. Biol. Chem.* **287**, 10 967–10 976.
- Zhang, H.F., Francl, L.J., Jordahl, J.G. and Meinhardt, S.W. (1997) Structural and physical properties of a necrosis-inducing toxin from *Pyrenophora tritici-repentis*. *Phytopathology*, **87**, 154–160.
- Zhang, W.J., Pedersen, C., Kwaaitaal, M., Gregersen, P.L., Morch, S.M., Hanisch, S., Kristensen, A., Fuglsang, A.T., Collinge, D.B. and Thordal-Christensen, H. (2012) Interaction of barley powdery mildew effector candidate CSEP0055 with the defence protein PR17c. *Mol. Plant Pathol.* **13**, 1110–1119.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Fig. S1 Plant assays showing activity of the recombinant ToxA proteins in the ToxA-sensitive tetraploid Langdon wheat. Second leaves of 2-week-old plants were infiltrated with ToxA at the estimated final concentrations (ng/μL) indicated on the right (highest concentration shown for the N102A mutant). Control plants were treated with the protein sample of the 'mock' strain shown in Fig. 3A (lane 3). Marks delimit the infiltrated area. Photographs were taken at 72 h post-infiltration (one representative leaf of each treatment is shown).

Fig. S2 Characterization of the purified recombinant PR-1-5 proteins. (A) Size exclusion chromatography analysis of the recombinant PR-1-5 (left) and PR-1-5-N141A (right) proteins. A single peak was detected for both proteins (samples were separated using Phenomenex Yarra SEC-2000 columns; Torrance, CA, USA). (B) Mass spectrum showing the tryptic fragment (peak, 41.85) containing the single-residue mutation site in the recombinant PR-1-5-N141A mutant protein. The amino acid (aa) sequence of the 18-aa fragment (corresponding positions of 138–155 in the PR-1-5 protein) is shown with the mutated residue highlighted in red [detailed tandem mass spectrometry (MS/MS) data are available on request].

Fig. S3 Plant assays showing the necrosis-promoting activity of the PR-1-5 protein in hexaploid wheat cultivars. ToxA-sensitive (Grandin, A) and insensitive (BR34, B) wheat plants were infiltrated with the recombinant ToxA and/or PR-1-5 proteins as indicated on the right. Photographs were taken at 48 h post-infiltration (one or three representative leaves of each treatment are shown).

Fig. S4 Expression patterns of *PR-1-1*, *ToxABP1* and *PCN* in comparison with that of *PR-1-5* (Fig. 5) in response to ToxA treatment. mRNA samples were collected from the second leaves of ToxA-insensitive (BR34) and ToxA-sensitive (Grandin) wheat plants that were 'mock'-infiltrated (–) or infiltrated with ToxA protein (+) at a final concentration of 1.0 ng/μL. Polymerase chain reaction (PCR) products were separated on ethidium bromide-stained 1% agarose gels. Numbers on the right indicate size (in base pairs) of the amplicons. Oligonucleotide sequences of the primers used for reverse transcriptase-polymerase chain reaction (RT-PCR): 5'-CCTTTTAGACACCGAACCAGGAAGT-3' (forward) and

5'-ATGGCGCCATATCGTCGCTTCT-3' (forward) and 5'-TTAATGCTTCAAGGGATAAGCGCT-3' (reverse) for TOXABP1 (Manning *et al.*, 2007), and 5'-ATGGCCGCCCTCTCCTCTGCA-3' (forward) and 5'-TTAGTTGACGGTGACCTTGCCGA-3' (reverse) for PCN (Tai *et al.*, 2007).

Fig. S5 Amino acid sequence alignment showing the residues consisting of the turning loop between β C and β D in the PR-1 domains of plant PR-1 proteins. Residues shown correspond to positions 135–150 of the PR-1-5 protein (GenBank accession number HQ541965). Arrows indicate residues aligned to the N141 site of PR-1-5 and its 'sister' protein PR-1-4 (underlined), both of which have a unique 'NNNL' motif (highlighted in yellow). Numbers on the right indicate the positions in the full-length

proteins. Alignments were generated using the MegAlign program (DNASTAR). Species abbreviations: At, *Arabidopsis thaliana*; Bn, *Brassica napus*; Cc, *Capsicum chinense*; Ca, *C. annuum*; Ew, *Eutrema wasabi*; Gm, *Glycine max*; Hv, *Hordeum vulgare*; Le, *Lycopersicon esculentum*; Md, *Malus \times domestica*; Nt, *Nicotiana tabacum*; Os, *Oryza sativa*; Sb, *Sorghum bicolor*; St, *Solanum tuberosum*; Ta, *Triticum aestivum*; Tm, *T. monococcum*; Zm, *Zea mays*. The 23 wheat (Ta) PR-1-like proteins (PR-1-1 to PR-1-23) have been reported previously by Lu *et al.* (2011).

Table S1 Polymerase chain reaction (PCR) primers used for the generation of yeast two-hybrid constructs.

Table S2 Polymerase chain reaction (PCR) primers used for the generation of *Pichia* expression constructs.