



Published in final edited form as:

Insect Biochem Mol Biol. 2019 August ; 111: 103182. doi:10.1016/j.ibmb.2019.103182.

Mosquito-fungus interactions and antifungal immunity

P. Tawidian¹, V. L. Rhodes², and K. Michel^{1,*}

¹Division of Biology, Kansas State University, 267 Chalmers Hall, Manhattan, KS 66506

²Missouri Southern State University, Biology Department, Reynolds Hall 220, 3950 E. Newman Rd., Joplin, MO 64801-1595 USA


Abstract

The mosquito immune system has evolved in the presence of continuous encounters with fungi that range from food to foes. Herein, we review the field of mosquito-fungal interactions, providing an overview of current knowledge and topics of interest. Mosquitoes encounter fungi in their aquatic and terrestrial habitats. Mosquito larvae are exposed to fungi on plant detritus, within the water column, and at the water surface. Adult mosquitoes are exposed to fungi during indoor and outdoor resting, blood and sugar feeding, mating, and oviposition. Fungi enter the mosquito body through different routes, including ingestion and through active or passive breaches in the cuticle. Oral uptake of fungi can be beneficial to mosquitoes, as yeasts hold nutritional value and support larval development. However, ingestion of or surface contact with fungal entomopathogens leads to colonization of the mosquito with often lethal consequences to the host. The mosquito immune system recognizes fungi and mounts cellular and humoral immune responses in the hemocoel, and possibly epithelial immune responses in the gut. These responses are regulated transcriptionally through multiple signal transduction pathways. Proteolytic protease cascades provide additional regulation of antifungal immunity. Together, these immune responses provide an efficient barrier to fungal infections, which need to be overcome by entomopathogens. Therefore, fungi constitute an excellent tool to examine the molecular underpinnings of mosquito immunity and to identify novel antifungal peptides. In addition, recent advances in mycobiome analyses can now be used to examine the contribution of fungi to various mosquito traits, including vector competence.

Graphical Abstract

*Corresponding author: kmichel@ksu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Keywords

mycobiota; entomopathogen; infection; melanization; humoral immunity

Introduction

Fungi hold nutritional value for larval and adult mosquitoes, and thus are mosquito prey (Asahina, 1964). Fungal commensals can form longer-term associations in the mosquito larval gut, with little to no impact on host survival (Tuzet and Manier, 1947). In contrast, some water molds (Chromista) and fungi utilize mosquitoes as their growth medium, most often with detrimental or lethal effects on their host (Braun, 1855; Galli-Valerio and Rochaz de Jongh, 1906; Eckstein, 1922; Roubaud and Toumanoff, 1930; Couch, 1935). The potential use of such fungal entomopathogens to control mosquito populations has largely driven the research on fungal-mosquito interactions over more than a century (reviewed in e.g. Christophers, 1952; Castillo and Roberts, 1980a,b; Jenkins 1964; Roberts, 1974; Lakon, 1919; Kanzok and Jacobs-Lorena, 2006; Scholte, 2005). While this holds true today (e.g. Lovett et al., 2019), mosquito microbiome studies have recently been extended to fungi (Kaufman et al., 2008; Muturi et al., 2016; Luis et al., 2019). These studies provide new insight into the range of mosquito-fungi encounters that may be fleeting, but may stimulate mosquito immunity through fungal surface molecules and/or secondary metabolites.

Entomopathogenic chromista and fungi infect mosquitoes through different routes, including active penetration of the body wall, through wounds, or through ingestion, dependent on the pathogen as well as the life stage of the mosquito (e.g. Clark et al., 1966, 1968; McCray et al., 1973; Greenfield et al., 2014). Most commonly, entomopathogens isolated from field-caught mosquitoes belong to the genus *Coelomomyces*, which are ascomycetes fungi in the order of Blastocladales (Castillo and Roberts, 1980; Couch and Umphlett, 1963). *Coelomomyces* fungi infect mosquito larvae through their aquatic zoospores. Zygospores attach to the larval integumental cuticle, and penetrate the host body wall by a penetration tube. Hyphal growth and sporangia formation occurs in the hemolymph, leading to larval death (reviewed in Araujo and Hughes, 2014; Scholte et al., 2004). However, their obligate two-host life cycle limits their use in mosquito control programs (Laird, 1967; Service, 1977). Currently, entomopathogens being proposed for mosquito control are ascomycetes fungi in the genera of *Beauveria* and *Metarhizium* in the order Hypocreales; Farenhorst et al., 2011; Lovett et al., 2019; Popko et al., 2018; Scholte et al., 2005). In general, fungi in these two genera infect mosquitoes via asexual spores called conidia, which germinate and produce penetration pegs that breach the host cuticle. Upon reaching the hemolymph, the

fungus proliferates and forms yeast-like blastospores that rapidly colonize the host leading to its death (reviewed in Scholte et al., 2004).

Recognition of largely unknown fungal and water mold-derived molecules by mosquito pathogen recognition receptors (PRRs), activates mosquito epithelial, cellular and humoral immune responses. Mosquito cellular immune responses include phagocytosis, encapsulation, and nodulation (Bartholomay and Michel, 2018; Hillyer and Strand, 2014). Humoral immune responses include complement-like reactions, melanization, and the expression of antimicrobial peptides (Blandin et al., 2008; Nakhleh et al., 2017; Rhodes and Michel, 2017). These immune responses are regulated through proteolytic activation of key immune factors and/or on a transcriptional levels by several signal transduction pathways including Toll pathway, the immune deficiency (IMD) pathway, Janus Kinase (JAK)-signal transducer and activator of transcription (STAT), and Jun N-terminal kinase (JNK) and mitogen-activated protein kinase (MAPK) p38 pathway (Nakhleh et al., 2017; Rhodes and Michel, 2017). The most commonly observed immune response against fungi and water molds is melanization (Brey et al., 1988; Coluzzi, 1966; Mc Innis and Zattau, 1982; Clark et al. 1968). Molecular insights of mosquito antifungal immune responses are driven by comparative invertebrate immunology (Al Souhail et al., 2016; Chen et al., 2010; Valanne et al., 2011), and experiments performed using a small number of fungal laboratory infection models (Vizioli et al., 2001a; Ramirez et al., 2019; Rhodes et al., 2018; Yassine et al., 2012).

This review will summarize the potential environmental and molecular interactions of mosquito adults and larvae with fungi and water molds. The first two sections will provide an overview of their reported environmental encounters, and detail the potential routes of infection. The sum of these interactions across the symbiotic continuum shapes the mosquito anti-fungal immune responses, which we describe in the final section of this review.

1. Mosquito-fungus encounters in the environment

Fungi are ubiquitous and found in all mosquito habitats (Goh and Hyde, 1996). In addition, the aquatic environment of mosquito larvae provides habitat for water molds. This close proximity leads to a myriad of possible interactions between mosquitoes and fungi as well as water molds throughout the mosquito's life history (Fig. 1). We mined the existing literature (see Supplementary Text for methods and references) and found records for 158 and 43 species of fungi and water molds, respectively having been observed in/isolated across 149 mosquito species (Tables S1 and S2, and references within). Only one third of these species have been isolated/observed in adult mosquitoes, reflecting the substantially lower number of fungus sampling efforts in adult mosquitoes compared to collections in immature life stages (Table S2). Two thirds of the isolated fungi belong to the orders Blastocladales (39 species, 38 of which in the genus *Coelomomyces*), Eurotiales (28 species, including 16 *Penicillium* sp. and ten *Aspergillus* sp.), Hypocreales (24 species across 14 genera including *Beauveria* and *Metarhizium*), and Saccharomycetales (17 species with more than half belonging to the genus *Candida*). Only one sixth of the isolated water molds are known to be pathogenic, while nearly two thirds of the isolated fungi are either opportunistic, facultative or obligate pathogens (Table S1). However, these numbers are likely underestimates of the true range of mosquito-fungi interactions. Amplicon-based sequencing of adults from eleven

mosquito species identified recently 347 and 1 species of fungi and water molds, respectively. Of these, 96 % were described for the first time in or on adult mosquitoes, and their impact on mosquito immunity is entirely unknown.

This section describes the possible encounters with fungi and water molds across the mosquito life history, facilitated by mosquito behavior in their aquatic and terrestrial environments.

1.1 Larva-fungus interactions through ingestion

Fungi and water molds in the aquatic environment can enter the mosquito body passively through diverse larval feeding behaviors, including collecting-filtering, as well as grazing on and shredding of decaying and living organic matter (Fish and Carpenter, 1982; Merritt et al., 1992; Yee et al., 2004). Upon uptake, they either are digested, pass through the intestine unharmed, or manage to stay and grow within the mosquito host.

Fungi act as food and provide nutrients for the development of mosquito larvae. An example is the baker's yeast, *Saccharomyces cerevisiae* that is commonly used for the rearing of mosquito larvae (Asahina, 1964). Seven individual yeast species that had been isolated previously from field-collected *Culex theileri* and *Cx. pipiens* larvae, supported the growth of *Cx. pipiens* larvae. This study thus confirmed the nutritional value of yeast ingestion by mosquito larvae (Steyn et al., 2016). Shift in fungal microbial populations on decaying oak leaf matter in the presence and absence of *Aedes triseriatus* larvae further suggests the ingestion of fungi through larval browsing and grazing (Fish and Carpenter, 1982; Kaufman et al., 2008).

Mosquito feeding behavior also leads to the establishment of commensals in the mosquito hindgut. At least four fungal species in the genus *Smittium* sp., belonging to the order of Harpellales, are able to attach and replicate in the hindgut of various mosquito species, with most often no effect on larval development or survival (Lopez-Lastra, 1997; Pereira et al., 2005; Sweeney, 1981; Tuzet and Manier, 1947; White et al., 2006).

During filter feeding and grazing, mosquito larvae do also ingest potential entomopathogenic fungi and water molds. The fungal entomopathogen *Culicinomyces clavispurus* causes infection in *Culex fatigans* larvae only upon ingestion of the fungal conidia (Sweeney, 1975). Likewise, *Smittium morbosum* infects mosquito larvae solely after oral uptake (Sato et al., 1989; Sweeney, 1981). *M. anisopliae* has deleterious effects on the larval hosts after oral uptake, which may be due to subsequent infection of the hemocoel, but at least in certain instances are due to toxicity in the gut (Cheng and Liu, 1990; Butt et al., 2013). The water molds *Leptolegnia chapmanii* and *Lagenidium giganteum* both infect the hemocoel of mosquito larvae after oral uptake of zoospores and by cuticular penetration of the body wall (Zattau and McInnis, 1987; McCray et al., 1973).

1.2. Larva-fungus interactions through contact

Entomopathogenic fungi and water molds commonly infect mosquito larvae through active cuticular penetration. Fungi in the genus *Coelomomyces*, including *C. opifexi* are only known to infect mosquito larvae via cuticle penetration (Wong and Pillai, 1980). *C.*

psorophorae spores infect only through attachment and cuticular penetration of the head, intersegmental regions, and the base of anal gills (Travland, 1979). In addition to aquatic fungal entomopathogens, soil-borne entomopathogens like *M. anisopliae* and *B. bassiana*, infect larvae of multiple mosquito species via direct penetration of the cuticle. Studies on *M. anisopliae* route of infection in mosquito larvae suggest that infection is mainly due to contact with spores at the water surface, mostly in and around the siphon (Crisan, 1971; Lacey et al., 1988). Likewise, dependent on formulation, *B. bassiana* conidia float on the water surface, and exposure preferentially happens around the perispiracular lobes (Clark et al., 1968).

1.3. Transstadial transmission of fungal communities

Transstadial transmission of bacterial microbiota from mosquito larvae to pupae and adults is limited, and has been reported in *Culex tarsalis*, *Cx. pipiens*, *An. gambiae*, and *Ae. aegypti* (Duguma et al., 2015; Moll et al., 2001). Studies conducted on yeast in *Culex* spp. have shown no transmission from larvae to pupae, or to adult stages (Díaz-Nieto et al., 2016; Steyn et al., 2016). These data suggest that transstadial transmission of fungi and water molds, especially if located solely in the midgut, does not occur and is not the source of fungal associations or infections in adult mosquitoes. One exception are the 36 species of *Coelomomyces*. While their persistence to adult mosquitoes occurs at a low frequency, transmission from infected larvae to pupae and adults has been observed repeatedly (Garland and Pillai, 1979; Lucarotti, 1987; Lucarotti and Andreadis, 1995).

1.4. Adult-fungus interactions through ingestion

Adult male and female mosquitoes commonly feed on plant nectars (Foster, 1995). These nectars contain many yeasts that likely are taken up by adult mosquitoes during feeding. The yeast nectar community is diverse and largely depends on inoculation through pollinator species including insects and birds (reviewed in Chappel and Fukami, 2018). Yeast community composition within the guts of pollinators overlaps with the yeast nectar community, strongly suggesting that such composition is largely driven by oral uptake and environmental filtering (Sandhu and Waraich, 1985). Several yeasts that have been identified in field-collected adult mosquitoes, including *C. parapsilosis* and *Hansenula* spp. in *Ae. triseriatus*, as well as *Pichia* spp. in *Aedes japonicus* (Bozic et al., 2017; Muturi et al., 2016; Ricci et al., 2011), overlap with nectar-associated yeasts. Most recently, amplicon-based sequencing of adult *Ae. albopictus* identified a large number of yeast species (Luis et al., 2019). Together, these studies support the notion that nectar-associated yeasts are regularly ingested during nectar feeding by adult mosquitoes (Fig. 1G). Similarly, yeast may be taken up during blood feeding (Fig. 1C). Fungal cultures from human blood samples and blood-fed female mosquitoes share several yeast species, including *C. parapsilosis*, *Rhodotorula* spp., *Saccharomyces* spp., and *Cryptococcus* spp. (Bille et al., 1982; Chang et al., 2001; Muturi et al., 2016).

1.5. Adult-fungus interactions through substrate contact

Adult mosquitoes commonly rest in- and outdoors, and thus are exposed continuously to fungi through contact with contaminated surfaces. Mosquitoes, resting in tree cavities or on flower and leaf surfaces, come in touch with a diverse community of microorganisms

including yeasts and filamentous fungi (Fig. 1. H, D, Levetin and Dorsey, 2006). Examples of such surface contact exposures is the occurrence of infections of adult *Culex pipiens* with the fungal entomopathogen *Entomophthora conglomerata* and *Entomophthora destruens* in wine cellars and natural caves, respectively (Novak, 1965; Weiser and Batko, 1966). This exposure route is also exploited for vector population control purposes. The provision of *M. anisopliae* and *B. bassiana*-impregnated netting, cloth, and mud panels as indoor resting sites increased the mortality of *An. gambiae* adults (Fig. 1A, Mnyone et al., 2010). Similarly, outdoor mosquito resting sites can be targeted using both entomopathogen species on the surface of clay pot traps, roof covers, and within resting stations like extra-domiciliary odor-baited stations (Luz et al., 2010; Lwetoijera et al., 2010; Mnyone et al., 2010). In addition, transfer of spores through contact with screens impregnated with *B. bassiana* in oviposition traps was used successfully to target female *Ae. aegypti* (Snetselaar et al., 2014).

Gravid female mosquitoes can locate suitable oviposition sites by the detection of secondary metabolites produced by certain fungi (Eneh et al., 2016; Geetha et al., 2003). During oviposition, these females are exposed to these and other soil-borne and/or aquatic fungi through direct contact (Fig. 1E). In how far these brief encounters contribute to fungal-mosquito interactions is currently unclear. The positive correlation of infection prevalence with number of ovipositions suggests that at least *Entomophthora conglomerata* may be able to infect *Culex pipiens* during oviposition (Kupriyanova, 1966). It is feasible that treatment of oviposition sites with fungal entomopathogens to control mosquito egg and early larval stages may also cause infection in adult females (Sousa et al., 2013).

1.6. Additional adult-fungus interactions

Mosquitoes can pass on spores that are attached to their own cuticle. Transfer of entomopathogenic fungal spores between individual mosquitoes through mating has been observed in several species. Infection of healthy male *An. gambiae* mosquitoes by *M. anisopliae* was observed post-mating with topically exposed female mosquitoes (Scholte et al., 2004a). Reduction in survival of healthy female *Ae. aegypti* mosquitoes post-mating was observed with both *M. anisopliae* and *B. bassiana*-exposed male mosquitoes, confirming conidial dissemination through mating in *Ae. aegypti* mosquitoes (Fig. 1B) (García-Munguía et al., 2011; Garza-Hernández et al., 2015; Reyes-Villanueva et al., 2011). As such, mosquitoes can be used to auto-disseminate fungal entomopathogens (Scholte et al., 2004a), and thus extend the reach of these biocontrol agents beyond the initial contact with the impregnated surface (Shah and Pell, 2003). Autodissemination may also extend to oviposition sites (Fig. 1E). A specialized case is the oviposition of resting sporangia of *Coelomomyces* sp., including *C. psorophorae* and *C. stegomyiae* by female mosquitoes. Infected females are sterile, but take blood meals, and actively oviposit sporangia, a behavior that contributes to the dissemination of the fungus to new aquatic environments (Lucarotti, 1987; Laird et al., 1992).

In addition to surfaces, indoor and outdoor air contains a highly diverse fungal spore community (Frohlich-Nowoisky et al., 2009), with some taxa also being found in mosquitoes (Guegan et al., 2018; Muturi et al., 2016). Conidia of *B. bassiana* can be air-

borne (Feng et al., 1994), and thus may come in contact with mosquitoes through air rather than surfaces (Clark et al., 1968).

2. Getting in: Colonization of mosquitoes by fungal and water mold symbionts

Out of the 158 species of fungi for which we found published evidence of isolation from mosquitoes, roughly half can establish longer-term interactions with their hosts. 67 species from six fungal orders are considered pathogens of mosquitoes, using the hemocoel and/or various tissues as growth medium (Section 1, Fig. 2, top and middle panel). Of these, only *Entomophthora sp.* and *Coelomomyces sp.* are obligate pathogens, while the rest are facultative pathogens that switch between pathogenic and saprophytic or even endophytic life styles. In addition, 22 species of fungi in the orders Eurotiales, Hypocreales and Mucorales are opportunistic pathogens, who themselves cannot actively invade the mosquito hemocoel, but can establish an infection if entering through breaches in the cuticle (Fig. 2, top panel). Five *Smittium* species (order Harpellales) are commensals of mosquitoes, attaching to and growing within the hindgut of mosquito larvae (see Section 2.2, Fig. 2, bottom panel).

Out of the 43 species of water molds that have been observed and/or isolated from mosquitoes, only five species (in the genera *Lagenidium*, *Leptolegnia* and *Saprolegnia*) are known facultative pathogens of mosquitoes. No commensal water molds have been described in mosquitoes. This section will briefly describe how some of these symbionts colonize their mosquito hosts, either through breaching the cuticle and/or through ingestion.

2.1. Active penetration of cuticle

As mentioned in Sections 1.2 and 1.5 above, insects often encounter entomopathogens through contact. Upon contact, colonization of the host is achieved through active penetration, a multistep process that somewhat varies dependent on fungal species (reviewed in Butt et al., 2016; Scholte et al., 2004; Lovett and St. Leger, 2017). In general, active penetration begins with the attachment of spores to the host epicuticle. Fungal spores germinate and form germ tubes that differentiate into specialized attachment organs, called appressoria, that then form a penetration tube, which penetrates the host cuticular layers through pressure exertion and secretion of cuticle-degrading enzymes. Upon reaching the hemolymph, the fungus proliferates as blastospores (yeast-like budded cells) or hyphal bodies (chains of budded cells). As the infection progresses, the insect host succumbs to the infection, either through fungal toxins or by starvation. In most fungal entomopathogens, hyphae break through the cuticle to produce either infective spores or resting structures on the insect cadaver (Fig. 2, top panel; recently reviewed in Lovett and St. Leger, 2017).

Infection by penetration is observed in mosquito larvae and adults, with histological examinations available for several species combinations of fungi/water molds and mosquitoes. Clark et al. (1968) describes the histological manifestation of *B. bassiana* infection in *Cx. pipiens* larvae, observing mycelial growth preferentially around the tracheal trunks, suggesting invasion in an around the trachea. Infection of *Cx. pipiens* larvae by *B.*

bassiana led to the disintegration and deformation of the epicuticle, midgut epithelium, and muscles (Benzina et al., 2018). Cuticle penetration has also been commonly observed in mosquito larval infections by *Coelomomyces* sp. (Shoukamy and Lucarotti, 1998; Wong and Pillai, 1980). *C. psorophorae* zygospore adhesion to *Culiseta inornata* larval cuticle is followed by germination and development of a narrow hyphal body that penetrates the host epidermal cells (Travland, 1979). An ultrastructural study of *C. stegomyiae*-infected *Ae. aegypti* larvae revealed hyphal growth in the muscles with deterioration of myofibrils, degeneration of midgut muscles, and fragmentation of the microvilli that line the malpighian tubules (Shoukamy et al., 2001).

The infection process of water molds is similar. Motile zoospores of *L. giganteum* attach to the cuticle of mosquito larvae, a germ tube penetrates the body wall, hyphae grow in the hemocoel, and start to produce sporangia. The sporangia develop exit tubes that grow out through the cuticle, ultimately releasing zoospore vesicles into the aquatic environment. In addition, resting structures in the form of sexual oospores can also be produced (Fetter-Lasko and Washino, 1983; Brey et al., 1988). Attachment of *L. giganteum* zoospores followed by penetration of the larval cuticle has been observed in *Aedes aegypti*, *An. gambiae*, and *Cx. pipiens* larvae (Brey et al., 1988; Golkar et al., 1993). Similarly, zoospores of *L. chapmanii* have been shown to aggregate, and then enters via penetration tubes the head, body, anal papillae, and intersegmental folds of *Ae. aegypti* and *Cx. quinquefasciatus* larvae (Lord and Fukuda, 1988; Zattau and McInnis, 1987).

Exposure of adult *An. stephensi* mosquitoes to *B. bassiana* impregnated filter papers, led to the accumulation and germination of fungal spores on the proboscis, tarsi, legs and wings. During early stages of infection, hyphae invaded tissues and organs, including compound eyes, brain, salivary glands, mouthparts, muscles, midgut, ovaries, and malpighian tubules (Ishii et al., 2017). *Conidiobolus coronatus* infects *Aedes taeniorhynchus* and *Cx. quinquefasciatus* adult mosquitoes through cuticle penetration of the intersegmental interstices, head, and dorsal thoracic regions. Histological studies on infected mosquitoes revealed the dissemination of hyphal bodies in hemocoel, muscles, fat body, and gonads (Lowe et al., 1968; Lowe and Kennel, 1972). Infection of adult *Aedes sierrensis* mosquitoes with *Tolypocladium cylindrosporum*, showed fungal hyphae and hyphal bodies exclusively in the thorax, suggesting active penetration through the cuticle of the thoracic spiracles (Soarés, 1982).

2.2. Ingestion of fungal spores

Fungal spore uptake through ingestion has been commonly observed in mosquito larvae. Ingestion of *C. clavosporus*, *B. bassiana*, *M. anisopliae*, and *Aspergillus clavatus* spores was reported for *Aedes*, *Culex*, and *Anopheles* mosquito larvae (Miranpuri and Khachatourians, 1991; Scholte et al., 2004b; Seye et al., 2009). Upon ingestion, the spores may remain in the midgut and release toxins that cause pathology and larval killing (Fig. 2, center panel). Cross sections of *Cx. pipiens* larvae during early stages of infection by *M. anisopliae* revealed the accumulation of intact fungal spores in the gut, confirming spore ingestion. Digestion of fungal spores was observed in cross section of dead larvae where large number of spores were disrupted (Crisan, 1971). Ingestion of *M. anisopliae* spores by *Cx. quinquefasciatus*

larvae induced larval mortality 24 hours post-exposure, suggesting the release of toxins during fungal spore digestion. Guts of infected larvae revealed the accumulation of non-germinated and partially digested fungal spores (Lacey et al., 1988). In neither study, oral uptake of *M. anisopliae* did result in disseminated fungal infection.

Other fungal entomopathogens spread from the midgut lumen throughout the mosquito body by penetration through the midgut epithelium (Fig. 2, center panel). For example, the conidia of *C. clavissporus* adhere to the gut walls of *Cx. fatigans* mosquito larvae, where they germinate and penetrate the gut epithelium (Sweeney, 1975). Ingestion of *B. bassiana* blastospores rather than conidia by *Ae. aegypti* larvae result in the spread of fungal spores throughout the larval body 24 hours post ingestion, with heavy colonization in the fore-, mid-, and hindgut (Miranpuri and Khachatourians, 1991). Exposure of *Cx. quinquefasciatus* larvae to *A. clavatus* led to the accumulation of fungal spores in the alimentary canal. Infection resulted in the destruction of gut cells and disruption of the gut epithelium in several locations, facilitating fungal dissemination in the mosquito body (Bawin et al., 2016). Ingestion *Fusarium oxysporum* spores by *Aedes detritus* and *Cx. pipiens* led to the colonization of the gut lumen and penetration of the epithelial lining (Hasan and Vago, 1972).

Several species in the genus *Smittium* establish symbiotic relationships due to attachment to the cuticular lining of the hindgut (Fig. 2, bottom panel; Lichtwardt, 1996). In most cases, the symbiotic relationship remains commensal, and the fungus is shed with every molt. Pathology can be induced by hyphal overgrowth causing complete blockage of the alimentary canal, as it is observed for *Smittium culisetae* (Williams and Lichtwardt, 1972). The only species within this genus that can be considered a facultative pathogen of mosquito larvae is *Smittium morbosum* (Coluzzi, 1966; Sweeney, 1981; Sato et al., 1989). Upon initial attachment to the hindgut, *Smittium morbosum* grows anteriorly, and hyphae penetrate the posterior midgut epithelium. Systemic infection is prevented by the mosquito immune system through strong melanization at the basal side of the midgut epithelium. However, this melanotic mass anchors the hyphae, and either allows transstadial transmission, or leads to incomplete molting, causing larval death. *S. morbosum*.

In addition to infection by cuticle penetration, as described in section 2.1, zoospores of the water mold *L. chapmanii* have been shown to be ingested by *Ae. aegypti* larvae (Pelizza et al., 2008; Zattau and McInnis, 1987). Upon ingestion, the zoospores germinated in the midgut, elongated and formed branched hyphae that penetrated the peritrophic matrix and midgut epithelium, reaching the hemocoel. Once in the hemocoel, infection proceeded similarly to infection initiated by cuticle penetration (Zattau and McInnis, 1987).

To the best of our knowledge, there is currently no evidence that adult mosquitoes commonly ingest entomopathogenic fungi, or that oral uptake of fungal entomopathogens results in adult mosquito infection.

2.3 Fungal entry through injury

Fungal infection of mosquitoes can be facilitated through injuries that breach the cuticle, as these allow entry of opportunistic pathogens (Fig. 2, top panel). The entomopathogenic

fungus *Pythium sp.* cannot infect mosquito larvae through penetration, but infects and kills mechanically injured *Aedes*, *Anopheles*, and *Culex* mosquito larvae (Clark et al., 1966). Experimental infections can also be achieved through intrathoracic conidial spore injection, as observed for *B. bassiana* in adult *An. gambiae* mosquitoes and *M. anisopliae* and *A. clavatus* in *Cx. quinquefasciatus* larvae (Bawin et al., 2014; Yassine et al., 2012). While this route of infection is perhaps occurs less frequently in the field, and is difficult to exploit for biological control, it can facilitate research of molecular mosquito-fungal interactions and antifungal immunity.

3. Antifungal immune mechanisms in mosquitoes

Once a water mold or a fungus has penetrated the mosquito cuticle and entered the hemocoel, it encounters both efficient cellular and humoral immune responses. Survival thus depends on its ability to largely avoid or eventually overcome these immune responses. Nevertheless, immune escape is incomplete. For example, *B. bassiana* is recognized, and its growth is limited by the mosquito immune system. The outcome of infection is thus strongly influenced by the immune competency of the host. Somewhat surprisingly, few data are available to date on the molecular interactions of mosquitoes with fungi. This section will provide a brief overview of the molecules and pathways currently known to be involved in fungus recognition and killing in various mosquito species.

3.1 Recognition

β -1,3-glucans are the known molecules on the surface of fungal cell walls that are recognized by insect immune systems (Lu and St. Leger, 2016). Studies in various insects have identified β -1,3-glucan recognition proteins (GRPs) and Gram-negative binding proteins (GNBPs) to bind to β -1,3-glucan and trigger downstream immune responses, including Toll pathway activation and/or melanization (Jiang et al., 2004; Matskevich et al., 2010; Ochiai and Ashida, 2000). The only molecule currently confirmed to bind β -1,3-glucan in mosquitoes is a GRP from *Armigeres subalbatus* (AsGRP). AsGRP binds to curdlan, an insoluble β -1,3-glucan polymer, and is required for melanization of some bacterial species (Wang et al., 2006, 2005). If AsGRP contributes to melanization of fungal species is unclear. GGBP2 was found to be transcriptionally upregulated in *An. gambiae* mosquitoes that were challenged by injection of dead *B. bassiana* conidia (Aguilar et al., 2005). Whether GGBP2 can directly bind to fungal surfaces however remains to be tested. The water mold *L. chapmanii* is also detected in mosquito larvae, as evidenced by localized immune responses to its surface (Zattau and McInnis, 1987). The cell wall of *L. chapmanii*, and that of all water molds, does not contain chitin, and it is unclear how and which molecules on its surface are responsible for immune recognition.

Thioester-containing proteins (TEPs) bind to many foreign surfaces and are critical for pathogen recognition and innate immunity in mosquitoes (Blandin and Levashina, 2004). The activation of the complement-like pathway in mosquitoes leads to the deposition of thioester-containing protein 1 (TEP1) on foreign surfaces (Fig. 3). TEP1 acts as an opsonin and promotes the phagocytosis of bacteria (Levashina et al., 2001; Moita et al., 2005) and lysis of the rodent malaria parasite, *Plasmodium berghei* (Blandin et al., 2004). Several

studies point to the involvement of TEPs in antifungal immunity in mosquitoes. Infection of *Ae. aegypti* mosquitoes with *B. bassiana* and *Cordyceps javanica* resulted in upregulation of *TEP22* in fat body and midgut of the infected mosquitoes (Ramirez et al., 2018a; 2019). Knockdown of *TEP22* increased the susceptibility of *Ae. aegypti* to *B. bassiana* infection, confirming the role of *TEP22* in antifungal immune response (Wang et al., 2015). In *An. gambiae*, *TEP1* binds to *B. bassiana* hyphal bodies (Yassine et al., 2012). Knockdown of *TEP1* decreased hyphal melanization and increased susceptibility of *An. gambiae* mosquitoes to infection by *B. bassiana* (Yassine et al., 2012). Together, these studies suggest that TEPs are important molecules for recognition of fungi in the mosquito hemocoel and critically contribute to the antifungal immune response in mosquitoes. However, the binding partner of TEPs on the surface of fungi, or any other surfaces awaits identification.

In addition to β -1,3-glucan, other sugar moieties on the fungal surface may be recognized. Neuraminic acid and/or sialic acid on the surface of *B. bassiana* blastospores are potential binding partners for lectins (Wanchoo et al., 2009). It remains to be seen whether mosquito-derived lectins can recognize these sugars and contribute to antifungal immunity or immunity against water molds.

3.2 Killing mechanisms

3.2.1 Cellular Immune Responses—Cellular immune responses, mediated by hemocytes, can kill fungal cells within the hemolymph in insects. Hemocytes phagocytize, encapsulate, and nodulate blastospores and hyphal bodies (Lavine and Strand, 2002). Phagocytosis is initiated by the recognition and binding of phagocytic cells to fungal particles, leading to particle engulfment through cytoskeleton modification and intracellular vesicular transport to the phagosome (Jiravanichpaisal et al., 2006). Phagocytosis of blastospores of the fungal species *M. anisopliae*, *B. bassiana*, *Cordyceps farinosa*, *Metarhizium rileyi*, *Cordyceps fumosorosea*, and *C. albicans* has been observed in several insect species (Hung et al., 1993; Hung and Boucias, 1992; Kawakami, 1965; Ouedraogo et al., 2003; Pendland and Boucias, 1996). In mosquitoes, granulocytes and less commonly prohemocytes exhibit phagocytic activity (Hillyer and Strand, 2014). Inoculation of *An. albimanus* with *S. cerevisiae* resulted in the melanization of yeast cells and encapsulation by plasmatocytes. However, phagocytosis of yeast cells was not observed (Hernández-Martínez et al., 2002).

Cellular immune responses have also been reported in response to water mold infections. *L. giganteum* increased transiently total hemocyte numbers in *Cx. quinquefasciatus* larvae, as well as causing a relative increase of granulocytes compared to over hemocyte cell types (Fei and Huai-en, 2001). This suggests that *L. giganteum* can stimulate cellular immunity in mosquitoes. Indeed, hemocytes in *Ae. aegypti* larvae form loose capsules on *L. giganteum* hyphae. In how far these cellular responses limit *L. giganteum* infection of mosquito larvae remains to be tested.

3.2.2 Antimicrobial Peptides (AMPs)—In addition to cellular immune responses, insects mount humoral immune responses against fungi, including the expression of antimicrobial peptides (AMPs, Fig. 3). Most insect AMPs are short amphipathic peptides

that are active against gram-positive and/or gram-negative bacteria (Boman, 2003; Tonk and Vilcinskis, 2017). All mosquito genomes annotated to date, encode members of the defensin, cecropin, dipterin, and gambicin AMP families (Bartholomay et al., 2010; Christophides et al., 2002; García Gil de Muñoz et al., 2008; Neafsey et al., 2015; Waterhouse et al., 2007). In addition, a putative *attacin* gene has been identified in *Ae. aegypti* and *An. gambiae* (Waterhouse et al., 2007). Of these, defensin A and C (DEF1), cecropin A (CEC1), and gambicin (GAM1) are currently implicated in antifungal immunity. Infection of *Cx. quinquefasciatus* larvae with *Metarhizium brunneum* conidia and blastospores upregulates *DEF1*, *CEC1*, and *GAM1* (Alkhaibari et al., 2018). Infection of adult *Ae. aegypti* with either *B. bassiana* or *C. javanica* upregulated *defensin A* and *C*, as well as *cecropin D* and *G* (Ramirez et al., 2019). Synthetic *An. gambiae* CEC1 exhibited antifungal activity against the yeasts *S. cerevisiae*, *Cryptococcus neoformans*, and *C. albicans*, and filamentous fungi belonging to *Aspergillus* and *Fusarium* genera (Delucca et al., 1997; Vizioli et al., 2000). Similarly, CEC1, purified from *Ae. aegypti* hemolymph exhibited antifungal activity against *C. albicans*, *C. neoformans*, *S. cerevisiae*, *Fusarium culmorum*, and *Neurospora crassa* (Lamberty et al., 2001; Lowenberger et al., 1999). DEF, isolated from female *An. gambiae* midgut tissues, inhibited hyphal growth of the filamentous fungi *Botrytis cinerea*, *F. oxysporum*, *F. culmorum*, and *N. crassa* (Vizioli et al., 2001b). GAM1, purified from *An. gambiae* 4a-3B cell conditioned media inhibited hyphal growth of *N. crassa* (Vizioli et al., 2001a). Beyond known AMPs, few studies have specifically searched for antifungal peptides (AFPs) in insects (Al Souhail et al., 2016; Faruck et al., 2016), and none have been conducted in mosquitoes.

3.2.3 Melanization—In addition to AMP expression, the arthropod-specific melanization immune response is commonly observed against fungi (Fig. 3). Aromatic amino acids are converted to eumelanin, which is deposited on surfaces that are recognized by the immune system as foreign. This acellular encapsulation is thought to kill pathogens through site-directed cytotoxicity and/or by blocking nutrient uptake (Nappi and Christensen, 2005; Nappi and Vass, 1993). A key enzyme in melanization is prophenoloxidase (PPO), which is expressed from nine to ten paralogous genes in all mosquito genomes annotated to date (Bartholomay et al., 2010; Neafsey et al., 2015; Waterhouse et al., 2007). Infection of *An. gambiae* larvae with *L. giganteum* triggered a heavy melanization response of hyphae two hours post-infection, while *Ae. aegypti* larvae had a lower melanization response of less than 10% of hyphae (Golkar et al., 1993). Melanization in *Ae. aegypti* larvae was observed at the entry points of *L. chapmanii* zoospores and along the hyphal growth into the coelomic cavity (Zattau and McInnis, 1987). Yeast cells, deposited into the hemocoel of mosquitoes trigger an efficient melanization response. *S. cerevisiae*, injected into the hemocoel of *An. albimanus* is unable to grow and establish an infection in the hemolymph, due to the melanization of the yeast cells (Hernández-Martínez et al., 2002). Likewise, *C. albicans* is efficiently melanized and killed in the hemolymph of *Cx. quinquefasciatus* (Da Silva et al., 2000). Melanization in response to fungal entomopathogen infection has been observed in larvae of multiple mosquito species. *Anopheles amicus*, *Anopheles annulipes*, *Cx. fatigans*, and *Ae. tarsalis* larvae form melanotic capsules around *Culicinomyces* at all sites of infection (Sweeney, 1975). The fungus is encapsulated on the cuticle of the larval foregut and hindgut, the fungal penetration

sites between gut and hemolymph, and inside the hemolymph. Cuticular pigmentation and darkening of the head capsule was observed in early larval instars of *An. stephensi* when exposed to *B. bassiana* (Prasad and Veerwal, 2010). The melanization response in *An. gambiae* confers partial resistance to *B. bassiana* infection, as fungal load and mosquito death rate increased when the melanization response was experimentally decreased (Yassine et al., 2012). The resistance to fungal infection was in part mediated by TEPI, as its depletion from mosquito hemolymph resulted in less deposition of phenoloxidase on the surface of *B. bassiana* hyphal bodies. This study thus provided first molecular insight into mosquito antifungal immunity beyond AFPs, providing additional evidence that TEPI functions as an opsonin for melanization (Yassine et al., 2012).

3.3 Regulation of antifungal immunity through signal transduction pathways

—Several signal transduction pathways contribute to mosquito immunity, including Toll, IMD, JAK-STAT, JNK, and p38 signal transduction pathways (Fig. 3, Bian et al., 2005; Frolet et al., 2006; Meister et al., 2009, 2005; Shin et al., 2005). Below we briefly summarize current knowledge of these signal transduction pathways with regards to their (i) activation by fungal infections, (ii) control of antifungal molecule expression, and (iii) impact on fungal infection outcome.

3.3.1 Toll Pathway—The Toll pathway is a major immune signaling pathway in insects, whose role in antifungal immunity is well documented (Lemaitre et al., 1996). Named after its transmembrane receptor Toll, it consists of an extracellular protease cascade that activates the Toll ligand Spatzle, and an intracellular signal transduction cascade that translocates an NF- κ B transcription factor into the nucleus (Hoffmann and Reichhart, 2002; Lemaitre, 2004; Rhodes and Michel, 2017; Valanne et al., 2011). In mosquitoes, the NF- κ B transcription factor downstream of the Toll pathway is REL1 (Barillas-Mury et al., 1996; Meister et al., 2005; Shin et al., 2005). *B. bassiana* infection activates the Toll pathway in *Ae. aegypti* and *An. gambiae* adult mosquitoes (Dong et al., 2012; Rhodes et al., 2018; Shin et al., 2006). *Cordyceps javanica* and *Beauveria brongniartii* also activate the Toll pathway in *Ae. aegypti* adult mosquitoes (Ramirez et al., 2018a). Manipulation of the pathway in mosquitoes strongly influences fungal infection outcome. In *Ae. aegypti*, knockdown of *Toll5A* and *Spz1C* decreased survival in *B. bassiana*-infected mosquitoes (Shin et al., 2006). Similarly, *Rel1* knockdown in *Ae. aegypti* and *An. gambiae* mosquitoes decreased mosquito survival to infection by *B. bassiana* (Bian et al., 2005; Rhodes et al., 2018; Shin et al., 2005). Knockdown of *Cactus*, a negative regulator of the Toll pathway increased survival of *An. gambiae* mosquito to infections with *B. bassiana* (Rhodes et al., 2018). The Toll pathway regulates the expression of several known AMPs with antifungal activity, including *DEF1*, *CEC1*, and *GAM1* (Barillas-Mury et al., 1996; Garver et al., 2009; Luna et al., 2006; Shin et al., 2005; Zhang et al., 2017). In addition, the Toll pathway regulates the basal expression levels of TEPI (Frolet et al., 2006). As the Toll pathway controls about 10% of the mosquito transcriptome, its influence on fungal infection likely goes beyond the regulation of these few antifungal factors.

3.3.2 IMD Pathway—The IMD pathway is a major immune signal transduction pathway in the gut and regulates malaria parasite killing in mosquitoes (Antonova et al., 2009; Chen

et al., 2012; Garver et al., 2009, 2012; Meister et al., 2005, 2009). Its downstream NF- κ B transcription factor in mosquitoes is called REL2 (Fig. 3, Antonova et al., 2009; Meister et al., 2005). Infection of *Ae. Aegypti* mosquitoes with *B. bassiana*, *B. brongniartii*, and *C. javanica* upregulated *Rel2* transcript levels in the midgut and fat body six days post-infection (Ramirez et al., 2018a; Ramirez et al., 2019), suggesting the activation of the IMD pathway upon fungal challenge.

The IMD pathway confers resistance to fungal infections in mosquitoes. Knockdown of *Rel2* in *B. bassiana* and *C. javanica*-infected *Ae. aegypti* adult mosquitoes decreased mosquito survival to infection and increased fungal load in infected mosquitoes (Ramirez et al., 2018b). Similar to the Toll pathway, the IMD pathway regulates the expression of the *DEF1*, *CEC1*, and *GAM1* (Antonova et al., 2009; Garver et al., 2009; Luna et al., 2006; Meister et al., 2005, 2009; Ramirez et al., 2018b; Zhang et al., 2017). Altered expression of these antifungal molecules may contribute to the antifungal resistance that is exerted through the IMD pathway.

3.3.3 JAK-STAT Pathway—In mosquitoes, the JAK-STAT pathway is involved in antiviral and anti-*Plasmodium* immunity (Carissimo et al., 2015; Gupta et al., 2009; Souza-Neto et al., 2009). The pathway is activated by ligand-binding to the domeless (Dome) receptor, and ultimately translocates the transcription factor Stat92E to the nucleus (Fig. 3, Zeidler and Bausek, 2013). Infection of adult *Ae. aegypti* with *B. bassiana* upregulated the expression of *Dome* and the STAT-regulated anti-dengue restriction factor 1 (*DVRF1*). Similarly, the expression of *STAT* was upregulated in *Ae. aegypti* mosquitoes infected by *B. bassiana*, *Be. brongniartii*, and *Isaria amoenerosea* (Ramirez et al., 2018b, 2018c). Knockdown of *Dome* or *DVRF1* decreased mosquito survival to *B. bassiana* infection (Dong et al., 2012). In *Ae. aegypti* cells, the JAK-STAT pathway upregulates *GAM1* expression in response to *C. albicans* exposure (Zhang et al., 2017), which may contribute to the JAK-STAT pathway-mediated antifungal immunity in mosquitoes.

3.3.3 MAPK Pathways—Less is known about the role of the JNK and p38 MAP-kinase signaling pathways in mosquito immunity including responses to fungal infection. In *Ae. aegypti* mosquitoes, infection with *C. albicans* upregulates MAP kinase kinase 4 (MAPK4), which is upstream to JNK and p38 kinases (Wu and Cho, 2014). Likewise, infection with *Be. brongniartii* upregulates the expression of *JNK* in the midgut and fat body (Ramirez et al., 2018a). It is unclear whether either of these pathways contribute to antifungal immunity in mosquitoes. However, in *Drosophila*, *p38* mutant flies exhibited increased susceptibility to infection by *B. bassiana* and *Aspergillus fumigatus* (Chen et al., 2010).

3.4 Regulation of antifungal immunity through protease cascades—In addition to transcriptional regulation, several mosquito humoral immune responses are regulated through proteolytic activation of key immune factors and enzymes, including TEPI and PPO (Nakhleh et al., 2017; Rhodes and Michel, 2017). Proteolytic activity is regulated through cascades of Clip-domain containing serine proteinases (CLIPs) and their proteolytically inactive homologs (clip-SPHs, most often CLIPAs). TEPI function is augmented by two clip-SPHs, SPCLIP1 and CLIPA2. SPCLIP1 is essential for TEPI deposition on the surface of microbes and acts as a positive regulator of the complement-like pathway (Povelones et

al., 2013). CLIPA2 acts as a negative regulator through inhibiting the production of active TEPI during infection (Yassine et al., 2014). Knockdown of CLIPA2 decreased susceptibility of *An. gambiae* to *B. bassiana* in a TEPI-dependent manner, confirming the role of TEPI in antifungal immunity (Kamareddine et al., 2016).

An. gambiae CLIPA8 is a positive regulator of phenoloxidase activation and required for melanization in response to *P. berghei* parasites, bacteria, and *B. bassiana* (Schnitger et al., 2007; Yassine et al., 2012). Knockdown of CLIPA8 abolished the melanization of *B. bassiana* hyphae, and decreased resistance to *B. bassiana* infection (Yassine et al., 2012). While several CLIPBs are required for PO activation in mosquitoes (An et al., 2011; Zhang et al., 2016), their role in antifungal immunity in mosquitoes is untested.

4. Summary and outlook

The mosquito immune system has evolved in the context of continuous encounters between mosquitoes and fungi ranging from food to foes. Many of these encounters indeed are beneficial to the mosquito. Yeast hold nutritional value for mosquito larvae (Asahina, 1964; Steyn et al., 2016), and may generate a hypoxic environment in the larval midgut required for mosquito development (Valzania et al., 2018). Encounters with potential opportunistic fungal pathogens may remain benign for the mosquito due to physical and physiological barriers, including the cuticle and digestive enzymes (Gillespie et al., 2000). However, an significant number of fungal entomopathogens can overcome barriers to infection and colonize the mosquito hemocoel with often lethal consequences to the host. It is thus not surprising to find all branches of the mosquito immune system to be engaged in antifungal immunity. The current body of literature describes some of these interactions on a histological, and to a much lesser extent molecular level.

Important knowledge gaps remain. (i) Out of the many potential mosquito-fungal encounters, little is known of the mosquito immune system's role in fungal interactions beyond defense against a small number of well-studied entomopathogens in adult infections. Comparative studies using multiple mosquito and fungal species combinations would be valuable to address whether a core of anti-fungal immunity mechanisms are engaged across mosquito species, and what biotic and abiotic factors influence the efficacy of these immune responses. Mosquito larval immunity is largely unexplored, and its contribution to antifungal defense is unclear. It is likely that exposure of the larval immune system impacts adult immunity. This is important to explore, as it could impact vector-borne pathogen susceptibility and transmission rates.

(ii) The molecular identity of antifungal molecules in both mosquito larvae and adults is largely unexplored. Naturally occurring AFPs constitute a numerous and structurally highly diverse group of peptides (van der Weerden et al., 2013), and it is likely that mosquitoes produce antifungals beyond canonical AMPs. Beyond knowledge gain, mosquito AFPs could have practical applications, including their use as biomarkers of exposure, and natural product fungicides (Rautenbach et al., 2016; van der Weerden et al., 2013).

(iii) During colonization, entomopathogenic fungi clearly interact with mosquito epithelia, including the epidermis, tracheal linings, and the gut epithelium. However, the role of epithelial immunity in limiting fungal colonization is unknown. Mosquito epithelial immune responses contribute to maintenance of a healthy microbiota and defense against vector-borne disease pathogens, and likely also limit fungal infections through similar mechanisms.

(iv) In laboratory studies, boosting mosquito basal immunity reduces fungal entomopathogen infections (Rhodes et al., 2018; Yassine et al., 2012). Different strains and species of fungal entomopathogens differ in their virulence against the same mosquito species (Blanford et al., 2012). Entomopathogen virulence is also influenced by abiotic factors, such as temperature (Heinig et al., 2015). However, in contrast to field and laboratory studies on variation in host susceptibility to vector-borne pathogens (Collins et al., 1986; Gubler, 1979; Gubler and Rosen, 1976; Huff, 1929, 1927; Niaré et al., 2002; Wallis et al., 1985), mosquito host genotype influence on fungal entomopathogen infection is, to our knowledge, unexplored. Such studies may help to identify non-canonical antifungal immune responses in mosquitoes. Such knowledge could further aid biocontrol programs to select for strains that even more readily overcome mosquito immunity.

(v) Our knowledge of mosquito-fungus interactions is mostly confined to entomopathogens, due to their potential for biological control of mosquitoes. In contrast to bacterial microbiomes, investigations of naturally occurring fungal associations with mosquito larvae and adults are just beginning. So far, a handful of studies have identified yeasts and filamentous fungi associated with mosquitoes. Future studies should explore whether the transient nature of these associations impact mosquito physiology, and shape the immune status of mosquitoes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Carol Sevin, Research librarian at Kansas State University, for helping us to retrieve literature on mosquito interactions with *B. bassiana* and *M. anisopliae*. Dr. K.M. is partially supported by the National Institutes of Health grant number R01AI140760, and P.T. is supported through the USDA-ARS Specific Cooperative Agreement 58-5430-4-022. In addition, this work was supported by the USDA National Institute of Food and Agriculture, Hatch project 1003627. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the funding agencies. This is contribution 19-190-J from the Kansas Agricultural Experiment Station.

References

- Aguilar R, Jedlicka AE, Mintz M, Mahairaki V, Scott AL, Dimopoulos G, 2005 Global gene expression analysis of *Anopheles gambiae* responses to microbial challenge. *Insect Biochemistry and Molecular Biology*, Genetic manipulation of insects 35, 709–719.
- Al Souhail Q, Hiromasa Y, Rahnamaeian M, Giraldo MC, Takahashi D, Valent B, Vilcinskas A, Kanost MR, 2016 Characterization and regulation of expression of an antifungal peptide from hemolymph of an insect, *Manduca sexta*. *Dev. Comp. Immunol*
- Alkhaibari AM, Maffei T, Bull JC, Butt TM, 2018 Combined use of the entomopathogenic fungus, *Metarhizium brunneum*, and the mosquito predator, *Toxorhynchites brevipalpis*, for control of

mosquito larvae: Is this a risky biocontrol strategy? *Journal of Invertebrate Pathology* 153, 38–50. [PubMed: 29425967]

- An C, Budd A, Kanost MR, Michel K, 2011 Characterization of a regulatory unit that controls melanization and affects longevity of mosquitoes. *Cellular and Molecular Life Sciences* 68, 1929–1939. [PubMed: 20953892]
- Antonova Y, Alvarez KS, Kim YJ, Kokoza V, Raikhel AS, 2009 The role of NF- κ B factor REL2 in the *Aedes aegypti* immune response. *Insect Biochemistry and Molecular Biology* 39, 303–314. [PubMed: 19552893]
- Asahina S, 1964 Food material and feeding procedures for mosquito larvae. *Bull World Health Organ* 31, 465–466. [PubMed: 14272445]
- Araújo JPM, Hughes DP, 2014 Diversity of entomopathogens Fungi: Which groups conquered the insect body?
- Barillas-Mury C, Charlesworth A, Gross I, Richman A, Hoffmann JA, Kafatos FC, 1996 Immune factor Gambif1, a new Rel family member from the human malaria vector, *Anopheles gambiae*. *Embo J* 15, 4691–4701. [PubMed: 8887560]
- Bartholomay LC, Michel K, 2018 Mosquito Immunobiology: The intersection of vector health and vector competence. *Annual Review of Entomology* 63, 145–167.
- Bartholomay LC, Waterhouse RM, Mayhew GF, Campbell CL, Michel K, Zou Z, Ramirez JL, Das S, Alvarez K, Arensburger P, Bryant B, Chapman SB, Dong Y, Erickson SM, Karunaratne SHPP, Kokoza V, Kodira CD, Pignatelli P, Shin SW, Vanlandingham DL, Atkinson PW, Birren B, Christophides GK, Clem RJ, Hemingway J, Higgs S, Megy K, Ranson H, Zdobnov EM, Raikhel AS, Christensen BM, Dimopoulos G, Muskavitch MAT, 2010 Pathogenomics of *Culex quinquefasciatus* and meta-analysis of infection responses to diverse pathogens. *Science* 330, 88–90. [PubMed: 20929811]
- Bawin T, Seye F, Boukraa S, Zimmer J-Y, Raharimalala FN, Ndiaye M, Compere P, Delvigne F, Francis F, 2016 Histopathological effects of *Aspergillus clavatus* (Ascomycota: Trichocomaceae) on larvae of the southern house mosquito, *Culex quinquefasciatus* (Diptera: Culicidae). *Fungal Biology* 120, 489–499. [PubMed: 27020151]
- Benzina F, Hamid S, Mohand-Kaci H, Bissaad F, Halouane F, 2018 Histological changes in the larvae of the domestic mosquito *Culex pipiens* treated with the entomopathogenic fungus *Beauveria bassiana*. *Scientific Research and Essays* 13, 1–10.
- Bian G, Shin SW, Cheon H-M, Kokoza V, Raikhel AS, 2005 Transgenic alteration of Toll immune pathway in the female mosquito *Aedes aegypti*. *Proceedings of the National Academy of Sciences* 102, 13568–13573.
- Bille J, Stockman L, Roberts GD, 1982 Detection of yeasts and filamentous fungi in blood cultures during a 10-year period (1972 to 1981). *Journal of clinical microbiology* 16, 968–970. [PubMed: 7153348]
- Blandin S, Levashina EA, 2004 Thioester-containing proteins and insect immunity. *Molecular Immunology, Innate Immunity* 40, 903–908.
- Blandin S, Shiao S-H, Moita LF, Janse CJ, Waters AP, Kafatos FC, Levashina EA, 2004 Complement-like protein TEPI is a determinant of vectorial capacity in the malaria vector *Anopheles gambiae*. *Cell* 116, 661–670. [PubMed: 15006349]
- Blandin SA, Marois E, Levashina EA, 2008 Antimalarial responses in *Anopheles gambiae*: from a complement-like protein to a complement-like pathway. *Cell Host & Microbe* 3, 364–374. [PubMed: 18541213]
- Blanford S, Jenkins NE, Read AF, Thomas MB, 2012 Evaluating the lethal and pre-lethal effects of a range of fungi against adult *Anopheles stephensi* mosquitoes. *Malaria Journal* 11, 365. [PubMed: 23126549]
- Boman HG, 2003 Antibacterial peptides: basic facts and emerging concepts. *J. Intern. Med* 254, 197–215. [PubMed: 12930229]
- Bozic J, Capone A, Pediconi D, Mensah P, Cappelli A, Valzano M, Mancini MV, Scuppa P, Martin E, Epis S, Rossi P, Favia G, Ricci I, 2017 Mosquitoes can harbour yeasts of clinical significance and contribute to their environmental dissemination. *Environ Microbiol Rep* 9, 642–648. [PubMed: 28714286]

- Braun A, 1855 Algarum unicellularium genera nova et minus cognita: praemissis observationibus de algis unicellularibus in genere. Engelmann.
- Brey PT, Lebrun RA, Papierok B, Ohayon H, Vennavalli S, Hafez J, 1988 Defense reactions by larvae of *Aedes aegypti* during infection by the aquatic fungus *Lagenidium giganteum* (Oomycete). *Cell Tissue Res.* 253, 245–250. [PubMed: 3416342]
- Butt TM, Greenfield BPJ, Greig C, Maffei TGG, Taylor JWD, Piasecka J, Dudley E, Abdulla A, Dubovskiy IM, Garrido-Jurado I, Quesada-Moraga E, Penny MW, Eastwood DC, 2013 *Metarhizium anisopliae* pathogenesis of mosquito larvae: a verdict of accidental death. *PLoS ONE* 8, e81686. [PubMed: 24349111]
- Butt TM, Coates CJ, Dubovskiy IM, Ratcliffe NA, 2016 Entomopathogenic fungi: new insights into host-pathogen interactions. *Adv. Genet* 94, 307–364. [PubMed: 27131329]
- Carissimo G, Pondeville E, McFarlane M, Dietrich I, Mitri C, Bischoff E, Antoniewski C, Bourgoquin C, Failloux A-B, Kohl A, Vernick KD, 2015 Antiviral immunity of *Anopheles gambiae* is highly compartmentalized, with distinct roles for RNA interference and gut microbiota. *Proceedings of the National Academy of Sciences* 112, E176–E185.
- Castillo JM, Roberts DW, 1980 *Coelomomyces* pathogens of Culicidae (mosquitos). *Bull World Health Organ* 58, 53–67. [PubMed: 20604461]
- Castillo JM, Roberts DW, 1980 Fungal pathogens, except *Coelomomyces*, of Culicidae (mosquitos). *Bull. World Health Organ.* 58 Suppl, 69–83. [PubMed: 20604462]
- Chang HC, Leaw SN, Huang AH, Wu TL, Chang TC, 2001 Rapid identification of yeasts in positive blood cultures by a multiplex PCR method. *Journal of Clinical Microbiology* 39, 3466–3471. [PubMed: 11574557]
- Chappell CR, Fukami T, 2018 Nectar yeasts: a natural microcosm for ecology. *Yeast* 35, 417–423. [PubMed: 29476620]
- Chen J, Xie C, Tian L, Hong L, Wu X, Han J, 2010 Participation of the p38 pathway in *Drosophila* host defense against pathogenic bacteria and fungi. *Proc Natl Acad Sci U S A* 107, 20774–20779. [PubMed: 21076039]
- Chen Y, Dong Y, Sandiford S, Dimopoulos G, 2012 Transcriptional mediators Kto and Skd are involved in the regulation of the IMD pathway and anti-*Plasmodium* defense in *Anopheles gambiae*. *PLOS ONE* 7, e45580. [PubMed: 23049816]
- Chang N-T, Liu S-D, 1990 Virulence of entomogenous fungus *Metarhizium anisopliae* var. *anisopliae* to *Aedes aegypti* and *Aedes albopictus* larvae. *Chinese Journal of Microbiology and Immunology* 23, 253–257. [PubMed: 2091906]
- Christophers SR, 1952 The recorded parasites of mosquitoes. *Rivista di parassitologia* 13, 21–28.
- Christophides GK, Zdobnov E, Barillas-Mury C, Birney E, Blandin S, Blass C, Brey PT, Collins FH, Danielli A, Dimopoulos G, Hetru C, Hoa NT, Hoffmann JA, Kanzok SM, Letunic I, Levashina EA, Loukeris TG, Lycett G, Meister S, Michel K, Moita LF, Muller H-M, Osta MA, Paskewitz SM, Reichhart J-M, Rzhetsky A, Troxler L, Vernick KD, Vlachou D, Volz J, von Mering C, Xu J, Zheng L, Bork P, Kafatos FC, 2002 Immunity-related genes and gene families in *Anopheles gambiae*. *Science* 298, 159–165. [PubMed: 12364793]
- Clark TB, Kellen WR, Fukuda T, Lindegren JE, 1968 Field and laboratory studies on the pathogenicity of the fungus *Beauveria bassiana* to three genera of mosquitoes. *J. Invertebr. Pathol* 11, 1–7. [PubMed: 5654770]
- Clark TB, Kellen WR, Lindegren JE, Sanders RD, 1966 *Pythium* sp. (Phycomycetes: Pythiales) pathogenic to mosquito larvae. *Journal of Invertebrate Pathology* 8, 351–354.
- Collins FH, Sakai RK, Vernick KD, Paskewitz S, Seeley DC, Miller LH, Collins WE, Campbell CC, Gwadz RW, 1986 Genetic selection of a Plasmodium-refractory strain of the malaria vector *An. gambiae*. *Science* 234, 607–610. [PubMed: 3532325]
- Coluzzi M, 1966 Experimental infections with *Rubetella* fungi in *Anopheles gambiae* and other mosquitoes. in: Corradetti A (Ed.), *Proceedings of the First International Congress of Parasitology Pergamon*, pp. 592–593.
- Couch JN, 1935 A New Saprophytic species of *Lagenidium*, with notes on other forms. *Mycologia* 27, 376–387.
- Couch JN, Umphlett CJ, 1963 *Coelomomyces* Infections, in: *Insect Pathology*. Elsevier, pp. 149–188.

- Crisan EV, 1971 Mechanism responsible for release of toxin by *Metarhizium* spores in mosquito larvae. *Journal of Invertebrate Pathology* 17, 260–264. [PubMed: 4396947]
- Da Silva JB, De Albuquerque CMR, De Araujo EC, Peixoto CA, Hurd H, 2000 Immune defense mechanisms of *Culex quinquefasciatus* (Diptera: Culicidae) against *Candida albicans* infection. *Journal of Invertebrate Pathology* 76, 257–262. [PubMed: 11112370]
- DeLucca AJ, Bland JM, Jacks TJ, Grimm C, Cleveland TE, & Walsh TJ, 1997 Fungicidal activity of cecropin A. *Antimicrobial agents and chemotherap.* 41, 481–483.
- Díaz-Nieto LM, D'Alessio C, Perotti MA, Beron CM, 2016 *Culex pipiens* development is greatly influenced by native bacteria and exogenous yeast. *PLOS ONE* 11, e0153133. [PubMed: 27055276]
- Dikgolz VE, Toledo AV, Topa PE, Lopez Lastra CC, 2005 Evaluation of histological techniques for the detection of fungal infections caused by *Leptolegnia chapmanii* (Oomycetes: Saprolegniales) in *Aedes aegypti* (Diptera: Culicidae) larvae. *Folia Microbiologica* 50, 125–127. [PubMed: 16110916]
- Dong Y, Morton JC, Ramirez JL, Souza-Neto JA, Dimopoulos G, 2012 The entomopathogenic fungus *Beauveria bassiana* activate toll and JAK-STAT pathway-controlled effector genes and anti-dengue activity in *Aedes aegypti*. *Insect Biochem. Mol. Biol.* 42, 126–132. [PubMed: 22198333]
- Duguma D, Hall MW, Rugman-Jones P, Stouthamer R, Terenius O, Neufeld JD, Walton WE, 2015 Developmental succession of the microbiome of *Culex* mosquitoes. *BMC Microbiology* 15, 140. [PubMed: 26205080]
- Ekstein F, 1922 Beitrage zur Erkenntnis der Stechmueckenparasiten. *Centralblatt fUr Bakteriologie, Parasitenkunde und Infektionskrankheiten.* 1. Abt., Originale. 88, 128–135.
- Eneh LK, Saijo H, Borg-Karlson A-K, Lindh JM, Rajarao GK, 2016 Cedrol, a malaria mosquito oviposition attractant is produced by fungi isolated from rhizomes of the grass *Cyperus rotundus*. *Malaria Journal* 15, 478. [PubMed: 27639972]
- Farenhorst M, Hilhorst A, Thomas MB, Knols BGJ, 2011 Development of fungal applications on netting substrates for malaria vector C=control. *J Med Entomol* 48, 305–313. [PubMed: 21485366]
- Faruck MO, Yusof F, Chowdhury S, 2016 An overview of antifungal peptides derived from insect. *Peptides, Invertebrate Neuropeptides XVI* 80, 80–88.
- Fei M, Huai-en B, 2001 Observation on hemocytes of *Culex pipiens quinquefasciatus* larvae infected by *Lagenidium giganteum*. *Zhongguo ji sheng chong xue yu ji sheng chong bing za zhi= Chinese journal of parasitology & parasitic diseases* 19, 330–332. [PubMed: 12572061]
- Feng MG, Poprawski TJ, Khachatourians GG, 1994 Production, formulation and application of the entomopathogenic fungus *Beauveria bassiana* for insect control: current status. *Biocontrol Science and Technology* 4, 3–34.
- Fetter-Lasko JL, Washino RK, 1983 In Situ Studies on Seasonality and Recycling Pattern in California of *Lagenidium giganteum* Couch, an Aquatic Fungal Pathogen of Mosquitoes. *Environ Entomol* 12, 635–640.
- Fish D, Carpenter SR, 1982 Leaf Litter and Larval Mosquito Dynamics in Tree-Hole Ecosystems. *Ecology* 63, 283–288.
- Foster WA, 1995 Mosquito sugar feeding and reproductive energetics. *Annual Review of Entomology* 40, 443–474.
- Frohlich-Nowoisky J, Pickersgill DA, Despres VR, Poschl U, 2009 High diversity of fungi in air particulate matter. *Proceedings of the National Academy of Sciences* 106, 12814–12819.
- Frolet C, Thoma M, Blandin S, Hoffmann JA, Levashina EA, 2006 Boosting NF- κ B-Dependent basal immunity of *Anopheles gambiae* aborts development of *Plasmodium berghei*. *Immunity* 25, 677–685. [PubMed: 17045818]
- Galli-Valerio B, Rochaz de Jongh J, 1906 Uber die Wirkung von *Aspergillus niger* und *A. glaucus* auf die Larven von *Culex* und *Anopheles*. 2, Mitteilung. *Centralblatt fur Bakteriologie, Parasitenkunde und Infektionskrankheiten.* 1. Abt, Originale. 40, 630–633.
- García Gil de Muñoz FL, Martínez-Barnette J, Lanz-Mendoza H, Rodríguez MH, Hernández-Hernández FC, 2008 Prostaglandin E2 modulates the expression of antimicrobial peptides in the

fat body and midgut of *Anopheles albimanus*. Archives of Insect Biochemistry and Physiology 68, 14–25. [PubMed: 18412259]

- García-Munguía AM, Garza-Hernández JA, Rebollar-Tellez EA, Rodríguez-Pérez MA, Reyes-Villanueva F, 2011 Transmission of *Beauveria bassiana* from male to female *Aedes aegypti* mosquitoes. Parasites & Vectors 4, 24. [PubMed: 21352560]
- Garland CD, Pillai JS, 1979 *Coelomomyces opifexi* Pillai & Smith (Coelomomycetaceae: Blastocladales) V. Development of the fungus in pupae and adults of *Aedes australis*. New Zealand Journal of Zoology 6, 279–283.
- Garver LS, Bahia AC, Das S, Souza-Neto JA, Shiao J, Dong Y, Dimopoulos G, 2012 *Anopheles* Imd pathway factors and effectors in infection intensity-dependent anti-*Plasmodium* action. PLoS Pathogens 8, e1002737. [PubMed: 22685401]
- Garver LS, Dong Y, Dimopoulos G, 2009 Caspar Controls Resistance to *Plasmodium falciparum* in Diverse Anopheline Species. PLoS Pathogens 5, e1000335. [PubMed: 19282971]
- Garza-Hernández JA, Reyes-Villanueva F, Russell TL, Braks MAH, Garcia-Munguía AM, Rodríguez-Pérez MA, 2015 Copulation Activity, Sperm Production and Conidia Transfer in *Aedes aegypti* Males Contaminated by *Metarhizium anisopliae*: A Biological Control Prospect. PLOS Neglected Tropical Diseases 9, e0004144. [PubMed: 26473490]
- Geetha I, Paily KP, Padmanaban V, Balaraman K, 2003 Oviposition response of the mosquito, *Culex quinquefasciatus* to the secondary metabolite(s) of the fungus, *Trichoderma viride*. Mem. Inst. Oswaldo Cruz 98, 223–226. [PubMed: 12764438]
- Gillespie JP, Bailey AM, Cobb B, Vilcinskas A, 2000 Fungi as elicitors of insect immune responses. Archives of Insect Biochemistry and Physiology 44, 49–68. [PubMed: 10861866]
- Goh TK, Hyde KD, 1996 Biodiversity of freshwater fungi. Journal of Industrial Microbiology & Biotechnology 17, 328–345.
- Golkar L, Lebrun RA, Ohayon H, Gounon P, Papierok B, Brey PT, 1993 Variation of larval susceptibility to *Lagenidium giganteum* in three mosquito species. Journal of Invertebrate Pathology 62, 1–8. [PubMed: 8104999]
- Greenfield Bethany P. J., Lord Alex M., Dudley Ed, Butt Tariq M., 2014 Conidia of the insect pathogenic fungus, *Metarhizium anisopliae*, fail to adhere to mosquito larval cuticle. Royal Society Open Science 1, 140193. [PubMed: 26064542]
- Gubler DJ, 1979 Variation in susceptibility to oral infection with dengue viruses among geographic strains of *Aedes aegypti*. Am. J. Trop. Med. Hyg 28, 1045–1052. [PubMed: 507282]
- Gubler DJ, Rosen L, 1976 Variation among geographic strains of *Aedes Albopictus* in susceptibility to infection with Dengue viruses. The American Journal of Tropical Medicine and Hygiene 25, 318–325. [PubMed: 1259091]
- Guegan M, Zouache K, Demichel C, Minard G, Tran Van V, Potier P, Mavingui P, Valiente Moro C, 2018 The mosquito holobiont: fresh insight into mosquito-microbiota interactions. Microbiome 6, 49. [PubMed: 29554951]
- Gupta L, Molina-Cruz A, Kumar S, Rodrigues J, Dixit R, Zamora RE, Barillas-Mury C, 2009 The STAT pathway mediates late-phase immunity against *Plasmodium* in the mosquito *Anopheles gambiae*. Cell Host & Microbe 5, 498–507. [PubMed: 19454353]
- Hasan S, Vago C, 1972 The pathogenicity of *Fusarium oxysporum* to mosquito larvae. Journal of Invertebrate Pathology 20, 268–271. [PubMed: 4655538]
- Heinig RL, Paaijmans KP, Hancock PA, Thomas MB, 2015 The potential for fungal biopesticides to reduce malaria transmission under diverse environmental conditions. J Appl Ecol 52, 1558–1566. [PubMed: 26792946]
- Hernández-Martínez S, Lanz H, Rodríguez MH, González-Ceron L, Tsutsumi V, 2002 Cellular-mediated reactions to foreign organisms inoculated into the hemocoel of *Anopheles albimanus* (Diptera: Culicidae). J Med Entomol 39, 61–69. [PubMed: 11931273]
- Hillyer JF, Strand MR, 2014 Mosquito hemocyte-mediated immune responses. Current Opinion in Insect Science, Vectors and medical and veterinary entomology 3, 14–21.
- Hoffmann JA, Reichhart J-M, 2002 *Drosophila* innate immunity: an evolutionary perspective. Nature Immunology 3, 121–126. [PubMed: 11812988]

- Huff CG, 1929 The effects of selection upon susceptibility to bird malaria in *Culex pipiens* Linn. *Annals of Tropical Medicine and Parasitology* 23, 427–442.
- Huff CGG, 1927 Studies on the infectivity of plasmodia of birds for mosquitoes, with special reference to the problem of immunity in the mosquito. *The American Journal of Hygiene* 7, 706–734.
- Hung S-Y, Boucias DG, 1992 Influence of *Beauveria bassiana* on the cellular defense response of the beet armyworm, *Spodoptera exigua*. *Journal of Invertebrate Pathology* 60, 152–158.
- Hung S-Y, Boucias DG, Vey AJ, 1993 Effect of *Beauveria bassiana* and *Candida albicans* on the Cellular Defense Response of *Spodoptera exigua*. *Journal of Invertebrate Pathology* 61, 179–187. [PubMed: 8463710]
- Impoinvil DE, Kongere JO, Foster WA, Njiru BN, Killeen GF, Githure JI, Beier JC, Hassanali A, Knols BGJ, 2004 Feeding and survival of the malaria vector *Anopheles gambiae* on plants growing in Kenya. *Medical and Veterinary Entomology* 18, 108–115. [PubMed: 15189235]
- Ishii M, Kanuka H, Badolo A, Sagnon N, Guelbeogo WM, Koike M, Aiuchi D, 2017 Proboscis infection route of *Beauveria bassiana* triggers early death of *Anopheles* mosquito. *Scientific Reports* 7, 3476. [PubMed: 28615623]
- Jenkins DW, 1964 Pathogens, parasites and predators of medically important arthropods. annotated list and bibliography. *Bulletin of the World Health Organization* 30, 5–150.
- Jiang H, Ma C, Lu Z-Q, Kanost MR, 2004 β -1,3-Glucan recognition protein-2 (β GRP-2) from *Manduca sexta*: an acute-phase protein that binds β -1,3-glucan and lipoteichoic acid to aggregate fungi and bacteria and stimulate prophenoloxidase activation. *Insect Biochemistry and Molecular Biology* 34, 89–100. [PubMed: 14976985]
- Jiravanichpaisal P, Lee BL, Soderhall K, 2006 Cell-mediated immunity in arthropods: Hematopoiesis, coagulation, melanization and opsonization. *Immunobiology* 211, 213–236. [PubMed: 16697916]
- Kamareddine L, Nakhleh J, Osta MA, 2016 Functional interaction between apolipoproteins and complement regulate the mosquito immune response to systemic infections. *Journal of Innate Immunity* 8, 314–326. [PubMed: 26950600]
- Kanzok SM, Jacobs-Lorena M, 2006 Entomopathogenic fungi as biological insecticides to control malaria. *Trends in Parasitology* 22, 49–51. [PubMed: 16377249]
- Kaufman MG, Chen S, Walker ED, 2008 Leaf-Associated Bacterial and Fungal Taxa Shifts in Response to Larvae of the Tree Hole Mosquito, *Ochlerotatus triseriatus*. *Microb Ecol* 55, 673–684. [PubMed: 17899246]
- Kawakami K, 1965 Phagocytosis in muscardine-diseased larvae of the silkworm, *Bombyx mori* (Linnaeus). *Journal of Invertebrate Pathology* 7, 203–208. [PubMed: 5828994]
- Kupriyanova ES, 1966 On an *Entomophthora* fungus parasitising mosquitoes of the complex of *Culex pipiens* L. *Zoologicheskii Zhurnal* 45.
- Lacey CM, Lacey LA, Roberts DR, 1988 Route of invasion and histopathology of *Metarhizium anisopliae* in *Culex quinquefasciatus*. *Journal of Invertebrate Pathology* 52, 108–118. [PubMed: 3418133]
- Laird M, Mogi M, Sota T, 1992 Northernmost occurrences of the protistan pathogen, *Coelomomyces stegomyiae* var. *stegomyiae*. *J Am Mosq Control Assoc* 8, 430–432. [PubMed: 1474392]
- Lakon G, 1919 Die Insektenfeinde aus der Familie der Entomophthoreen. *Zeitschrift für Angewandte Entomologie* 5, 161–216.
- Lamberty M, Zachary D, Lanot R, Bordereau C, Robert A, Hoffmann JA, Bulet P, 2001 Constitutive expression of a cysteine-rich antifungal and a linear antibacterial peptide in a termite insect. *J. Biol. Chem* 276, 4085–4092. [PubMed: 11053427]
- Lavine MD, Strand MR, 2002 Insect hemocytes and their role in immunity. *Insect Biochemistry and Molecular Biology, Recent Progress in Insect Molecular Biology* 32, 1295–1309.
- Lemaitre B, 2004b The road to Toll. *Nature Reviews Immunology* 4, 521–527.
- Lemaitre B, Nicolas E, Michaut L, Reichhart J-M, Hoffmann JA, 1996 The dorsoventral regulatory gene cassette *spätzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell* 86, 973–983. [PubMed: 8808632]
- Levashina EA, Moita LF, Blandin S, Vriend G, Lagueux M, Kafatos FC, 2001 Conserved role of a complement-like protein in phagocytosis revealed by dsRNA knockout in cultured cells of the mosquito, *Anopheles gambiae*. *Cell* 104, 709–718. [PubMed: 11257225]

- Levetin E, Dorsey K, 2006 Contribution of leaf surface fungi to the air spora. *Aerobiologia* 22, 3–12.
- Lopez Lastra CC, 1997 Primera cita de *Smittium culisetae* Y. *S. culicis* (Trichomycetes: Harpellales) en larvas de mosquitos (Diptera: Culicidae) de la republica Argentina. *Bol. Soc. Argent. Bot* 33, 43–46.
- López Lastra CC, Scorsetti AC, Marti GA, Garcia JJ, 2004 Host range and specificity of an Argentinean isolate of the aquatic fungus *Leptolegnia chapmanii* (Oomycetes: Saprolegniales), a pathogen of mosquito larvae (Diptera: Culicidae). *Mycopathologia* 158, 311–315. [PubMed: 15702269]
- Lord JC, Fukuda T, 1988 An ultrastructural study of the invasion of *Culex quinquefasciatus* larvae by *Leptolegnia chapmanii* (Oomycetes: Saprolegniales). *Mycopathologia* 104, 67–73. [PubMed: 3221914]
- Lovett B, Bilgo E, Millogo SA, Ouattarra AK, Sare I, Gnambani EJ, Dabire RK, Diabate A, St. Leger RJ, 2019 Transgenic *Metarhizium* rapidly kills mosquitoes in a malaria-endemic region of Burkina Faso. *Science* 364, 894–897. [PubMed: 31147521]
- Lovett B, Leger RJS, 2017 The Insect Pathogens. *Microbiology Spectrum* 5, 1–19.
- Lowe RE, Rumbaugh RG, Patterson RS, 1968 *Entomophthora coronata* as a pathogen of mosquitoes. *Journal of Invertebrate Pathology* 11.
- Lowe RE, Kennel EW, 1972 Pathogenicity of the fungus *Entomophthora coronata* *Culex pipiens quinquefasciatus* and *Aedes taeniorhynchus*. *Mosquito news*. 32, 614–620.
- Lowenberger C, Charlet M, Vizioli J, Kamal S, Richman A, Christensen BM, Bulet P, 1999 Antimicrobial activity spectrum, cDNA cloning, and mRNA expression of a newly isolated member of the Cecropin family from the mosquito vector *Aedes aegypti*. *Journal of Biological Chemistry* 274, 20092–20097. [PubMed: 10400619]
- Lu H-L, St. Leger RJ, 2016 Chapter Seven - Insect immunity to entomopathogenic fungi, in: Lovett B, St. Leger Raymond J. (Eds.), *Advances in Genetics, Genetics and Molecular Biology of Entomopathogenic Fungi*. Academic Press 94, 251–285.
- Lucarotti CJ, & Andreadis TG, 1995 Reproductive strategies and adaptations for survival among obligatory microsporidian and fungal parasites of mosquitoes: a comparative analysis of *Amblyospora* and *Coelomomyces*. *J Am Mosq Control. Assoc* 11, 111–121. [PubMed: 7616177]
- Lucarotti CJ, 1987 *Coelomomyces stegomyiae* Infection in Adult *Aedes aegypti*. *Mycologia* 79, 362–369.
- Luis P, Vallon L, Tran F-H, Hugoni M, Tran-Van V, Mavingui P, Minard G, Moro CV, 2019 *Aedes albopictus* mosquitoes host a locally structured mycobiota with evidence of reduced fungal diversity in invasive populations. *Fungal Ecology* 39, 257–266.
- Luna C, Hoa NT, Lin H, Zhang L, Nguyen HLA, Kanzok SM, Zheng L, 2006 Expression of immune responsive genes in cell lines from two different Anopheline species. *Insect Molecular Biology* 15, 721–729. [PubMed: 17201765]
- Luz C, Mnyone LL, Sangusangu R, Lyimo IN, Rocha LFN, Humber RA, Russell TL, 2010 A new resting trap to sample fungus-infected mosquitoes, and the pathogenicity of *Lecanicillium muscarium* to culicid adults. *Acta Tropica* 116, 105–107. [PubMed: 20452324]
- Lwetoijera DW, Sumaye RD, Madumla EP, Kavishe DR, Mnyone LL, Russell TL, Okumu FO, 2010 An extra-domiciliary method of delivering entomopathogenic fungus, *Metarhizium anisopliae* IP 46 for controlling adult populations of the malaria vector, *Anopheles arabiensis*. *Parasites & Vectors* 3, 18. [PubMed: 20233423]
- Matskevich AA, Quintin J, Ferrandon D, 2010 The *Drosophila* PRR GGBP3 assembles effector complexes involved in antifungal defenses independently of its Toll-pathway activation function. *European Journal of Immunology* 40, 1244–1254. [PubMed: 20201042]
- Mccray EM, Umphlett CJ, Fay RW, 1973 Laboratory studies on a new fungal pathogen of mosquitoes. *Mosquito News* 33, 54–60.
- Mc Innis T, Zattau WC, 1982 Experimental infection of mosquito larvae by a species of the aquatic fungus *Leptolegnia*. *Journal of Invertebrate Pathology* 39, 98–104.
- Meister S, Agianian B, Turlure F, Relogio A, Morlais I, Kafatos FC, Christophides GK, 2009 *Anopheles gambiae* PGRPLC-mediated defense against bacteria modulates infections with malaria parasites. *PLoS Pathogens*. 5, e1000542. [PubMed: 19662170]

- Meister S, Kanzok SM, Zheng X. -I., Luna C, Li T-R, Hoa NT, Clayton JR, White KP, Kafatos FC, Christophides GK, Zheng L, 2005 Immune signaling pathways regulating bacterial and malaria parasite infection of the mosquito *Anopheles gambiae*. *Proceedings of the National Academy of Sciences* 102, 11420–11425.
- Merritt RW, Dadd RH, Walker ED, 1992 Feeding Behavior, Natural Food, and Nutritional Relationships of Larval Mosquitoes. *Annual Review of Entomology* 37, 349–374. [10.1146/annurev.en.37.010192.002025](https://doi.org/10.1146/annurev.en.37.010192.002025)
- Meyling NV, Eilenberg J, 2007 Ecology of the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* in temperate agroecosystems: Potential for conservation biological control. *Biological Control* 43, 145–155.
- Miranpuri GS, Khachatourians GG, 1991 Infection sites of the entomopathogenic fungus *Beauveria bassiana* in the larvae of the mosquito *Aedes aegypti*. *Entomologia Experimentalis et Applicata* 59, 19–27.
- Mnyone LL, Kirby MJ, Lwetoijera DW, Mpingwa MW, Simfukwe ET, Knols BG, Takken W, Russell TL, 2010 Tools for delivering entomopathogenic fungi to malaria mosquitoes: effects of delivery surfaces on fungal efficacy and persistence. *Malaria Journal* 9, 246. [PubMed: 20799967]
- Moita LF, Wang-Sattler R, Michel K, Zimmermann T, Blandin S, Levashina EA, Kafatos FC, 2005 In vivo identification of novel regulators and conserved pathways of phagocytosis in *An. gambiae*. *Immunity* 23, 65–73. [PubMed: 16039580]
- Moll RM, Romoser WS, Modrakowski MC, Moncayo AC, Lerdthusnee K, 2001 Meconial peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae) metamorphosis. *J Med Entomol* 38, 29–32. [PubMed: 11268687]
- Muturi EJ, Bara JJ, Rooney AP, Hansen AK, 2016 Midgut fungal and bacterial microbiota of *Aedes triseriatus* and *Aedes japonicus* shift in response to La Crosse virus infection. *Molecular Ecology* 25, 4075–4090. [PubMed: 27357374]
- Nakhleh J, El Moussawi L, Osta MA, 2017 Chapter Three - The melanization response in insect immunity, in: Ligoxygakis P (Ed.), *Advances in Insect Physiology, Insect Immunity*. Academic Press 52, 83–109.
- Nappi AJ, Christensen BM, 2005 Melanogenesis and associated cytotoxic reactions: Applications to insect innate immunity. *Insect Biochemistry and Molecular Biology* 35, 443–459. [PubMed: 15804578]
- Nappi AJ, Vass E, 1993 Melanogenesis and the Generation of Cytotoxic Molecules During Insect Cellular Immune Reactions. *Pigment Cell Research* 6, 117–126. [PubMed: 8234196]
- Neafsey DE, Waterhouse RM, Abai MR, Aganezov SS, Alekseyev MA, Allen JE, Amon J, Arca B, Arensburger P, Artemov G, Assour LA, Basseri H, Berlin A, Birren BW, Blandin SA, Brockman AI, Burkot TR, Burt A, Chan CS, Chauve C, Chiu JC, Christensen M, Costantini C, Davidson VLM, Deligianni E, Dottorini T, Dritsou V, Gabriel SB, Guelbeogo WM, Hall AB, Han MV, Hlaing T, Hughes DST, Jenkins AM, Jiang X, Jungreis I, Kakani EG, Kamali M, Kempainen P, Kennedy RC, Kirmizoglou IK, Koekemoer LL, Laban N, Langridge N, Lawniczak MKN, Lirakis M, Lobo NF, Lowy E, MacCallum RM, Mao C, Maslen G, Mbogo C, McCarthy J, Michel K, Mitchell SN, Moore W, Murphy KA, Naumenko AN, Nolan T, Novoa EM, O’Loughlin S, Oranganje C, Oshaghi MA, Pakpour N, Papathanos PA, Peery AN, Povelones M, Prakash A, Price DP, Rajaraman A, Reimer LJ, Rinker DC, Rokas A, Russell TL, Sagnon N, Sharakhova MV, Shea T, Simao FA, Simard F, Slotman MA, Somboon P, Stegny V, Struchiner CJ, Thomas GWC, Tojo M, Topalis P, Tubio JMC, Unger MF, Vontas J, Walton C, Wilding CS, Willis JH, Wu Y-C, Yan G, Zdobnov EM, Zhou X, Catteruccia F, Christophides GK, Collins FH, Cornman RS, Crisanti A, Donnelly MJ, Emrich SJ, Fontaine MC, Gelbart W, Hahn MW, Hansen IA, Howell PI, Kafatos FC, Kellis M, Lawson D, Louis C, Luckhart S, Muskavitch MAT, Ribeiro JM, Riehle MA, Sharakhov IV, Tu Z, Zwiebel LJ, Besansky NJ, 2015 Highly evolvable malaria vectors: The genomes of 16 *Anopheles* mosquitoes. *Science* 347, 1258522–1258522. [PubMed: 25554792]
- Niaré O, Markianos K, Volz J, Oduol F, Touré A, Bagayoko M, Sangaré D, Traoré SF, Wang R, Blass C, Dolo G, Bouaré M, Kafatos FCC, Kruglyak L, Touré YT, Vernick KDD, Niare O, Markianos K, Volz J, Oduol F, Toure A, Bagayoko M, Sangare D, Traore SF, Wang R, Blass C, Dolo G, Bouare M, Kafatos FCC, Kruglyak L, Toure YT, Vernick KDD, 2002 Genetic loci affecting

- resistance to human malaria parasites in a West African mosquito vector population. *Science* 298, 213–216. [PubMed: 12364806]
- Novák D, 1965 Zum Auftreten der Mykosen bei Stechmücken in Mähren (Diptera: Culicidae). 1 15, 135–137.
- Ochiai M, Ashida M, 2000 The binding domain and the cDNA cloning of β -1,3-glucan recognition protein from the silkworm, *Bombyx mori*. *J. Biol. Chem* 275, 4995–5002. [PubMed: 10671539]
- Ouedraogo RM, Cusson M, Goettel MS, Brodeur J, 2003 Inhibition of fungal growth in thermoregulating locusts, *Locusta migratoria*, infected by the fungus *Metarhizium anisopliae* var acridum. *Journal of Invertebrate Pathology* 82, 103–109. [PubMed: 12623310]
- Pelizza SA, López Lastra CC, Becnel JJ, Humber RA, García JJ, 2008 Further research on the production, longevity and infectivity of the zoospores of *Leptolegnia chapmanii* Seymour (Oomycota: Peronosporomycetes). *Journal of Invertebrate Pathology* 98, 314–319. [PubMed: 18511067]
- Pendland JC, Boucias DG, 1996 Phagocytosis of lectin-opsonized fungal cells and endocytosis of the ligand by insect *Spodoptera exigua* granular hemocytes: an ultrastructural and immunocytochemical study. *Cell and Tissue Research* 285, 57–67.
- Pereira E. da S., Ferreira RLM, Hamada N, Lichtwardt RW, 2005 Trichomycete fungi (Zygomycota) associated with mosquito larvae (Diptera: Culicidae) in natural and artificial habitats in Manaus, AM Brazil. *Neotropical Entomology* 34, 325–329.
- Popko DA, Henke JA, Mullens BA, Walton WE, 2018 Evaluation of Two Entomopathogenic Fungi for Control of *Culex quinquefasciatus* (Diptera: Culicidae) in Underground Storm Drains in the Coachella Valley, California, United States. *J Med Entomol* 55, 654–665. [PubMed: 29294059]
- Povelones M, Bhagavatula L, Yassine H, Tan LA, Upton LM, Osta MA, Christophides GK, 2013 The CLIP-Domain Serine Protease Homolog SPCLIP1 Regulates Complement Recruitment to Microbial Surfaces in the Malaria Mosquito *Anopheles gambiae*. *PLOS Pathogens* 9, e1003623. [PubMed: 24039584]
- Prasad A, Veerwal B, 2010 Biotoxicity of entomopathogenic fungus *Beauveria bassiana* (Balsamo) vuillemin, against early instars of anopheline mosquitoes. *Journal of Herbal Medicine*. 4, 181–188.
- Ramirez JL, Dunlap CA, Muturi EJ, Barletta ABF, Rooney AP, 2018a Entomopathogenic fungal infection leads to temporospatial modulation of the mosquito immune system. *PLoS Negl Trop Dis*. 12, e0006433. [PubMed: 29684026]
- Ramirez JL, Muturi EJ, Barletta ABF, Rooney AP, 2018b The *Aedes aegypti* IMD pathway is a critical component of the mosquito antifungal immune response. *Developmental & Comparative Immunology*. In press.
- Ramirez JL, Muturi EJ, Dunlap C, Rooney AP, 2018c Strain-specific pathogenicity and subversion of phenoloxidase activity in the mosquito *Aedes aegypti* by members of the fungal entomopathogenic genus *Isaria*. *Sci Rep*. 8, 9896. [PubMed: 29967469]
- Ramirez JL, Muturi EJ, Barletta ABF, Rooney AP, 2019 The *Aedes aegypti* IMD pathway is a critical component of the mosquito antifungal immune response. *Developmental & Comparative Immunology* 95, 1–9. [PubMed: 30582948]
- Rautenbach M, Troskie AM, Vosloo JA, 2016 Antifungal peptides: To be or not to be membrane active. *Biochimie* 130, 132–145. [PubMed: 27234616]
- Reyes-Villanueva F, Garza-Hernandez JA, Garcia-Munguia AM, Tamez-Guerra P, Howard AF, Rodriguez-Perez MA, 2011 Dissemination of *Metarhizium anisopliae* of low and high virulence by mating behavior in *Aedes aegypti*. *Parasites & Vectors* 4, 171. [PubMed: 21906283]
- Rhodes VL, Thomas MB, Michel K, 2018 The interplay between dose and immune system activation determines fungal infection outcome in the African malaria mosquito, *Anopheles gambiae*. *Developmental & Comparative Immunology* 85, 125–133. [PubMed: 29649553]
- Rhodes VLM, Michel K, 2017 Modulation of Mosquito Immune Defenses as a Control Strategy, in: *Arthropod Vector: Controller of Disease Transmission*, Elsevier. 1, 59–89.
- Ricci I, Mosca M, Valzano M, Damiani C, Scuppa P, Rossi P, Crotti E, Cappelli A, Ulissi U, Capone A, Esposito F, Alma A, Mandrioli M, Sacchi L, Bandi C, Daffonchio D, Favia G, 2011 Different

- mosquito species host *Wickerhamomyces anomalus* (*Pichia anomala*): perspectives on vector-borne diseases symbiotic control. *Antonie van Leeuwenhoek*. 99, 43–50. [PubMed: 21113816]
- Roberts D, 1974 Fungal Infections of Mosquitoes. *Le Controle des Moustiques/Mosquito Control* 143–193.
- Roubaud E, Toumanoff, 1930 Essais d'infection experimentale de larves de culicides par quelques champignons entomophytes. 23, 1025–1027.
- Sandhu DK, Waraich MK, 1985 Yeasts associated with pollinating bees and flower nectar. *Microbial Ecology*. 11, 51–58. [PubMed: 24221239]
- Sato H (Tokyo U. of A. and T., Shimada N, Aoki J, 1989 Light and electron microscopy of *Smittium morbosum* (Trichomycetes), newly recorded from Japan. *Transactions of the Mycological Society of Japan* (Japan).
- Saunders GA, Washburn JO, Egerter DE, Anderson JR, 1988 Pathogenicity of fungi isolated from field-collected larvae of the western treehole mosquito, *Aedes sierrensis* (Diptera: Culicidae). *Journal of Invertebrate Pathology* 52, 360–363. [PubMed: 3183417]
- Schnitger AKD, Kafatos FC, Osta MA, 2007 The melanization reaction is not required for survival of *Anopheles gambiae* mosquitoes after bacterial infections. *Journal of Biological Chemistry*. 282, 21884–21888. [PubMed: 17537726]
- Scholte E-J, 2005 An Entomopathogenic Fungus for Control of Adult African Malaria Mosquitoes. *Science*. 308, 1641–1642. [PubMed: 15947190]
- Scholte E-J, Knols BG, Takken W, 2004a Autodissemination of the entomopathogenic fungus *Metarhizium anisopliae* amongst adults of the malaria vector *Anopheles gambiae* s.s. *Malaria Journal*. 3, 45. [PubMed: 15566626]
- Scholte E-J, Knols BGJ, Samson RA, Takken W, 2004b Entomopathogenic fungi for mosquito control: a review. *J. Insect Sci*. 4, 19. [PubMed: 15861235]
- Schrank A, Vainstein MH, 2010 *Metarhizium anisopliae* enzymes and toxins. *Toxicon* 56, 1267–1274. [PubMed: 20298710]
- Seye F, Faye O, Ndiaye M, Njie E, Afoutou JM, 2009 Pathogenicity of the fungus, *Aspergillus clavatus*, isolated from the locust, *Oedaleus senegalensis*, against larvae of the mosquitoes *Aedes aegypti*, *Anopheles gambiae* and *Culex quinquefasciatus*. *J Insect Sci*. 9.
- Shah PA, Pell JK, 2003 Entomopathogenic fungi as biological control agents. *Applied Microbiology and Biotechnology* 61, 413–423. [PubMed: 12764556]
- Shin SW, Bian G, Raikhel AS, 2006 A Toll Receptor and a Cytokine, Toll5A and Spz1C, are involved in Toll antifungal immune signaling in the mosquito *Aedes aegypti*. *Journal of Biological Chemistry* 281, 39388–39395. [PubMed: 17068331]
- Shin SW, Kokoza V, Bian G, Cheon H-M, Kim YJ, Raikhel AS, 2005 REL1, a Homologue of *Drosophila* Dorsal, regulates Toll antifungal immune pathway in the female mosquito *Aedes aegypti*. *Journal of Biological Chemistry* 280, 16499–16507. [PubMed: 15722339]
- Shoukamy MA, Abdelzaher HMA, Shahin AAB, 2001 Ultrastructural changes in the muscles, midgut, hemopoietic organ, imaginal discs and malpighian tubules of the mosquito *Aedes aegypti* larvae infected by the fungus *Coelomomyces stegomyiae*. *Mycopathologia* 149, 99–106. [PubMed: 11265168]
- Shoukamy MA, Lucarotti CJ, 1998 Pathology of *Coelomomyces stegomyiae* in Larval *Aedes aegypti*. *Mycologia* 90, 559–564.
- Snetselaar J, Andriessen R, Suer RA, Osinga AJ, Knols BG, Farenhorst M, 2014 Development and evaluation of a novel contamination device that targets multiple life-stages of *Aedes aegypti*. *Parasit Vectors*. 7, 200. [PubMed: 24766772]
- Soarés GG, 1982 Pathogenesis of infection by the hyphomycetous fungus, *Tolyposcladium cylindrosporium* in *Aedes sierrensis* and *Culex tarsalis* [Dip.: Culicidae]. *Entomophaga*. 27, 283–299.
- Sousa NA, Lobo LS, Rodrigues J, Luz C, 2013 New insights on the effectiveness of *Metarhizium anisopliae* formulation and application against *Aedes aegypti* eggs. *Letters in Applied Microbiology*. 57, 193–199. [PubMed: 23638865]

- Souza-Neto JA, Sim S, Dimopoulos G, 2009 An evolutionary conserved function of the JAK-STAT pathway in anti-dengue defense. *Proceedings of the National Academy of Sciences*. 106, 17841–17846.
- Steyn A, Roets F, Botha A, 2016 Yeasts associated with *Culex pipiens* and *Culex theileri* mosquito larvae and the effect of selected yeast strains on the ontogeny of *Culex pipiens*. *Microbial Ecology*. 71, 747–760. [PubMed: 26573833]
- Sur B, Bihari V, Sharma A, Basu SK, 1999 Survey of termite-inhabited soil and mosquito breeding sites in Lucknow, India for potential mycopathogens of *Anopheles stephensi*. *Mycopathologia* 144, 77–80.
- Sweeney A, 1975 The Mode of Infection of the Insect Pathogenic Fungus *Culicinomyces* in Larvae of the Mosquito *Culex Fatigans*. *Australian Journal of Zoology*. 23, 49.
- Sweeney AW, 1981 An undescribed species of *Smittium* (Trichomycetes) pathogenic to mosquito larvae in Australia. *Transactions of the British Mycological Society* 77, 55–60.
- Tajedin L, Hashemi J, Abaei M, Hosseinpour L, Rafei F, Basseri H, 2009 Study on fungal flora in the midgut of the larva and adult of the different populations of the malaria vector *Anopheles stephensi*. *Iran J Arthropod Borne Dis*. 3, 36–40.
- Tonk M, Vilcinskis A, 2017 The Medical Potential of Antimicrobial Peptides from Insects. *Curr Top Med Chem*. 17, 554–575. [PubMed: 27411327]
- Travland LB, 1979 Initiation of infection of mosquito larvae (*Culiseta inornata*) by *Coelomomycespsorophorae*. *Journal of Invertebrate Pathology*. 33, 95–105.
- Tuzet O, Manier JF, 1947 *Orphella culici* n. sp., entophyte parasite du rectum des larves de *Culex hortensis* Fcbl. *Comptes rendus des seances de la Societe de biologie et des ses filiales* 225, 264–265
- Valanne S, Wang J-H, Ramet M, 2011 The *Drosophila* Toll Signaling Pathway. *The Journal of Immunology*. 186, 649–656. [PubMed: 21209287]
- Valzania L, Martinson VG, Harrison RE, Boyd BM, Coon KL, Brown MR, Strand MR, 2018 Both living bacteria and eukaryotes in the mosquito gut promote growth of larvae. *PLOS Neglected Tropical Diseases*. 12, e0006638. [PubMed: 29979680]
- van der Weerden NL, Bleackley MR, Anderson MA, 2013 Properties and mechanisms of action of naturally occurring antifungal peptides. *Cellular and Molecular Life Sciences*. 70, 3545–3570. [PubMed: 23381653]
- Vizioli J, Bulet P, Charlet M, Lowenberger C, Blass C, Muller H-M, Dimopoulos G, Hoffmann J, Kafatos FC, Richman A, 2000 Cloning and analysis of a cecropin gene from the malaria vector mosquito, *Anopheles gambiae*. *Insect Molecular Biology*. 9, 75–84. [PubMed: 10672074]
- Vizioli J, Bulet P, Hoffmann JA, Kafatos FC, Muller H-M, Dimopoulos G, 2001a Gambicin: A novel immune responsive antimicrobial peptide from the malaria vector *Anopheles gambiae*. *Proc Natl Acad Sci U S A*. 98, 12630–12635. 10.1073/pnas.221466798 [PubMed: 11606751]
- Vizioli J, Richman AM, Uttenweiler-Joseph S, Blass C, Bulet P, 2001b The defending peptide of the malaria vector mosquito *Anopheles gambiae*: antimicrobial activities and expression in adult mosquitoes. *Insect Biochemistry and Molecular Biology*. 31, 241–248. [PubMed: 11167093]
- Wallis GP, Aitken THG, Amato GD, Tabachnick AJ, 1985 Selection for susceptibility and refractoriness of *Aedes aegypti* to oral infection with yellow fever virus. *Am J Trop Med Hyg*. 34, 1225–1231. [PubMed: 3834805]
- Wanchoo A, Lewis MW, Keyhani NO, 2009 Lectin mapping reveals stage-specific display of surface carbohydrates in in vitro and haemolymph-derived cells of the entomopathogenic fungus *Beauveria bassiana*. *Microbiology*. 155, 3121–3133. [PubMed: 19608611]
- Wang C, Wang S, 2017 Insect Pathogenic Fungi: Genomics, Molecular Interactions, and Genetic Improvements. *Annual Review of Entomology*. 62, 73–90.
- Wang X, Fuchs JF, Infanger L-C, Rocheleau TA, Hillyer JF, Chen C-C, Christensen BM, 2005 Mosquito innate immunity: involvement of β 1,3-glucan recognition protein in melanotic encapsulation immune responses in *Armigeres subalbatus*. *Molecular and Biochemical Parasitology*. 139, 65–73. [PubMed: 15610820]

- Wang X, Rocheleau TA, Fuchs JF, Christensen BM, 2006 Beta 1, 3-glucan recognition protein from the mosquito, *Armigeres subalbatus*, is involved in the recognition of distinct types of bacteria in innate immune responses. *Cellular Microbiology*. 8, 1581–1590. [PubMed: 16984413]
- Wang Y-H, Hu Y, Xing L-S, Jiang H, Hu S-N, Raikhel AS, Zou Z, 2015 A critical role for CLSP2 in the modulation of antifungal immune response in mosquitoes. *PLoS Pathog.* 11, e1004931. [PubMed: 26057557]
- Waterhouse RM, Kriventseva EV, Meister S, Xi Z, Alvarez KS, Bartholomay LC, Barillas-Mury C, Bian G, Blandin S, Christensen BM, Dong Y, Jiang H, Kanost MR, Koutsos AC, Levashina EA, Li J, Ligoxygakis P, MacCallum RM, Mayhew GF, Mendes A, Michel K, Osta MA, Paskewitz S, Shin SW, Vlachou D, Wang L, Wei W, Zheng L, Zou Z, Severson DW, Raikhel AS, Kafatos FC, Dimopoulos G, Zdobnov EM, Christophides GK, 2007 Evolutionary Dynamics of Immune-Related Genes and Pathways in Disease-Vector Mosquitoes. *Science*. 316, 1738–1743. [PubMed: 17588928]
- Weiser J, Batko A, 1966 A new parasite of *Culex pipiens* L., *Entomophthora destruens* sp. nov. (Phycomycetes, Entomophthoraceae). *Folia Parasitologica* 13.
- Williams MC, Lichtwardt RW, 1972 Infection of *Aedes aegypti* Larvae by Axenic Cultures of the Fungal Genus *Smittium* (Trichomycetes). *American Journal of Botany* 59, 189–193.
- White MM, Siri A, Lichtwardt RW, 2006 Trichomycete insect symbionts in Great Smoky Mountains National Park and vicinity. *Mycologia* 98, 333–352. [PubMed: 16894978]
- Wong TL, Pillai JS, 1978 *Coelomomyces opifexi* Pillai & Smith (Coelomomycetaceae: Blastocladiiales) IV. Host range and relative susceptibility of *Aedes australis* and *Opifex fuscus* larvae. *New Zealand Journal of Zoology* 5, 807–810.
- Wong TL, Pillai JS, 1980 *Coelomomyces opifexi* Pillai & Smith (Coelomomycetaceae: Blastocladiiales) VI. Observations on the mode of entry into *Aedes australis* larvae. *New Zealand Journal of Zoology* 7, 135–139.
- Wu RC-C, Cho W-L, 2014 Cloning and characterization of microbial activated *Aedes aegypti* MEK4 (AaMEK4): influences of noncatalytic domains on enzymatic activity. *Insect Molecular Biology*. 23, 644–655. [PubMed: 25039995]
- Yassine H, Kamareddine L, Chamat S, Christophides GK, Osta MA, 2014 A serine protease homolog negatively regulates TEP1 consumption in systemic infections of the malaria vector *Anopheles gambiae*. *JIN*. 6, 806–818.
- Yassine H, Kamareddine L, Osta MA, 2012 The mosquito melanization response is implicated in defense against the entomopathogenic fungus *Beauveria bassiana*. *PLOS Pathogens*. 8, e1003029. [PubMed: 23166497]
- Yee DA, Kesavaraju B, Juliano SA, 2004 Larval feeding behavior of three co-occurring species of container mosquitoes. *J Vector Ecol.* 29, 315–322. [PubMed: 15707290]
- Zattau WC, McInnis T, 1987 Life cycle and mode of infection of *Leptolegnia chapmanii* (Oomycetes) parasitizing *Aedes aegypti*. *Journal of Invertebrate Pathology* 50, 134–145. [PubMed: 3624894]
- Zeidler MP, Bausek N, 2013 The *Drosophila* JAK-STAT pathway. *JAK-STAT*. 2, e25353. [PubMed: 24069564]
- Zhang Rudian, Zhu Y, Pang X, Xiao X, Zhang Renli, Cheng G, 2017 Regulation of Antimicrobial Peptides in *Aedes aegypti* Aag2 Cells. *Front. Cell. Infect. Microbiol.* 7, 22. [PubMed: 28217557]
- Zhang X, An C, Sprigg K, Michel K, 2016 CLIPB8 is part of the prophenoloxidase activation system in *Anopheles gambiae* mosquitoes. *Insect Biochemistry and Molecular Biology*. 71, 106–115. [PubMed: 26926112]

Highlights

1. Mosquito-fungus interactions occur in all life stages.
2. Fungal entomopathogens are taken up passively or enter the mosquito actively.
3. Fungal infections activate all branches of the mosquito immune system.
4. New technology enables research of mosquito-fungus interactions beyond biocontrol.




Figure 1: The multitude of potential fungus-mosquito encounters.

Adult mosquitoes are exposed to fungi on a multitude of surfaces during indoor and outdoor resting (A, D, H), blood and sugar feeding (C, G). Additional fungal encounters can occur during mating (B) and oviposition (E). Mosquito larvae in their aquatic environment are exposed to fungi on plant detritus, within the water column, and at the water surface (F).




Figure 2: Entry routes of fungal entomopathogens into their mosquito hosts.

Entomopathogens enter mosquito larvae and adults through two distinct routes. Top panel: Fungal and water mold asexual spores attach to the mosquito cuticle and penetrate actively via penetration pegs. Opportunistic fungal pathogens can gain entry through wound sites. Center panel: Ingested spores may be degraded through digestive enzymes in the larval midgut, resulting in toxin release. Some spores may germinate and penetrate through the midgut epithelium, subsequently disseminating throughout the larval body. It is unclear, whether this occurs in adult mosquitoes. Bottom panel: Trichospores of *Smittium sp.* attach to the cuticular lining of the hindgut in mosquito larvae, germinate and grow locally. Hyphae of *Smittium morbosum* can grow anteriorly, and penetrate the posterior midgut epithelium, where they are melanized.




Figure 3: Immune modules that regulate mosquito antifungal immunity. Fungi that penetrate epithelia and reach the hemolymph, as indicated by the germinating blastospore drawing, are attacked by multiple antifungal molecules. These molecules can be expression products of signal transduction pathways, including the Toll, IMD, JAK/STAT and MAPK pathways. Alternatively, these molecules are proteolytic activation products of humoral killing modules, circulating in the hemolymph, including PO and TEPI, which are products of the melanization and complement-like pathways, respectively. Proteins are bolded if their role in mosquito antifungal immunity is supported experimentally. Other key factors of the mosquito immune modules are listed in gray.