



# HHS Public Access

Author manuscript

*Curr Immunol Rev.* Author manuscript; available in PMC 2020 January 01.

Published in final edited form as:

*Curr Immunol Rev.* 2019 ; 15(1): 63–75. doi:10.2174/1573395514666180528081204.

## Mucosal Immunity in HIV/SIV Infection: T Cells, B Cells and Beyond

Barbara L. Shacklett<sup>\*,1,2</sup>

<sup>1</sup>Department of Medical Microbiology and Immunology

<sup>2</sup>Division of Infectious Diseases, Department of Medicine, School of Medicine, University of California, Davis, CA 95616

### Abstract

As our understanding of mucosal immunity increases, it is becoming clear that the host response to HIV-1 is more complex and nuanced than originally believed. The mucosal landscape is populated with a variety of specialized cell types whose functions include combating infectious agents while preserving commensal microbiota, maintaining barrier integrity, and ensuring immune homeostasis. Advances in multiparameter flow cytometry, gene expression analysis and bioinformatics have allowed more detailed characterization of these cell types and their roles in host defense than was previously possible. This review provides an overview of existing literature on immunity to HIV-1 and SIVmac in mucosal tissues of the female reproductive tract and the gastrointestinal tract, focusing on major effector cell populations and briefly summarizing new information on tissue resident memory T cells, T<sub>reg</sub>, Th17, Th22 and innate lymphocytes (ILC), subsets that have been studied primarily in the gastrointestinal mucosa.

### Keywords

HIV-1; SIV; mucosa; gut; T-cell; adaptive; innate

## 1. INTRODUCTION

Mucosal tissues house a majority of the body's lymphocytes, including the CD4<sup>+</sup> T cells that serve as the primary targets for HIV-1 infection. The reproductive and gastrointestinal mucosal tissues are the major portals of entry for HIV-1. Over the past decades, numerous studies have explored the immune responses occurring at these tissue sites, with results underscoring the concept that the balance between inflammation and protective immunity is established at or near these portals of entry. While many aspects of mucosal immunity remain to be fully elucidated, these efforts have revealed important new insights into the innate and adaptive immune cells housed within mucosal tissues, including trafficking

\*Corresponding Author: Barbara L. Shacklett, PhD, Professor, Dept. of Medical Microbiology and Immunology, 3327 Tupper Hall, 1 Shields Avenue, Davis, CA 95616, Tel: (530) 752-6785, Fax: (530) 752-8692, blshacklett@ucdavis.edu.

Conflict of Interest

The author is aware of no conflict of interest that could be perceived to bias this work.

patterns, regulation of effector functions, induction of memory, and functional plasticity. What follows is a summary of this literature, with an emphasis on recent developments.

## 2. THE FEMALE REPRODUCTIVE TRACT

### 2.1. The Female Reproductive Tract: Which Sites Are Most Important for Immune Defense?

The vast majority of HIV-1 infections result from sexual contact via the genital tract or the anorectal canal. Women account for roughly half of the estimated 36-37 million persons living with HIV-1 infection worldwide. HIV-1 acquisition via the female reproductive tract (FRT) is believed to be less efficient than transmission via the anorectal mucosa, an interpretation supported by experimental infection studies in rhesus macaques [1]. The lower FRT, which includes the vagina and ectocervix, is shielded from infection by a relatively thick, stratified squamous (type II) epithelium [2]. In contrast, the upper FRT, which includes the endocervix and uterine endometrium, is lined with simple columnar (type I) epithelium. These two tissues meet at the cervical transformation zone, an area rich in potential HIV-1 target cells such as CD4<sup>+</sup> T cells and antigen-presenting cells [3]. Cervical mucus, which helps protect the upper tract, varies in volume, composition and physical characteristics throughout the menstrual cycle [4], becoming a thick “mucus plug” during pregnancy [5].

The majority of HIV-1 infections in women are believed to occur *via* the lower FRT, a concept supported by experimental inoculation studies of rhesus macaques with simian immunodeficiency virus (SIVmac) [6, 7]. Hysterectomized macaques are susceptible to productive infection following intravaginal exposure [8], and a randomized clinical trial revealed no reduction in HIV-1 acquisition among women using a diaphragm (which blocks access to the upper FRT) compared to controls [9]. Proposed mechanisms by which HIV-1 may enter the body through the lower FRT include disruption of the mucosal surface through micro-breaches occurring during sexual intercourse; disruption due to inflammation or ulceration associated with other sexually transmitted infections; uptake of virus by Langerhans' cells or dendritic cells within or directly below the epithelium; and direct infection of CD4<sup>+</sup> T cells or macrophages within the epithelium [10–14]. In a recent study, CCR6<sup>+</sup> CD4<sup>+</sup> T cells of the Th17 lineage were identified as the primary targets of SIV during vaginal transmission [15], extending previous studies showing high susceptibility of this subset to HIV-1 infection *in vitro* [16].

Although HIV-1 transmission clearly can and does occur via the lower FRT, susceptible target cells are present throughout the upper and lower tract [14, 17]. The extent to which the upper FRT serves as a target and/or reservoir for HIV-1 replication is not known. CD4<sup>+</sup> T cells in the upper FRT express CCR5 and exhibit an activated memory phenotype [17]. In experimental infection studies of rhesus macaques using SIVmac, tissues of the upper tract, including the ovary, were shown to become infected following intravaginal exposure [18]. *In vitro* studies have also demonstrated susceptibility of ovarian CD4<sup>+</sup> T cells to infection with laboratory-adapted HIV-1 strains [19]. However, imaging studies in women undergoing simulated intercourse, utilizing radiolabeled surrogates for cell-free and cell-associated virus, did not reveal migration of the surrogates to the upper FRT [20]. *In vitro*, endometrial

macrophages and T cells are permissive to HIV-1 infection; however, endometrial and decidual macrophages express the HIV-1 restriction factor SAMHD1, which may restrict their susceptibility to productive infection *in vivo* [21, 22]. To date, few studies have addressed the extent to which CD4<sup>+</sup> T cells in the upper FRT are infected *in vivo* [23]. In summary, then, further studies are needed to fully address the role of the upper FRT in HIV-1 transmission, replication and pathogenesis.

## 2.2. Antigen-Specific T-cell Responses in the FRT

Comprehensive studies of immune responses in the human FRT present significant logistical challenges, and are therefore rare in the literature. However, several groups have investigated adaptive responses in rhesus macaques experimentally infected with SIVmac, and to a lesser extent in HIV-1-infected women. Mucosal CD8<sup>+</sup> T-cell responses to SIVmac emerge in the cervicovaginal mucosa with kinetics that are “too little and too late” to prevent viral dissemination to draining lymph nodes [24]. Coupled with observations from murine lymphocytic choriomeningitis virus infection, this finding has suggested that vaccine-mediated induction of a sizable population of HIV-1-specific T cells in the FRT might provide protection against vaginal exposure [25, 26]. However, it has proven challenging to induce sufficiently large populations of antigen-specific T cells at mucosal ‘front lines’ to allow direct testing of this hypothesis.

During chronic infection, HIV-1-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells are detected in the cervix and vagina. In early studies, SIVmac-specific cytotoxic T cells (CTL) were identified in vaginal tissues of infected rhesus macaques [27], and HIV-1-specific CTL activity was detected by <sup>51</sup>Cr release assay in polyclonally expanded cervical T cells from HIV-1-positive women [28, 29]. Comparison of MHC restriction, TCR CDR3 region sequences, and epitopes recognized by CTL from cervix and blood revealed that certain T-cell clones were indeed found at both sites, suggesting that some HIV-1-specific T-cell clones are widely distributed throughout the body rather than restricted to tissue residency [28, 29].

A series of more recent studies, conducted among South African women with HIV-1, focused on cervical CD8<sup>+</sup> T-cell responses, their relationship to cervical inflammation, and their role in limiting viral shedding in cervical secretions [30–32]. Gumbi and colleagues [33] studied the relationship between local inflammation and HIV-1 Gag-specific responses in CD8<sup>+</sup> T cells from cervical cytobrush. They detected no relationship between the magnitude of cervical responses and virus shedding in genital secretions, implying that cervical T-cell responses are often ineffective at containing viral replication. Women who were shedding virus in cervical secretions had higher concentrations of proinflammatory cytokines (TNF $\alpha$ , IL-1 $\beta$ , IL-6, and IL-8) in secretions than non-shedders, suggesting that genital inflammation may promote HIV-1 replication and shedding within the cervical mucosa [33].

T-cell “polyfunctionality”, i.e., the ability to respond to T-cell receptor stimulation by producing multiple effector cytokines, chemokines and cytolytic granule constituents, has been associated with control of HIV-1 replication in Elite Controllers, who maintain low to undetectable viral load without antiretroviral therapy [34, 35]. Investigating cervical HIV-1 Gag-specific T cells in a South African cohort, Bere and colleagues [36] found that these

cells were predominantly monofunctional. Polyfunctional cervical T cells were sometimes detected in women with relatively high blood CD4<sup>+</sup> T-cell counts and low plasma viral loads, but the presence of polyfunctional T cells did not prevent HIV-1 shedding from the genital tract [36].

Nkwanyana and colleagues [37] explored the phenotype of cervical T cells from HIV-1 infected women. CD4<sup>+</sup> T cells were relatively depleted, most CD8<sup>+</sup> T cells had an effector memory phenotype, and T cell counts were positively correlated with concentrations of proinflammatory cytokines [37]. Mkhize and colleagues [38] studied the effects of highly active antiretroviral therapy (HAART) on cervical CD4<sup>+</sup> T-cell reconstitution and HIV-1 viral load suppression in cervical secretions. Their work revealed a significant association between genital HIV-1 shedding and viral RNA levels in blood, both of which were well suppressed in HAART-compliant women. However, HIV-1-specific CD8<sup>+</sup> T-cell responses in blood were greatly reduced following HAART, while those in the cervix were maintained. This finding may suggest that suboptimal drug concentrations were present in the lower FRT, allowing for low-level or intermittent viral persistence that would stimulate memory T-cell responses [38].

### 2.3. Reproductive Hormones and Adaptive Immunity in the FRT

Reproductive hormones impact many features of the FRT, including the composition, thickness and abundance of genital mucus [39]; the composition of vaginal microbiota [39]; the thickness of the vaginal epithelium in nonhuman primates [40]; and the presence and activity of immune effector cells and molecules of innate immunity [41]. The effects of hormonal contraceptives on the FRT, and implications for HIV-1 susceptibility, have become the subject of considerable study and debate. Two recent meta-analyses concluded that women using the long-lasting injectable contraceptive depot-medroxyprogesterone acetate (DMPA) face a 40-50% greater risk of HIV-1 acquisition compared to controls [42, 43]. The mechanisms underlying this enhanced risk are incompletely understood, but may include increased target cell density; increased HIV-1 coreceptor expression; thinning of the epithelial barrier; and/or changes in the vaginal microbiome [44–47]. The question of enhanced susceptibility promises to remain a topic of active investigation, with important implications for public health policy.

Tissues of the upper and lower FRT respond differently to cyclic hormonal fluctuations during the menstrual cycle [48, 49]. In the lower tract (ectocervix and vagina), cytotoxic T-cell activity remains relatively constant throughout the menstrual cycle. However, levels of immunoglobulins (IgG and IgA) in cervical secretions decline after ovulation, rebounding near the end of the cycle [41]. Similar patterns have been observed for soluble factors of innate immunity (e.g., lactoferrin, SLPI and HBD2), suggesting that the post-ovulatory phase, termed the secretory or luteal phase, represents a window of vulnerability to HIV-1 and other sexually transmitted pathogens for the lower reproductive tract [48, 49].

In the upper FRT, adaptive T-cell responses vary through the menstrual cycle: CD8<sup>+</sup> T cells isolated from proliferative phase endometrial tissue show strong cytotoxic activity in <sup>51</sup>Cr release assays; however, T cells isolated during the secretory phase show reduced activity [48, 49]. Thus, antigen-specific T-cell responses are suppressed in the upper tract during the

secretory phase, reinforcing the notion of a “window of opportunity” for pathogens. Other changes affect the distribution of endometrial T cells: during the proliferative phase, CD8<sup>+</sup> T cells localize to lymphoid aggregates located in the lamina basalis of the uterine endometrium [50, 51]. The function of these structures is not known, but they may serve to promote retention of resident memory T cells during menstrual shedding [41]. HIV-1-specific T-cell responses in the upper FRT have not been extensively characterized [52], nor has the impact of the menstrual cycle on these responses been fully explored to date.

### 3. THE GASTROINTESTINAL TRACT

#### 3.1. The Gastrointestinal Tract as a Portal of HIV-1 Entry

The gastrointestinal (GI) tract is the largest lymphoid organ in the body, as well as a major physical and immunological barrier against infectious pathogens. In the early years of the HIV/AIDS epidemic, diarrhea and wasting syndromes were among the most commonly reported features of the disease, although the reasons for this were initially unclear. Research later revealed that the GI tract houses CD4<sup>+</sup> T-cell subsets that are highly susceptible to HIV-1 infection, and are among the first cells in the body to be depleted regardless of the route of HIV-1 transmission [53]. Subsequent work provided evidence linking acute HIV-1 infection to gut epithelial damage and microbial translocation, contributing to the generalized immune activation associated with HIV-1 disease [54, 55].

Transmission studies in rhesus macaques suggest that anal intercourse is significantly more likely than vaginal intercourse to result in HIV-1 acquisition by the receptive partner[1]. The upper anorectal canal is lined with simple columnar (type I) epithelium; the lower canal is lined with stratified squamous (type II) epithelium. These are separated by the dentate (or pectinate) line and a narrow zone of transitional epithelium [56]. HIV-1 target cells are abundant in the distal GI tract [57–59]. During receptive anal intercourse, HIV-1 transmission may occur through several mechanisms: epithelial microabrasions or breaches; transcytosis across epithelial cells; or virion capture by mucosal dendritic cells (reviewed in [60]). Recent studies have used SPECT/CT imaging of radiolabeled surrogates to model dissemination of viral particles in the anorectal canal during simulated intercourse. Radiolabeled surrogates reached peak levels at 10-20 cm from the anal verge, near the rectosigmoid junction [61]. Although under-studied in this context, the anorectal mucosa should not be overlooked as a potential portal of entry for HIV-1 in heterosexual women. A recent meta-analysis suggested significant prevalence and frequency of anal intercourse among heterosexual women in a wide range of cultures and environments [62, 63].

The susceptibility of the GI tract to HIV-1 infection, along with the relative accessibility of tissue sampling in this compartment, have made possible a wide range of detailed studies designed to elucidate the role of individual immune cell subsets in the host response to HIV/SIV. The remainder of this section presents a summary of major findings, organized by immune cell type.

### 3.2. Gastrointestinal CD8<sup>+</sup> T Cells and HIV-1 Infection

The literature describing CD8<sup>+</sup> T-cell responses to HIV-1 infection is considerable; however, the vast majority of studies have focused on T cells isolated from blood. Early mucosal studies of individuals with chronic HIV-1 infection who were also seropositive for cytomegalovirus (CMV) revealed that HIV-1-specific CD8<sup>+</sup> T cells could be detected in rectal and duodenal biopsies, while CMV-specific CD8<sup>+</sup> T cells, though abundant in blood, were rare in these tissues [64–66]. Polyclonally expanded rectal CD8<sup>+</sup> T cells were demonstrated to kill HIV-1 antigen-pulsed target cells in <sup>51</sup>Cr release assays [29, 65]. Mapping of HIV-1-specific CD8<sup>+</sup> T-cell epitope recognition, using pooled peptides, revealed that mucosal responses tended to mirror those in blood, at least in antigenic specificity [67]. With the advent of multiparameter flow cytometry, protocols were developed to measure T-cell production of multiple cytokines, chemokines, and cytolytic granule constituents. These approaches prompted attempts to identify immunologic correlates of viral clearance. Studies of individuals with chronic HIV-1 infection and a range of viral loads and CD4<sup>+</sup> T-cell counts revealed that strong, polyfunctional HIV-1 Gag-specific responses in rectal mucosa were frequently associated with well-preserved mucosal CD4<sup>+</sup> T-cell populations and low plasma viral load [68, 69]. Individuals meeting the definition of ‘HIV controllers’ had significant preservation of mucosal CD4<sup>+</sup> T cells, as well as strong, polyfunctional HIV Gag-specific CD8<sup>+</sup> [34, 70] and CD4<sup>+</sup> [71] T-cell responses in rectal mucosa. Many, but not all of these controllers possessed the MHC class I alleles HLA-B\*57 and/or B\*27. Taken together, these findings supported the widely held interpretation, later confirmed by genome-wide association studies [72, 73], that MHC-restricted T-cell responses account for a significant component of the “elite controller” phenomenon, and further suggested that T-cell populations residing in the GI tract contribute to this immune control [34].

**3.2.1. Are Gastrointestinal CD8<sup>+</sup> T Cells “Dysfunctional” for Cytotoxicity?—** MHC class I restricted, CD8<sup>+</sup> T cells, which release cytolytic granules upon T-cell receptor (TCR) stimulation, have generally been considered to act primarily by killing infected host cells. Accordingly, the terms CD8<sup>+</sup> T cell and cytotoxic T cell (CTL) are often used interchangeably. However, close examination of gastrointestinal CD8<sup>+</sup> T cells revealed that these cells contain significantly less perforin and granzyme B than their blood counterparts, regardless of HIV-1 infection status [74]. This deficit was most evident in rectal effector memory (T<sub>EM</sub>) and terminally differentiated effector (T<sub>EMRA</sub>) CD8<sup>+</sup> T-cell subsets, whose blood counterparts contain the highest levels of perforin and granzyme B [75]. Furthermore, rectal CD8<sup>+</sup> T cells from both healthy and HIV-1-infected individuals were significantly less able than blood CD8<sup>+</sup> T cells to kill GFP-labeled P815 target cells in a redirected lysis assay [75]. Finally, two transcription factors whose expression is required for perforin expression and cytotoxicity, T-bet and Eomesodermin (Eomes), are weakly expressed by rectal CD8<sup>+</sup> T cells [75]. These observations are consistent with the recently described tissue resident memory subset (T<sub>RM</sub>), described below. Taken together, these findings suggest that the predominant antiviral functions of rectal CD8<sup>+</sup> T cells involve release of HIV-1-blocking chemokines and immunomodulatory cytokines, with cytotoxic killing of infected CD4<sup>+</sup> T cells playing a secondary role [75]. We speculate that tissue-specific limitation on cytotoxic activity (i.e., low perforin and granzyme expression) might prove beneficial to the host by limiting mucosal T-cell depletion and tissue damage; however, in the case of chronic viral



infection such as HIV-1, the inability to fully eradicate infected cells likely also contributes to the role of the GI tract as a viral reservoir. It is tempting to suggest that approaches designed to stimulate more effective mucosal cytotoxic T-cell responses, either by increasing their abundance or by “programming” them for greater cytotoxic capacity, might provide more effective first-line protection against mucosal infections, and/or promote better pathogen clearance from tissues in the context of chronic infection [26, 75]

### 3.2.2. Tissue Resident Memory T Cells ( $T_{RM}$ ): A Unique Role in Antiviral Host Defense

—Until recently the dominant paradigm for T-cell differentiation suggested that activated effector T cells migrate continuously throughout the host, searching for and responding to local tissue infection. However, extensive studies in mice revealed that effectors entering certain tissues, including the gastrointestinal tract, are induced to differentiate into “tissue resident” cells by exposure to locally produced cytokines including TGF $\beta$ , IL-15, IL-33 and TNF $\alpha$  [76–83].  $T_{RM}$  exhibit a distinct cell surface phenotype, with expression of  $\alpha E\beta 7$  integrin/CD103, CD69, and downregulation of CD62L, CCR7 and S1PR1 [84]. In contrast to short-lived effector cells, which express T-bet, and lymphoid-localized central memory cells, which express Eomesodermin,  $T_{RM}$  are typically T-bet<sup>Low</sup> and Eomes<sup>Neg</sup>. This corresponds to the phenotype previously observed to be abundant in rectal mucosal CD8<sup>+</sup> T cells from HIV-1<sup>+</sup> and healthy individuals [75, 82, 85].

Given that  $T_{RM}$  in mouse models play important roles in triggering innate immune responses and recruiting other immune cells to sites of infection, it will be important to clarify the contribution of  $T_{RM}$  to HIV-1 clearance in humans. As stated above, a more nuanced understanding of the role of mucosal CD8<sup>+</sup> T cells in fighting viral infection may inform attempts to purge mucosal tissues of residual HIV-1 replication, particularly in the context of therapeutic vaccination and the ‘HIV Cure’ initiative.

### 3.3. Gastrointestinal CD4<sup>+</sup>T Cells and HIV Infection

As discussed extensively elsewhere, gastrointestinal CD4<sup>+</sup> T cells are a major target for HIV/SIV infection and depletion during the acute phase of infection [86–89]. This depletion occurs rapidly, regardless of the initial infection route [53]. The mechanisms driving this phenomenon include direct viral infection [88] and apoptosis of uninfected bystander cells [87], as well as disruption of mucosal homing pathways involving integrin and chemokine expression [90, 91]. Initiation of antiretroviral therapy during acute HIV infection can partially preserve and/or restore mucosal CD4<sup>+</sup> T-cell populations; specific findings have varied by study [92–95].

#### 3.3.1. Th17 Cells

—Th17 cells appear to be preferentially infected and depleted in the GI tract during acute HIV/SIV infection [96]. In addition to IL-17, these CD4<sup>+</sup> T cells produce IL-22, IL-21, and are important for maintaining epithelial integrity and defense against extracellular bacteria and fungi [96]. In humans, Th17 differentiation requires transcription factor ROR $\gamma$ t and exposure to polarizing cytokines including IL-23 [97]. Th17 cells are preferentially lost in hosts with pathogenic lentivirus infection (HIV-1 infected humans; SIVmac infected rhesus macaques), but not from hosts with nonpathogenic infection (sooty

mangabeys infected with SIVsm; African green monkeys) [96]. Individuals characterized as HIV-1 longterm nonprogressors also have intact mucosal Th17 populations [98].

Early loss of intestinal Th17 cells is associated with impairment of the gut epithelial barrier, promoting microbial translocation and chronic immune activation. In macaques, loss of Th17 cells as a result of SIVmac infection is associated with reduced immune control of *Salmonella enterica* serovar Typhimurium [99]. However, initiation of HAART during early acute HIV-1 infection (Fiebig I/II stages) leads to relative preservation of mucosal Th17 cells and reversal of systemic immune activation [95].

Using cells isolated from human blood to explore Th17-polarized subsets in greater detail than could be attempted with mucosal biopsies, Gosselin and colleagues [100,101] demonstrated that Th17-like cells with the phenotypes CCR4<sup>+</sup>/CCR6<sup>+</sup> and CXCR3<sup>+</sup>/CCR6<sup>+</sup> are highly permissive to HIV-1 infection *in vitro*. In colonic mucosa and blood of HIV-1-infected individuals on ART with undetectable plasma viremia, memory CD4<sup>+</sup> T cells expressing CCR6 were found to be enriched for replication-competent HIV DNA [102]. CD4<sup>+</sup> T cells activated under Th17-polarizing conditions reportedly have diminished expression of RNAses that can inhibit HIV-1 replication [103]; this may partly explain the increased susceptibility of such cells to HIV/SIV infection.

**3.3.2. Regulatory T Cells**—Regulatory T cells (T<sub>reg</sub>) limit activation and effector functions of multiple immune cell types [104]. Their role is to restrict potentially harmful functions of autoreactive cells that have escaped central tolerance, as well as to limit ‘collateral damage’ that can arise as a consequence of normal pathogen-specific immune responses [104]. T<sub>reg</sub> are identified by expression of the high-affinity interleukin-2 (IL-2) receptor alpha chain, CD25, and the transcription factor Foxp3. In chronic infections such as HIV-1, which require long-term commitment of adaptive immune ‘resources’, T<sub>reg</sub> may inappropriately dampen the host response, favoring pathogen persistence. Similar to the classical Th1/Th2 paradigm describing a reciprocal relationship between two CD4<sup>+</sup> T-cell lineages with complementary functions, T<sub>reg</sub> and Th17 are derived from a common progenitor and their differentiation is determined by cytokines produced by antigen-presenting cells in response to microbial products [105].

Studying HIV-1<sup>+</sup> controllers and non-controllers, Shaw and colleagues [106] found that the frequency of T<sub>reg</sub> in rectal mucosa positively correlated with plasma viral load and expression of T-cell activation markers. Thus, HIV-1-positive individuals with high viral load and immune activation had high T<sub>reg</sub> frequencies in mucosa. Mucosal T<sub>reg</sub> in chronic HIV-1 infection maintained their capacity to suppress proliferation of autologous non-T<sub>reg</sub> cells [106], suggesting that T<sub>reg</sub> may contribute to suboptimal HIV-1-specific T-cell responses and, by extension, to viral persistence. In a cohort of Colombian HIV-positive subjects, HAART treatment significantly decreased, but did not normalize, T<sub>reg</sub> frequency in rectal mucosa compared to seronegatives [107]. Exploring the mechanisms driving T<sub>reg</sub> expansion in tissues of SIV-infected rhesus macaques, Presicce and colleagues [108] found that mature myeloid dendritic cells (mDCs) from spleen and mesenteric lymph nodes efficiently induced ‘conversion’ of autologous non-T<sub>reg</sub> to Foxp3<sup>+</sup> T<sub>reg</sub>.



**3.3.3. The T<sub>reg</sub>/Th17 Balance in the Gastrointestinal Tract**—In nonhuman primate models of lentiviral infection, disease progression is associated with induction of T<sub>reg</sub> and loss of Th17 cells; thus, the T<sub>reg</sub>/Th17 balance is skewed towards T<sub>reg</sub> [96, 109]. This development is strongly associated with increased systemic immune activation: Th17 are preserved in nonpathogenic SIV infection, but T<sub>reg</sub> are expanded and Th17 lost in progressive SIV disease [96, 109, 110]. The enzyme indoleamine 2,3-dioxygenase (IDO) plays a critical role in regulating the T<sub>reg</sub>/Th17 balance. Expressed by certain antigen-presenting dendritic cells, IDO is upregulated by interferons and TLR agonists, which may be triggered by numerous infectious and inflammatory conditions [111]. Increased IDO activity leads to higher levels of tryptophan catabolites, inducing Foxp3 expression and T<sub>reg</sub> development, and suppressing the Th17 lineage [112]. This dysregulation contributes to the persistent inflammatory state observed in tissues during chronic HIV-1 disease [113, 114], which is only partially reversed by antiretroviral therapy (ART) [114]. Conversely, loss of CD103<sup>+</sup> mucosal dendritic cells (DC), a subset that can induce naïve T cells to express *RORc*, has been associated with loss of mucosal lymphocytes producing IL-17 and IL-22 in the SIVmac model [115]. Altered T-cell homing patterns also contribute to the Th17/T<sub>reg</sub> imbalance: expression of chemokine CCL20 by small intestine epithelial cells is perturbed in chronic HIV-1 infection, including in ART-treated individuals. This leads to impaired homing of CD4<sup>+</sup> T cells expressing the chemokine receptor CCR6, specifically Th17 cells, to the small intestine, and the proportion of gut T<sub>reg</sub> is correspondingly increased [116].

Intriguingly, administration of a lactobacillus probiotic along with IL-21 and ART during chronic SIV infection led to reduced IDO activity and expansion of intestinal Th17 cells, suggesting that such approaches might help restore mucosal immune homeostasis during chronic HIV-1 infection [117, 118].

**3.3.4. Th22 Cells**—The cytokine IL-22 promotes innate defense against mucosal pathogens, is important for maintaining mucosal barrier integrity, and promotes epithelial modeling and repair [119]. Although IL-22 is produced by Th17 cells, a distinct subset of CD4<sup>+</sup> T cells found in epithelia of the skin and GI tract produces IL-22 in the absence of IFN $\gamma$  and IL-17; this subset has been designated Th22 [97, 120–122]. Th22 cells share certain common surface markers with Th17 cells (e.g., CCR4 and CCR6), but also express CCR10 and the aryl hydrocarbon receptor [97, 120–122]. Like Th17, Th22 cells also express CCR5 and are depleted during HIV-1 infection, and their loss is associated with a loss of epithelial barrier function and increased microbial translocation [123]. Evidence from *in vitro* studies and the SIVmac model revealed that the Th17 and Th22 subsets act cooperatively to maintain mucosal barrier functions [124]. Also, SIVmac-infected animals with high colorectal Th17 and Th22 numbers and function had significantly lower levels of SIV DNA, both during ART and after ART interruption, than animals with fewer and/or less functional Th17 and Th22 cells [125]. Thus, colorectal IL-17 and -22 producing CD4<sup>+</sup> T cells may be regulators and predictors of HIV/SIV viral persistence [125].

### 3.4. Gastrointestinal B Cells

Although B cells are quite abundant in mucosal tissues, few reports have addressed the effects of HIV-1 infection on mucosal B-cell populations. Hypergammaglobulinemia and

polyclonal B-cell activation were reported early in the HIV-1 epidemic [126, 127], and increased B-cell apoptosis has been detected in lymphoid tissues [128]. Histology has revealed loss of germinal center structure in lymph nodes during acute HIV-1 infection, a finding that was extended to germinal centers in terminal ileum Peyer's patches, with 88% of follicles exhibiting B- or T-cell apoptosis and follicular lysis [129].

There is a paucity of HIV-1-specific IgA production in plasma and mucosal secretions in response to HIV-1 and SIV infection [130–132]; the reasons for this are currently unclear [133]. One hypothesis relates to the ability of HIV-1 Nef to perturb B-cell signaling and block immunoglobulin class switching [134, 135]. The proposed mechanism involves the formation of actin-driven cellular conduits that allow Nef to move from infected macrophages and dendritic cells into uninfected germinal center B cells. Accumulation of Nef in germinal centers has been associated with reduced expression of activation-induced cytidine deaminase (AID), an enzyme required for class switching [135].

Plasmablasts are cycling (Ki67<sup>+</sup>) B cells that secrete immunoglobulins; they represent approximately 1 to 3% of circulating B cells in healthy individuals and are less differentiated than the mature antibody-secreting cells known as plasma cells [136]. In healthy individuals, blood plasmablasts produce mainly IgA [137]. In HIV-1 infection, the number and percentage of circulating plasmablasts are elevated compared to controls, and the proportion that are IgA-secreting is reduced [136]. However, most of these circulating plasmablasts likely arise from polyclonal B-cell activation and are not HIV-1-specific, which may partially explain why the antibody response fails to control HIV [136].

The bias towards IgG-secreting plasma cells reported in HIV-1 disease is not unique to HIV-1 infection, but also reportedly occurs in other conditions, such as Crohn's disease and chronic granulomatous disease (CGD), that lead to mucosal inflammation [138]. When intestinal inflammation is present, IgG-secreting plasma cells express increased CXCR4 and decreased CCR10; and the frequency of CXCR4<sup>+</sup>/IgG<sup>+</sup> plasma cells in the GI mucosa is associated with the severity of the inflammation [138]. These findings suggest that mucosal B cells may play an important role in mucosal immune cell homeostasis and the response to gut inflammation.

### 3.5. Beyond T and B Cells: Innate Effector Subsets

**3.5.1. Natural Killer (NK) Cells**—Although there have been extensive studies of NK cells in peripheral blood in the setting of HIV infection, few studies have focused on the role of mucosal NK cells. Mucosal NK cells are often characterized as belonging to the ILC1 group of innate lymphocytes [139], described in the section below. Mela and colleagues [140] reported depletion of NK subsets in the colonic lamina propria of viremic HIV<sup>+</sup> subjects, and partial reconstitution of these cells following ART in subjects with undetectable viremia. Taborda and colleagues [141] detected reduced expression of the activation marker CD69 on colonic NK cells in a cohort of Colombian HIV controllers compared to typical progressors. Sips and colleagues [142] identified two distinct populations of colonic NKp46<sup>+</sup> NK cells: the first and more abundant subset in both HIV<sup>+</sup> and HIV<sup>-</sup> individuals was localized to the HIV- intraepithelial region; the second was localized to the lamina propria. There was a general trend towards decreased frequencies of

both subsets in untreated HIV<sup>+</sup> individuals with high viral HIV-/loads, as compared to HIV<sup>-</sup> subjects. Intriguingly, HIV controllers expressing protective KIR/HLA genotypes showed a trend towards higher numbers of intraepithelial NK cells than controllers lacking protective genotypes, suggesting a possible role for these cells in immune control of HIV [142].

Despite species-specific differences, given the difficulties inherent in obtaining fresh mucosal tissue from human subjects, several groups have focused on elucidating mucosal NK cells in the SIVmac model system [143, 144]. Reeves and colleagues [145] demonstrated that chronic SIVmac infection induces expansion of blood NK cells expressing  $\alpha 4\beta 7$  integrin and lacking CCR7; these cells are presumably trafficking towards the gastrointestinal mucosa and away from lymph nodes. Two distinct NK subsets were described in SIV-infected macaques: NKG2A<sup>+</sup> NK cells, cytotoxic and IFN $\gamma$ -producing, were distributed widely in blood and tissues; in contrast, NKp44<sup>+</sup> NK cells, non-cytotoxic and predominantly cytokine-producing (IL-22 and IL-17), were restricted to mucosal tissues [146]. Subsequent work led to re-classification of the NKp44<sup>+</sup> subset as ILC3 cells [147]. In chronic SIV infection, this NKp44<sup>+</sup> population (now designated ILC3) was depleted from mucosal tissues and/or adopted a modified functional profile, with increased IFN $\gamma$ , reduced IL-17 production and increased cytotoxic potential [146, 148]. These functional alterations were correlated with increased mucosal expression of inflammatory mediators such as IDO [146]. ART increased the frequency of rectal NKG2A<sup>+</sup> NK and peripheral blood CD16<sup>+</sup> NK cells, but did not restore a normal frequency of rectal NKp44<sup>+</sup> cells [148].

NK cells expressing CXCR6 are recruited to the liver via the chemokine CXCL16. A recent study by Evans and colleagues [149] suggested that translocation of microbial products during chronic SIV infection leads to increased production of CXCL16 by liver myeloid dendritic cells (mDC). This, in turn, leads to recruitment of hypercytotoxic NK cells that contribute to local inflammation and liver damage.

**3.5.2. Innate Lymphocytes (ILC)**—Until recently, little was known of the relatively rare populations of mucosal ILCs and how they respond to infection. Within the past 5 years, several studies have elucidated the various subsets of ILCs and their distinct roles in tissue repair and host defense. Although they lack a T- cell receptor, ILCs share phenotypic and functional properties with classical, MHC-restricted T- cell subsets; this similarity is reflected in their nomenclature. ILCs have been grouped into three categories: ILC1, 2 and 3. Human ILC1 share many features with Th1 cells and/or tissue resident memory T cells [139, 150], such as production of IFN $\gamma$  in response to IL-12 and IL-15, intraepithelial location and evidence of TGF $\beta$  imprinting, and have been described by some as an innate counterpart of T<sub>RM</sub> [151]. NK cells fall within the ILC1 designation [139]. ILC2, with properties similar to Th2 T cells, express transcription factor GATA-3 and secrete IL-5 and IL-13. ILC2 play important roles in the host response to helminthes, as well as in the pathogenesis of allergy and asthma [152, 153]. ILC3, analogous to Th17 cells, express ROR $\gamma$ t, and secrete cytokines IL-17 and IL-22. ILC3 are important for the host response to extracellular bacteria and for regulation of microbiota [154]. Group 3 ILCs also encompass lymphoid tissue inducer cells (LTi), which are essential for the development of lymphoid tissues during embryogenesis, and continue to regulate lymphoid architecture after birth [155–158].

Despite certain functional similarities to CD4<sup>+</sup> T-cell subsets, ILCs do not express receptors for HIV/SIV and are not infectable by either virus [159]. Nevertheless, ILC3 are progressively depleted from the GI tract in SIVmac-infected rhesus macaques [147, 160, 161] and in humanized mice infected with HIV-1 [162]. ILC3 in rhesus macaques, identified as CD3<sup>-</sup>/NKp44<sup>+</sup> and producing IL-17, undergo significant depletion in colonic and jejunal lamina propria during acute infection [147]. Apoptosis appears to be the primary mechanism driving depletion, as these cells increase their expression of activated caspase-3 by greater than 100-fold by 14 days post-infection [147]. Although NKp44<sup>+</sup> ILCs are normally noncytotoxic, under inflammatory conditions they can acquire cytotoxic potential [146]; during chronic SIV infection these cells reportedly increased intracellular perforin expression by 4-fold in colon, jejunum and mesenteric lymph nodes, modulating their phenotype towards that of the ILC1 subset [147].

Studying ILC dynamics in peripheral blood, Kløverpris and colleagues [163] demonstrated rapid loss of all ILC subsets during acute HIV-1 infection concomitant with an increase in plasma I-FABP, a marker of immune activation. This depletion appeared to be mediated by apoptotic pathways, a finding supported by *in vitro* studies showing that rhesus ILC3 could be induced to undergo apoptosis by microbial products signaling through the TLR2 and/or TLR4 pathways [160]. Blood ILCs could be preserved by initiation of ART during acute, but not chronic, HIV-1 infection [163]. Much work remains in order to fully elucidate the significance of ILC subsets, their role in intestinal homeostasis, and their capacity for plasticity, as well as clarification of differences in ILC dynamics between the SIVmac/rhesus macaque model and human HIV-1 infection.

**3.5.3. NKT and Mucosa-associated Invariant T Cells (MAIT)**—NKT and MAIT cells are T cells with limited receptor diversity that are considered to exhibit properties of both the innate and adaptive immune system: like adaptive immune cells (e.g., conventional T and B cells), they express antigen receptors generated through V(D)J recombination; however, like ILCs, they recognize a limited range of foreign antigens and respond quickly to challenge [164, 165]. NKT cells express certain receptors typical of NK cells, including CD161 in humans and NK1.1 in mice. In humans, most NKT cells express a T-cell receptor (TCR) that includes V $\alpha$ 24 co-expressed with J $\alpha$ 18 and V $\beta$ 11 [164]. Unlike conventional T cells, whose TCRs recognize short peptides presented by MHC class I or II, NKT cells recognize glycolipid antigens presented by the nonclassical MHC-like molecule CD1d. These glycolipids include the model antigen  $\alpha$ -galactosylceramide ( $\alpha$ GalCer);  $\alpha$ -glycuronylceramides from the cell walls of certain Gram-negative, LPS-negative bacteria; and the mammalian glycosphingolipid isoglobotrihexosylceramide [164]. CD1d is expressed on antigen-presenting cells, including DC and B cells. Upon TCR stimulation, NKT cells respond rapidly by producing IFN $\gamma$  and IL-4. A role for NKT cells has been suggested in a wide range of conditions including autoimmune diseases, allergy, atherosclerosis, and infectious diseases including bacterial, parasitic and viral infections [164]. Accordingly, while these cells do not directly respond to HIV-1 antigens, their loss and/or dysregulation in HIV-1 infection may have important consequences for the immune system.

NKT cells can be subdivided into CD4<sup>+</sup> and CD4<sup>-</sup> subsets with differential expression of tissue homing receptors; both subsets frequently express CXCR4 and CCR5 [166].

Peripheral CD4<sup>+</sup> NKT cells, particularly those expressing CD62L, are lost in viremic HIV-1 infection [166–168], but may be expanded upon treatment with IL-2 [169]. Reconstitution of blood NKT is delayed after effective ART [170], and those NKT cells that persist during chronic HIV infection display functional impairment and elevated expression of the exhaustion marker PD-1 [171]. In addition, the HIV-1 protein Vpu can interfere with dendritic cell-mediated presentation of lipid antigens to NKT cells [172]. In the GI tract, HIV infection leads to preferential loss of CD4<sup>+</sup> NKT cells, and the extent of NKT loss is related to the level of systemic immune activation [173].

Like NKT cells, human mucosa-associated invariant T cells (MAIT) express CD161 and a semi-invariant T-cell receptor, in this case usually encoded by V $\alpha$ 7.2 and V $\beta$ 2 or 13. The majority of human MAIT cells are CD8<sup>+</sup>, expressing either CD8 $\alpha\alpha$  or  $\alpha\beta$ , with minor populations of CD4<sup>+</sup> or double negative MAITs [174]. MAIT cells are present at relatively high frequency in mucosal tissues, including the GI tract and liver, and recognize microbial vitamin metabolites presented by the nonclassical MHC-like molecule MR1 [175]. Upon stimulation, MAIT cells express both proinflammatory and tissue protective cytokines including IFN $\gamma$ , TNF $\alpha$ , IL-17 and IL-22.

Although MAIT cells are predominantly CD8<sup>+</sup>, rather than CD4<sup>+</sup>, and therefore not directly infected by HIV, MAIT cells are significantly depleted from blood in viremic HIV-1 infection [174, 176–179], with an apparently slower decline in colorectal mucosa [174]. The functionality of remaining MAITs is reduced, with decreased cytokine production and cytotoxicity in response to bacterial antigen [180]; functionality is partially restored following suppressive ART, while MAIT numbers are not [174, 181]. While the underlying cause of MAIT cell loss in HIV-1 infection is not known, data suggest a mechanism driven by persistent exposure to microbial antigens, resulting in exhaustion and apoptosis [181]. There have been some discrepancies between reports in the kinetics of MAIT cell loss and subsequent repopulation on ART, likely due to differences in subject characteristics, mucosal sampling sites, and markers used to identify MAIT cells [174, 176, 182]. Studies in rhesus macaques revealed reduced MAIT cell frequencies in blood, mesenteric lymph nodes, and bronchoalveolar lavage of SIV<sub>mac</sub>-infected macaques compared to uninfected controls [183]. However, no significant differences between groups were detected in jejunum or liver, and increased expression of Ki67 in SIV<sub>mac</sub>-infected macaques suggested greater MAIT cell proliferation and turnover compared to uninfected macaques in this particular study [183]. Further work is needed to fully elucidate the role of MAIT cells in HIV/SIV infection, the mechanism(s) driving their loss and repopulation, and consequences of their loss or dysfunction for immune control of other microbes.

**3.5.4. Gamma Delta T Cells**—In humans, GI IELs are predominantly CD8<sup>+</sup>, with the majority expressing the TCR  $\alpha$  and  $\beta$  chains [184]. A minority of human intestinal IEL are  $\gamma\delta$  T cells, although this subset is far more abundant in mice [185, 186]. Nevertheless,  $\gamma\delta$  T cells, also present in other mucosal tissues and skin, play important roles in immune regulation, tumor surveillance, and innate immune responses to pathogens. They can recognize three sets of stimuli: TCR ligands, including microbial and endogenous phosphoantigens; stress-induced ligands that engage activating natural killer receptors (NKR) such as NKG2D; and/or pathogen-associated molecular patterns (PAMPs) that are

recognized by pattern recognition receptors (PRR) such as Toll-like receptors and Dectin-1 [187]. TCR  $\gamma\delta$  cells can produce a wide range of cytokines, including those typically associated with Th1 (IFN $\gamma$ ), Th2 (IL-4, and IL-13), and Th17 (IL-17) subsets; they can also exhibit cytotoxicity mediated by FAS, TRAIL-R, and the perforin/granzyme pathway [187]. This broad range of functions can be explained by the existence of multiple TCR- $\gamma\delta$  subsets, by developmental programming, and to some extent by functional plasticity. In human peripheral blood, the V $\gamma$ 9V $\delta$ 2 (also referred to as V $\gamma$ 2V $\delta$ 2 [188]) variable regions predominate; however, in the GI the dominant population expresses V $\delta$ 1 [187]. V $\delta$ 1 T cells recognize primarily MICA and MICB, which are stress-induced molecules expressed on intestinal epithelial cells [189]. In HIV infection, absolute numbers of TCR- $\gamma\delta$  cells are increased, and activated V $\delta$ 1-expressing cells are expanded in blood and lungs, causing an inversion of the normal V $\delta$ 1/V $\delta$ 2 ratio in blood [190].

Investigating mucosal  $\gamma\delta$  in HIV-1 infected adults, Brandtzaeg, Nilssen and colleagues [185,191] reported that the  $\gamma\delta$  T-cell subset in duodenal mucosa was enriched in HIV<sup>+</sup> subjects compared to controls, both in percentage and absolute number per unit area. Most were negative for CD8 and >90% expressed the V $\delta$ 1/J $\delta$ 1 TCR segments. Their abundance in the HIV<sup>+</sup> group was inversely related to serum markers of immune activation such as neopterin and  $\beta$ 2 microglobulin, and was lowest in individuals with advanced disease. Subsequently, Poles and colleagues [192] demonstrated expansion of V $\delta$ 1 and contraction of V $\delta$ 2 populations in both rectosigmoid mucosa and blood of HIV<sup>+</sup> individuals; these changes were detected during acute infection, persisted into the chronic phase, and were not reversed by HAART. Although the mechanisms driving these changes are not entirely clear, and most  $\gamma\delta$  T cells do not express CD4, V $\gamma$ 2V $\delta$ 2 T cells that express CCR5 and  $\alpha$ 4 $\beta$ 7 integrin can reportedly be induced to undergo caspase-mediated cell death following binding to soluble or cell-associated gp120 from CCR5-tropic HIV [188].

In a nonhuman primate model of nonprogressive lentiviral disease, sooty mangabey monkeys infected with SIV<sub>mac</sub> did not develop the inversion of the normal V $\delta$ 1/V $\delta$ 2 ratio that is typically observed in blood of HIV<sup>+</sup> individuals [193]. In contrast, rhesus macaques infected with pathogenic SIV<sub>smE543</sub> did exhibit SIV-associated inversion of the V $\delta$ 1/V $\delta$ 2 ratio in blood and tissues, corresponding to expansion of the V $\delta$ 1 subset. Macaque  $\gamma\delta$  T cells were not infected by SIV *in vivo*, and the stimulus responsible for V $\delta$ 1 expansion appeared to be microbial translocation [194]. Intriguingly, peripheral V $\gamma$ 9V $\delta$ 2 T cells were recently reported as a previously unrecognized reservoir of latent HIV infection [195]. The proposed mechanism involves upregulation of CD4 and CCR5 due to pathological immune activation in early HIV infection [195].

#### 4. CONCLUSIONS AND FUTURE DIRECTIONS

Mucosal tissues play a major role in the transmission and pathogenesis of HIV infection, and these tissues house a variety of novel immune cell subsets that are phenotypically and functionally distinct from their counterparts in peripheral blood. While the challenges inherent in tissue sampling have prevented full exploration of mucosal tissues in the past, new technologies in imaging, gene and protein expression analysis and bioinformatics, are now poised to elucidate these novel cell subsets and their interactions in exciting new ways.



New approaches for measuring viral persistence *in situ* will enhance our understanding of host-virus interactions in tissues. Much work remains in order to fully understand the nature of innate and adaptive immunity in tissues, the trafficking patterns of mucosal immune cells, the concept of ‘tissue residency’, and the plasticity of immune cell subsets. Perhaps the greatest challenge remains that of utilizing this information to develop novel preventive and therapeutic strategies that will ultimately lead to eradication of HIV/AIDS.

## Acknowledgments

The author wishes to thank colleagues and collaborators at the University of California at Davis and the University of California at San Francisco, as well as current and past laboratory members, for their contributions to work described in this review, and regrets that many excellent articles could not be cited here due to space limitations. The author’s laboratory is supported by NIH/NIAID R01 AI057020 and NIH/NIDDK R01 DK108350.

## List of Abbreviations

<b>SAMHD1</b>	Sterile alpha motif domain and histidine-aspartic domain-containing protein 1
<b>SLPI</b>	Secretory leukocyte protease inhibitor
<b>HBD2</b>	Human beta-defensin 2
<b>MICA, MICB</b>	MHC class I polypeptide-related sequences A and B

## REFERENCES

- [1]. Chenine AL, Siddappa NB, Kramer VG, et al. Relative transmissibility of an R5 clade C simian-human immunodeficiency virus across different mucosae in macaques parallels the relative risks of sexual HIV-1 transmission in humans via different routes. *J Infect Dis* 2010; 201(8):1155–63. [PubMed: 20214475]
- [2]. Iwasaki A Antiviral immune responses in the genital tract: clues for vaccines. *Nat Rev Immunol* 2010; 10(10):699–711. [PubMed: 20829886]
- [3]. Pudney J, Quayle AJ, Anderson DJ. Immunological microenvironments in the human vagina and cervix: mediators of cellular immunity are concentrated in the cervical transformation zone. *Biol Reprod* 2005; 73(6):1253–63. [PubMed: 16093359]
- [4]. Grande G, Milardi D, Vincenzoni F, et al. Proteomic characterization of the qualitative and quantitative differences in cervical mucus composition during the menstrual cycle. *Mol Biosyst* 2015; 11(6):1717–25. [PubMed: 25959140]
- [5]. Lee DC, Hassan SS, Romero R, et al. Protein profiling underscores immunological functions of uterine cervical mucus plug in human pregnancy. *J Proteomics* 2011; 74(6):817–28. [PubMed: 21362502]
- [6]. Hu J, Gardner MB, Miller CJ. Simian immunodeficiency virus rapidly penetrates the cervicovaginal mucosa after intravaginal inoculation and infects intraepithelial dendritic cells. *J Virol* 2000; 74(13):6087–95. [PubMed: 10846092]
- [7]. Zhang Z, Schuler T, Zupancic M, et al. Sexual transmission and propagation of SIV and HIV in resting and activated CD4+ T cells. *Science* 1999; 286(5443):1353–7. [PubMed: 10558989]
- [8]. Miller CJ, Alexander NJ, Vogel P, Anderson J, Marx PA. Mechanism of genital transmission of SIV: a hypothesis based on transmission studies and the location of SIV in the genital tract of chronically infected female rhesus macaques. *J Med Primatol* 1992; 21(2–3):64–8. [PubMed: 1433268]

- [9]. Padian NS, van der Straten A, Ramjee G, et al. Diaphragm and lubricant gel for prevention of HIV acquisition in southern African women: a randomised controlled trial. *Lancet* 2007; 370(9583): 251–61. [PubMed: 17631387]
- [10]. Ballweber L, Robinson B, Kreger A., et al. Vaginal Langerhans cells nonproductively transporting HIV-1 mediate infection of T cells. *J Virol* 2011; 85(24):13443–7. [PubMed: 21976645]
- [11]. Hladik F, McElrath MJ. Setting the stage: host invasion by HIV. *Nat Rev Immunol* 2008; 8(6): 447–57. [PubMed: 18469831]
- [12]. Hladik F, Sakchalathorn P, Ballweber L, et al. Initial events in establishing vaginal entry and infection by human immunodeficiency virus type-1. *Immunity* 2007; 26(2):257–70. [PubMed: 17306567]
- [13]. Shen R, Kappes JC, Smythies LE, Richter HE, Novak L, Smith PD. Vaginal myeloid dendritic cells transmit founder HIV-1. *J Virol* 2014; 88(13):7683–8. [PubMed: 24741097]
- [14]. Shen R, Richter HE, Smith PD. Interactions between HIV-1 and mucosal cells in the female reproductive tract. *Am J Reprod Immunol* 2014; 71(6):608–17. [PubMed: 24689653]
- [15]. Stieh DJ, Matias E, Xu H, et al. Th17 cells are preferentially infected very early after vaginal transmission of SIV in macaques. *Cell Host Microbe* 2016; 19(4):529–40. [PubMed: 27078070]
- [16]. Rodriguez-Garcia M, Barr FD, Crist SG, Fahey JV, Wira CR. Phenotype and susceptibility to HIV infection of CD4+ Th17 cells in the human female reproductive tract. *Mucosal Immunol* 2014; 7(6):1375–85. [PubMed: 24759207]
- [17]. Shanmugasundaram U, Critchfield JW, Pannell J., et al. Phenotype and functionality of CD4+ and CD8+ T cells in the upper reproductive tract of healthy premenopausal women. *Am J Reprod Immunol* 2014; 71(2):95–108. [PubMed: 24313954]
- [18]. Stieh DJ, Maric D, Kelley ZL, et al. Vaginal challenge with an SIV-based dual reporter system reveals that infection can occur throughout the upper and lower female reproductive tract. *PLoS Pathog* 2014; 10(10):e1004440. [PubMed: 25299616]
- [19]. Shen Z, Rodriguez-Garcia M, Ochsenbauer C, Wira CR. Characterization of immune cells and infection by HIV in human ovarian tissues. *Am J Reprod Immunol* 2017; 78(1):e12687.
- [20]. Louissaint NA, Fuchs EJ, Bakshi RP., et al. Distribution of cell-free and cell-associated HIV surrogates in the female genital tract after simulated vaginal intercourse. *J Infect Dis* 2012; 205(5):725–32. [PubMed: 22279121]
- [21]. El Costa H, Quillay H, Marlin R, et al. The local environment orchestrates mucosal decidual macrophage differentiation and substantially inhibits HIV-1 replication. *Mucosal Immunol* 2016; 9(3):634–46. [PubMed: 26349662]
- [22]. Quillay H, El Costa H, Marlin R, , et al. Distinct characteristics of endometrial and decidual macrophages and regulation of their permissivity to HIV-1 infection by SAMHD1. *J Virol* 2015; 89(2):1329–39. [PubMed: 25392215]
- [23]. Rahangdale L, De Paris K, Kashuba AD., et al. Immunologic, virologic, and pharmacologic characterization of the female upper genital tract in HIV-infected women. *J Acquir Immune Defic Syndr* 2015; 68(4):420–4. [PubMed: 25501615]
- [24]. Reynolds MR, Rakasz E, Skinner PJ, et al. CD8+ T-lymphocyte response to major immunodominant epitopes after vaginal exposure to simian immunodeficiency virus: too late and too little. *J Virol* 2005; 79(14):9228–35. [PubMed: 15994817]
- [25]. Li Q, Skinner PJ, Ha SJ, et al. Visualizing antigen-specific and infected cells in situ predicts outcomes in early viral infection. *Science* 2009; 323(5922):1726–9. [PubMed: 19325114]
- [26]. Masopust D Developing an HIV cytotoxic T-lymphocyte vaccine: issues of CD8 T-cell quantity, quality and location. *J Intern Med* 2009; 265(1):125–37. [PubMed: 19093965]
- [27]. Lohman BL, Miller CJ, McChesney MB. Antiviral cytotoxic T lymphocytes in vaginal mucosa of simian immunodeficiency virus-infected rhesus macaques. *J Immunol* 1995; 155(12):5855–60. [PubMed: 7499875]
- [28]. Musey L, Ding Y, Cao J, et al. Ontogeny and specificities of mucosal and blood human immunodeficiency virus type 1-specific CD8(+) cytotoxic T lymphocytes. *J Virol* 2003; 77(1): 291–300. [PubMed: 12477834]

- [29]. Musey L, Hu Y, Eckert L, Christensen M, Karchmer T, McElrath MJ. HIV-1 induces cytotoxic T lymphocytes in the cervix of infected women. *J Exp Med* 1997; 185(2):293–303. [PubMed: 9016878]
- [30]. Bere A, Denny L, Burgers WA, Passmore JA. Polyclonal expansion of cervical cytobrush-derived T cells to investigate HIV-specific responses in the female genital tract. *Immunology* 2010; 130(1):23–33. [PubMed: 20201983]
- [31]. Bere A, Denny L, Hanekom W, Burgers WA, Passmore JA. Comparison of polyclonal expansion methods to improve the recovery of cervical cytobrush-derived T cells from the female genital tract of HIV-infected women. *J Immunol Methods* 2010; 354(1–2):68–79. [PubMed: 20149794]
- [32]. McKinnon LR, Hughes SM, De Rosa SC, et al. Optimizing viable leukocyte sampling from the female genital tract for clinical trials: an international multi-site study. *PLoS One* 2014; 9(1):e85675. [PubMed: 24454917]
- [33]. Gumbi PP, Nkwanyana NN, Bere A, et al. Impact of mucosal inflammation on cervical human immunodeficiency virus (HIV-1)-specific CD8 T-cell responses in the female genital tract during chronic HIV infection. *J Virol* 2008; 82(17):8529–36. [PubMed: 18562528]
- [34]. Ferre AL, Hunt PW, Critchfield JW, et al. Mucosal immune responses to HIV-1 in elite controllers: a potential correlate of immune control. *Blood* 2009; 113(17):3978–89. [PubMed: 19109229]
- [35]. Makedonas G, Betts MR. Polyfunctional analysis of human t cell responses: importance in vaccine immunogenicity and natural infection. *Springer Semin Immunopathol* 2006; 28(3):209–19. [PubMed: 16932955]
- [36]. Bere A, Denny L, Naicker P, Burgers WA, Passmore JA. HIV-specific T-cell responses detected in the genital tract of chronically HIV-infected women are largely monofunctional. *Immunology* 2013; 139(3):342–51. [PubMed: 23374084]
- [37]. Nkwanyana NN, Gumbi PP, Roberts L, et al. Impact of human immunodeficiency virus 1 infection and inflammation on the composition and yield of cervical mononuclear cells in the female genital tract. *Immunology* 2009; 128(1 Suppl):e746–57. [PubMed: 19740336]
- [38]. Mkhize NN, Gumbi PP, Liebenberg LJ, et al. Persistence of genital tract T cell responses in HIV-infected women on highly active antiretroviral therapy. *J Virol* 2010; 84(20):10765–72. [PubMed: 20686039]
- [39]. Eschenbach DA, Thwin SS, Patton DL, et al. Influence of the normal menstrual cycle on vaginal tissue, discharge, and microflora. *Clin Infect Dis* 2000; 30(6):901–7. [PubMed: 10852812]
- [40]. Poonia B, Walter L, Dufour J, Harrison R, Marx PA, Veazey RS. Cyclic changes in the vaginal epithelium of normal rhesus macaques. *J Endocrinol* 2006; 190(3):829–35. [PubMed: 17003283]
- [41]. Wira CR, Rodriguez-Garcia M, Patel MV. The role of sex hormones in immune protection of the female reproductive tract. *Nat Rev Immunol* 2015; 15(4):217–30. [PubMed: 25743222]
- [42]. Morrison CS, Chen PL, Kwok C, et al. Hormonal contraception and the risk of HIV acquisition: an individual participant data meta-analysis. *PLoS Med* 2015; 12(1):e1001778. [PubMed: 25612136]
- [43]. Ralph LJ, McCoy SI, Shiu K, Padian NS. Hormonal contraceptive use and women’s risk of HIV acquisition: a meta-analysis of observational studies. *Lancet Infect Dis* 2015; 15(2):181–9. [PubMed: 25578825]
- [44]. Achilles SL, Creinin MD, Stoner KA, Chen BA, Meyn L, Hillier SL. Changes in genital tract immune cell populations after initiation of intrauterine contraception. *Am J Obstet Gynecol* 2014; 211(5):489 e1–9. [PubMed: 24834865]
- [45]. Achilles SL, Hillier SL. The complexity of contraceptives: understanding their impact on genital immune cells and vaginal microbiota. *AIDS* 2013; 27 Suppl 1:S5–15. [PubMed: 24088684]
- [46]. Byrne EH, Anahtar MN, Cohen KE, et al. Association between injectable progestin-only contraceptives and HIV acquisition and HIV target cell frequency in the female genital tract in South African women: a prospective cohort study. *Lancet Infect Dis* 2016; 16(4):441–8. [PubMed: 26723758]
- [47]. Smith-McCune KK, Hilton JF, Shanmugasundaram U, et al. Effects of depot-medroxyprogesterone acetate on the immune microenvironment of the human cervix and

- endometrium: implications for HIV susceptibility. *Mucosal Immunol* 2017; doi: 10.1038/mi.2016.121.
- [48]. Wira CR, Fahey JV. A new strategy to understand how HIV infects women: identification of a window of vulnerability during the menstrual cycle. *AIDS* 2008; 22(15):1909–17. [PubMed: 18784454]
- [49]. Wira CR, Fahey JV, Rodriguez-Garcia M, Shen Z, Patel MV. Regulation of mucosal immunity in the female reproductive tract: the role of sex hormones in immune protection against sexually transmitted pathogens. *Am J Reprod Immunol* 2014; 72(2):236–58. [PubMed: 24734774]
- [50]. White HD, Yeaman GR, Givan AL, Wira CR. Mucosal immunity in the human female reproductive tract: cytotoxic T lymphocyte function in the cervix and vagina of premenopausal and postmenopausal women. *Am J Reprod Immunol* 1997; 37(1):30–8. [PubMed: 9138451]
- [51]. Yeaman GR, Guyre PM, Fanger MW, et al. Unique CD8+ T cell-rich lymphoid aggregates in human uterine endometrium. *J Leukoc Biol* 1997; 61(4):427–35. [PubMed: 9103229]
- [52]. White HD, Musey LK, Andrews MM, et al. Human immunodeficiency virus-specific and CD3-redredirected cytotoxic T lymphocyte activity in the human female reproductive tract: lack of correlation between mucosa and peripheral blood. *J Infect Dis* 2001; 183(6):977–83. [PubMed: 11237817]
- [53]. Veazey RS, DeMaria M, Chalifoux LV, et al. Gastrointestinal tract as a major site of CD4+ T cell depletion and viral replication in SIV infection. *Science* 1998; 280(5362):427–31. [PubMed: 9545219]
- [54]. Brenchley JM, Douek DC. The mucosal barrier and immune activation in HIV pathogenesis. *Curr Opin HIV AIDS* 2008; 3(3):356–61. [PubMed: 19372990]
- [55]. Brenchley JM, Price DA, Schacker TW, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med* 2006; 12(12):1365–71. [PubMed: 17115046]
- [56]. Fritsch H, Zehm S, Illig R, Moser P, Aigner F. New insights into the development and differentiation of the human anorectal epithelia. Are there clinical consequences? *Int J Colorectal Dis* 2010; 25(10):1231–42. [PubMed: 20563874]
- [57]. McElrath MJ, Smythe K, Randolph-Habecker J, et al. Comprehensive assessment of HIV target cells in the distal human gut suggests increasing HIV susceptibility toward the anus. *J Acquir Immune Defic Syndr* 2013; 63(3):263–71. [PubMed: 23392465]
- [58]. Poles MA, Elliott J, Taing P, Anton PA, Chen IS. A preponderance of CCR5(+) CXCR4(+) mononuclear cells enhances gastrointestinal mucosal susceptibility to human immunodeficiency virus type 1 infection. *J Virol* 2001; 75(18):8390–9. [PubMed: 11507184]
- [59]. Preza GC, Tanner K, Elliott J, Yang OO, Anton PA, Ochoa MT. Antigen-presenting cell candidates for HIV-1 transmission in human distal colonic mucosa defined by CD207 dendritic cells and CD209 macrophages. *AIDS Res Hum Retroviruses* 2014; 30(3):241–9. [PubMed: 24134315]
- [60]. Shacklett BL, Anton PA. HIV infection and gut mucosal immune function: updates on pathogenesis with implications for management and intervention. *Curr Infect Dis Rep* 2010; 12(1):19–27. [PubMed: 20174448]
- [61]. Louissaint NA, Nimmagadda S, Fuchs EJ, et al. Distribution of cell-free and cell-associated HIV surrogates in the colon after simulated receptive anal intercourse in men who have sex with men. *J Acquir Immune Defic Syndr* 2012; 59(1):10–7. [PubMed: 21937920]
- [62]. Baggaley RF, White RG, Boily MC. HIV transmission risk through anal intercourse: systematic review, meta-analysis and implications for HIV prevention. *Int J Epidemiol* 2010; 39(4):1048–63. [PubMed: 20406794]
- [63]. Gorbach PM, Manhart LE, Hess KL, Stoner BP, Martin DH, Holmes KK. Anal intercourse among young heterosexuals in three sexually transmitted disease clinics in the United States. *Sex Transm Dis* 2009; 36(4):193–8. [PubMed: 19265740]
- [64]. Shacklett BL, Beadle TJ, Pacheco PAG, et al. Isolation of cytomegalovirus-specific cytotoxic T-lymphocytes from gut-associated lymphoid tissue (GALT) of HIV type 1-infected subjects. *AIDS Res Hum Retroviruses* 2000; 16(12):1157–62. [PubMed: 10954891]

- [65]. Shacklett BL, Beadle TJ, Pacheco PA, et al. Characterization of HIV-1-specific cytotoxic T lymphocytes expressing the mucosal lymphocyte integrin CD103 in rectal and duodenal lymphoid tissue of HIV-1-infected subjects. *Virology* 2000; 270(2):317–27. [PubMed: 10792991]
- [66]. Shacklett BL, Cox CA, Sandberg JK, Stollman NH, Jacobson MA, Nixon DF. Trafficking of human immunodeficiency virus type 1-specific CD8+ T cells to gut-associated lymphoid tissue during chronic infection. *J Virol* 2003; 77(10):5621–31. [PubMed: 12719554]
- [67]. Ibarrondo FJ, Anton PA, Fuerst M, et al. Parallel human immunodeficiency virus type 1-specific CD8+ T-lymphocyte responses in blood and mucosa during chronic infection. *J Virol* 2005; 79(7):4289–97. [PubMed: 15767429]
- [68]. Critchfield JW, Young DH, Hayes TL, et al. Magnitude and complexity of rectal mucosa HIV-1-specific CD8+ T-cell responses during chronic infection reflect clinical status. *PLoS One* 2008; 3(10):e3577. [PubMed: 18974782]
- [69]. Critchfield JW, Lemongello D, Walker DH, et al. Multifunctional human immunodeficiency virus (HIV) gag-specific CD8+ T-cell responses in rectal mucosa and peripheral blood mononuclear cells during chronic HIV type 1 infection. *J Virol* 2007; 81(11):5460–71. [PubMed: 17344302]
- [70]. Ferre AL, Lemongello D, Hunt PW, et al. Immunodominant HIV-specific CD8+ T-cell responses are common to blood and gastrointestinal mucosa, and Gag-specific responses dominate in rectal mucosa of HIV controllers. *J Virol* 2010; 84(19):10354–65. [PubMed: 20668079]
- [71]. Ferre AL, Hunt PW, McConnell DH, et al. HIV controllers with HLA-DRB1\*13 and HLA-DQB1\*06 alleles have strong, polyfunctional mucosal CD4+ T-cell responses. *J Virol* 2010; 84(21):11020–9. [PubMed: 20719952]
- [72]. Fellay J, Ge D, Shianna KV, et al. Common genetic variation and the control of HIV-1 in humans. *PLoS Genet* 2009; 5(12):e1000791. [PubMed: 20041166]
- [73]. International HIVCS, Pereyra F, Jia X, et al. The major genetic determinants of HIV-1 control affect HLA class I peptide presentation. *Science* 2010; 330(6010):1551–7. [PubMed: 21051598]
- [74]. Shacklett BL, Cox CA, Quigley MF, et al. Abundant expression of granzyme A, but not perforin, in granules of CD8+ T cells in GALT: implications for immune control of HIV-1 infection. *J Immunol* 2004; 173(1):641–8. [PubMed: 15210827]
- [75]. Kiniry BE, Ganesh A, Critchfield JW, et al. Predominance of weakly cytotoxic, Tbet<sup>Low</sup>Eomes<sup>Neg</sup> CD8+ T-cells in human gastrointestinal mucosa: implications for HIV infection. *Mucosal Immunol* 2017; 10(4):1008–20. [PubMed: 27827375]
- [76]. Casey KA, Fraser KA, Schenkel JM, et al. Antigen-independent differentiation and maintenance of effector-like resident memory T cells in tissues. *J Immunol* 2012; 188(10):4866–75. [PubMed: 22504644]
- [77]. Rosato PC, Beura LK, Masopust D. Tissue resident memory T cells and viral immunity. *Curr Opin Virol* 2017; 22:44–50. [PubMed: 27987416]
- [78]. Schenkel JM, Fraser KA, Beura LK, Pauken KE, Vezys V, Masopust D. T cell memory. Resident memory CD8 T cells trigger protective innate and adaptive immune responses. *Science* 2014; 346(6205):98–101. [PubMed: 25170049]
- [79]. Steinert EM, Schenkel JM, Fraser KA, et al. Quantifying memory CD8 T cells reveals regionalization of immunosurveillance. *Cell* 2015; 161(4):737–49. [PubMed: 25957682]
- [80]. Mackay LK, Rahimpour A, Ma JZ, et al. The developmental pathway for CD103(+) CD8+ tissue-resident memory T cells of skin. *Nat Immunol* 2013; 14(12):1294–301. [PubMed: 24162776]
- [81]. Mackay LK, Stock AT, Ma JZ, et al. Long-lived epithelial immunity by tissue-resident memory T (TRM) cells in the absence of persisting local antigen presentation. *Proc Natl Acad Sci U S A* 2012; 109(18):7037–42. [PubMed: 22509047]
- [82]. Mackay LK, Wynne-Jones E, Freestone D, et al. T-box transcription factors combine with the cytokines TGF-beta and IL-15 to control tissue-resident memory T cell fate. *Immunity* 2015; 43(6):1101–11. [PubMed: 26682984]
- [83]. Carbone FR, Mackay LK, Heath WR, Gebhardt T. Distinct resident and recirculating memory T cell subsets in non-lymphoid tissues. *Curr Opin Immunol* 2013; 25(3):329–33. [PubMed: 23746791]



- [84]. Mueller SN, Mackay LK. Tissue-resident memory T cells: local specialists in immune defence. *Nat Rev Immunol* 2016; 16(2):79–89. [PubMed: 26688350]
- [85]. Mackay LK, Kallies A. Transcriptional regulation of tissue-resident lymphocytes. *Trends Immunol* 2017; 38(2):94–103. [PubMed: 27939451]
- [86]. Guadalupe M, Reay E, Sankaran S, et al. Severe CD4+ T-cell depletion in gut lymphoid tissue during primary human immunodeficiency virus type 1 infection and substantial delay in restoration following highly active antiretroviral therapy. *J Virol* 2003; 77(21):11708–17. [PubMed: 14557656]
- [87]. Li Q, Duan L, Estes JD, et al. Peak SIV replication in resting memory CD4+ T cells depletes gut lamina propria CD4+ T cells. *Nature* 2005; 434(7037):1148–52. [PubMed: 15793562]
- [88]. Mattapallil JJ, Douek DC, Hill B, Nishimura Y, Martin M, Roederer M. Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV infection. *Nature* 2005; 434(7037):1093–7. [PubMed: 15793563]
- [89]. Mehandru S, Poles MA, Tenner-Racz K, et al. Primary HIV-1 infection is associated with preferential depletion of CD4+ T lymphocytes from effector sites in the gastrointestinal tract. *J Exp Med* 2004; 200(6):761–70. [PubMed: 15365095]
- [90]. Girard A, Vergnon-Miszczycha D, Depince-Berger AE, et al. Brief Report: A high rate of beta7+ gut-homing lymphocytes in HIV-infected immunological nonresponders is associated With poor CD4 T-cell recovery during suppressive HAART. *J Acquir Immune Defic Syndr* 2016; 72(3): 259–65. [PubMed: 27306505]
- [91]. Mavigner M, Cazabat M, Dubois M, et al. Altered CD4+ T cell homing to the gut impairs mucosal immune reconstitution in treated HIV-infected individuals. *J Clin Invest* 2012; 122(1): 62–9. [PubMed: 22156200]
- [92]. Allers K, Puyskens A, Epple HJ, et al. The effect of timing of antiretroviral therapy on CD4+ T-cell reconstitution in the intestine of HIV-infected patients. *Mucosal Immunol* 2016; 9(1):265–74. [PubMed: 26129649]
- [93]. Deleage C, Schuetz A, Alvord WG, et al. Impact of early cART in the gut during acute HIV infection. *JCI Insight* 2016; 1(10):e87065. [PubMed: 27446990]
- [94]. Kok A, Hocqueloux L, Hocini H, et al. Early initiation of combined antiretroviral therapy preserves immune function in the gut of HIV-infected patients. *Mucosal Immunol* 2015; 8(1): 127–40. [PubMed: 24985081]
- [95]. Schuetz A, Deleage C, Sereti I, et al. Initiation of ART during early acute HIV infection preserves mucosal Th17 function and reverses HIV-related immune activation. *PLoS Pathog* 2014; 10(12):e1004543. [PubMed: 25503054]
- [96]. Cecchinato V, Franchini G. Th17 cells in pathogenic simian immunodeficiency virus infection of macaques. *Curr Opin HIV AIDS* 2010; 5(2):141–5. [PubMed: 20543591]
- [97]. Sallusto F, Zielinski CE, Lanzavecchia A. Human Th17 subsets. *Eur J Immunol* 2012; 42(9): 2215–20. [PubMed: 22949319]
- [98]. Ciccone EJ, Greenwald JH, Lee PI, et al. CD4+ T cells, including Th17 and cycling subsets, are intact in the gut mucosa of HIV-1-infected long-term nonprogressors. *J Virol* 2011; 85(12):5880–8. [PubMed: 21471231]
- [99]. Raffatellu M, Santos RL, Verhoeven DE, et al. Simian immunodeficiency virus-induced mucosal interleukin-17 deficiency promotes Salmonella dissemination from the gut. *Nat Med* 2008; 14(4): 421–8. [PubMed: 18376406]
- [100]. Gosselin A, Monteiro P, Chomont N, et al. Peripheral blood CCR4+CCR6+ and CXCR3+CCR6+CD4+ T cells are highly permissive to HIV-1 infection. *J Immunol* 2010; 184(3):1604–16. [PubMed: 20042588]
- [101]. Wacleche VS, Goulet JP, Gosselin A, et al. New insights into the heterogeneity of Th17 subsets contributing to HIV-1 persistence during antiretroviral therapy. *Retrovirology* 2016; 13(1):59. [PubMed: 27553844]
- [102]. Gosselin A, Wiche Salinas TR, Planas D, et al. HIV persists in CCR6+CD4+ T cells from colon and blood during antiretroviral therapy. *AIDS* 2017; 31(1):35–48. [PubMed: 27835617]



- [103]. Christensen-Quick A, Lafferty M, Sun L, Marchionni L, DeVico A, Garzino-Demo A. Human Th17 cells lack HIV-inhibitory RNases and are highly permissive to productive HIV infection. *J Virol* 2016; 90(17):7833–47. [PubMed: 27334595]
- [104]. Belkaid Y, Tarbell K. Regulatory T cells in the control of host-microorganism interactions. *Annu Rev Immunol* 2009; 27:551–89. [PubMed: 19302048]
- [105]. Kanwar B, Favre D, McCune JM. Th17 and regulatory T cells: implications for AIDS pathogenesis. *Curr Opin HIV AIDS* 2010; 5(2):151–7. [PubMed: 20543593]
- [106]. Shaw JM, Hunt PW, Critchfield JW, et al. Increased frequency of regulatory T cells accompanies increased immune activation in rectal mucosae of HIV-positive noncontrollers. *J Virol* 2011; 85(21):11422–34. [PubMed: 21880771]
- [107]. Rueda CM, Velilla PA, Chougnet CA, Rugeles MT. Incomplete normalization of regulatory T-cell frequency in the gut mucosa of Colombian HIV-infected patients receiving long-term antiretroviral treatment. *PLoS One* 2013; 8(8):e71062. [PubMed: 23967152]
- [108]. Presicce P, Shaw JM, Miller CJ, Shacklett BL, Chougnet CA. Myeloid dendritic cells isolated from tissues of SIV-infected Rhesus macaques promote the induction of regulatory T cells. *AIDS* 2012; 26(3):263–73. [PubMed: 22095196]
- [109]. Moreno-Fernandez ME, Presicce P, Chougnet CA. Homeostasis and function of regulatory T cells in HIV/SIV infection. *J Virol* 2012; 86(19):10262–9. [PubMed: 22811537]
- [110]. Favre D, Lederer S, Kanwar B, et al. Critical loss of the balance between Th17 and T regulatory cell populations in pathogenic SIV infection. *PLoS Pathog* 2009; 5(2):e1000295. [PubMed: 19214220]
- [111]. Mellor AL, Munn DH. IDO expression by dendritic cells: tolerance and tryptophan catabolism. *Nat Rev Immunol* 2004; 4(10):762–74. [PubMed: 15459668]
- [112]. Baban B, Chandler PR, Sharma MD, et al. IDO activates regulatory T cells and blocks their conversion into Th17-like T cells. *J Immunol* 2009; 183(4):2475–83. [PubMed: 19635913]
- [113]. Favre D, Mold J, Hunt PW, et al. Tryptophan catabolism by indoleamine 2,3-dioxygenase 1 alters the balance of TH17 to regulatory T cells in HIV disease. *Sci Transl Med* 2010; 2(32):32ra36.
- [114]. Jenabian MA, El-Far M, Vyboh K, et al. Immunosuppressive tryptophan catabolism and gut mucosal dysfunction following early HIV infection. *J Infect Dis* 2015; 212(3):355–66. [PubMed: 25616404]
- [115]. Klatt NR, Estes JD, Sun X, et al. Loss of mucosal CD103+ DCs and IL-17+ and IL-22+ lymphocytes is associated with mucosal damage in SIV infection. *Mucosal Immunol* 2012; 5(6):646–57. [PubMed: 22643849]
- [116]. Loiseau C, Requena M, Mavigner M, et al. CCR6(–) regulatory T cells blunt the restoration of gut Th17 cells along the CCR6-CCL20 axis in treated HIV-1-infected individuals. *Mucosal Immunol* 2016; 9(5):1137–50. [PubMed: 26883727]
- [117]. Ortiz AM, Klase ZA, DiNapoli SR, et al. IL-21 and probiotic therapy improve Th17 frequencies, microbial translocation, and microbiome in ARV-treated, SIV-infected macaques. *Mucosal Immunol* 2016; 9(2):458–67. [PubMed: 26286233]
- [118]. Vujkovic-Cvijin I, Swainson LA, Chu SN, et al. Gut-resident lactobacillus abundance associates with IDO1 inhibition and Th17 dynamics in SIV-infected macaques. *Cell Rep* 2015; 13(8):1589–97. [PubMed: 26586432]
- [119]. Sonnenberg GF, Fouser LA, Artis D. Border patrol: regulation of immunity, inflammation and tissue homeostasis at barrier surfaces by IL-22. *Nat Immunol* 2011; 12(5):383–90. [PubMed: 21502992]
- [120]. Duhon T, Geiger R, Jarrossay D, Lanzavecchia A, Sallusto F. Production of interleukin 22 but not interleukin 17 by a subset of human skin-homing memory T cells. *Nat Immunol* 2009; 10(8):857–63. [PubMed: 19578369]
- [121]. Eyerich S, Eyerich K, Pennino D, et al. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* 2009; 119(12):3573–85. [PubMed: 19920355]

- [122]. Trifari S, Kaplan CD, Tran EH, Crellin NK, Spits H. Identification of a human helper T cell population that has abundant production of interleukin 22 and is distinct from T(H)-17, T(H)1 and T(H)2 cells. *Nat Immunol* 2009; 10(8):864–71. [PubMed: 19578368]
- [123]. Canary LA, Vinton CL, Morcock DR, et al. Rate of AIDS progression is associated with gastrointestinal dysfunction in simian immunodeficiency virus-infected pigtail macaques. *J Immunol* 2013; 190(6):2959–65. [PubMed: 23401593]
- [124]. Xu H, Wang X, Veazey RS. Th17 cells coordinate with Th22 cells in maintaining homeostasis of intestinal tissues and both are depleted in SIV-infected macaques. *J AIDS Clin Res* 2014; 5(5): 302. [PubMed: 25364618]
- [125]. Ryan ES, Micci L, Fromentin R, et al. Loss of function of intestinal IL-17 and IL-22 producing cells contributes to inflammation and viral persistence in SIV-infected rhesus macaques. *PLoS Pathog* 2016; 12(2):e1005412. [PubMed: 26829644]
- [126]. Lane HC, Masur H, Edgar LC, Whalen G, Rook AH, Fauci AS. Abnormalities of B-cell activation and immunoregulation in patients with the acquired immunodeficiency syndrome. *N Engl J Med* 1983; 309(8):453–8. [PubMed: 6224088]
- [127]. Shirai A, Cosentino M, Leitman-Klinman SF, Klinman DM. Human immunodeficiency virus infection induces both polyclonal and virus-specific B cell activation. *J Clin Invest* 1992; 89(2): 561–6. [PubMed: 1737846]
- [128]. Moir S, Fauci AS. Pathogenic mechanisms of B-lymphocyte dysfunction in HIV disease. *J Allergy Clin Immunol* 2008; 122(1):12–9; quiz 20–1. [PubMed: 18547629]
- [129]. Levesque MC, Moody MA, Hwang KK, et al. Polyclonal B cell differentiation and loss of gastrointestinal tract germinal centers in the earliest stages of HIV-1 infection. *PLoS Med* 2009; 6(7):e1000107. [PubMed: 19582166]
- [130]. Kotler DP, Scholes JV, Tierney AR. Intestinal plasma cell alterations in acquired immunodeficiency syndrome. *Dig Dis Sci* 1987; 32(2):129–38. [PubMed: 3542444]
- [131]. Raux M, Finkielstzajn L, Salmon-Ceron D, et al. Comparison of the distribution of IgG and IgA antibodies in serum and various mucosal fluids of HIV type 1-infected subjects. *AIDS Res Hum Retroviruses* 1999; 15(15):1365–76. [PubMed: 10515152]
- [132]. Schneider T, Zippel T, Schmidt W, Zeitz M, Ullrich R. Secretory immunity in HIV infection. *Pathobiology* 1998; 66(3–4):131–8. [PubMed: 9693313]
- [133]. Chaoul N, Burelout C, Peruchon S, et al. Default in plasma and intestinal IgA responses during acute infection by simian immunodeficiency virus. *Retrovirology* 2012; 9:43. [PubMed: 22632376]
- [134]. Qiao X, He B, Chiu A, Knowles DM, Chadburn A, Cerutti A. Human immunodeficiency virus 1 Nef suppresses CD40-dependent immunoglobulin class switching in bystander B cells. *Nat Immunol* 2006; 7(3):302–10. [PubMed: 16429138]
- [135]. Xu W, Santini PA, Sullivan JS, et al. HIV-1 evades virus-specific IgG2 and IgA responses by targeting systemic and intestinal B cells via long-range intercellular conduits. *Nat Immunol* 2009; 10(9):1008–17. [PubMed: 19648924]
- [136]. Buckner CM, Moir S, Ho J, et al. Characterization of plasmablasts in the blood of HIV-infected viremic individuals: evidence for nonspecific immune activation. *J Virol* 2013; 87(10):5800–11. [PubMed: 23487459]
- [137]. Mei HE, Yoshida T, Sime W, et al. Blood-borne human plasma cells in steady state are derived from mucosal immune responses. *Blood* 2009; 113(11):2461–9. [PubMed: 18987362]
- [138]. Buckner CM, Moir S, Kardava L, et al. CXCR4/IgG-expressing plasma cells are associated with human gastrointestinal tissue inflammation. *J Allergy Clin Immunol* 2014; 133(6):1676–85 e5. [PubMed: 24373354]
- [139]. Walker JA, Barlow JL, McKenzie AN. Innate lymphoid cells--how did we miss them? *Nat Rev Immunol* 2013; 13(2):75–87. [PubMed: 23292121]
- [140]. Mela CM, Steel A, Lindsay J, Gazzard BG, Gotch FM, Goodier MR. Depletion of natural killer cells in the colonic lamina propria of viraemic HIV-1-infected individuals. *AIDS* 2007; 21(16): 2177–82. [PubMed: 18090044]

- [141]. Taborda NA, Gonzalez SM, Alvarez CM, Correa LA, Montoya CJ, Rugeles MT. Higher frequency of NK and CD4+ T-cells in mucosa and potent cytotoxic response in HIV controllers. *PLoS One* 2015; 10(8):e0136292. [PubMed: 26291824]
- [142]. Sips M, Sciaranghella G, Diefenbach T, et al. Altered distribution of mucosal NK cells during HIV infection. *Mucosal Immunol* 2012; 5(1):30–40. [PubMed: 21993602]
- [143]. Hong HS, Rajakumar PA, Billingsley JM, Reeves RK, Johnson RP. No monkey business: why studying NK cells in non-human primates pays off. *Front Immunol* 2013; 4:32. [PubMed: 23423644]
- [144]. Webster RL, Johnson RP. Delineation of multiple subpopulations of natural killer cells in rhesus macaques. *Immunology* 2005; 115(2):206–14. [PubMed: 15885126]
- [145]. Reeves RK, Evans TI, Gillis J, Johnson RP. Simian immunodeficiency virus infection induces expansion of alpha4beta7+ and cytotoxic CD56+ NK cells. *J Virol* 2010; 84(17):8959–63. [PubMed: 20554780]
- [146]. Reeves RK, Rajakumar PA, Evans TI, et al. Gut inflammation and indoleamine deoxygenase inhibit IL-17 production and promote cytotoxic potential in NKp44+ mucosal NK cells during SIV infection. *Blood* 2011; 118(12):3321–30. [PubMed: 21791421]
- [147]. Li H, Richert-Spuhler LE, Evans TI, et al. Hypercytotoxicity and rapid loss of NKp44+ innate lymphoid cells during acute SIV infection. *PLoS Pathog* 2014; 10(12):e1004551. [PubMed: 25503264]
- [148]. Liyanage NP, Gordon SN, Doster MN, et al. Antiretroviral therapy partly reverses the systemic and mucosal distribution of NK cell subsets that is altered by SIVmac(2)(5)(1) infection of macaques. *Virology* 2014; 450–451:359–68.
- [149]. Evans TI, Li H, Schafer JL, et al. SIV-induced translocation of bacterial products in the liver mobilizes myeloid dendritic and natural killer cells associated with liver damage. *J Infect Dis* 2016; 213(3):361–9. [PubMed: 26238685]
- [150]. Fuchs A, Vermi W, Lee JS, et al. Intraepithelial type 1 innate lymphoid cells are a unique subset of IL-12- and IL-15-responsive IFN $\gamma$ -producing cells. *Immunity* 2013; 38(4):769–81. [PubMed: 23453631]
- [151]. Fuchs A, Colonna M. Innate lymphoid cells in homeostasis, infection, chronic inflammation and tumors of the gastrointestinal tract. *Curr Opin Gastroenterol* 2013; 29(6):581–7. [PubMed: 24100718]
- [152]. Mjosberg J, Bernink J, Golebski K, et al. The transcription factor GATA3 is essential for the function of human type 2 innate lymphoid cells. *Immunity* 2012; 37(4):649–59. [PubMed: 23063330]
- [153]. Nussbaum JC, Van Dyken SJ, von Moltke J, et al. Type 2 innate lymphoid cells control eosinophil homeostasis. *Nature* 2013; 502(7470):245–8. [PubMed: 24037376]
- [154]. Serafini N, Voshenrich CA, Di Santo JP. Transcriptional regulation of innate lymphoid cell fate. *Nat Rev Immunol* 2015; 15(7):415–28. [PubMed: 26065585]
- [155]. Hepworth MR, Monticelli LA, Fung TC, et al. Innate lymphoid cells regulate CD4+ T-cell responses to intestinal commensal bacteria. *Nature* 2013; 498(7452):113–7. [PubMed: 23698371]
- [156]. Magri G, Miyajima M, Bascones S, et al. Innate lymphoid cells integrate stromal and immunological signals to enhance antibody production by splenic marginal zone B cells. *Nat Immunol* 2014; 15(4):354–64. [PubMed: 24562309]
- [157]. Qiu J, Guo X, Chen ZM, et al. Group 3 innate lymphoid cells inhibit T-cell-mediated intestinal inflammation through aryl hydrocarbon receptor signaling and regulation of microflora. *Immunity* 2013; 39(2):386–99. [PubMed: 23954130]
- [158]. van de Pavert SA, Ferreira M, Domingues RG, et al. Maternal retinoids control type 3 innate lymphoid cells and set the offspring immunity. *Nature* 2014; 508(7494):123–7. [PubMed: 24670648]
- [159]. Mudd JC, Brenchley JM. ILC you later: early and irreparable loss of innate lymphocytes in HIV infection. *Immunity* 2016; 44(2):216–8. [PubMed: 26885853]
- [160]. Xu H, Wang X, Lackner AA, Veazey RS. Type 3 innate lymphoid cell depletion is mediated by TLRs in lymphoid tissues of simian immunodeficiency virus-infected macaques. *FASEB J* 2015; 29(12):5072–80. [PubMed: 26283536]

- [161]. Xu H, Wang X, Liu DX, Moroney-Rasmussen T, Lackner AA, Veazey RS. IL-17-producing innate lymphoid cells are restricted to mucosal tissues and are depleted in SIV-infected macaques. *Mucosal Immunol* 2012; 5(6):658–69. [PubMed: 22669579]
- [162]. Zhang Z, Cheng L, Zhao J, et al. Plasmacytoid dendritic cells promote HIV-1-induced group 3 innate lymphoid cell depletion. *J Clin Invest* 2015; 125(9):3692–703. [PubMed: 26301812]
- [163]. Klooverpris HN, Kazer SW, Mjosberg J, et al. Innate lymphoid cells are depleted irreversibly during acute HIV-1 infection in the absence of viral suppression. *Immunity* 2016; 44(2):391–405. [PubMed: 26850658]
- [164]. Bendelac A, Savage PB, Teyton L. The biology of NKT cells. *Annu Rev Immunol* 2007; 25:297–336. [PubMed: 17150027]
- [165]. Sandberg JK, Ljunggren HG. Development and function of CD1d-restricted NKT cells: influence of sphingolipids, SAP and sex. *Trends Immunol* 2005; 26(7):347–9. [PubMed: 15925541]
- [166]. Sandberg JK, Fast NM, Palacios EH, et al. Selective loss of innate CD4(+) V alpha 24 natural killer T cells in human immunodeficiency virus infection. *J Virol* 2002; 76(15):7528–34. [PubMed: 12097565]
- [167]. Motsinger A, Haas DW, Stanic AK, Van Kaer L, Joyce S, Unutmaz D. CD1d-restricted human natural killer T cells are highly susceptible to human immunodeficiency virus 1 infection. *J Exp Med* 2002; 195(7):869–79. [PubMed: 11927631]
- [168]. van der Vliet HJ, von Blomberg BM, Hazenberg MD, et al. Selective decrease in circulating V alpha 24+V beta 11+ NKT cells during HIV type 1 infection. *J Immunol* 2002; 168(3):1490–5. [PubMed: 11801694]
- [169]. Moll M, Snyder-Cappione J, Spotts G, Hecht FM, Sandberg JK, Nixon DF. Expansion of CD1d-restricted NKT cells in patients with primary HIV-1 infection treated with interleukin-2. *Blood* 2006; 107(8):3081–3. [PubMed: 16368878]
- [170]. Yang OO, Wilson SB, Hultin LE, et al. Delayed reconstitution of CD4+ iNKT cells after effective HIV type 1 therapy. *AIDS Res Hum Retroviruses* 2007; 23(7):913–22. [PubMed: 17678476]
- [171]. Moll M, Kuylensstierna C, Gonzalez VD, et al. Severe functional impairment and elevated PD-1 expression in CD1d-restricted NKT cells retained during chronic HIV-1 infection. *Eur J Immunol* 2009; 39(3):902–11. [PubMed: 19197939]
- [172]. Moll M, Andersson SK, Smed-Sorensen A, Sandberg JK. Inhibition of lipid antigen presentation in dendritic cells by HIV-1 Vpu interference with CD1d recycling from endosomal compartments. *Blood* 2010; 116(11):1876–84. [PubMed: 20530791]
- [173]. Ibarondo FJ, Wilson SB, Hultin LE, et al. Preferential depletion of gut CD4-expressing iNKT cells contributes to systemic immune activation in HIV-1 infection. *Mucosal Immunol* 2013; 6(3):591–600. [PubMed: 23149661]
- [174]. Leeansyah E, Ganesh A, Quigley MF, et al. Activation, exhaustion, and persistent decline of the antimicrobial MR1-restricted MAIT-cell population in chronic HIV-1 infection. *Blood* 2013; 121(7):1124–35. [PubMed: 23243281]
- [175]. Kjer-Nielsen L, Patel O, Corbett AJ, et al. MR1 presents microbial vitamin B metabolites to MAIT cells. *Nature* 2012; 491(7426):717–23. [PubMed: 23051753]
- [176]. Cosgrove C, Ussher JE, Rauch A, et al. Early and nonreversible decrease of CD161+ MAIT cells in HIV infection. *Blood* 2013; 121(6):951–61. [PubMed: 23255555]
- [177]. Fernandez CS, Amarasena T, Kelleher AD, et al. MAIT cells are depleted early but retain functional cytokine expression in HIV infection. *Immunol Cell Biol* 2015; 93(2):177–88. [PubMed: 25348935]
- [178]. Eberhard JM, Hartjen P, Kummer S, et al. CD161+ MAIT cells are severely reduced in peripheral blood and lymph nodes of HIV-infected individuals independently of disease progression. *PLoS One* 2014; 9(11):e111323. [PubMed: 25369333]
- [179]. Khaitan A, Kilberg M, Kravietz A, et al. HIV-infected children have lower frequencies of CD8+ mucosal-associated invariant T (MAIT) cells that correlate with innate, Th17 and Th22 cell subsets. *PLoS One* 2016; 11(8):e0161786. [PubMed: 27560150]

- [180]. Leeansyah E, Svard J, Dias J, et al. Arming of MAIT cell cytolytic antimicrobial activity is induced by IL-7 and defective in HIV-1 infection. *PLoS Pathog* 2015; 11(8):e1005072. [PubMed: 26295709]
- [181]. Sandberg JK, Dias J, Shacklett BL, Leeansyah E. Will loss of your MAITs weaken your HAART [corrected]? *AIDS* 2013; 27(16):2501–4. [PubMed: 23595154]
- [182]. Greathead L, Metcalf R, Gazzard B, Gotch F, Steel A, Kelleher P. CD8+/CD161++ mucosal-associated invariant T-cell levels in the colon are restored on long-term antiretroviral therapy and correlate with CD8+ T-cell immune activation. *AIDS* 2014; 28(11):1690–2. [PubMed: 24911351]
- [183]. Vinton C, Wu F, Rossjohn J, et al. Mucosa-associated invariant T cells are systemically depleted in simian immunodeficiency virus-infected rhesus macaques. *J Virol* 2016; 90(9):4520–9. [PubMed: 26912615]
- [184]. Brandtzaeg P, Bosnes V, Halstensen TS, Scott H, Sollid LM, Valnes KN. T lymphocytes in human gut epithelium preferentially express the alpha/beta antigen receptor and are often CD45/UCHL1-positive. *Scand J Immunol* 1989; 30(1):123–8. [PubMed: 2526965]
- [185]. Nilssen DE, Muller F, Oktedalen O, et al. Intraepithelial gamma/delta T cells in duodenal mucosa are related to the immune state and survival time in AIDS. *J Virol* 1996; 70(6):3545–50. [PubMed: 8648688]
- [186]. Kagnoff MF. Current concepts in mucosal immunity. III. Ontogeny and function of gamma delta T cells in the intestine. *Am J Physiol* 1998; 274(3 Pt 1):G455–8. [PubMed: 9530144]
- [187]. Bonneville M, O'Brien RL, Born WK. Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity. *Nat Rev Immunol* 2010; 10(7):467–78. [PubMed: 20539306]
- [188]. Li H, Pauza CD. HIV envelope-mediated, CCR5/ $\alpha$ 4 $\beta$ 7-dependent killing of CD4-negative gammadelta T cells which are lost during progression to AIDS. *Blood* 2011; 118(22):5824–31. [PubMed: 21926353]
- [189]. Groh V, Steinle A, Bauer S, Spies T. Recognition of stress-induced MHC molecules by intestinal epithelial gammadelta T cells. *Science* 1998; 279(5357):1737–40. [PubMed: 9497295]
- [190]. Sciammas R, Bluestone JA. TCRgammadelta cells and viruses. *Microbes Infect Inst Pasteur* 1999; 1(3):203–12.
- [191]. Nilssen DE, Brandtzaeg P. Intraepithelial  $\gamma\delta$  T cells remain increased in the duodenum of AIDS patients despite antiretroviral treatment. *PLoS One* 2012; 7(1):e29066. [PubMed: 22238587]
- [192]. Poles MA, Barsoum S, Yu W, et al. Human immunodeficiency virus type 1 induces persistent changes in mucosal and blood gammadelta T cells despite suppressive therapy. *J Virol* 2003; 77(19):10456–67. [PubMed: 12970431]
- [193]. Kosub DA, Lehrman G, Milush JM, et al. Gamma/Delta T-cell functional responses differ after pathogenic human immunodeficiency virus and nonpathogenic simian immunodeficiency virus infections. *J Virol* 2008; 82(3):1155–65. [PubMed: 18045946]
- [194]. Harris LD, Klatt NR, Vinton C, et al. Mechanisms underlying gammadelta T-cell subset perturbations in SIV-infected Asian rhesus macaques. *Blood* 2010; 116(20):4148–57. [PubMed: 20660793]
- [195]. Soriano-Sarabia N, Archin NM, Bateson R, et al. Peripheral V $\gamma$ 9V $\delta$ 2T cells are a novel reservoir of latent HIV infection. *PLoS Pathog* 2015; 11(10):e1005201. [PubMed: 26473478]