



Published in final edited form as:

Chem Commun (Camb). 2019 July 09; 55(56): 8150–8153. doi:10.1039/c9cc02830g.

Gold Redox Catalysis for Cyclization/Arylation of Allylic Oximes: Synthesis of Isoxazoline Derivatives

Abiola Azeez Jimoh^a, Seyedmorteza Hosseyni^a, Xiaohan Ye^a, Lukasz Wojtas^a, Yong Hu^b, and Xiaodong Shi^a

^aDepartment of Chemistry, University of South Florida, Tampa, FL 33620, USA

^bDepartment of Neonatology, Shanghai Children's Hospital, Shanghai Jiao Tong University, Shanghai, Shanghai 200040, China

Abstract

Base-assisted diazonium activation has been employed to promote gold (I)/(III) redox catalysis toward allylic oxime cyclization/aryl coupling. Functional isoxazolines were prepared with good to excellent yields, while the alternative photoactivation method provided trace amounts of the isoxazoline products. This study further broadens the scope of gold redox chemistry.

Homogenous gold catalysis has been successfully applied in the areas of alkyne, allene, and alkene activation for effective C-C and C-heteroatom bond formation.¹ However, due to the challenge associated with a high oxidation potential of gold, gold redox catalysis has not been well explored until recently.² Typically, strong oxidants such as PIDA or selectfluor have been used in oxidizing Au(I) to Au(III).³ The discovery of diazonium salts as oxidizing agents for gold redox catalysis has recently stimulated the interest of many researchers.⁴ As initially reported by Glorius in 2013, followed by Toste, Shin, Alcaide, Hashimi, and others, photoactivation of diazonium salts lead to the oxidation of Au(I) to Ar-Au(III) intermediates, which promote sequential C-C multiple bonds activation followed by reductive elimination (Scheme 1A).⁵ It is generally believed that the resulting aryl radical from photoactivated diazonium salts initiates gold oxidation.⁶ These seminal works significantly advanced gold catalysis research since they offered effective reactive species (aryl diazonium) as oxidants in promoting the gold-redox cycle.

Our group recently reported a Lewis base-promoted gold oxidation process using diazonium salts.⁷ In those reports, inorganic bases were found to be adequate to facilitate diazonium salt activation under thermal conditions (40–60 °C). Herein, we report the successful cyclization of allylic oximes for the preparation of isoxazoline derivatives under mild reaction conditions. On the other hand, allylic oximes were found to undergo significant decomposition under photoconditions, giving trace amounts of the desired product.

xmshi@usf.edu.

†Electronic supplementary information (ESI) available: Experimental procedures and spectroscopic data of new compounds. CCDC 1521359–1521361.

Conflicts of Interest

There are no conflicts to declare

Due to the strong interest in developing a new strategy for heterocycle synthesis under mild conditions, we were curious about the possibility of using the gold-redox strategy for constructing the isoxazoline motif through an allylic oxime cyclization method (Scheme 1B). Notably, isoxazolines and their derivatives are a versatile class of compounds that can be found in numerous natural products and biologically active compounds.⁸ However, due to the poor substrate stability toward transition metal catalysts, there are only limited successful examples reported regarding oxime cyclization.⁹ In 2010, Loh's group reported the first example of Pd-catalyzed allylic oxime cyclization for the dioxygenation of an alkene (O₂ as both oxidant and the second hydroxyl source).¹⁰ Later, Chen's group reported Pd catalyzed allylic oxime cyclization with aryl halides.¹¹ More recently, Zhu and coworkers reported the Pd-catalyzed homoallylic oxime cyclization for the synthesis of 5-methylisoxazoles.¹² Unlike Pd catalysis, gold redox chemistry with aryldiazonium salts does not undergo oxidative addition through an aryl halide pathway. Therefore, gold redox catalysis will offer greater orthogonal functional group tolerance compared to traditional Pd catalysis. One general concern is whether oxime could survive under gold redox conditions with radical labile N-O and C=N bonds. In fact, when treating allyl oxime **1a** with diazonium **3a** under photocatalytic conditions, no desired product **2a** was observed although **1a** was completely consumed within two hours. With an interest in exploring this useful transformation with gold-redox catalysis, we further evaluated the reactivity of **1a** in the presence of diazonium salt **3a**. The results are summarized in Figure 1.

Although with reactive N-O and C=C bonds, oxime **1a** showed good stability while exposed to light; giving less than 5% decomposition under blue LED initiation. However, upon treatment with diazonium salt **3a** (1:1 mixture of **1a** and **3a**), rapid decomposition occurred at room temperature and 100% of **1a** was consumed with no clear identifiable products observed when exposed to either blue LED or daylight. Slower decomposition was observed (100% decomposition, 5 h) when treating the mixture under darkness. Notably, ¹⁹F NMR studies showed that most of the diazonium salt was consumed (entries 1–4). Unfortunately, these results ruled out the possibility of applying diazonium salt in gold redox catalysis under photo-activation conditions.

As discussed above, our group had initiated a chemical activation approach as an alternative for diazonium salt promoted gold redox catalysis. We argued that a base will assist the in-situ N₂ extrusion process to promote the oxidation of Au(I) to Au(III)-Ar, thereby, slowing down the radical generation which decomposes the oxime.^{7,13} With this in mind, we found that addition of 2 eq. Li₂CO₃, did slow down **1a** decomposition: 10% conversion of **1a** in 6 h under daylight (entry 5). Meanwhile, ¹⁹F NMR studies showed that only a little amount of diazonium salt was consumed in the presence of Li₂CO₃. This further suggests that Li₂CO₃ could potentially prevent the diazonium salt from decomposition to the aryl radical. Encouraged by this result, we evaluated the proposed cyclization-arylation under various conditions, including the choices of catalysts, solvents, bases, and reaction temperatures. Eventually, 7.5% PPh₃AuCl, 5 eq. Li₂CO₃ in CH₃CN as solvent was identified as the optimal conditions, giving the desired product **2a** in 78% NMR yield (74% isolated yield). Comparison of various reaction conditions is summarized in Table 1.

As shown above, the gold catalyst was found to be crucial for the allylic oxime cyclization since no conversion of the substrate was observed without gold (entry 2). Conducting the reaction in the absence of a base gave complete conversion with no appreciable amount of the desired product. Li_2CO_3 (5 eq) was identified as the optimal choice for this transformation (entries 5–6). Notably, although crude ^1H NMR of the reaction mixture indicated a very ‘clean’ reaction, the overall yield was around 75%. Meanwhile, the addition of excess diazonium salt was found to lead to lower yield (entry 9). Also, acetonitrile was the only solvent suitable for this transformation; switching to other solvents like 1,2-dichloroethane or toluene gave no desired product. This is due to the poor solubility of the diazonium salt in these solvents, even at elevated temperature. To emphasize the uniqueness of our chemical activation method, photochemical conditions were used for the cyclization in the presence of diazonium salt. The substrate readily decomposed in the presence of diazonium salts under this method. The standard photoactivation conditions using gold and $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$ in degassed methanol did not provide a significant amount of desired isoxazoline as confirmed by ^1H NMR analysis of the crude reaction mixture. This result suggests that the photoactivation condition is unsuitable for the oxime substrates in the presence of diazonium salt. With the optimal reaction condition in hand, we applied it to a series of allylic oximes (Scheme 2).

As shown in Scheme 2, both aromatic and aliphatic substrates were evaluated for the reaction. Good yields and high reaction efficiency could be observed when aryl groups were used (**2a-2j**). The presence of electron withdrawing groups appeared to be more favorable with respect to the isolated yield (**2e-2g**). Also, the naphthyl group afforded a good yield (**2i**). In the case of electron donating substituent **2h**, a modest yield was achieved. Compared to **2j**, the more sterically hindered **2b** provided a lower yield, suggesting that sterics played a role. However, aliphatic substrate gave a much lower yield for this reaction (**2k**). Also, the homoallylic oxime was found to be viable for this reaction giving a good yield of 5,6-dihydro-4H-1,2-oxazine derivative (**2l**). This is a significant step towards broadening the scope of this reaction; owing to the fact that oxazines are also known to be a highly desirable class of heterocycles. The 7 or 8 membered-ring analogs could not be observed under the reported reaction conditions.

Furthermore, a series of diazonium salts were investigated (Scheme 3); electron poor aryl diazonium salts could also be enlisted, giving the desired products in high yields (**4a-4h**). The iodo-substituted diazonium salt **4e** was also successfully applied with modest yield. This result highlights the orthogonal reactivity under gold catalysis, compared with previously reported Pd catalysis conditions. However, electron rich diazonium salts failed to give the desired product due to the instability of the diazonium species.

Although the substituted isoxazolines are in themselves useful compounds, we carried out their reduction to the corresponding hydroxyl ketones (Scheme 4A). Isoxazoline **2a** was subjected to reductive N-O bond cleavage followed by subsequent hydrolysis of the resulting imine. This reductive ring opening was achieved up to 82% yield. To further demonstrate the synthetic utility, we synthesized the *p*-iodo substituted aromatic substrate **6**, which was further subjected to our reaction conditions. The desired isoxazoline **7** was obtained with 72% yield (Scheme 4B). Notably, when treating **6** under previously reported Pd-catalyzed

conditions, no desired product was observed with 100% conversion. The fast oxidative addition of the C-I bond led to the side reaction pathway. This result clearly indicated the orthogonal catalytic reactivity between Pd and Au. The following Sonogashira reaction of **7** afforded alkyne substituted aromatic isoxazoline **8**, allowing an extended synthetic handle.

In conclusion, gold-redox chemistry has been applied for the cyclization/functionalization of allylic oximes in the presence of aryl diazonium salts. This transformation gives access to aryl functionalized 2-isoxazolines in good to excellent yields. Moreover, we believe the apparent compatibility of Au redox chemistry with oximes will provide an alternate route towards the synthesis of isoxazoline derivatives.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the NSF (CHE-1665122), NIH (1R01GM120240-01) and NSFC (21228204) for financial support.

References

- (1) (a). Hashmi ASK and Hutchings GJ, *Angew. Chem., Int. Ed.*, 2006, 45, 7896;(b)Widenhofer RA and Han XQ, *Eur. J. Org. Chem.*, 2006, 4555;(c)Gorin DJ and Toste FD, *Nature*, 2007, 446, 395; [PubMed: 17377576] (d)Gorin DJ, Sherry BD and Toste FD, *Chem. Rev.*, 2008, 108, 3351; [PubMed: 18652511] (e)Hashmi ASK, *Angew. Chem., Int. Ed.*, 2010, 49, 5232;(f)Sengupta S and Shi XD, *Chemcatchem*, 2010, 2, 609;(g)Aubert C, Fensterbank L, Garcia P, Malacria M and Simonneau A, *Chem. Rev.*, 2011, 111, 1954; [PubMed: 21391568] (h)Wegner HA and Auzias M, *Angew. Chem., Int. Ed.*, 2011, 50, 8236;(i)Hashmi ASK, *Acc. Chem. Res.*, 2014, 47, 864; [PubMed: 24533563] (j)Wang YM, Lackner AD and Toste FD, *Acc. Chem. Res.*, 2014, 47, 889; [PubMed: 24228794] (k)Zhang LM, *Acc. Chem. Res.*, 2014, 47, 877. [PubMed: 24428596] (l)Jia MQ and Bandini M. *ACS Catal*, 2015, 5, 1638;(m)Dorel R and Echavarren AM, *Chem. Rev.*, 2015, 115, 9028; [PubMed: 25844920] (n)Asiri AM and Hashmi ASK, *Chem. Soc. Rev.*, 2016, 45, 4471. [PubMed: 27385433]
- (2). Bratsch SG, *J. Phys. Chem Ref. Data*, 1989, 18, 1.
- (3). (a)For review, see: Hopkinson MN, Gee AD and Gouverneur V, *Chem. Eur. J.*, 2011, 17, 8248; [PubMed: 21678513] (b)Joost M, Amgoune A and Bourissou D, *Angew. Chem., Int. Ed.*, 2015, 54, 15022;(c)Miró J and del Pozo C, *Chem. Rev.*, 2016, 116, 11924; [PubMed: 27548659] (d)Silva F, Tierno AF and Wengryniuk SE, *Molecules*, 2017, 22.(a)For selected examples, see: Zhang G, Peng Y, Cui L and Zhang L, *Angew. Chem., Int. Ed.*, 2009, 48, 3112;(b)Tkatchouk E, Mankad NP, Benitez D, Goddard WA and Toste FD, *J. Am. Chem. Soc.*, 2011, 133, 14293; [PubMed: 21861448] (c)Leyva-Perez A, Domenech A, Al-Resayes SI and Corma A, *ACS Catal*, 2012, 2, 121;(d)Peng H, Xi Y, Ronaghi N, Dong B, Akhmedov NG and Shi X, *J. Am. Chem. Soc.*, 2014, 136, 13174; [PubMed: 25184690] (e)Levin MD and Toste FD, *Angew. Chem., Int. Ed.*, 2014, 53, 6211;(f)Hofer M, Genoux A, Kumar R and Nevado C, *Angew. Chem., Int. Ed.*, 2017, 56, 1021;(g)Wang J, Zhang S, Xu C, Wojtas L, Akhmedov NG, Chen H and Shi X, *Angew. Chem., Int. Ed.*, 2018, 57, 6915.
- (4). (a)For review, see: Hopkinson MN, Tlahuext-Aca A and Glorius F, *Acc. Chem. Res.*, 2016, 49, 2261; [PubMed: 27610939] (b)Akram MO, Banerjee S, Saswade SS, Bedi V and Patil NT, *Chem. Commun*, 2018, 54, 11069.
- (5). (a)For selected examples see: Sahoo B, Hopkinson MN and Glorius F, *J. Am. Chem. Soc.*, 2013, 135, 5505; [PubMed: 23565980] (b)Tlahuext-Aca, Hopkinson MN, Sahoo B and Glorius F, *Chem. Sci*, 2016, 7, 89; [PubMed: 29861968] (c)Shu XZ, Zhang M, He Y, Frei H and Toste FD,

J. Am. Chem. Soc, 2014, 136, 5844; [PubMed: 24730447] (d)Kawai H, Wolf WJ, DiPasquale AG, Winston MS and Toste FD, J. Am. Chem. Soc, 2016, 138, 587; [PubMed: 26744765] (e)Kim S, Rojas-Martin J and Toste FD, Chem. Sci, 2016, 7, 85; [PubMed: 29861967] (f)Huang L, Rudolph M, Rominger F and Hashmi ASK, Angew. Chem., Int. Ed, 2016, 55, 4808;(g)Xie J, Sekine K, Witzel S, Krämer P, Rudolph M, Rominger F and Hashmi ASK, Angew. Chem., Int. Ed, 2018, 57, 16648;(h)Patil DV, Yun H and Shin S, Adv. Synth. Catal, 2015, 357, 2622; (i)Cornilleau T, Hermange P and Fouquet E, Chem. Commun, 2016, 52, 10040;(j)Qu CH, Zhang SL, Du HB and Zhu CJ, Chem. Commun, 2016, 52, 14400.(k)Um J, Yun H and Shin S, Org. Lett, 2016, 18, 484; [PubMed: 26761155] (l)Akram MO, Mali PS and Patil NT, Org. Lett, 2017, 19, 3075; [PubMed: 28590758] (m)Gauchot V, Sutherland DR and Lee AL, Chem. Sci, 2017, 8, 2885. [PubMed: 28553527] (n)Alcaide B, Almendros P, Busto E and Luna A, Adv. Synth. Catal, 2016, 358, 1526(o)Alcaide B, Almendros P, Busto E and Lázaro-Milla C, J.Org. Chem, 2017, 82, 2177. [PubMed: 28150493] (p)Alcaide B, Almendros P, Aparicio B, Lázaro-Milla C, Luna A and Faza ON, Adv. Synth. Catal, 2017, 359, 2789.

- (6). Huang L, Rominger F, Rudolph M and Hashmi ASK, Chem. Commun, 2016, 52, 6435.
- (7) (a). Cai R, Lu M, Aguilera EY, Xi Y, Akhmedov NG, Petersen JL, Chen H, Shi X, Angew. Chem. Int. Ed 2015, 54, 8772;(b)Peng H, Cai R, Xu C, Chen H and Shi X, Chem. Sci, 2016, 7, 6190; [PubMed: 30034759] (c)Dong B, Peng H, Motika SE and Shi X, Chem. Eur. J, 2017, 23, 11093. [PubMed: 28603854]
- (8) (a). Pohjakallio A, Pihko PM and Laitinen UM, Chem. Eur. J, 2010, 16, 11325; [PubMed: 20726019] (b)Landais Y, Robert F, Godineau E, Huet L and Likhite N, Tetrahedron, 2013, 69, 10073;(c)Gassel M, Wolf C, Noack S, Williams H and Ilg T, Insect Biochem. Mol. Biol, 2014, 45, 111; [PubMed: 24365472] (d)Shoop WL, Hartline EJ, Gould BR, Waddell ME, McDowell RG, Kinney JB, Lahm GP, Long JK, Xu M, Wagerle T, Jones GS, Dietrich RF, Cordova D, Schroeder ME, Rhoades DF, Benner EA and Confalone PN, Vet. Parasitol, 2014, 201, 179; [PubMed: 24631502] (e)Ismail T, Shafi S, Swam S, Sidiq T, Khajuria A, Rouf A, Yadav M, Saikam V, Singh PP, Alam MS, Islam N, Sharma K and Kumar HMS, Eur. J. Med. Chem, 2016, 123, 90; [PubMed: 27474926] (f)Puttaswamy N, Kumar GSP, Al-Ghorbani M, Vigneshwaran V, Prabhakar BT and Khanum SA, Eur. J. Med. Chem, 2016, 114, 153; [PubMed: 26974382] (g)Ma H, Stone VN, Wang HQ, Kellogg GE, Xu P and Zhang Y, Bioorg. Med. Chem. Lett, 2017, 27, 3840; [PubMed: 28668193] (h)Park SW, Kim SH, Song J, Park GY, Kim D, Nam TG and Hong KB, Beilstein J. Org. Chem, 2018, 14, 1028; [PubMed: 29977375] (i)Prasad SS and Baskaran S, J. Org. Chem, 2018, 83, 1558. [PubMed: 29313687]
- (9). (a)For selected examples, see: House HO, Manning DT, Melillo DG, Lee LF, Haynes OR and Wilkes BE, J. Org. Chem, 1976, 41, 855;(b)Bates RW and Sa-Ei K, Org. Lett, 2002, 4, 4225; [PubMed: 12443064] (c)Moriyama K, Izumisawa Y and Togo H, J. Org. Chem, 2011, 76, 7249; [PubMed: 21823624] (d)Zhang W, Su Y, Wang K-H, Wu L, Chang B, Shi Y, Huang D and Hu Y, Org. Lett, 2017, 19, 37.
- (10). Zhu M-K, Zhao J-F and Loh T-P, J. Am. Chem. Soc, 2010, 132, 6284. [PubMed: 20394368]
- (11). Jiang D, Peng J and Chen Y, Org. Lett, 2008, 10, 1695. [PubMed: 18393515]
- (12). Dong K-Y, Qin H-T, Bao X-X, Liu F and Zhu C, Org. Lett, 2014, 16, 5266 [PubMed: 25255195]
- (13) (a). Kindt S, Wicht K and Heinrich MR, Org. Lett, 2015, 17, 6122; [PubMed: 26636470] (b)Xia Z and Zhu Q, Org. Lett, 2013, 15, 4110; [PubMed: 23919834] (c)Hartmann M, Li Y and Studer A, J. Am. Chem. Soc, 2012, 134, 16516. [PubMed: 23009829]








Figure 1.
Allylic oxime reactivity with diazonium salt

(A) Gold redox catalysis using diazonium salt as oxidant**(B)** Challenge in oxime cyclization: radical-labile substrates

Scheme 1.
Cyclization of Oximes


**Scheme 2.**

Substrate scope for allylic oxime. General reaction conditions: **1a** (0.33 mmol), [Au] (0.025 mmol), diazonium salt (0.66 mmol), and base (1.65 mmol) in CH_3CN (0.4 M), 60 °C; b Isolated yield.


**Scheme 3.**

Substrate scope for diazonium salt. General reaction conditions: 1a (0.33 mmol), [Au] (0.025 mmol), diazonium salt (0.66 mmol), and base (1.65 mmol) in CH_3CN (0.4 M), 60 °C; b Isolated yield.

a) Reduction of Isoxazoline




b) orthogonal reactivity



Scheme 4.
Isoxazoline Reactivity

Table 1

Selected initial optimization reactions



Entry	Reaction conditions varies from standard ^a	convn 1a	yield ^b 2a	yield ^b 2a'
1	None	100%	78%	0%
2	No [Au]	0%	-	-
3	No [ArN ₂][BF ₄]	18%	-	-
4	No Base	100%	trace	-
5	NaHCO ₃ 2 eq.	95%	58%	0%
6	Li ₂ CO ₃ 2 eq.	100%	74%	0%
7	5 % PPh ₃ AuCl	90%	51%	0%
8	5 eq. [ArN ₂][BF ₄]	100%	65%	-
9 ^c	No [Au], UVA or Blue LED, degassed MeOH	100%	0%	0%
10 ^c	10% PPh ₃ AuCl, degassed MeOH or MeCN, 12W blue LEDs, RT	100%	<5%	0%
11 ^c	10% PPh ₃ AuCl, 2.5% [Ru(bpy) ₃](PF ₆) ₂ degassed MeOH or MeCN, 12 W blue LEDs, RT	100%	<5%	0%

^a General reaction conditions: 1a (0.33 mmol), [Au] (0.025 mmol) diazonium salt (0.66 mmol) and base (1.65 mmol) in CH₃CN (0.4 M), 60 °C

^b Yield was based on ¹HNMR with 1,3,5-trimethoxybenzene as internal standard.

^c Solvents were degassed using the Freeze-Pump-Thaw method.