

Expression and prognostic role of IKBKE and TBK1 in stage I non-small cell lung cancer

This article was published in the following Dove Press journal:
Cancer Management and Research

Xin Wang^{1,2}

Feifei Teng²

Jie Lu³

Dianbin Mu⁴

Jianbo Zhang⁴

Jinming Yu^{1,2}

¹Department of Oncology, Renmin Hospital of Wuhan University, Wuhan, Hubei 430060, People's Republic of China;

²Department of Radiation Oncology, Shandong Cancer Hospital Affiliated to Shandong University, Shandong Academy of Medical Sciences, Jinan, Shandong 250117, People's Republic of China;

³Department of Neurosurgery, Shandong Province Qianfoshan Hospital of Shandong University, Jinan, Shandong 250014, People's Republic of China; ⁴Department of Pathology, Shandong Cancer Hospital Affiliated to Shandong University, Shandong Academy of Medical Sciences, Jinan, Shandong 250117, People's Republic of China

Background: The inhibitors of nuclear factor kappa-B kinase subunit epsilon (*IKBKE*) and TANK-binding kinase 1 (*TBK1*) are important members of the nonclassical IKK family that share the kinase domain. They are important oncogenes for activation of several signaling pathways in several tumors. This study aims to explore the expression of *IKBKE* and *TBK1* and their prognostic role in stage I non-small cell lung cancer (NSCLC).

Patients and methods: A total of 142 surgically resected stage I NSCLC patients were enrolled and immunohistochemistry of *IKBKE* and *TBK1* was performed.

Results: *IKBKE* and *TBK1* were expressed in 121 (85.2%) and 114 (80.3%) of stage I NSCLC patients respectively. *IKBKE* expression was significantly associated with *TBK1* expression ($P=0.004$). Furthermore, multivariate regression analyses showed there was a significant relationship between patients with risk factors, the recurrence pattern of metastasis and *IKBKE*+/*TBK1*+ co-expression ($P=0.032$ and $P=0.022$, respectively). In Kaplan–Meier survival curve analyses, the *IKBKE*+/*TBK1*+ co-expression subgroup was significantly associated with poor overall survival ($P=0.014$).

Conclusions: This is the first study to investigate the relationship between *IKBKE* and *TBK1* expression and clinicopathologic characteristics in stage I NSCLC patients. *IKBKE*+/*TBK1*+ co-expression was significantly obvious in patients with risk factors and with recurrence pattern of distant metastasis. Furthermore, *IKBKE*+/*TBK1*+ is also an effective prognostic predictor for poor overall survival.

Keywords: *IKBKE*, *TBK1*, NSCLC, prognosis, cancer

Introduction

Lung cancer is the leading cause of cancer-related death worldwide. Non-small cell lung cancer (NSCLC) accounts for 80–85% of all lung cancer cases.^{1,2} The primary treatment for stage I NSCLC is surgical resection, but at least 30% of patients experienced recurrence within 5 years.³ Adjuvant chemotherapy is recommended for patients with risk factors, such as poor histologic differentiation, vascular invasion, wedge resection, tumors >4 cm, incomplete lymph node sampling, and visceral pleural invasion.⁴ Even after receiving adjuvant chemotherapy, the 5-year survival rate is only 60–90% due to local recurrence or distant metastasis.^{5,6} Therefore, we defined high-risk patients as tending to local recurrence or distant metastasis. It is challenging to screen true high-risk patients for adjuvant chemotherapy, and to avoid unnecessary adjuvant chemotherapy in low-risk patients to reduce the survival reduction caused by treatment-related toxicity.

The IKK family contains five protein factors. *IKK α* , *IKK β* , and *IKK γ* belong to canonical group, and the other two (*IKBKE* and *TBK1*) belong to non-canonical

Correspondence: Jinming Yu
Department of Oncology, Renmin Hospital of Wuhan University, Wuhan, Hubei 430060, People's Republic of China
Tel +86 6762 6971
Email sdyujinming@163.com

group.⁷ The amino acid sequence analysis confirmed that IKBKE has 67% homology with TBK1.⁸ IKBKE is highly expressed in human normal pancreatic tissue, thyroid tissue, spleen, and peripheral blood leukocytes, and can be up-regulated by multiple cytokines such as tumor necrosis factor alpha (TNF- α), IL-1, IL-6, and interferon- γ .⁹ TBK1 is constitutively expressed in immune cells, brain, lungs, gastrointestinal tract, and reproductive organs.¹⁰ Studies have demonstrated that IKBKE and TBK1 need to bind to scaffold proteins to form a protease complex to activate downstream protein factors, thereby activating the *NF- κ B* pathway. TNAK (TNF receptor-associated factor family member-associated NF- κ B activator), NAPI (NAK associated protein 1), and SINTBAD (similar to NAP and TBK1 adaptor) are the main scaffold proteins.^{11–17} Several studies have shown that IKBKE was overexpressed in breast cancer, glioma, gastric cancer, pancreatic cancer, ovarian cancer, renal cancer, and lung cancer and was associated with poor prognosis.^{18–24} IKBKE and TBK1 may play important roles in tumorigenesis, proliferation, and angiogenesis.^{25,26} Nevertheless, the association between IKBKE and TBK1 expression and clinicopathological characteristics and its role in the prognosis of stage I NSCLC has not been reported.

Patients and methods

Patient selection

A total of 142 tissue samples were retrospectively collected from patients who were pathologically diagnosed with stage I NSCLC between January 2004 and June 2012 in Shandong Cancer hospital. All of them received anatomical segmentectomy or standard lobectomy via open thoracotomy and did not receive other treatments except for surgery. This retrospective study was approved by the ethics committee at Shandong Cancer Hospital and Institute. All patients have provided written informed consent, and this study was conducted in accordance with the Declaration of Helsinki.

Immunohistochemistry

Tumor tissue samples were obtained from 142 surgically resected stage I NSCLC patients. Surgically resected specimens were fixed in formalin and embedded in paraffin. Then, 4- μ m paraffin-embedded sections were dewaxed and heated to 95°C for 20 mins for antigen retrieval in 10 mmol/L citrate buffer. Then, the sections were cleared of endogenous peroxidase activity and were blocked with 5% BSA for 30 mins

in room temperature (RT). Then, all sections were incubated at 4°C overnight with primary anti-*IKBKE* (ab7891, Abcam, IHC 1:100, Cambridge, MA) or anti-*TBK1* (ab109735, Abcam, IHC 1:400, Cambridge, MA). Then, the specimens were incubated with biotinylated secondary antibodies for 1 hr in RT. After incubation with ABC-peroxidase for 1 hr then all sections were colored using a DAB Kit (ZSGB-BIO, People's Republic of China) and counterstained with hematoxylin. All IHC results were evaluated by two pathologists. Double-blind readings were performed by two pathologists at the unknown tumor grade, and 10 fields were randomly collected from each pathological section, and the percentage of positive cells and the intensity of staining were scored: 1) coloring cells accounted for <5% cell count is 0; 5–25% is 1 point; 26–50% is 2 points; 51–75% is 3 points; and >75% is 4 points; 2) no staining is 0 points; the light yellow is 1 point; the brown and yellow is 2 points; the sepia is 3 points; 3) the two points are multiplied to form a positive grade: 0 is negative (-), 1–4 is weakly positive (+), 5–8 is positive (++) and 9–12 is strongly positive (+++).

Statistical analysis

The categorical variables were tested using Pearson's chi-squared tests or Fisher's exact test, and continuous variables were expressed as means \pm SD and were analyzed using independent *t*-test. The significant ($P<0.05$) clinicopathological characteristics in univariate analysis were enrolled in multivariate logistic regression analysis, and multivariate logistic regression analysis was used to confirm the independent clinicopathological characteristics which were associated with IKBKE and TBK1 expression. The Kaplan–Meier was used to analyze progression-free survival (PFS) and OS, and the survival curves of four subgroups were compared by log-rank test. All analysis was two-sided, and $P<0.05$ was considered statistically significant. Statistical analysis was performed using SPSS (version 20.0).

Results

Expression of IKBKE and TBK1 and their association in stage I NSCLC patients

Among the 142 enrolled stage I NSCLC patients, 48 (33.8%) patients were squamous cell carcinoma, and 65 patients (45.8%) were adenocarcinoma. Twenty-nine patients were other types of NSCLC including 8 large cell carcinoma, 5 atypical carcinoid tumor, 5 pleomorphic carcinoma, 9 adenosquamous carcinoma, and 2 mucinous epidermoid carcinoma. IKBKE and TBK1 expression was

found in 85.2% (n=121) and 80.3% (n=114) of stage I NSCLC patients, respectively. The association between IKBKE and TBK1 expression and clinicopathological characteristics is summarized in Table 1. IKBKE and TBK1 expression was associated with risk factors ($P=0.021$ and $P=0.014$). From Table 1 we found patients with risk factors showed increased IKBKE and TBK1 expression compared to patients who had no risk factors. IKBKE expression was observed in 91.3% and 77.4% of patients who had risk factors compared to patients who had no risk factors. Besides, TBK1 expression was observed in 87.5% and 71.0% in these two subgroups. In addition, TBK1 expression was significantly associated with recurrence pattern of metastasis compared to local recurrence ($P<0.001$) and TBK1 expression was found in 89.6% of patients with the recurrence pattern of metastasis compared to 64.0% of patients with the recurrence pattern of local recurrence. We further explored the correlation between IKBKE expression and TBK1 expression and found there was a significant correlation between IKBKE and TBK1 expression ($P=0.004$), as 84.3% (102/121) of tumors with IKBKE positive expression simultaneously showed TBK1 positive expression (Table 2).

IKBKE and TBK1 co-expression in stage I NSCLC patients

We divided all patients into four subgroups according to IKBKE and TBK1 expression as follows: IKBKE-/TBK1- (n=9, 6.3%); IKBKE+/TBK1- (n=19, 13.4%); IKBKE-/TBK1+ (n=12, 8.5%); IKBKE+/TBK1+ (n=102, 71.8%). The representative examples of four subgroups are shown in Figure 1. We then explored the association of IKBKE and TBK1 co-expression and clinicopathological characteristics (Table 3). Patients with risk factors and patients with the recurrence pattern of metastasis had more frequency in the IKBKE+/TBK1+ subgroup than in other three subgroups ($P=0.042$ and $P=0.004$, respectively). In addition, we did not find significant association between other clinicopathological characteristics and the expression of IKBKE and TBK1. In multivariate regression analysis of clinicopathological characteristics for IKBKE and TBK1 co-expression, we found patients with risk factors and the recurrence pattern of metastasis were significantly related to IKBKE+/TBK1+ co-expression ($P=0.032$ and $P=0.022$, respectively). The adjusted OR of patients with risk factors was 1.633, with a 95% CI 1.043–2.557 when compared with patients with no risk factors. The adjusted

OR of patients with recurrence pattern of metastasis was 1.670, with a 95% CI 1.078–2.587 when compared with patients with recurrence pattern of local recurrence (Table 4).

The prognostic significance of IKBKE and TBK1 co-expression in stage I NSCLC patients

The median follow-up period for OS and PFS was 1,442 days (range 488–3,239 days) and 1,021 days (range 107–2,989 days), respectively. During the follow-up period, the recurrence rate was 89.4% (n=127). Survival was analyzed using the Kaplan–Meier method with stratification of IKBKE and TBK1 expression. The results showed that IKBKE+/TBK1+ co-expression subgroup was significantly associated with poor OS (median 1,351 days; 95% CI 1,178–1,565; $P=0.014$) than those in IKBKE-/TBK1- subgroup (median 2,378 days; 95% CI 1,924–4,291), IKBKE+/TBK1- subgroup (median 1,554 days; 95% CI 959–2,148), and IKBKE-/TBK1+ subgroup (median 1,643 days; 95% CI 1,529–1,836). However, the IKBKE+/TBK1+ co-expression subgroup did not show significant association with poor PFS than the other three subgroups ($P=0.109$), although the Kaplan–Meier survival curve showed this tendency (Figure 2).

Discussion

Our study tried to investigate IKBKE and TBK1 expression in stage I NSCLC and the association between their expression and clinicopathological characteristics. We found that IKBKE and TBK1 expressed in 85.2% and 80.3% of stage I NSCLC, respectively, and IKBKE and TBK1 co-expressed in 71.8% of all cases. Identifying high-risk patients who need postoperative adjuvant chemotherapy to improve OS and reduce the toxicity of unnecessary chemotherapy is a key element in the whole treatment management of early-stage NSCLC. Our study demonstrated that IKBKE and TBK1 might play an important role in tumorigenesis and proliferation in stage I NSCLC. IKBKE and TBK1 may be good molecular targets for combination therapy.

Targeted therapy is the treatment of precise targeting molecular “targets” by small molecule drugs that specifically bind to these targets. Protein kinases are important signal messengers to modulate cell life activities and play an important role in cell proliferation, survival, apoptosis, metabolism, transcription, and differentiation. IKBKE and

Table 1 The association of IKBKE and TBK1 expression and clinicopathological characteristics

Variable	IKBKE expression			TBK1 expression		
	Total cases N=142(%)	Positive N=121(85.2%)	Negative N=21(14.8%)	Positive N=114(80.3%)	Negative N=28(19.7%)	P
Gender						
Male	91 (64.1%)	77 (84.6%)	14 (15.4%)	72(79.1%)	19(20.9%)	0.642
Female	51 (35.9%)	44 (86.3%)	7 (13.7%)	42(82.4%)	9(17.6%)	
Age						
≤60	67 (47.2%)	61 (91.0%)	6 (9.0%)	55(82.1%)	12(17.9%)	0.609
>60	75 (52.8%)	60 (80.0%)	15 (20.0%)	59(78.7%)	16(21.3%)	
Differentiation						
Poor	37 (26.1%)	31 (83.8%)	6 (16.2%)	29(78.4%)	8(21.6%)	0.940
Moderate	62 (43.7%)	51 (82.3%)	11 (17.7%)	50(80.6%)	12(19.4%)	
Well	43 (30.2%)	39 (90.7%)	4 (9.3%)	35(81.4%)	8(18.6%)	
Histologic type						
Squamous carcinoma	48 (33.8%)	43 (89.6%)	5 (10.4%)	37(77.1%)	11(22.9%)	0.723
Adenocarcinoma	65 (45.8%)	54 (83.1%)	11 (16.9%)	54(83.1%)	11(16.9%)	
Large cell carcinoma	8 (5.6%)	7 (87.5%)	1 (12.5%)	6(75.0%)	2(25.0%)	
Atypical carcinoid tumor	5 (3.5%)	4 (80.0%)	1 (20.0%)	4(80.0%)	1(20.0%)	
Pleomorphic carcinoma	5 (3.5%)	5 (100.0%)	0 (0%)	4(80.0%)	1(20.0%)	
Adenosquamous carcinoma	9 (6.4%)	6 (66.7%)	3 (33.3%)	8(88.9%)	1(11.1%)	
Mucinous epidermoid carcinoma	2 (1.4%)	2 (100.0%)	0 (0%)	1(50.0%)	1(50.0%)	
Risk factors						
Yes	80 (56.3%)	73 (91.3%)	7 (8.7%)	70(87.5%)	10(12.5%)	0.014
No	62 (43.7%)	48 (77.4%)	14 (22.6%)	44(71.0%)	18(29.0%)	
Recurrence pattern						
Local	50 (38.7%)	41 (82.0%)	9 (18.0%)	32(64.0%)	18(36.0%)	<0.001
Metastasis	77 (54.2%)	68 (88.3%)	9 (11.7%)	69(89.6%)	8(10.4%)	

Table 2 The association between IKBKE and TBK1 expression

Variable	IKBKE expression		P
	Positive (n=121)	Negative (n=21)	
TBK1 expression			0.004
Positive (n=114)	102 (84.3%)	12 (57.1%)	
Negative (n=28)	19 (15.7%)	9 (42.9%)	

TBK1 are serine/threonine protein kinases which is part of IKK family.⁹ Ma et al demonstrated the activity of TBK1 is strongly promoted by phosphorylation at Ser 172.²⁷ In breast cancer, as an oncogene, IKBKE is regulated by K63-linked polyubiquitination at lysine 30 and lysine 401 and controlled by the cIAP1/cIAP2/TRAF2 E3 ligase complex ubiquitinates itself.²⁸ Although IKBKE and TBK1 were characterized as activators of *NF- κ B* pathway, several studies indicated non-canonical IKKs (IKBKE and TBK1) were sufficient, but not essential for *NF- κ B* activation.^{18,29} *NF- κ B* pathway plays a pivotal role in the development of various tumors. In addition to *NF- κ B* pathway, IKBKE and TBK1 also activated AKT by direct phosphorylation of AKT to promote cell proliferation and survival.³⁰ In addition, Saxton et al³¹ found mechanistic target of rapamycin (mTOR) complex 1 (mTORC1) controls metabolic processes and phosphorylates key substrates to promote oncoprotein-induced cell proliferation and survival. Yu et al³² showed TBK1 can regulate AKT-mTORC1 signaling axis. Bodur et al³³ recently demonstrated TBK1 can directly phosphorylate

mTORC1 at Ser 2159 to promote its tumorigenic activity. Recent studies have confirmed that IKBKE can directly phosphorylate signal transducers and activators of transcription 3 (STAT3), knocking down IKBKE can significantly reduce the content of STAT3 and phosphorylated STAT3, and affect the proliferation and progression of lymphoma through STAT3 signaling pathway.³⁴ Furthermore, several studies have explored the tumorigenic function of IKBKE and TBK1 in various cancers. Qin et al³⁵ indicated silencing IKBKE using synthetic siRNA can reduce focus formation potential and clonogenicity in human breast cancer cells. Li et al³⁶ verified that IKBKE overexpression in glioma is positively correlated to the grade of glioma. Silencing IKBKE can significantly inhibit tumor cell proliferation and invasion, and lead to cell cycle arrest. Lu et al demonstrated IKBKE plays an important role in regulating cell proliferation, invasion and epithelial-mesenchymal transition of glioblastoma cells in vitro and in vivo.³⁷ Besides, Niederberger et al³⁸ found IKBKE/TBK1-sensitive acute myeloid leukemia (AML) cells tend to have MYC oncogenic signatures and inhibition of IKBKE/TBK1 can downregulate the MYC oncogenic pathway in AML cell lines. Accumulating evidence indicates the oncogenic potential of IKBKE and TBK1 is an important basis for them as targets in targeted therapy. Recent studies have shown that CYT387, a small molecule inhibitor that is traditionally thought to inhibit JAK/STAT signal transduction pathway, could inhibit IKBKE kinase activity and become a newly discovered small inhibitor of IKBKE. Experiments demonstrated CYT387 could significantly




Figure 1 Representative examples of the four subgroups. A and B, E and F, G and H were moderately differentiated squamous cell carcinoma. C and D were large cell lung cancer. IKBKE and TBK were cytoplasmic stain. A and B: IKBKE+/TBK1+. C and D: IKBKE+/TBK1-. E and F: IKBKE-/TBK1+. G and H: IKBKE-/TBK1-.

Table 3 The association of IKBKE and TBK1 co-expression and clinicopathological characteristics

Variable	IKBKE-/TBK1 - IKBKE+/TBK- IKBKE-/TBK+		IKBKE+/TBK1 +		P
	N=9 (6.3%)	N=19 (13.4%)	N=12 (8.5%)	N=102 (71.8%)	
Gender					0.961
Male	6 (6.6%)	13 (14.3%)	8 (8.8%)	64 (70.3%)	
Female	3 (5.9%)	6 (11.8%)	4 (7.8%)	38 (74.5%)	
Age					0.114
≤60	1 (1.5%)	11 (16.4%)	5 (7.5%)	50 (74.6%)	
>60	8 (10.7%)	8 (10.7%)	7 (9.3%)	52 (69.3%)	
Differentiation					0.937
Poor	3 (8.1%)	5 (13.5%)	3 (8.1%)	26 (70.3%)	
Moderate	4 (6.4%)	8 (12.9%)	7 (11.3%)	43 (69.4%)	
Well	2 (4.7%)	6 (14.0%)	2 (4.6%)	33 (76.7%)	
Histologic type					0.548
Squamous carcinoma	4 (8.3%)	7 (14.6%)	1 (2.1%)	36 (75.0%)	
Adenocarcinoma	3 (4.6%)	8 (12.3%)	8 (12.3%)	46 (70.8%)	
Large cell carcinoma	1 (12.5%)	1 (12.5%)	0 (0%)	6 (75.0%)	
Atypical carcinoid tumor	0 (0%)	1 (20.0%)	1 (20.0%)	3 (60.0%)	
Pleomorphic carcinoma	0 (0%)	1 (20.0%)	0 (0%)	4 (80.0%)	
Adenosquamous carcinoma	1 (11.1%)	0 (0%)	2 (22.2%)	6 (66.7%)	
Mucinous epidermoid carcinoma	0 (0%)	1 (50.0%)	0 (0%)	1 (50.0%)	
Risk factors					
Yes	3 (3.7%)	7 (8.8%)	5 (6.2%)	65 (81.3%)	
No	6 (9.7%)	12 (19.3%)	7 (11.3%)	37 (59.7%)	
Recurrence pattern					0.004
Local	6 (12.0%)	12 (24.0%)	3 (6.0%)	29 (58.0%)	
Metastasis	1 (1.3%)	7 (9.1%)	8 (10.4%)	61 (79.2%)	

Table 4 Multivariate regression analysis of clinicopathological characteristics for IKBKE and TBK1 co-expression

Variables	P	OR	95% CI
Risk factors	0.032	1.633	1.043–2.557
Recurrence pattern	0.022	1.670	1.078–2.587

inhibit the proliferation of NSCLC lines in vitro, block the development and progression of NSCLC induced by IKBKE and TBK1 mediated KRAS-driven tumorigenesis.³⁹ Considering the high positive expression rate of IKBKE and TBK1 in stage I NSCLC, targeted therapy targeting IKBKE and TBK1 might be a potential strategy to suppress cancer development.

In our study, we explored the correlation between the following clinicopathological characteristics, including gender, age, differentiation, histologic type, risk factors, and recurrence pattern and IKBKE/TBK1 expression status. We found IKBKE+/TBK1+ expressed in 71.8% of all cases, suggesting that IKBKE and TBK1 may play a pivotal role in NSCLC initiation. The multivariate regression analysis was applied to select prediction characteristics for IKBKE+/TBK1+ expression in stage I NSCLC and the multivariate analysis demonstrated patients with risk factors and the recurrence pattern of metastasis were significantly related to IKBKE+/TBK1+ co-expression ($P=0.032$ and $P=0.022$, respectively).

In addition, we did not find significant association between other clinicopathological characteristics and the expression of IKBKE and TBK1. The stage I NSCLC patients with risk factors were optimal candidates to receive adjuvant chemotherapy, whereas the locoregional or systemic recurrence was still high in these patients.⁵ Even in stage I NSCLC patients with no risk factors, the locoregional or systemic recurrence also could not be ignored. Therefore, how to explore effective treatments to prolong the OS time in high-risk patients remains challenging. Our research provides a new perspective for screening high-risk patients. We found stage I NSCLC patients with risk factors were significantly related to IKBKE+/TBK1+ co-expression ($P=0.032$). Furthermore, we found the recurrence pattern of metastasis was also significantly related to IKBKE+/TBK1+ co-expression ($P=0.022$). Recently, studies have shown IKBKE and TBK1 could phosphorylate AKT to activate AKT pathway,^{30,40} and AKT1 regulates pathological angiogenesis, vascular maturation, and permeability.⁴¹ All these may explain why the patients with IKBKE+/TBK1+ co-expression were prone to recurrence with metastasis in our study and the IKBKE+/TBK1+ subgroup has shorter OS than other subgroups ($P=0.014$), whereas we did not find statistically significant PFS differences between the four subgroups ($P=0.109$). Li et al showed IKBKE




Figure 2 The Kaplan–Meier overall survival (A) and progression free survival (B) curves for stage I NSCLC according to IKBKE and TBK1 co-expression stratification. **Abbreviations:** OS, overall survival; PFS, progression free survival.

and TBK1 dual inhibitors could suppress AKT activation and reduce VEGF expression, leading to impaired angiogenesis and inhibition of tumor growth.⁴² Furthermore, several studies have shown that the correlation between high expression of IKBKE and chemotherapy resistance. Guo et al²² verified that IKBKE is upregulated on mRNA and protein levels in 63 ovarian cancer tissues of 96 ovarian cancer patients, and elevated IKBKE obviously enhances the ability of ovarian cancer cells to resist cisplatin. Besides, Zhang et al⁴³ demonstrated that PLK4 decreases temozolomide sensitivity by regulating the IKBKE/NF- κ B axis which indicates IKBKE could influence glioblastoma proliferation and chemosensitivity. In addition to ovarian cancer and glioblastoma, Guo et al²⁴ reported that the expression of IKBKE was increased in 98 cases of NSCLC, and the sensitivity of NSCLC cell lines to chemotherapy drugs was increased after silencing IKBKE, suggesting that IKBKE plays a key role in lung cancer and tumor resistance. Zhang et al⁴⁴ also demonstrated that TBK1 contributes to tumor invasion and might be a driving factor for metastatic spread of breast cancer. Thus, IKBKE/TBK1 dual inhibitors combined with adjuvant chemotherapy may be a potential strategy for high-risk stage I NSCLC patients.

Apparently, our study has several limitations that should be considered. The retrospective character of this study exists inevitable bias. Considering the limited amount of tissue, we cannot test as many as possible relevant proteins. Moreover, when the patient relapsed, we were unable to obtain the specimen, but only analyzed the specimen that was surgically removed at the initial diagnosis, which limited our understanding of the changes in tumor molecular characteristics.

In conclusion, the current study provides further insights for tumor targeted therapy and the landscape of kinase proteins IKBKE and TBK1 in various cancers. To our knowledge, our study is the first study to investigate the relationships between expression of IKBKE and TBK1 and clinicopathologic characteristics in stage I NSCLC patients and their predictive significance on patients' survival. We found IKBKE+/TBK1+ co-expression was significantly obvious in patients with risk factors and the recurrence pattern of metastasis, and is an effective prognostic predictor for poor OS. Of note, further investigation is required to identify optimal patients who likely obtain benefit from IKBKE and TBK1 dual inhibitors therapy.

Ethical statement

All procedures performed in studies involving human participants were in accordance with ethical standards of the institutional and national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Acknowledgment

This work was financially supported by a research grant from the Construction and Verification of Precise Medical Diagnosis and Treatment System for Non-small Cell Lung Cancer (2016CYJS01A03).

Disclosure

The authors declare that they have no conflicts of interest in this work.

References

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin*. 2013;63:11–30. doi:10.3322/caac.21166
2. D'Addario G, Felip E; Group EGW. Non-small-cell lung cancer: ESMO clinical recommendations for diagnosis, treatment and follow-up. *Ann Oncol*. 2009;20:68–70. doi:10.1093/annonc/mdp132
3. Santarpia M, Altavilla G, Pitini V, Rosell R. Personalized treatment of early-stage non-small-cell lung cancer: the challenging role of EGFR inhibitors. *Future Oncol*. 2015;11:1259–1274. doi:10.2217/fon.14.320
4. Maeda R, Yoshida J, Ishii G, Hishida T, Nishimura M, Nagai K. Risk factors for tumor recurrence in patients with early-stage (stage I and II) non-small cell lung cancer: patient selection criteria for adjuvant chemotherapy according to the seventh edition TNM classification. *Chest*. 2011;140:1494–1502. doi:10.1378/chest.10-3279
5. Custodio AB, Gonzalez-Larriba JL, Bobokova J, et al. Prognostic and predictive markers of benefit from adjuvant chemotherapy in early-stage non-small cell lung cancer. *J Thorac Oncol*. 2009;4:891–910. doi:10.1097/JTO.0b013e3181a4b8fb
6. Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-cell lung cancer. *N Engl J Med*. 2005;352:2589–2597. doi:10.1056/NEJMoa043623
7. Verhelst K, Verstrepen L, Carpentier I, et al. IkappaB kinase epsilon (IKKepsilon): a therapeutic target in inflammation and cancer. *Biochem Pharmacol*. 2013;85:873–880. doi:10.1016/j.bcp.2013.01.007
8. Chien Y, Kim S, Bumeister R, et al. RalB GTPase-mediated activation of the I κ B family kinase TBK1 couples innate immune signaling to tumor cell survival. *Cell*. 2006;127(1):0–170. doi:10.1016/j.cell.2006.08.034
9. Shimada T, Kawai T, Takeda K, et al. IKK-i, a novel lipopolysaccharide-inducible kinase that is related to IkappaB kinases. *Int Immunol*. 1999;11:1357–1362. doi:10.1093/intimm/11.8.1357
10. Larabi A, Dev JM, Ng SL, et al. Crystal structure and mechanism of activation of TANK-binding kinase 1. *Cell Rep*. 2013;3:734–746. doi:10.1016/j.celrep.2013.01.034
11. Kawai T, Akira S. Signaling to NF-kappaB by Toll-like receptors. *Trends Mol Med*. 2007;13(11):460–469. doi:10.1016/j.molmed.2007.09.002
12. Chau TL, Gioia R, Gatot J-S, et al. Are the IKKs and IKK-related kinases TBK1 and IKK- γ similarly activated? *Trends Biochem Sci*. 2008;33(4):171–180. doi:10.1016/j.tibs.2008.01.002

13. Pomerantz JL, Baltimore D. NF-kappaB activation by a signaling complex containing TRAF2, TANK and TBK1, a novel IKK-related kinase. *Embo J*. 1999;18(23):6694–6704. doi:10.1093/emboj/18.23.6694
14. Nomura F, Kawai T, Nakanishi K, Akira S. NF-kappaB activation through IKK-i-dependent I-TRAF/TANK phosphorylation. *Genes Cells*. 2000;5(3):191–202.
15. Gatot JS, Gioia R, Chau TL, et al. Lipopolysaccharide-mediated interferon regulatory factor activation involves TBK1-IKKε-dependent Lys63-linked polyubiquitination and phosphorylation of TANK/I-TRAF. *J Biol Chem*. 2007;282(43):31131. doi:10.1074/jbc.M701690200
16. Fujita F, Taniguchi Y, Kato T, et al. Identification of NAP1, a regulatory subunit of IκB kinase-related kinases that potentiates NF-κB signaling. *Mol Cell Biol*. 2003;23(21):7780–7793. doi:10.1128/mcb.23.21.7780-7793.2003
17. Ryzhakov G, Randow F. SINTBAD, a novel component of innate antiviral immunity, shares a TBK1-binding domain with NAP1 and TANK. *Embo J*. 2007;26(13):3180–3190. doi:10.1038/sj.emboj.7601743
18. Eddy SF, Guo S, Demicco EG, et al. Inducible IκappaB kinase/IκappaB kinase epsilon expression is induced by CK2 and promotes aberrant nuclear factor-kappaB activation in breast cancer cells. *Cancer Res*. 2005;65(24):11375. doi:10.1158/0008-5472.CAN-04-4557
19. Guan H, Zhang H, Cai J, et al. IKBKE is over-expressed in glioma and contributes to resistance of glioma cells to apoptosis via activating NF-κB. *J Pathol*. 2011;223(3):436–445. doi:10.1002/path.2815
20. Lee SE, Hong M, Cho J, et al. IKKε and TBK1 expression in gastric cancer. *Oncotarget*. 2016;8(10):16233–16242.
21. Cheng A, Guo J, Henderson-Jackson E, Kim D, Malafa M, Coppola D. Ikb kinase ε expression in pancreatic ductal adenocarcinoma. *Am J Clin Pathol*. 2011;136(1):60–66. doi:10.1309/AJCP2JJGYNIUAS2V
22. Guo JP, Shu SK, He L, et al. Deregulation of IKBKE is associated with tumor progression, poor prognosis, and cisplatin resistance in ovarian cancer. *Am J Pathol*. 2009;175(1):324–333. doi:10.2353/ajpath.2009.080767
23. Ghatalia P, Yang ES, Lasseigne BN, et al. Kinase gene expression profiling of metastatic clear cell renal cell carcinoma tissue identifies potential new therapeutic targets. *PLoS One*. 2016;11(8):e0160924. doi:10.1371/journal.pone.0160924
24. Guo J, Kim D, Gao J, et al. IKBKE is induced by STAT3 and tobacco carcinogen and determines chemosensitivity in non-small cell lung cancer. *Oncogene*. 2013;32(2):151–159. doi:10.1038/onc.2012.39
25. Seo SI, Song SY, Kang MR, et al. Immunohistochemical analysis of NF-kappaB signaling proteins IKKepsilon, p50/p105, p52/p100 and RelA in prostate cancers. *Apmis*. 2009;117:623–628. doi:10.1111/j.1600-0463.2009.02506.x
26. Kim JY, Welsh EA, Oguz U, et al. Dissection of TBK1 signaling via phosphoproteomics in lung cancer cells. *Proc Natl Acad Sci USA*. 2013;110:12414–12419. doi:10.1073/pnas.1220674110
27. Ma X, Helgason E, Phung QT, et al. Molecular basis of tank-binding kinase 1 activation by transautophosphorylation. *Proc Natl Acad Sci USA*. 2012;109:9378–9383. doi:10.1073/pnas.1121552109
28. Zhou AY, Shen RR, Kim E, et al. IKK^γ-mediated tumorigenesis requires K63-linked polyubiquitination by a cIAP1/cIAP2/TRAF2 E3 ubiquitin ligase complex. *Cell Rep*. 2013;3:724–733. doi:10.1016/j.celrep.2013.02.004
29. Rokavec M, Luo JL. The transient and constitutive inflammatory signaling in tumorigenesis. *Cell Cycle*. 2012;11:2587–2588. doi:10.4161/cc.21139
30. Gulen MF, Bulek K, Xiao H, et al. Inactivation of the enzyme GSK3α by the kinase IKKi promotes AKT-mTOR signaling pathway that mediates interleukin-1-induced Th17 cell maintenance. *Immunity*. 2012;37(5):800–812. doi:10.1016/j.immuni.2012.08.019
31. Saxton RA, Sabatini DM. mTOR signaling in growth, metabolism, and disease. *Cell*. 2017;168:960–976. doi:10.1016/j.cell.2017.02.004
32. Yu J, Zhou X, Chang M, et al. Regulation of T-cell activation and migration by the kinase TBK1 during neuroinflammation. *Nat Commun*. 2015;6:6074. doi:10.1038/ncomms7074
33. Bodur C, Kazyken D, Huang K, et al. The IKK-related kinase TBK1 activates mTORC1 directly in response to growth factors and innate immune agonists. *Embo J*. 2018;37:19–38. doi:10.15252/embj.201696164
34. Zhang H, Chen L, Cai SH, Cheng H. Identification of TBK1 and IKKε, the non-canonical IκB kinases, as crucial pro-survival factors in HTLV-1-transformed T lymphocytes. *Leuk Res*. 2016;46:37–44. doi:10.1016/j.leukres.2016.04.012
35. Qin B, Cheng K, Gite S, Ostendorff HP, Narod S, Rothschild KJ. Silencing of the IKKε gene by siRNA inhibits invasiveness and growth of breast cancer cells. *Breast Cancer Res*. 2010;12(5):R74. doi:10.1186/bcr2722
36. Li H, Chen L, Zhang A, et al. Silencing of IKKε using siRNA inhibits proliferation and invasion of glioma cells in vitro and in vivo. *Int J Oncol*. 2012;41(1):169–178. doi:10.3892/ijo.2012.1452
37. Lu J, Yang Y, Guo G, et al. Ikbke regulates cell proliferation and epithelial-mesenchymal transition of human malignant glioma via the hippo pathway. *Oncotarget*. 2017;8(30):49502–49514. doi:10.18632/oncotarget.17738
38. Niederberger E, Möser C, Kynast K, et al. The non-canonical IjB kinafound IKBKE/TBK1-sensitive acute myeloid leukemia (AML) cells tend to have MYC ses IKKε and TBK1 as potential targets for the development of novel therapeutic drugs. *Curr Mol Med*. 2012;13:7.
39. Zhu Z, Aref AR, Cohoon TJ, et al. Inhibition of KRAS-driven tumorigenicity by interruption of an autocrine cytokine circuit. *Cancer Discov*. 2014. doi:10.1158/2159-8290.CD-13-0646
40. Guo JP, Coppola D, Cheng JQ. IKBKE protein activates AKT independent of phosphatidylinositol 3-kinase/PDK1/mTORC2 and the pleckstrin homology domain to sustain malignant transformation. *J Biol Chem*. 2011;286:37389–37398. doi:10.1074/jbc.M111.287433
41. Chen J, Somanath PR, Razorenova O, et al. AKT1 regulates pathological angiogenesis, vascular maturation and permeability in vivo. *Nat Med*. 2005;11(11):1188–1196. doi:10.1038/nm1307
42. Li J, Huang J, Jeong JH, et al. Selective TBK1/IKKi dual inhibitors with anticancer potency. *Int J Cancer*. 2014;134(8):1972–1980. doi:10.1002/ijc.28507
43. Zhang Z, Wang Z, Huang K, et al. PLK4 is a determinant of temozolomide sensitivity through phosphorylation of IKBKE in glioblastoma. *Cancer Lett*. 2019;443:91–107. doi:10.1016/j.canlet.2018.11.034
44. Zhang Y, Unnithan RVM, Hamidi A, et al. TANK-binding kinase 1 is a mediator of platelet-induced EMT in mammary carcinoma cell. *Faseb J*. 2019;fj201801936RRR. doi:10.1096/fj.201801936RRR

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>