



Published in final edited form as:

Chem Commun (Camb). 2019 July 23; 55(60): 8820–8823. doi:10.1039/c9cc03498f.

Multi-component Heteroarene Couplings via Polarity-reversed Radical Cascades

Jeremy M. Lear^{†,a}, J. Quentin Buquoi^{†,a}, Xin Gu^a, Kui Pan^a, Darsheed N. Mustafa^a, David A. Nagib^a

^aDepartment of Chemistry and Biochemistry, The Ohio State University, Columbus, OH 43210, United States.

Abstract

A multi-component radical addition strategy enables difunctionalization of alkenes with heteroarenes and a variety of radical precursors, including N_3 , $P(O)R_2$, and CF_3 . This unified approach for coupling diverse classes of electrophilic radicals and heteroarenes to vinyl ethers allows for direct, vicinal C-C as well as C-N, C-P, and C-R_f bond formation.

Graphical Abstract



A unified strategy enables multi-component radical addition cascades to couple alkenes, heteroarenes, and various radicals, including N_3 , $P(O)R_2$, and CF_3 .

The unique reactivity and selectivity accessible via radical addition to alkenes has significantly enabled the synthesis of valuable medicines and materials.¹ Specifically, in the realm of drug discovery, certain privileged elements and architectures (e.g. N, P, F, heteroarenes) are frequently incorporated into a medicinal candidate to improve its pharmacological properties.² Among the most common radical reactions used to introduce heteroarenes in this arena is the Minisci reaction.^{3,4} This two-component reaction (Fig 1a) is best suited for coupling nucleophilic radicals (e.g. α -oxy) to electrophilic heteroarenes (e.g. pyridine, quinoline). Conversely, the polarity-mismatched⁵ coupling of radicals and heteroarenes that are both electron-deficient is typically disfavored – with rare exceptions.⁶ However, Minisci also demonstrated that a polarity-reversing strategy could bring together three components in a radical cascade (Fig 1b).⁷ This underutilized approach chemoselectively combines an *in situ* generated electrophilic radical with an electron-rich

[†]These authors contributed equally.

Electronic Supplementary Information (ESI) available: Procedures, Characterization, and Spectroscopic data of all new compounds. See DOI: 10.1039/x0xx00000x

Conflicts of interest

There are no conflicts to declare

alkene in the presence of an electron-deficient heteroarene. The resulting nucleophilic radical then combines with the protonated heteroarene to afford a difunctionalized product after subsequent oxidation. In a pioneering example, Minisci generated α -carbonyl radicals from acetone using $\text{Ag}/\text{S}_2\text{O}_8$ to initiate this cascade.⁷ Later, refluxing peroxides were employed to enable the perfluoroalkyl radical ($\bullet\text{R}_f$) variant.⁸

Recent efforts have focused on extending this approach to incorporate other radical partners – and to do so under milder conditions that are more suitable to medicinal applications. For example, Barriault developed a bimetallic, gold-catalyst to incorporate α -carbonyl radicals ($\bullet\text{R}_{\text{EWG}}$).⁹ Herzon and Baran reported hydropyridylations ($\bullet\text{H}$) with Co and Fe complexes.¹⁰ Liu developed an azido variant ($\bullet\text{N}_3$) with iodanes, and Chu reported a photocatalytic version shortly thereafter.¹¹ Other photolytic methods for incorporating trifluoromethyl ($\bullet\text{CF}_3$) and phosphonyl radicals ($\bullet\text{P}(\text{O})\text{R}_2$) have been developed by Hong, Matsunaga, and us.¹² However, in looking at the divergent conditions required to perform each of these specific reactions, we questioned if a single, unified strategy could be developed to enable the incorporation of multiple classes of electrophilic radicals within this cascade. The advantage of such an approach would be its utility as a robust synthetic tool to rapidly access complex molecules with vicinal substituted functionality – especially for applications in medicinal chemistry.

Towards our goal of a unified strategy, we sought to employ hypervalent iodanes, which are robust precursors for numerous electrophilic radicals (Fig 1c).¹³ In particular, we hypothesized azido, phosphonyl, and trifluoromethyl radicals could all be accessed by iodane-scission reactions, as shown in Fig 2. In all three cases, a neutral or anionic reagent combines with a λ_3 -iodane either directly or otherwise to form an electrophilic radical. Azido radicals **A** are formed via direct ligand displacement of $\text{PhI}(\text{OAc})_2$ by TMSN_3 , followed by homolysis of the weak N-I bond.¹⁴ Conversely, we proposed that phosphonyl radicals **B** may be formed indirectly by $\text{H}\bullet$ abstraction of the weak P-H in $\text{Ph}_2\text{P}(\text{O})\text{H}$ by $\bullet\text{N}_3$.^{14,15} Finally, trifluoromethyl radicals **C** may be generated by ligand displacement of $\text{PhI}(\text{OTFA})_2$ with Langlois' reagent NaSO_2CF_3 .¹⁶

To our delight, upon subjecting these three, distinct radicalprecursors (TMSN_3 , $\text{Ph}_2\text{P}(\text{O})\text{H}$, and NaSO_2CF_3) to an electron-rich alkene (ethyl vinyl ether) and heteroarene (phenanthridine) in the presence of $\text{PhI}(\text{OAc})_2$ or $\text{PhI}(\text{OTFA})_2$, all three classes of cascade reactions were accessible (**1–3**; 60–92%). As shown in Fig 3, the development of these three reactions were largely enabled by the identity of the oxidant, as well as its rate of addition. For example, in the azido variant, an initial addition of $\text{PhI}(\text{OAc})_2$ affords only 20% yield (entry 1), likely due to rapid formation of $\bullet\text{N}_3$ in too high of concentration. Alternatively, slow addition via syringe pump allows for efficient formation of the azido product **1** (entry 2, 92% yield). This strongly beneficial effect of slow addition was also seen for other heteroarene traps (see SI).

Next, we investigated the phosphonyl cascade reaction. As expected, direct displacement of both $\text{PhI}(\text{OAc})_2$ and $\text{PhI}(\text{OTFA})_2$ by diphenyl phosphine oxide is inefficient, with up to 14% yield observed only in the case of slow oxidant addition (entries 3–5). However, addition of

TMSN₃ as an H-atom transfer mediator,¹⁴ enables efficient formation of phosphonyl product **2** (entry 6, 90% yield).

Finally, we investigated the trifluoromethyl radical-mediated variant of this cascade. Knowing that NaSO₂CF₃ does not efficiently displace the ligands of PhI(OAc)₂ at room temp, we used the more reactive oxidant, PhI(OTFA)₂. Interestingly, in the event of slow oxidant addition, slow rates of both iodane substitution and ensuing formation of •CF₃ resulted in low reactivity (entries 7–9). Thus, for the CF₃ cascade reaction, we found that addition of the reagent at t = 0 is optimal, affording trifluoromethyl product **3** (entry 10, 60% yield).

With a unified, iodane-mediated strategy in hand for heteroaryl difunctionalization of alkenes by a polarity-reversing radical cascade, we wished to investigate the generality of these three classes of reactions. In the azido three-component coupling (Fig. 4), we were pleased to find the phenanthridine-based protocol could be extended to include several other electronically diverse heteroarenes. For example, both isoquinolines (**4–10**) and quinolines (**11–12**), bearing various functional groups, including ketones, esters, amides, and halides, were tolerated. Additionally, in place of ethyl vinyl ether, 2,3-dihydrofuran could be used as the alkene partner to afford azido furan **13** as a single diastereomer (> 20:1 d.r.). Unfortunately, non-etheral alkenes were inefficient coupling partners – providing diminished reactivity in all three radical cascades.¹⁷

Due to the importance of phosphorus motifs in materials and medicines, we then turned our attention to exploring the scope of the phosphonyl three-component coupling (Fig. 5). As in the case of the azide variant, the incorporation of diphenyl phosphine oxide through phosphinylalkylation of heteroarenes was efficient and synthetically generalizable. Similarly, both isoquinolines (**14–20**, **23**) and quinolines (**21–22**) were tolerated. Synthetic utility was again demonstrated by the incorporation of various functional groups, including ketones, esters, amides, and halides. Notably, aryl bromide **20**, which may undergo deleterious halide abstraction by phosphonyl radicals, was tolerated. And as before, both the 2- and 4-positions of quinoline are susceptible to efficient addition when the other position is blocked with a substituent. To our delight, furan derivative **23** was also accessed from the cyclic precursor in high diastereoselectivity (> 20:1 d.r.).

Finally, given the ubiquity of trifluoromethyl groups in medicine, we also sought to explore the generality of the trifluoromethyl three-component coupling (Fig. 6). Despite the use of the more reactive oxidant, PhI(OTFA)₂, the reaction scope appears similarly general to the other two cascade reaction classes. To test this reaction in more complex and challenging cases, ketones and halides were included in the heteroarene component at varying positions without issue (**24–27**). Similarly, a cyclic alkene was employed – affording a single diastereomer of the fluoroalkylated heteroarene (> 20:1 d.r.).

In conclusion, we have demonstrated the development of a unified, modular strategy for three-component couplings to access alkyl heteroarenes in which three contiguous carbons contain biologically valuable substitution (N, P, R_f, O, pyridine). The synthetically facile approach employs hypervalent iodine reagents to generate azido, phosphonyl, and

trifluoromethyl radicals, which chemo- and regio-selectively combine with enol ethers and heteroarenes. The mildness of the platform permits wide functional group tolerance and enables the synthesis of diverse classes of heteroarenes with medicinal relevance.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Institutes of Health (NIH R35 GM119812), National Science Foundation (NSF CAREER 1654656), and the United States Air Force (J.Q.B.) for financial support.

References

1. Hart DJ, Science, 1984, 223, 883–887; M. Yan, J. C. Lo, J. T. Edwards and P. S. Baran, J. Am. Chem. Soc., 2016, 138, 12692–12714. [PubMed: 17781610]
2. Smith BR, Eastman CM and Njardarson JT, J. Med. Chem, 2014, 57, 9764–9773; E. Vitaku, D. T. Smith and J. T. Njardarson, J. Med. Chem., 2014, 57, 10257–10274. [PubMed: 25255063]
3. Duncton MAJ, Med. Chem. Commun, 2011, 2, 1135–1161; 10.1002/anie.201900977.
4. Recent examples of the Minisci reaction: Devari S and Shah BA, Chem. Commun, 2016, 52, 1490–1493; M. C. Quattrini, S. Fujii, K. Yamada, T. Fukuyama, D. Ravelli, M. Fagnoni and I. Ryu, Chem. Commun., 2017, 53, 2335–2338; L. Zhang, G. Zhang, Y. Li, S. Wang and A. Lei, Chem. Commun., 2018, 54, 5744–5747; A. H. Jatoi, G. G. Pawar, F. Robert and Y. Landais, Chem. Commun., 2019, 55, 466–469.
5. Roberts BP, Chem. Soc. Rev, 1999, 28, 25–35; F. De Vleeschouwer, V. Van Speybroeck, M. Waroquier, P. Geerlings and F. De Proft, Org. Lett., 2007, 9, 2721–2724.
6. Nagib DA and MacMillan DWC, Nature, 2011, 480, 224–228; Y. Ji, T. Brueckl, R. D. Baxter, Y. Fujiwara, I. B. Seiple, S. Su, D. G. Blackmond and P. S. Baran, Proc. Natl. Acad. Sci. U. S. A., 2011, 108, 14411–14415; Y. Fujiwara, J. A. Dixon, F. O'hara, E. D. Funder, D. D. Dixon, R. A. Rodriguez, R. D. Baxter, B. Herlé, N. Sach, M. R. Collins, Y. Ishihara and P. S. Baran, Nature, 2012, 492, 95–99. [PubMed: 22158245]
7. Citterio A, Gentile A and Minisci F, Tetrahedron Lett, 1982, 23, 5587–5590.
8. 18 Antonietti F, Mele A, Minisci F, Punta C, Recupero F and Fontana F, J. Fluor. Chem, 2004, 125, 205–211.
9. McCallum T and Barriault L, Chem. Sci, 2016, 7, 4754–4758. [PubMed: 30155127]
10. Ma X and Herzon SB, J. Am. Chem. Soc, 2016, 138, 8718–8721; J. C. Lo, D. Kim, C.-M. Pan, J. T. Edwards, Y. Yabe, J. Gui, T. Qin, S. Gutiérrez, J. Giacoboni, M. W. Smith, P. L. Holland and P. S. Baran, J. Am. Chem. Soc., 2017, 139, 2484–2503; X. Ma, H. Dang, J. A. Rose, P. Rablen and S. B. Herzon, J. Am. Chem. Soc., 2017, 139, 5998–6007. [PubMed: 27384921]
11. Liu Z and Liu Z-Q, Org. Lett, 2017, 19, 5649–5652; J. Chen, S. Zhu, J. Qin and L. Chu, Chem. Commun., 2019, 55, 2336–2339. [PubMed: 28961001]
12. He Y-T, Kang D, Kim I and Hong S, Green Chem, 2018, 20, 5209–5214; J. Q. Buquoi, J. M. Lear, X. Gu, and D. A. Nagib. ACS Catal., 2019, DOI: 10.1021/acscatal.9b01580 ; Y. Kumagai, N. Murakami, F. Kamiyama, R. Tanaka, T. Yoshino, M. Kojima, and S. Matsunaga, Org. Lett., 2019, DOI: 10.1021/acscatal.9b01580.1021/acs.orglett.9b01015; Y. Kumagai, N. Murakami, F. Kamiyama, R. Tanaka, T. Yoshino, M. Kojima, and S. Matsunaga, Org. Lett., 2019, DOI: 10.1021/acs.orglett.9b01015 .
13. Togo H and Katohgi M, Synlett, 2001, 2001, 0565–0581; A. Yoshimura and V. V. Zhdankin, Chem. Rev., 2016, 116, 3328–3435.
14. Fontana F, Minisci F, Yong MY and Lihua Z, Tetrahedron Lett, 1993, 34, 2517–2520; K. Matcha and A. P. Antonchick, Angew. Chem. Int. Ed., 2013, 52, 2082–2086; A. P. Antonchick and L.



Burgmann, *Angew. Chem. Int. Ed.*, 2013, 52, 3267–3271; K. Matcha, R. Narayan and A. P. Antonchick, *Angew. Chem. Int. Ed.*, 2013, 52, 7985–7989.

15. Williams RH and Hamilton LA, *J. Am. Chem. Soc.*, 1952, 74, 5418–5420; S. Deprèle and J.-L. Montchamp, *J. Org. Chem.*, 2001, 66, 6745–6755.
16. Yang Y-D, Iwamoto K, Tokunaga E and Shibata N, *Chem. Commun*, 2013, 49, 5510–5512. L. Shi, X. Yang, Y. Wang, H. Yang and H. Fu, *Adv. Synth. Catal.*, 2014, 356, 1021–1028.
17. Non-ethereal alkenes afford the following yields for each reaction: N₃ (53%), P (35%), and CF₃ (10%). A likely explanation is the decreased nucleophilicity of the intermediate, non-ketyl, alkyl radical is less suited to heteroarene addition than to competing pathways (e.g. oxidation or H-atom transfer).

a. Minisci reaction - a polarity-matched, two-component coupling



b. Polarity-reversing radical cascade for three-component coupling



c. Unified strategy for polarity-reversed radical couplings




Fig. 1.
Design of a unified strategy for polarity-reversal cascades





Fig. 2.
Diverse electrophilic radicals from hypervalent iodanes




entry	R• precursor	oxidant	addition	product	yield
1	TMSN ₃	PhI(OAc) ₂	t = 0	1	20%
2	TMSN₃	PhI(OAc)₂	2 mL / hr	1	92%
3	Ph ₂ P(O)H	PhI(OAc) ₂	t = 0	2	0%
4	Ph ₂ P(O)H	PhI(OTFA) ₂	t = 0	2	0%
5	Ph ₂ P(O)H	PhI(OAc) ₂	2 mL / hr	2	14%
6	Ph₂P(O)H	TMSN₃, PhI(OAc)₂	2 mL / hr	2	90%
7	NaSO ₂ CF ₃	PhI(OTFA) ₂	2 mL / hr	3	1%
8	NaSO ₂ CF ₃	PhI(OTFA) ₂	3 mL / hr	3	35%
9	NaSO ₂ CF ₃	PhI(OTFA) ₂	6 mL / hr	3	51%
10	NaSO₂CF₃	PhI(OTFA)₂	t = 0	3	60%

Fig. 3.

Development of three radical cascade reaction classes ^a

^a **N₃**: Heteroarene (0.1 mmol), TFA (1 eq), TMSN₃ (2 eq), alkene (4 eq), PhI(OAc)₂ (2 eq), CH₂Cl₂ (0.05 M), 23 °C, 30 min. **P**: Heteroarene (0.1 mmol), TFA (2 eq), TMSN₃ (1 eq), phosphine oxide (6 eq), alkene (2 eq), PhI(OAc)₂ (2 equiv), CH₂Cl₂ (0.05 M), 23 °C, 30 min. **CF₃**: Heteroarene (0.1 mmol), TFA (1 eq), NaSO₂CF₃ (2 eq), alkene (2 eq), PhI(OTFA)₂ (2 eq), MeCN (0.1 M), 23 °C, 45 min. Yields determined by NMR.

•N₃

azido alkylation



R = H, **4**, 58%
Me, **5**, 64%










**6**, 80%**7**, 66%**8**, 91%**9**, 79%**10**, 75%**11**, 59%**12**, 93%**13**, 72%, > 20:1 d.r.

Fig. 4.
Azido three-component coupling



•P

phosphonyl alkylation





Fig. 5.
Phosphonyl three-component coupling

•CF₃

trifluoromethyl alkylation




**24**, 65%R = OBz, **25**, 52%
Br, **26**, 42%**27**, 48%
> 20:1 d.r.

Fig. 6.
Trifluoromethyl three-component coupling