



# HHS Public Access

Author manuscript

*Bioorg Chem.* Author manuscript; available in PMC 2020 August 01.

Published in final edited form as:

*Bioorg Chem.* 2019 August ; 89: 103014. doi:10.1016/j.bioorg.2019.103014.

## Phototoxicity of 7-Oxycoumarins with keratinocytes in culture

Christophe Guillon<sup>1,\*</sup>, Yi-Hua Jan<sup>2</sup>, Diane E. Heck<sup>3</sup>, Thomas M. Mariano<sup>2</sup>, Robert D. Rapp<sup>1</sup>, Michele Jetter<sup>1</sup>, Keith Kardos<sup>1</sup>, Marilyn Whittemore<sup>4</sup>, Eric Akyea<sup>1</sup>, Ivan Jabin<sup>5</sup>, Jeffrey D. Laskin<sup>2</sup>, Ned D. Heindel<sup>1</sup>

<sup>1</sup>Department of Chemistry, Lehigh University, Bethlehem, PA 18015 USA

<sup>2</sup>Department of Environmental and Occupational Health, Rutgers University School of Public Health, Piscataway, NJ 08854 USA

<sup>3</sup>Department of Environmental Science, New York Medical College, Valhalla, NY, 10595 USA

<sup>4</sup>Buckman Laboratories, 1256 N. McLean Blvd, Memphis, TN 38108 USA

<sup>5</sup>Laboratoire de Chimie Organique, Université Libre de Bruxelles, B-1050 Brussels, BELGIUM

### Abstract


Seventy-one 7-oxycoumarins, 66 synthesized and 5 commercially sourced, were tested for their ability to inhibit growth in murine PAM212 keratinocytes. Forty-nine compounds from the library demonstrated light-induced lethality. None was toxic in the absence of UVA light. Structure-activity correlations indicate that the ability of the compounds to inhibit cell growth was dependent not only on their physiochemical characteristics, but also on their ability to absorb UVA light. Relative lipophilicity was an important factor as was electron density in the pyrone ring. Coumarins with electron withdrawing moieties – cyano and fluoro at C<sub>3</sub> – were considerably less active while those with bromines or iodine at that location displayed enhanced activity. Coumarins that were found to inhibit keratinocyte growth were also tested for photo-induced DNA plasmid nicking. A concentration-dependent alteration in migration on neutral gels caused by nicking was observed.

### Graphical Abstract

Forty-nine members of a set of seventy-one structurally-diverse 7-oxycoumarins bearing alkyl, alkenyl, alkynyl, halo, nitro, cyano, trifluoromethyl, and amino residues were found to display photo-induced toxicity (UVA light) against murine PAM212 keratinocytes in culture.

\*Corresponding author's chg3@lehigh.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Structure-activity effects were evident and depended on halogen content and ring position, electron-withdrawing moieties at position R<sub>1</sub>, and relative lipophilicity of the alkenyl and alkynyl side chains at R<sub>3</sub> and R<sub>4</sub>. A photo-induced DNA plasmid nicking was observed in phototoxic members of the set.

### Keywords

7-hydroxycoumarins; 7-oxycoumarins; furocoumarins; psoralens; methoxsalen; 8-MOP; phototoxicity; PAM212 keratinocytes; DNA photo-damage

### Introduction:

Simple monomeric 7-hydroxycoumarins (or their glycosides) as well as linear fused-ring furocoumarins, a.k.a. psoralens, are widely distributed in many dicotyledonous plant families, including the *Apiaceae* (*Umbelliferae*), *Asteraceae*, *Fabiaceae*, *Moraceae*, *Rosaceae*, *Rubiaceae*, *Rutaceae* and *Solanaceae* as well as in monocotyledonous plants, such as *Gramineae*, palms, lilies, and orchids<sup>(1)</sup> In traditional and in established medicine, these compounds have found broad pharmaceutical utility as anti-microbials, immune-modulators, anti-inflammatories, therapeutics for hyperproliferative skin diseases, anti-virals, and cancer chemotherapeutics.<sup>(2,3)</sup>

8-Methoxypsoralen, a.k.a., methoxsalen or 8-MOP, and many similar furocoumarins, have an especially long and extensive literature as phototoxin/phototherapeutics.<sup>(4-6)</sup> However, the story on phototoxicity of the 7-hydroxycoumarins is more limited and ambiguous. Several researchers have concluded that “simple coumarins (umbelliferone), show no phototoxic effects”<sup>(4, 7-9)</sup> But, nevertheless, a number of coumarins – substituted with electron donating groups – were reported to be photo-labile and undergoing a 2+2 cycloaddition (dimerization) at the C<sub>3</sub>-C<sub>4</sub> double bond<sup>(10)</sup> Mousavi demonstrated a modest phototoxicity (IC<sub>50</sub> 476 μM)

for umbelliferone (7-hydroxycoumarin) against the MCF-7 breast carcinoma cell line<sup>(11)</sup> In addition, Goard has shown that a conjugate containing 6-bromo-7-hydroxycoumarin, attached at C<sub>4</sub> to the antibiotic anisomycin, served as a photo-activated inhibitor of protein biosynthesis better than did the parent antibiotic itself against CHO or HEK293 cells<sup>(12)</sup> Indeed, some coumarins can display phototoxic effects while others can display a structure-specific avidity for DNA binding although not by intercalation. Coumarin, esculetin, and a set of pyridyl-substituted coumarins formed reversible complexes with DNA; physical measurements supported their binding loci to be in the minor groove by a weak hydrogen-bonding association of the DNA with the coumarin carbonyl<sup>(13-15)</sup> Most psoralens, however, intercalate DNA and photo-crosslink at the pyrimidine bases<sup>(16)</sup> Thus, coumarins and psoralens exert their pharmacology by different mechanisms.

The psoralens and the 7-hydroxycoumarins are related in several ways. Virtually all the laboratory synthetic pathways to the linear furocoumarins pass through the functionalized 7-hydroxycoumarins as the penultimate precursors of the resulting three-ring psoralens<sup>(16)</sup> Nature also uses these same 7-hydroxycoumarins on the biosynthetic pathway to the natural product psoralens<sup>(16, 17)</sup> Besides sharing structural features and a similar synthetic pathway, the psoralens and the 7-hydroxycoumarins also share comparable absorption and qualitatively similar fluorescence behaviors. Methoxsalen, which absorbs at 220, 250, and 305 nm, can be excited at 250 nm, and thereafter gives a weak fluorescence emission at 505 nm with a quantum yield of 0.002<sup>(18)</sup> Similarly, 7-hydroxycoumarin also has broad absorption bands at 220, 255, and 295 with a  $\lambda_{\text{max}}$  of 330 nm and weak shoulder end-absorption at 370 nm. However, when excited at 330 nm it emits such intense blue fluorescence at 455 nm (with an impressive quantum yield of 0.81) that the compound finds wide use as a fluorescent whitener in laundry detergents<sup>(19-21)</sup> Because of their acidic phenolic -OH, 7-hydroxycoumarins show a range of pH-dependent excitations and emissions with quantum yields approaching unity above pH 9.4<sup>(21)</sup>. The psoralens show no such pH-dependency.

Because the phenolic-capped O-acylated and the O-glycosylated 7-hydroxycoumarins – like the corresponding psoralens – display almost no fluorescence, they can serve as useful fluorogenic substrates for quantifying enzymatic activity (hydrolytic and glycolytic).<sup>(22-23)</sup> Thus, enzymatic liberation of the highly fluorescent free phenol provides a readily measured marker; molecular orbital calculations have been advanced to explain this phenomenon<sup>(24)</sup>

Intrigued by the structural commonalities, the spectral dissimilarities and the ambiguous reports on the photobiology of umbelliferones, a study of the phototoxic effects of seventy-one 7-hydroxycoumarins, their ethers and esters, (see Figure 1 and Table 1) was undertaken. It is well recognized that ultraviolet light irradiation in combination with chemical photosensitizers, such as psoralens, has considerable biological activity in the skin. In addition to causing erythema, sunburn, and tanning, the photoactivated psoralens cause alterations in keratinocyte growth and differentiation<sup>(25)</sup> Indeed, because of these biological responses, the psoralens have been used to treat many skin disorders, particularly, eczema and psoriasis<sup>(26)</sup> Many derivatives of psoralens have been evaluated in phototoxicity assays using keratinocyte growth inhibition to evaluate biological activity. In the present studies we used a mouse keratinocyte cell line to evaluate phototoxicity of a number of structural

analogs of the 7-oxycoumarins. PAM212 cells in culture are a common mouse keratinocyte line often used to investigate mechanisms of drug action since results with these cells can lead directly to *in vivo* mouse models and subsequently to human keratinocytes as lead pharmaceuticals are advanced in preclinical development.

## Results and Discussion:

While coumarins including the 7-hydroxy compounds, their ethers and alkylated analogs, have shown broad promise as anti-inflammatories, anti-tumor, anti-HIV, anticoagulant, antimicrobial, antioxidant, hepatoprotective, anti-pyretic, anti-hypertensive, and anti-convulsant agents, they nevertheless display complex pharmacologies, which are truly difficult to correlate.<sup>(27)</sup> The problem is that for any given compound library, no validated quantitative structure-linked comparative principle has yet to be found although logP, electronic surface analysis, pKa, and size have all been considered. In coumarin papers, structure-activity relations are phrased in general terms and even then, any given set displays many exceptions. For example, “for TNF- $\alpha$  inhibitory effects...the C<sub>6</sub> position must contain a halogen or methoxy...hydrophobic groups at C<sub>3</sub> are favored...small aliphatic groups at C<sub>4</sub> are optimal.”<sup>(27)</sup> Similar general statements are reported for anti-HIV coumarins<sup>(28)</sup> and for anti-fungal coumarins.<sup>(29)</sup> For instance, Kostova noted the anti-HIV activities for substituents on C<sub>4</sub>, were “methyl>H>propyl>benzyl” but he similarly noted that “to describe structure-activity relationships on the potent anti-HIV coumarin derivatives is not an easy task.” The anti-fungal SARs for a series of 24 diverse coumarins were equally problematic. Guerra reported that all ethers of 6-hydroxycoumarin were inactive in his anti-fungal assay (the parent phenol did have modest activity) but the ethers of 7-hydroxycoumarin were active and with declining effect as the 7-alkoxy chain lengthened.<sup>(29)</sup> No other correlations were made.

In the present studies, PAM212 mouse keratinocytes were used to evaluate the biological activity of the coumarins.<sup>(30)</sup> In this assay the PAM212 cells were treated with the coumarins and then exposed to UVA light. Control cells were treated with the coumarin without UVA light. The cells were then allowed to grow for 4-5 days and then analyzed for growth. Figure 1 shows typical keratinocyte growth inhibition experiments with several of the coumarins. Structure activity analysis for the photobiological activity of the coumarins is shown in Table 1. Without UVA light treatment, no significant growth inhibition by the coumarins is evident. However, following UVA light treatment, there is a marked inhibition of cell growth. Table 1 summarizes the concentration of the coumarins inhibiting growth by 50% (IC<sub>50</sub>) for the compounds in Figure 1 as well as other coumarins in the data set compounds (1–71).

Based on these data (Table 1), lipophilicity is an important factor. Compare for example the parent 7-hydroxycoumarins and their corresponding 7-alkoxy ethers.

Dimethylumbelliferone, (**12**) logP = 2.1 displayed a modest IC<sub>50</sub> of 100  $\mu$ M while all its ethers (**2** to **11**) displayed substantially improved activity. The lower chain length ethers, compounds **2** to **9**, (whose cLogP values fell between 3.2 and 3.9 were considerably more active than were the longer chain length ethers (**10** and **11**) for which cLogPs of > 5.8 reflected an enhanced – and apparently less optimal – lipophilicity.

Furthermore, when the alkenyl or alkyl moiety was rearranged from the 7-position (where it was an ether) onto either ring positions 3- or 6- (e.g., **19–24**), all of these 7-hydroxycoumarins were more active ( $IC_{50}$ 's 1–21  $\mu M$ ) than the parent dimethylumbelliferone (**12**) from which they were synthetically derived. In addition, all these 3- or 6- alkylated 7-hydroxycoumarins were more lipophilic than **12** with their cLogPs falling between 3.6 and 4.6.

Similarly, in the 6-iodo series, the parent phenol, 6-iododimethylumbelliferone (**16** with  $clogP = 3.38$ ) possessed an  $IC_{50}$  of 18  $\mu M$  – much more photo-active than its nor-iodo, the dimethylumbelliferone (**12**,  $logP = 2.10$ ) – but even here its more lipophilic ethers the 6-iodo-7-alkoxy derivatives (viz., **13–15**) with  $clogPs$  of 4.3 to 5.4, were all more active. The 4,8-dimethyl-6-nitro-7-hydroxycoumarin (**18**) is a member of the family of acidic ortho-nitrophenols (in general  $pK_a$ 's average 7.2 for o-nitrophenols and  $clogPs$  average 1.7), which are extensively ionized at physiological pH and thus might be expected to display poor membrane permeability. At a  $cLogP$  of 2.39 compound **18** possesses no photocytotoxicity up to its solubility maximum of 100  $\mu M$ . On the other hand, the 4,8-dimethyl-6-amino-7-allyloxycoumarin (**17**), a member of the family of weakly basic aromatic ortho amino ethers, is unionized at physiological pH and in this case with  $clogP = 2.94$  displays an  $IC_{50}$  of 5  $\mu M$ .

In the 8-iodo series (**27–30**) the parent phenol, 4-methyl-8-iodo-7-hydroxycoumarin (**27**,  $cLogP = 2.93$ ) was inactive while its more lipophilic 7-alkoxy variants (**28**, **29**, and **30** with  $cLogPs$  of 4.81, 4.16, and 3.10 respectively) had  $IC_{50}$ 's of 22, 50, and 70  $\mu M$  respectively. Here the activity of the alkoxy compounds does increase with the increasing lipophilicity as provided by the carbon chain but since no analog has more than a five-carbon side chain a maximum lipophilicity on the Hansch curve has not been attained.

In keeping with the trend that the alkoxy groups are more photo-active than their parent phenols, as noted umbelliferone (7-hydroxycoumarin = **1**) showed no activity in the screen; but its O-methyl ether (7-methoxycoumarin = **32**) had an  $IC_{50}$  of 50  $\mu M$  while its bis-methyl ether (7,8-dimethoxycoumarin = **31**) had an  $IC_{50}$  of 33  $\mu M$ .

The two families of fluoro-substituted coumarins constitute diverse classes. The first of these, those bearing a trifluoromethyl at  $R_2$ , position 4 in traditional coumarin numbering, (e.g., **33–47**), were largely inactive. The two parent phenols, compound **37** (4-trifluoromethyl-8-methyl-7-hydroxycoumarin) and the nor-methyl compound **47** (4-trifluoromethyl-7-hydroxycoumarin), were inactive at their solubility maxima. The traditional alkyl, alkenyl, or alkynyl ethers of **47** (e.g., **36**, **39**, **41**, **42**, and **46**) which in the other 7-hydroxy structural families did impart significant activity over their less active parent phenols, were equally inactive in this particular 4-trifluoromethyl family (the one with no 8-methyl moiety). However, three analogs of phenol **37** (the 4-trifluoromethyl-8-methyl-7-hydroxy system), were active (viz., **33–35**). Is this observed activity an effect of increased lipophilicity or is it derived from an electronic effect? The answer is probably both since two of the active derivatives (**33** with  $cLogP = 4.09$  and **35** with  $cLogP = 4.20$ ) were indeed much more lipophilic than **37** ( $cLogP = 3.00$ ), the inactive parent phenol, but the 7-O-acetonyl analog (**34**) displayed equivalent lipophilicity ( $clogP = 2.50$ ) to the parent and yet a

substantially enhanced 40  $\mu\text{M}$  activity. There may be an explanation for the activity ( $\text{IC}_{50} = 20\mu\text{M}$ ) seen in **33** with its pendant chloro-containing side chain. Platz has claimed an augmented photo-induced anti-viral activity for sensitizable heterocyclics with halogen-bearing side chains<sup>(44)</sup>.

The second family of fluoro-substituted coumarins, compounds **67** to **70** bearing an F at R<sup>1</sup>, position 3 in the traditional coumarin numbering, was similarly an inactive class. Not only the parent phenol (3-fluoro-4,8-dimethyl-7-hydroxycoumarin, **68**) but also the O-allyl and C-allyl analogs displayed no activity up to their solubility maxima. One exception, compound **67**, with a 2,3-dibromopropyl side chain, was modestly active at 120  $\mu\text{M}$  probably by a free radical effect to be discussed later.

We are not the first to observe that fluoro-attachment can be a deactivating structural feature in otherwise promising umbelliferones and other coumarins as candidate therapeutics. A  $\text{CF}_3$  attachment at ring positions 3 or 4 (R<sub>1</sub> and R<sub>2</sub> in the generalized coumarin structure, see **Figure** on Table 1) was found to abolish otherwise significant anti-inflammatory activity in a study of more than 50 diverse coumarins<sup>(27)</sup> In addition, fluorine attachment to the ring carbons of 7-hydroxycoumarins has been found to impart a photostability, an inertness to photo-reactivity and a resistance to fluorescence quenching; these properties all being problematic to the successful use of non-fluorinated umbelliferones as fluorophores in biological systems<sup>(22)</sup> The photo-inactivity of the two types of fluoro coumarins in this study is therefore to be expected.

Not surprisingly, the 3-cyano coumarins (*e.g.*, **62-66**) show an activity profile similar to the trifluoromethyl and fluoro coumarins. The cyano, the trifluoromethyl, and the fluoro groups are all electronically-deactivating entities with closely related electron withdrawing sigmas of 0.56, 0.43, and 0.34<sup>(45)</sup> Very few members of these three electron withdrawing families display phototoxicity and those that do are the side-chain brominated analogs. As in the 3-fluoro class (see **67**) so in the 3-cyano class only **62** and **63** ( $\text{IC}_{50}$ 's 3 and 48  $\mu\text{M}$ ), with their 2,3-dibromopropyl side chains, demonstrated the photosensitization effect.

One unique structure-activity correlation is that all fourteen 3-bromo- and 3-iodo-coumarins (**49-60** and **71**) in this study displayed significant photosensitizing effects with eight of the set having  $\text{IC}_{50}$ 's of 10  $\mu\text{M}$  or less. The heavy-halogen class of coumarins is the most photo-active. The significant phototoxic activity observed in the 3-brominated /iodinated coumarins (*viz.* **49-61** and **71**) as well as in those derived from dibrominating the C<sub>6</sub> (R<sub>3</sub>) allyl (*viz.* **49, 60, 62, 63, and 67**), may arise from the well-known photo-homolytic cleavage of these labile carbon-halogen bonds<sup>(46)</sup> Not only is there a synthetically useful photo-radical cleavage which uses the 3-brominated/iodinated coumarins as starting materials for a coupling reaction<sup>(47)</sup> but also such 3-bromo/iodo species have demonstrated potent photo-viricidal properties<sup>(48)</sup> Park and colleagues advanced a theory in which 3-bromoumbelliferone – with a 7-alkoxy side chain to provide affinity to a viral nucleic acid strand – was photolyzed to a free radical at C<sub>3</sub> which subsequently effected a cleavage of that strand<sup>(48)</sup> While other non-fluoro halogens (Cl, Br, and I) attached at C<sub>3</sub> on the coumarin ring were claimed efficacious as photo-activated radical producers<sup>(48)</sup>, Platz also noted that halogens on an aliphatic side chain off the coumarin ring were equally efficacious

as light activated anti-viral agents<sup>(44)</sup> In the set of coumarins (see **49**, **60**, **62**, **63** and **67**), it was demonstrated that the 2,3-dibromopropyl side chain substantially augmented the phototoxicity of these dibromoalkyl compounds over that of close structural analogs lacking that moiety.

In summary, we established structural trends within the set of 71 coumarin derivatives presented herein. None of the compounds displayed growth inhibitory activity in the absence of light but 49 of them did in the presence of UVA light. Lipophilicity played a key role in the activity of those compounds. Within structural subsets, it was observed that the more lipophilic derivatives were the most active ones as long as the lipophilicity of said compounds did not exceed a value beyond which the activity started to diminish again. The addition of one or multiple halogens, such as bromo, iodo and chloro, resulted in enhanced activity; whether those substitutions occurred at the traditional ring positions (C<sub>3</sub>, C<sub>6</sub> and C<sub>8</sub>) or on a more distal side chain. However, fluorine substitution, in the form of a single fluorine atom or a trifluoromethyl group, typically resulted in partial or complete loss of activity.

As psoralens are known to intercalate into DNA, and a few psoralens are known to bind to the minor groove of DNA<sup>(13-15)</sup>, we next determined if the photo-activated coumarins had the capacity to bind and damage DNA. For these studies, we used a plasmid DNA nicking and unwinding assay<sup>(30)</sup> Results obtained in this study demonstrate that there is a structure-specific photolytic interaction of coumarins and DNA. If the candidate coumarin is able to nick and linearize the plasmid upon light exposure, the change in electrophoretic migration of the altered DNA is readily evident, (see examples in Figure 2 and Table 2). A set of five structurally diverse coumarins was selected from a group that was found to strongly inhibit keratinocyte growth (see Figure 1) but which ranged from halogenated variants, to 7-hydroxyls to 7-allyloxys. There was little or no damage to the plasmid DNA by any of these compounds in the absence of UVA. However, with each of the coumarins (**14**, **48**, **58**, **62** and **71**), UVA light treatment caused concentration-dependent alteration in migration in neutral gels caused by nicking, which resulted in a faster migrating supercoiled plasmid when compared to slower migrating linear or open circular forms of the plasmid. Generally similar results were evident when the plasmid was treated with the coumarins and UVA light and then analyzed in alkaline gels (**14**, **48** and **71**). These data indicate that the photo-activated coumarins can modify the DNA plasmid. A direct structure activity correlation between the phototoxicity results (Table 1) and the plasmid nicking/unwinding results (Figure 2 and Table 2) would not be expected, nor was it observed. The phototoxicity study is a whole cell assay with attendant complications induced by membrane transport, the presence of growth media, and metabolism, while photo nicking is performed extracellularly in simple buffer solution. The C<sub>7</sub> hydroxyl molecules **71** and **69** as well as the pre-hydroxyl (the readily hydrolyzed C<sub>7</sub> acetoxy molecule) **48** all displayed nearly five times greater linearization of the plasmid compared to the 7-allyloxy ethers **14** and **58** although the latter were among the most phototoxic compounds of the set .

## Experimental:

### Materials, Methods and Instrumentation:

The 7-oxycoumarins were purchased or synthesized as indicated on Table 1, and in the following Syntheses.  $^1\text{H}$  NMR spectra were obtained on a Bruker 500MHz nuclear magnetic resonance spectrometer in  $\text{CDCl}_3$  or  $\text{DMSO-d}_6$ . All solvents and reagents were of the highest purity commercially available. Combustion analyses were within  $\pm 0.4$  of theoretical values (deviations are noted in the text) and NMR spectra (see below) were in accord with expected resonance shifts. The cLogPs reported herein were obtained from the Chemical Properties calculator in ChemDraw Professional version 17.0. If a measured logP is known, that value is used in the discussion.

### Syntheses:

Coumarins from commercial sources or published syntheses are noted in Table 1 by the name of the supplier or by reference to a published article. The letter S means the indicated coumarin was synthesized for this study. In some cases, in which the published synthesis lacked experimental details or physical properties, we have repeated that synthesis and the newly measured properties are reported below. Compounds not present in the text below (**8**, **9**, **31**, **32**, **50** and **61-70**) were prepared according to methods described in the references displayed in Table 1 for those compounds. In addition, the synthetic methods for all previously unreported coumarins are described below.

**4,8-Dimethyl-7-[(3-methyl-2-buten-1-yl)oxy]coumarin (3)**, **4,8-dimethyl-3-(3-methyl-2-buten-1-yl)-7-hydroxycoumarin (19)**, and **4,8-dimethyl-6-(3-methyl-2-buten-1-yl)-7-hydroxycoumarin (20)** were all produced in the etherification of **4,8-dimethyl-7-hydroxycoumarin (12)** when 1.9 g (10.0 mmol) of the latter compound were dissolved in 49 mL of aqueous sodium hydroxide (containing 0.40 g or 10.0 mmol) and reacted with 1-bromo-3-methyl-2-butene (1.5 g, 10.0 mmol) over 14 hr of vigorous stirring at ambient temperature. The precipitate which had formed was removed by filtration, washed on the filter with 20% aqueous sodium hydroxide, and the base-soluble phases combined. As expected the base insoluble compound (0.44 g, 17%) was the expected butenyl ether, **4,8-dimethyl-7-[(3-methyl-2-buten-1-yl)oxy]coumarin (3)**, mp = 93-94°C, recrystallized from MeOH). Elemental analysis and  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ ) supported the proposed structure:  $\delta$  (ppm) 2.17 and 2.44 (two s, each 3H, ring methyls), 1.68 and 1.83 (two d by second order splitting, each 3H,  $=\text{C}(\text{CH}_3)_2$ ), 4.69 (m, 2H,  $-\text{OCH}_2-$ ), 5.40 (m, 1H,  $-\text{CH}=\text{C}$ ), 6.33 (s, 1H,  $\text{C}_3\text{-H}$ ), 7.43 (d, 1H,  $\text{C}_5\text{-H}$ ,  $J = 8.4$  Hz), and 6.90 (d, 1H,  $\text{C}_6\text{-H}$ ,  $J = 8.4$  Hz).

Neutralization with 6 N hydrochloric acid of the aqueous basic phase precipitated an orange oil which was separated by flash chromatography (silica gel, chloroform:ethyl acetate 95:5 elutant) into three phenols – unreacted starting material (not recovered) and two different C-alkylated phenols. The first eluted fraction (0.55 g, 21%, mp = 199-200°C, recrystallized from MeOH), elemental analysis within 0.4% of theoretical and  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) supported the proposed structure of **4,8-dimethyl-3-(3-methyl-2-buten-1-yl)-7-hydroxycoumarin (19)**:  $\delta$  (ppm) 2.32 and 2.36 (two s, each 3H, ring methyls), 1.68 (d, 3H,  $J = 1.2$  Hz,  $=\text{C}(\text{CH}_3)_2$ ), 1.78 (s, 3H,  $=\text{C}(\text{CH}_3)$ ), 3.35 (d, 2H,  $J = 6.5$  Hz,  $-\text{CH}_2-$ ), 5.08 (dd,



1H, J = 6.5 Hz, J' = 1.2 Hz, -CH=), 5.86 (br s, 1H, -OH, exchanges with D<sub>2</sub>O), 6.88 (d, 1H, J = 8.5 Hz, C<sub>6</sub>-H), and 7.27 ppm (d, 1H, J = 8.5 Hz, C<sub>5</sub>-H). The second component to elute was isolated as off-white short needles (0.31 g, 12%, mp = 165-167°C, recrystallized from MeOH), elemental analysis within 0.3% of theoretical and <sup>1</sup>H NMR (CDCl<sub>3</sub>) supported the proposed structure of **4,8-dimethyl-6-(3-methyl-2-buten-1-yl)-7-hydroxycoumarin (20)**: δ (ppm) 2.36 (s, 3H, C<sub>8</sub>-CH<sub>3</sub>), 2.39 (d, 3H, J = 1.2 Hz, C<sub>4</sub>-CH<sub>3</sub> coupled to C<sub>3</sub>-H), 1.81 (two overlapping closely spaced s, 6H, =C(CH<sub>3</sub>)<sub>2</sub>), 3.42 (d, 2H, J = 6.8 Hz, -CH<sub>2</sub>-), 5.30 (m, 1H, J = 6.8 Hz, J' = 1.2 Hz, -CH=), 5.95 (br s, 1H, phenolic -OH exchanges with D<sub>2</sub>O), 6.12 (q, 1H, J = 1.2 Hz, C<sub>3</sub>-H coupled to C<sub>4</sub>-CH<sub>3</sub>), and 7.18 ppm (s, 1H, C<sub>5</sub>-H).

**4,8-dimethyl-6-allyl-7-hydroxycoumarin (21)**, mp = 168-170°C and **4,8-dimethyl-6-allyl-7-acetoxycoumarin (24)**, mp = 149.5-150°C were prepared as reported.<sup>(34)</sup>

**Preparation of 4,8-dimethyl-7-[(2-buten-1-yl)oxy]coumarin (7) and 4,8-dimethyl-6-(2-buten-1-yl)-7-hydroxycoumarin (23)**. To a solution of 0.95 g (5.0 mmol) of **4,8-dimethyl-7-hydroxycoumarin (12)**, in 30 mL of aqueous sodium hydroxide containing 0.2 g (5 mmol) of the base, was added 1-chloro-2-butene (0.68 g, 5 mmol). The resulting solution was stirred at room temperature for 18 h, acidified with 10% hydrochloric acid, chilled, and filtered. The dried crude product was subjected to flash chromatography (silica gel, 99:1 methylene chloride:2-propanol elutant) with the first product fraction identified as **4,8-dimethyl-7-[(2-buten-1-yl)oxy]coumarin (7)**, 0.44 g, 36%, mp = 114-116°C, recrystallized from methanol/ethyl acetate and the second product fraction identified as **4,8-dimethyl-6-(2-buten-1-yl)-7-hydroxycoumarin (23)**, 0.47 g, 39%, mp = 158-160°C. Alternatively, **7** can be prepared without contamination by **23** if the acetone, potassium carbonate phenolic alkylation of **12** is used.<sup>(49)</sup>

**4,8-Dimethyl-6-(3-buten-2-yl)-7-hydroxycoumarin (22)** was prepared by the Claisen rearrangement of 0.40 g (16 mmol) of **4,8-dimethyl-7-[(2-buten-1-yl)oxy]coumarin (7)** refluxed for eight hours under a dry nitrogen stream in 30 mL of N,N-diethylaniline. The cooled reaction contents were poured into 40 mL of chloroform and extracted 3X with 30 mL each of 25% hydrochloric acid. The organic layer was washed with 30 mL of water, dried over anhydrous sodium sulfate, and evaporated *in vacuo* to a yellow-brown solid, which was crystallized from methanol to a fluffy beige solid (0.31 g, 78% yield, mp = 175-176°C). Elemental analysis was within ± 0.1% of theoretical and <sup>1</sup>H NMR confirmed the structure as **4,8-dimethyl-6-(3-buten-2-yl)-7-hydroxycoumarin (22)**. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ (ppm) 1.41 (d, 3H, J = 6.8 Hz, C-CH<sub>3</sub>), 2.32 (s, 3H, C<sub>8</sub>-CH<sub>3</sub>), 2.42 (d, 3H, J = 0.98 Hz, C<sub>4</sub>-CH<sub>3</sub>), 3.80 (m, 1H, -CH<sub>2</sub>CH<sub>3</sub>), 5.15-5.35 (m, 2H, =CH<sub>2</sub>), 6.12 (q, 1H, J = 0.98 Hz, C<sub>3</sub>-H), 7.24 (s, 1H, C<sub>5</sub>-H).

These **4,8-dimethyl-7-alkoxycoumarins (2, 4, 5, 6, 10 and 11)** were prepared by alkylation of **4,8-dimethyl-7-hydroxycoumarin (12)** (Regis Chemical Company) by the general method for phenolic etherification.<sup>(49)</sup> The requisite 7-hydroxycoumarin (1.0 mmol), the corresponding alkyl/alkenyl/alkynyl bromide (1.5 mmol), 50 mL of anhydrous acetone, and 3 g of potassium carbonate were stirred vigorously at reflux for 16 hr, chilled, filtered, and the solid residue washed on the filter with a minimum of boiling methanol. The organic phase (acetone, methanol and coumarin ether product) was evaporated and the crude solid

trituated with 20% aqueous sodium hydroxide to remove any unreacted 7-hydroxycoumarin. The dried organic solid was crystallized from methanol to yield the pure ethers in 45-79% yields with elemental analyses ( $\pm 0.4\%$  of theory) and  $^1\text{H NMR}$  ( $\text{CDCl}_3$ ) in accord with the expected: **(2)** mp = 145 -146°C, **(4)** mp 83-84°C, **(5)** mp 106-108°C, **(6)** mp 103-104°C, **(10)** mp = 87.0-88.5°C, and **(11)** mp = 80-82°C.

**Dimethyl-7-allyloxycoumarin (26)** was prepared as described by Kaufman.<sup>(36)</sup>

**4,8-Dimethyl-3-bromo-6-[3-(1,3-dioxoisindolin-2-yl)prop-1-en-1-yl]-7-acetoxycoumarin (59)** was prepared by the heating with vigorous stirring at 100°C of 1.85 g (10 mmol) of potassium phthalimide in 50 mL of dimethylacetamide with 2.55 g (5 mmol) of **4,8-dimethyl-3-bromo-6-(2,3-dibromopropyl)-7-acetoxycoumarin (60)**. The latter was prepared by our published method<sup>(41)</sup> At the end of five hours the reaction medium was chilled to room temperature in an ice-water bath and diluted with 300 mL of cold water to precipitate the organic materials. These were filtered, dried, taken up in methylene chloride and flash chromatographed. **59** was isolated (0.64 g, 26% yield, mp = 231-232 °C, recrystallized from  $\text{CH}_2\text{Cl}_2$ /hexane).  $^1\text{H NMR}$  ( $\text{CDCl}_3$ ) confirmed the structure:  $\delta$  (ppm) 2.22 and 2.37 (two s, each 3H,  $\text{C}_4\text{-CH}_3$  and  $\text{C}_8\text{-CH}_3$ ), 2.59 (s, 3H, -OAc), 4.44 (dd, 2H, - $\text{CH}_2\text{-N}$ ), 6.22 and 6.63 (two m, each 1H, side chain - $\text{CH}=\text{CH}$ -), 7.53 (s, 1H,  $\text{C}_5\text{-H}$ ), 7.73 and 7.85 ppm (two m, each 2H, phthalimido Ar-H).

**4,8-dimethyl-7-hydroxy-6-iodocoumarin (16): 4,8-dimethyl-7-hydroxycoumarin (12,** 10.0 g, 52.0 mmol) was dissolved in 80 mL of dioxane with 200 mL of concentrated aqueous ammonia subsequently added. A solution prepared from 13.6 g (53.5 mmol) of iodine in 500 mL of aqueous 5% KI was added dropwise with vigorous stirring over a two-hour period. The greenish color of the solution gradually disappeared and a yellow precipitate formed. The pH was adjusted to 3.0 with hydrochloric acid, cooled, and filtered to remove the product, which was taken up in a minimum of ethyl acetate and washed with water ( $3 \times 50$  ml). Evaporation of the ethyl acetate gave 15.7 g (94%) of the title compound which was recrystallized from MeOH to analytical purity: elemental analysis (C,H), mp = 219-220°C.<sup>(33)</sup>  $^1\text{H NMR}$  ( $\text{DMSO-d}_6$ ) confirmed that iodination had taken place on  $\text{C}_6$ :  $\delta$  (ppm) 2.25 and 2.36 (two s, each 3H,  $\text{CH}_3$ 's), 6.13 (s, 1H,  $\text{C}_3\text{-H}$ ), 7.84 (s, 1H,  $\text{C}_5\text{-H}$ ), 10.11 (br s, 1H, OH, exchangeable with  $\text{D}_2\text{O}$ ).

**4-Methyl-8-iodo-7-hydroxycoumarin (27)** was prepared as described in 35% yield, mp as reported 254-256°C.<sup>(32)</sup>

**4-Methyl-7-alkoxy-8-iodocoumarins, (28-30)**. A solution of **27** (3.87 g, 12.8 mmol), 2 g of 1-chloro-3-methyl-2-butene (19.1 mmol), 8.0 g potassium carbonate (58 mmol), and 200 mL of acetone was stirred at reflux for four hrs. After filtration the potassium carbonate was washed on the filter three times with hot acetone. Combined filtrates and washings were evaporated and the product, **4-methyl-7-[(3-methyl-2-buten-1-yl)oxy]-8-iodocoumarin (28)**, isolated as white microneedles (2.60 g, 55% yield, mp = 151.5-153°C, recrystallized from ethyl acetate). Elemental analysis (C,H,I within  $\pm 0.2\%$  of theory) and  $^1\text{H NMR}$  ( $\text{CDCl}_3$ ) confirmed the structure:  $\delta$  (ppm) 1.77 and 1.79 (two s, each 3H,  $=\text{C}(\text{CH}_3)_2$ ), 2.42 (d, 3H,  $\text{J} = 1.2$  Hz,  $\text{C}_4\text{-CH}_3$ ), 4.65 (d, 2H,  $\text{J} = 6.8$ , - $\text{OCH}_2$ ), 5.56 (dt, 1H,  $\text{J} = 6.8$ ,  $\text{J}' = 1.1$ , -

CH=), 6.13 (d, 1H, J = 1.2, C<sub>3</sub>-H), 6.83 (d, 1H, J = 8.8, C<sub>6</sub>-H), 7.51 (d, 1H, J = 8.8, C<sub>5</sub>-H). Similarly, with n-propyl bromide as the alkylating agent **4-methyl-7-n-propoxy-8-iodocoumarin (29)** was obtained (49% yield, mp = 129-130°C, recrystallized from ethanol). Elemental analysis (C,H,I within ± 0.15% of theoretical) and <sup>1</sup>H NMR (CDCl<sub>3</sub>) confirmed the assigned structure: δ (ppm) 1.13 (t, 3H, J = 7.33, -CH<sub>3</sub>), 1.85 (m, 2H, -CH<sub>2</sub>-), 2.41 (s, 3H, C<sub>4</sub>-CH<sub>3</sub>), 4.10 (t, 2H, J = 6.36, -OCH<sub>2</sub>-), 6.14 (s, 1H, C<sub>3</sub>-H), 6.77 (d, 1H, J = 8.8, C<sub>6</sub>-H), 7.54 (d, 1H, J = 8.8, C<sub>5</sub>-H).

Similarly, by the above method using methyl iodide as the alkylating agent, a 54% yield of **4-methyl-7-methoxy-8-iodocoumarin (30)** was obtained (mp = 194-195°C, recrystallized from ethanol). Elemental analysis (C,H,I within ± 0.1% of theoretical) and <sup>1</sup>H NMR (CDCl<sub>3</sub>) confirming of the assigned structure: δ (ppm) 2.42 (s, 3H, C<sub>4</sub>-CH<sub>3</sub>), 4.00 (s, 3H, -OCH<sub>3</sub>), 6.15 (s, 1H, C<sub>3</sub>-H), 6.82 (d, 1H, J = 8.8, C<sub>6</sub>-H), 7.57 (d, 1H, J = 8.8, C<sub>5</sub>-H).

**Syntheses of (13-15). 4,8-dimethyl-6-iodo-7-alkoxycoumarins** The allyl, 3-methyl-2-buten-1-yl, and n-propyl ethers were prepared by the general method of Vyas<sup>(49)</sup> from condensation of **4,8-dimethyl-6-iodo-7-hydroxycoumarin (16)** and the corresponding bromides (*viz.* allyl bromide, 3-methyl-2-butenyl bromide, and n-propyl bromide) in dry acetone over potassium carbonate in yields of 60-85% (recrystallization EtOH). Elemental analyses (C,H) and <sup>1</sup>H NMR were in accord with theoretical values and respective mp's of the purified ethers were (**13**) 158-159.5°C, (**14**) 146-147°C, and (**15**) 124.5-126°C.

**Syntheses of (52-57): 8-methyl-3-bromo-7-alkoxycoumarins.** The allyl, 3-methyl-2-buten-1-yl, n-propyl, propargyl, and methyl ethers were prepared by the general method.<sup>(49)</sup> The parent coumarin, **8-methyl-3-bromo-7-hydroxycoumarin (57)**, mp = 209-211°C, was prepared as described<sup>(42)</sup> and subsequently condensed with these corresponding bromides (*viz.* allyl, 3-methyl-2-butenyl, n-propyl, propargyl, and methyl bromide) in dry acetone over potassium carbonate in yields of 68-90%. Elemental analyses (C,H) and <sup>1</sup>H NMR were in accord with theoretical values and the mp's of the respective purified ethers were (**52**) 143-143.5°C, (**53**) 131.5-133°C, (**54**) 139.5-140°C, (**55**) 181-182°C, and (**56**) 154 - 155°C.

(**58**) **4,8-dimethyl-3-bromo-7-allyloxycoumarin** was prepared in 52% yield by allyl bromide alkylation in dry acetone/potassium carbonate of **4,8-dimethyl-3-bromo-7-hydroxycoumarin (51)** by the general method.<sup>(49)</sup> The starting material (**51**) was prepared as reported, mp = 269-271°C.<sup>(42)</sup> The title compound (**58**) appeared from MeOH as tiny white needles (mp = 147-148°C). <sup>1</sup>H NMR and elemental analysis were in accord with theoretical values.

(**37**) **4-trifluoromethyl-8-methyl-7-hydroxycoumarin** and (**47**) **4-trifluoromethyl-7-hydroxycoumarin**, were prepared as described<sup>(50)</sup> and had melting points (197-198°C and 186-188°C respectively) and <sup>1</sup>H NMRs in accord with those expected.

**Synthesis of (34, 35, 38, 44, and 45), 4-trifluoromethyl-8-methyl-7-alkoxycoumarins.** The 2-ketopropyl, n-propyl, propargyl, allyl, and methyl ethers were prepared by the general method<sup>(49)</sup> from condensation of **4-trifluoromethyl-8-methyl-7-hydroxycoumarin (37)** with chloroacetone, n-propyl bromide, propargyl bromide, allyl bromide and methyl iodide

in yields of 31 to 56% (recrystallization from EtOH). Elemental analyses (C,H) and  $^1\text{H}$  NMR were in accord with theoretical values and respective mp's of the purified ethers were **(34)** 143-144.5°C, **(35)** 112-113°C, **(38)** 160.5-161.5°C, **(44)** 114.5-115.5°C, and **(45)** 109.5-111°C.

Synthesis of **(36, 39, 41, 42, and 46)**. These **4-trifluoromethyl-7-alkoxycoumarins** were prepared by etherification of **4-trifluoromethyl-7-hydroxycoumarin (47)** with allyl bromide, n-propyl bromide, methyl iodide, propargyl bromide, and chloroacetone in 28 to 58% yields (recrystallized from EtOH) with elemental analyses (C,H within  $\pm 0.3\%$  of theory) and  $^1\text{H}$  NMR in accord with theory. The respective mp's of the purified ethers were **(36)** 76-77°C, **(39)** 97-98°C, **(41)** 112.5-113.5°C, **(42)** 110.5-112°C, and **(46)** 158-159°C.

Syntheses of **(33, 43 and 40)**. These 6-allyl-substituted coumarins were prepared by the Claisen rearrangement of the corresponding 7-hydroxy ethers. The rearrangement procedure in N,N-diethylaniline as described above for preparation of compound **(22)** was followed. First, **4-trifluoromethyl-8-methyl-7-hydroxycoumarin (37)** was etherified by Vyas's method<sup>(49)</sup> with 2-chloroallyl chloride (also known as 2,3-dichloropropene) and with allyl chloride respectively. The crude ethers were evaporated to dryness from the acetone solvent used to prepare them and each was solubilized in N,N-diethylaniline (five times the weight of aniline per unit weight of allyl ether) and heated at reflux under nitrogen for 12 hours. Work-up and product isolation as described for **(22)** gave 24% yield of chloroallyl **(33)**, light tan needles, mp 182-183°C, with elemental analysis (C,H,F,C1 within  $\pm 0.3\%$  of theory) and  $^1\text{H}$  NMR in accord with theory. The same procedure for the 7-allyloxy compound **(44)** gave 34% of the rearranged 6-allyl-7-hydroxy **(43)** as off-white crystals, mp 174-175°C. Elemental analysis (C,H,F) was within  $\pm 0.3\%$  and  $^1\text{H}$  NMR was in accord with theory. Acetylation of **(43)** with acetic anhydride/sodium acetate gave 58% of the O-acetyl coumarin, compound **(40)**, mp 151-152°C (recrystallization from MeOH). Analytical data confirmed structure and purity.

**(18) 4,8-dimethyl-6-nitro-7-hydroxycoumarin. 4,8-Dimethyl-7-hydroxycoumarin (12,** 7.50 g, 39.4 mmol) was dissolved in 75 mL concentrated sulfuric acid at room temperature, chilled to  $-20^\circ\text{C}$ , and treated to the addition of a pre-chilled nitrating mixture (3 mL concentrated  $\text{HNO}_3$  added to 9 mL concentrated  $\text{H}_2\text{SO}_4$ ). Stirring was continued for three hrs at  $-20^\circ\text{C}$  with the mixture allowed to warm before pouring into ice. Bright yellow crystals were filtered, washed with water and dried to recover 7.50 g (81% yield) crude solid. The product **(18)** was recrystallized from ethanol to give yellow green crystals with mp = 229.5-231.5°C. C,H,N analysis was within  $\pm 0.25\%$  of theoretical.  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ ):  $\delta$  (ppm) 2.29 and 2.41 (two s, 3H each,  $\text{C}_4\text{-CH}_3$  and  $\text{C}_8\text{-CH}_3$ ), 6.37 (s, 1H,  $\text{C}_3\text{H}$ ), 8.19 (s, 1H,  $\text{C}_5\text{H}$ ), 11.30 (br s, 1H, -OH).  $^{13}\text{C}$  NMR ( $\text{DMSO-d}_6$ ):  $\delta$  (ppm) 8.75, 18.00, 112.45, 112.93, 115.12, 119.90, 132.80, 152.91, 154.18, 155.10, 158.96.

**(17) 4,8-Dimethyl-6-amino-7-allyloxycoumarin. 4,8-Dimethyl-6-nitro-7-hydroxycoumarin (18,** 4.70 g, 20 mmol) was added to 40 mL dry acetone with  $\text{K}_2\text{CO}_3$  (2.76 g, 20 mmol) in an oven dried flask. Allyl bromide (2.90 g, 24 mmol) in 40 mL of acetone was added slowly with heating and stirring. After five hrs of vigorous agitation at reflux the mixture was filtered and the solid on the filter washed with boiling acetone.

Evaporation of the filtrate gave 3.30 g (60%) of crude **4,8-dimethyl-6-nitro-7-allyloxy coumarin** which was employed directly in the next step. To a round bottom flask was added finely ground **4,8-dimethyl-6-nitro-7-allyloxy coumarin** (3.00 g, 11 mmol), tin (3.24 g, 27.3 mmol), SnCl<sub>2</sub> (3.04 g, 16.0 mmol) and 2 mL concentrated HCl in 200 mL ethanol. The mixture was stirred overnight during which all tin dissolved. Most of the ethanol was removed *in vacuo*. As the solution was allowed to cool, a gel formed which was filtered to recover a small amount of pearlized crystals. Diethyl ether was added to the filtrate and a precipitate formed which was filtered, and rinsed with ether to remove SnCl<sub>2</sub>. The solids were dried and extracted with hot ethanol on the funnel of a filtration flask. Evaporation of the ethanol gave a crude product that was recrystallized from ethanol to provide 2.20 g (82%) of the **4,8-dimethyl-6-amino-7-allyloxy coumarin**, (**17**). The bright mustard yellow crystals had a mp of 145.5-147.5°C. C,H,N analysis within ± 0.30% of theoretical. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ (ppm) 2.28 and 2.49 (two s, each 3H, C<sub>4</sub>-CH<sub>3</sub> and C<sub>8</sub>-CH<sub>3</sub>), 4.54 (d, 2H, -O-CH<sub>2</sub>-), 5.20 (br s, 2H, -NH<sub>2</sub>), 5.28 (d, 1H, =CH), 5.48 (d, 1H, =CH'), 6.16 (m, 1H, -CH=), 6.39 (s, 1H, C<sub>3</sub>-H), 7.58 (s, 1H, C<sub>5</sub>-H). <sup>13</sup>C NMR (DMSO-d<sub>6</sub>): δ (ppm) 9.43, 18.14, 74.57, 113.64, 115.13, 116.00, 118.69, 120.09, 125.52, 133.48, 147.0, 151.49, 152.50, 159.62.

**(71) 3-Iodo-4,8-dimethyl-7-hydroxy coumarin**. Finely pulverized **4,8-dimethyl-7-hydroxy coumarin** (**12**) (3.00 g, 15.7 mmol) was added to 150 mL concentrated acetic acid with subsequent dropwise addition of ICl (5.35 g, 33.0 mmol). This mixture was heated at 50°C with stirring overnight. The mixture was cooled and the crystals collected by filtration. The yield was 3.22 g (65%). The white crystals were purified by recrystallization from methanol, with mp = 228-230°C. C,H,N analysis was within + 0.10% of theoretical. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ (ppm) 2.14 and 2.58 (two s, each 3H, C<sub>4</sub>-CH<sub>3</sub> and C<sub>8</sub>-CH<sub>3</sub>), 6.84 (d, 1H, C<sub>6</sub>-H, J = 8.7 Hz), 7.55 (d, 1H, C<sub>5</sub>-H, J = 8.7 Hz), 10.53 (br s, 1H, -OH).

## Biological Evaluation – Experimental Methods:

### Growth inhibition assays:

Assays to measure keratinocyte growth inhibition were performed as previously described.<sup>(30)</sup> Briefly, PAM212 mouse keratinocytes (25,000 cells per well) were inoculated into 6-well Falcon plastic tissue culture dishes in Dulbecco's Modified Eagle's medium supplemented with 10% newborn calf serum in an incubator containing 5% carbon dioxide at 37°C. After 24 hr, the medium was changed to fresh growth medium supplemented with control medium or medium containing increasing concentrations of the coumarins. The medium contained 10% serum and 0.2% DMSO which facilitated the solubilization of the coumarins. After an additional 30 min, the uncovered culture plates were exposed to UVA light (320-400 nm) emitted from a bank of four BLB fluorescence light tubes (F40 BL, Sylvania) placed approximately 10 cm above the cell culture plates. In these experiments, the incident light on the culture plates was 2.4 mW/cm<sup>2</sup> as measured with an International Light UV radiometer, Model IL442A. The cells were exposed to 1.28 J/cm<sup>2</sup> of UVA. After exposing the cells to UVA light, the cell culture medium was drained and the cells refed with fresh growth medium. After incubating the cells for 4-5 days, cells on the plates were

removed with trypsin and counted with a Coulter Counter. Results (see Table 1 and Figure 1) were presented as percent inhibition of cell growth when compared to controls.

### Plasmid DNA nicking and unwinding assays:

Assays to evaluate the effects of coumarins on plasmid DNA were performed as previously described.<sup>(30)</sup> Briefly, 10  $\mu\text{L}$  reaction mixtures were prepared that contained 75 ng of plasmid pZeoSV DNA, 0.1  $\mu\text{L}$  coumarin derivative solution (prepared as a 100  $\times$  concentrate in DMSO) and TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8). The reaction volume and components were scaled up several-fold so that multiple aliquots of 10  $\mu\text{L}$  could be removed for analysis in agarose gels before and after UVA light treatment. Treatment with UVA light was performed in V-bottomed 96-well plates as described above for the cell cultures. Samples of the reaction mixtures containing the plasmids were then analyzed by electrophoresis using 1.2% agarose gels either in neutral gel buffer (40 mM Tris base, 20 mM acetic acid, 1 mM EDTA) or under denaturing conditions. To denature the DNA, samples were first mixed with 2 M NaOH supplemented with 0.1 M EDTA and heated in a water bath at 90°C for 1 min. After cooling on ice, DNA samples were loaded on 1.2% alkaline agarose gels prepared and electrophoresed in alkaline gel buffer (50 mM Tris base, 45 mM boric acid, 30 mM NaOH, 1 mM EDTA). After electrophoresis, gels were washed and stained with ethidium bromide (0.5  $\mu\text{g}/\text{mL}$ ). Gels were then photographed (see examples in Figure 2) with an Eagle Eye II digital documentation system (Stratagene, San Diego, CA). Bands shown in Figure 2 were then subjected to image analysis to calculate the percent of plasmid unwinding (see Table 2) which ranged from 0.2 to 99% in the set chosen.

### Conclusions:

Forty-nine 7-oxycoumarins from a library of 71 structural variants demonstrated phototoxicity against PAM212 mouse keratinocytes. None of the compounds was phototoxic in the absence of UVA light. These data indicate that the ability of the compounds to inhibit cell growth was dependent not only on their physicochemical characteristics, but also on their ability to absorb UVA light. Based on the results, definite structure-activity correlations could be made. Relative lipophilicity was an important factor especially among structurally similar families of coumarins. In most substituted oxycoumarins, the 7-alkoxys were more active than the 7-hydroxy compounds. Coumarins with electron withdrawing moieties – cyano and fluoro at C<sub>3</sub> – were considerably less active while those with bromines or iodine at that location displayed enhanced activity. The electron withdrawing trifluoromethyl at C<sub>4</sub> was generally a deactivating functionality while brominated alkyl side chains substantially improved activity.

Of interest were the findings that photocytotoxic coumarins also possessed the ability to photo-damage plasmid DNA albeit not in the same rank order of their activity as inhibitors of keratinocyte growth. There is no evidence that the two-ring coumarins, unlike the three-ring furocoumarins (psoralens), intercalate DNA prior to manifesting phototoxicity, but outer groove DNA-affinity achieves the same proximity necessary for DNA binding. It thus appears that binding of the coumarins to DNA is sufficient to initiate photo-damage following exposure to UVA light. This includes both DNA strand nicking and unwinding. Taken together, our data indicates that the DNA in keratinocytes is likely to be an important

target for the coumarins. Further studies are needed to characterize coumarin-induced DNA photo-damage and its role in regulating keratinocyte cell growth.

## Acknowledgments:

This work was funded in part by the National Institutes of Health CounterACT Program through the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIH Grant number U54AR055073). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the federal government.

## References

- (1). Weinmann I, (1997), History of the development and applications of coumarin and coumarin-related compounds In: O'Kennedy R & Thornes RD, eds. *Coumarins – Biology, Applications and Mode of Action*, John Wiley & Sons Ltd, Chichester, UK, pp. 1–22.
- (2). Ojala T, (2001), Biological screening of plant coumarins, Doctoral dissertation, University of Helsinki, Finland, 22 2001. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.407.8611&rep=rep1&type=pdf>
- (3). Bruneton J, (1999), *Pharmacognosy, Phytochemistry, Medicinal Plants*, 2nd Ed. Intercept Ltd, Hampshire, UK, pp. 263–277.
- (4). Asif M, (2015), Pharmacological activities and phytochemistry of various plant containing coumarin derivatives. *Curr. Sci. Perspectives*, 1(3): 77–90.
- (5). Averbeck D, (1989), Recent advances in psoralen phototoxicity mechanism. *Photochem. Photobio.* 50(6): 859–882. DOI: 10.1111/j.1751-1097.1989.tb02917.x
- (6). Laskin JD, Lee E, Yurkow EJ, Laskin DL, Gallo MA, (1985), A possible mechanism of psoralen phototoxicity not involving direct interaction with DNA. *Proc. Natl. Acad. Sci. USA*, 82(18): 6158–6162. DOI: 10.1073/pnas.82.18.6158 [PubMed: 3862124]
- (7). Stansbury J, (2011), Photosensitizing herbs: skin diseases and cancers. *DDNR, Anti-aging, dermatology, oncology*, 15 2011, pp. 1–11. <http://ndnr.com/oncology/photosensitizing-coumarins-for-skin-diseases-and-cancers/>
- (8). Nigg HN, Nordby HE, Beier RC, Dillman A, Macias C, Hansen RC, (1993), Phototoxic coumarins in limes. *FoodChem. Toxicol.* 31(5): 331–335. DOI: 10.1016/0278-6915(93)90187-4
- (9). Ojala T, Vuorela P, Kiviranta J, Vuorela H, Hiltunen R, (1999), A bioassay using *Artemia salina* for detecting phototoxicity of plant coumarins. *Planta Med.*, 65(8): 715–718. DOI: 10.1055/s-1999-14049 [PubMed: 10630112]
- (10). Jivaramonaikul W, Rashatasakhon P, Wanichwecharungruang S, (2010), UVA absorption and photostability of coumarins. *Photochem, Photobiol, Sci*, 9(8): 1120–1125. DOI: 10.1039/C0PP00057D [PubMed: 20544120]
- (11). Mousavi SH, Davari AS, Iranshahi M, et al., (2015), Comparative analysis of the cytotoxic effects of 7-prenyloxycoumarin compounds and herniarin on MCF-7 cell line. *Avicenna J. Phytomed*, 5(6): 520–530. [PubMed: 26693409]
- (12). Goard M, Aakalu G, Fedorya OD, et al., (2005), Light-mediated inhibition of protein synthesis. *Chem. & Biol*, 12(6): 685–693. DOI: 10.1016/i.chembiol.2005.04.018 [PubMed: 15975514]
- (13). Sarwar T, Rehman SU, Husain MA, Ishqi HM, Tabish M, (2015), Interaction of coumarin with calf thymus DNA: deciphering the mode of binding by *in vitro* studies. *Int. J. Biol. Macromol*, 73: 9–16. DOI: 10.1016/j.ijbiomac.2014.10.017 [PubMed: 25453293]
- (14). Sarwar T, Husain MA, Rehman SU, Ishqi HM, Tabish M, (2015), Multi-spectroscopic and molecular modelling studies on the interaction of esculetin with calf thymus DNA. *Mol. Biosyst*, 11(2): 522–531. DOI: 10.1039/C4MB00636D [PubMed: 25424306]
- (15). Pivetta T, Valletta E, Ferino G, Isaia F, Pani A, Vascellari S, Castellano C, Demartin F, Cabiddu MG, Cadoni E, (2017), Novel coumarins and related copper complexes with biological activity: DNA binding, molecular docking and *in vitro* antiproliferative activity. *J. Inorg. Biochem*, 177:101–109. DOI: 10.1016/j.jinorgbio.2017.09.013 [PubMed: 28946026]
- (16). Bisagni E, (1992), Syntheses of psoralens and analogues. *J. Photochem. Photobio. B: Biol*, 14: 23–46. DOI: 10.1016/1011-1344(92)85081-5

- (17). Matern U, Luer P and Kreusch D, (1999), Biosynthesis of coumarins In: Barton D, Nakanishi K, Meth-Cohn O and Sankawa U (eds.): *Comprehensive Natural Products Chemistry*, Vol. 1, Polyketides and Other Secondary Metabolites Including Fatty Acids and Their Derivatives. Elsevier Science Ltd, Oxford, UK pp. 623–637.
- (18). Yeagers E, Augenstein L, (1965), Absorption and emission spectra of psoralen and 8-methoxypsoralen in powders and in solutions. *J. Invest. Derm*, 44(3): 181–186. DOI: 10.1038/jid.1965.32 [PubMed: 14275694]
- (19). Mattoo BN, (1956), Absorption and fluorescence spectra of coumarins. *Trans. Faraday Soc*, 52: 1184–1194. DOI: 10.1039/TF9565201184
- (20). Fink DW, Koehler WR, (1970), pH effects on fluorescence of umbelliferone. *Anal. Chem*, 42(9): 990–993. DOI: 10.1021/ac60291a034
- (21). Zhang J, Wei Y, Liu C, (2011), Fluorescence quantum yield and ionization constant of umbelliferone, *Chem. Bull*, 74: 957–960.
- (22). Gee KR, Haugland RP, Sun W-C, (1998), Derivatives of 6,8-difluoro-7-hydroxycoumarin. U. S. Patent 5,830,912 3 11 1998.
- (23). Invitrogen, (2010), *Molecular probes handbook: a guide to fluorescent probes and labeling technologies*. Chapter 10, 11th Edition (2010); Invitrogen publication by Thermo Fisher Scientific.
- (24). Abu-Eittah RH, El-Tawil BAH, (1985), The electronic absorption spectra of some coumarins. A molecular orbital treatment. *Can. J. Chem*, 63: 1173–1176. DOI: 10.1139/v85-200
- (25). Laskin JD, (1994), Cellular and molecular mechanisms in photochemical sensitization: studies on the mechanism of action of psoralens. *Food Chem. Toxicol*, 32(2): 1191–1127. DOI: 10.1016/0278-6915(94)90172-4
- (26). Racz E, Prens EP, (2015), Phototherapy and photochemotherapy for psoriasis. *Dermatol. Clin*, 33(1): 79–89. DOI: 10.1016/j.det.2014.09.007 [PubMed: 25412785]
- (27). Kayal G, Jain K, Malviya S, Kharia A, (2014), Comparative SAR of synthetic coumarin derivatives for their anti-inflammatory activity. *Int. J. Pharm. Sci. Res*, 5(9): 3577–3583. DOI: 10.13040/IJPSR.0975-8232.5(9).3577-84
- (28). Kostova I, Raleva S, Genova P, Argirova R, (2006), Structure-Activity Relationships of Synthetic Coumarins as HIV-1 Inhibitors. *Bioinorg. Chem. Appl*, 2006 (68274): 1–9. DOI: 10.1155/BCA/2006/68274
- (29). Guerra FQ, de Araujo RS, de Sousa JP, de Pereira FO, Mendoza-Junior FJ, Barbosa-Filho JM, de Oliveira Lima E, (2013), Evaluation of Antifungal Activity and Mode of Action of New Coumarin Derivative, 7-Hydroxy-6-nitro-2H-1-benzopyran-2-one, against *Aspergillus* spp. *Evid Based Complement Alternat. Med* 2015(925096): 8 pages. DOI: 10.1155/2015/925096.
- (30). Mariano TM, Vetrano AM, Gentile SL, Heck DE, Whittmore MS, Guillon CD, Jabin I, Rapp RD, Heindel ND, Laskin JD, (2002), Cell-impermeant pyridinium derivatives of psoralens as inhibitors of keratinocyte growth, *Biochem. Pharmacol*, 63(1): 31–39. DOI: 10.1016/S0006-2952(01)00855-3 [PubMed: 11754871]
- (31). Gupta VN, (1957), Nuclear methylation of coumarins. *J. Scientific Industrial Res*, 16B: 257–62.
- (32). Rodighiero P, Manzini P, Pastorini G, Bordin F, Guitto A, (1987), Synthesis of methyl derivatives of 8-desmethylxanthyletine and 8-desmethylseseline, potential antiproliferative agents. *J. Heterocyclic Chem*, 24: 485–488. DOI: 10.1002/jhet.5570240234
- (33). Yan H, Yin W, Liu P, Liu J, et al., (2014), Palladium-catalyzed synthesis of 6-allylcoumarins using organotin reagents as multicoupling organometallic nucleophiles. *Appl. Organometallic Chem*, 28(10): 747–749. DOI: 10.1002/aoc.3188
- (34). Rangaswami S and Seshadri TR, (1938), Fixation of the aromatic double bonds in the coumarin ring system. *Proc. - Indian Acad. Sci*, 7A: 8–12.
- (35). Kaufman KD, (1961), New synthesis of methylsubstituted psoralens and isopsoralens. *Journal of Organic Chemistry*, 26, 117–121. DOI:10.1021/jo01060a028
- (36). Kaufman KD, (1965), Preparation of alkylpsoralens. U.S. Patent P 17 81965.
- (37). Bissell ER, Larson DK, Croudace MC, (1981), Some 7-substituted-4-(trifluoromethyl)coumarins. *Journal of Chemical and Engineering Data*. 1981, 26(3), 348–350. DOI: 10.1021/je00025a041



- (38). Goldfarb DS, (2009), Method using lifespan-altering compounds for altering the lifespan of eukaryotic organisms and screening for such compounds. U. S. Patent (Appl. Pub.), 20090163545 A1.
- (39). Whalley WB, (1951), Organic fluoro compounds. VI. Some (trifluoromethyl)chromones and – coumarins. *Journal of the Chemical Society*, 3235–3238. DOI: 10.1039/JR9510003235
- (40). Heindel ND, Laskin JD, Heck DE, Rapp RD, Whittemore MS, McNeel TE, Jabin I (2001), Amino- and mercurio-substituted 4',5'-dihydropsoalens and therapeutic uses thereof. U. S. Patent 6,255,324B1, 3 7 2001.
- (41). Jabin I, Heindel ND, Rapp RD, Laskin JD, (2000), Synthetic approaches to 3-substituted-5'- (N-pyridiniummethyl)-4',5'-dihydropsoalens. *J. Heterocyclic Chem*, 37: 31–34. DOI:10.1002/jhet.5570370106
- (42). Chilin A, Confente A, Pastorini G, Guiott A, (2002), Synthesis of furo[3.2 g][1,4]benzoxazin-3-ones, new psoralen isosters, *Eur. J. Org. Chem*, 2002: 1937–1940. DOI: 10.1002/1099-0690(200206)2002:12<1937::AID-EJOC1937>3.0.CO;2-W
- (43). Takaoka A, Ibrahim MK, Kagaruki SRF, Ishikawa N, (1985), Synthesis of monofluoro heterocycles using fluoroolefins as starting materials. *Nippon Kagaku Kaishi*. (11), 2169–2176. DOI:10.1246/nikkashi.1985.2169
- (44). Platz MS, Goodrich RP Jr, Yerram N, (1995), Method of inactivation of viral and bacterial blood contaminants. U. S. Patent 5,418,130, 23 5 1995.
- (45). Leffler JE, Grunwald E, (1963), *Rates and Equilibria of Organic Reactions*. Wiley, New York, Table 7.1.
- (46). Reid ST, (1990), Photoelimination, Chapter 7 in Bryce-Smith D, Gilbert A. (eds), *Photochemistry*, Vol. 21, Royal Soc. of Chem. publishers, Cambridge, UK, p. 472.
- (47). Meng JB, Shen M-G, Fu D-C, Gao Z-H, et al., (1990), A photochemical synthesis of 3-aryl coumarins. *Synthesis*, 1990(8): 719–721. DOI: 10.1055/s-1990-26993
- (48). Park SC, Goodrich RP Jr., Yerram N, et al., (1996), Photoinactivation of viral and bacterial blood contaminants using halogenated coumarins. U. S. Patent 5,516,629, 14 5 1996.
- (49). Vyas GN, Shah NM, (1951), Quinacetophenone monomethyl ether. *Organic Syntheses*, 31, 90–92. DOI: DOI:10.15227/orgsyn.031.0090
- (50). Bayer V, Pastor R, Cambon A, (1982), Synthesis and spectra study of 4-(perfluoroalkyl)-coumarins. *J. Fluorine Chem*, 20(2): 187–202. DOI: 10.1016/S0022-1139(00)82207-1

**Highlights**


Polyfunctionalized non-toxic coumarins displayed a photo-induced toxicity against cultured keratinocytes. This toxicity may arise from a light-catalyzed nicking of DNA.

Author Manuscript

Author Manuscript


Author Manuscript

Author Manuscript



**Figure 1.**


Effects of coumarins on keratinocyte growth with and without UVA light. PAM212 cells were grown in tissue culture dishes and treated with increasing concentrations of the coumarins. After 30 min at 37°C, cells were treated with (closed symbols) or without (open symbols) UVA light. Cells were then refed with fresh growth medium. After 4-5 days, cells were removed from the dishes and counted on a Coulter Counter. Data are presented as percent inhibition of cell growth relative to controls. Data represent means  $\pm$  SEM ( $n = 3$ ). In a typical experiment, control plates contained  $5.5 \times 10^4$  cells.




**Figure 2.** Ability of coumarins and UVA light to damage plasmid DNA. Plasmid DNA was treated with increasing concentrations of the coumarins. Lane L shows products from the digestion of the plasmid with restriction endonuclease EcoRI to linearize the DNA (100 ng); in native gels, the linearized DNA migrates as one double-stranded band, whereas, in denaturing gels, it migrates as two well-separated single strands. Lane S shows untreated plasmid DNA (75 ng); in native gels, the intact plasmid DNA migrates mainly as one supercoiled double-stranded band, whereas in denaturing gels, it migrates as two closely migrating single strands. Plasmid was analyzed in neutral (panels A and B) or alkaline (panels C and D) agarose gels. Plasmid samples were treated with (panels B and D) or without (panels A and C) UVA light.

Table 1.


Effects of 7-Oxycoumarins plus UVA light on the growth of PAM212 keratinocytes





Cpd	Source	Melting point (°C)	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	IC <sub>50</sub> (μM)
1	A	228-230 <sup>(*)</sup>	H	H	H	-OH	H	>100
2	S	145-146	H	Me	H	-OC(Me) <sub>2</sub> C≡CH	Me	4
3	S	93-94	H	Me	H	-OCH <sub>2</sub> CH=CMe <sub>2</sub>	Me	10
4	S	83-84	H	Me	H	-O(CH <sub>2</sub> ) <sub>5</sub> Me	Me	10
5	S	106-108	H	Me	H	-O(CH <sub>2</sub> ) <sub>3</sub> Me	Me	10
6	S	103-104	H	Me	H	-O(CH <sub>2</sub> ) <sub>2</sub> Me	Me	15
7	S	114-116	H	Me	H	-OCH <sub>2</sub> CH=CHMe	Me	33
8	S	166-167 <sup>(31)</sup>	H	Me	H	-OMe	Me	40
9	S	139-141 <sup>(32)</sup>	H	Me	H	-OCH <sub>2</sub> C≡CH	Me	41
10	S	87-88.5	H	Me	H	-OCH <sub>2</sub> C≡C(CH <sub>2</sub> ) <sub>4</sub> Me	Me	60
11	S	80-82	H	Me	H	-O(CH <sub>2</sub> ) <sub>7</sub> Me	Me	60
12	B	260.5-262	H	Me	H	-OH	Me	100
13	S	158-159.5 <sup>(33)</sup>	H	Me	I	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	1
14	S	146-147	H	Me	I	-OCH <sub>2</sub> CH=C(Me) <sub>2</sub>	Me	5
15	S	124.5-126	H	Me	I	-O(CH <sub>2</sub> ) <sub>2</sub> Me	Me	10
16	S	219-220 <sup>(33)</sup>	H	Me	I	-OH	Me	18
17	S	145.5-147.5	H	Me	NH <sub>2</sub>	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	5
18	S	229.5-231.5	H	Me	NO <sub>2</sub>	-OH	Me	>100
19	S	199-200	-CH <sub>2</sub> CH=C(Me) <sub>2</sub>	Me	H	-OH	Me	12
20	S	165-167	H	Me	-CH <sub>2</sub> CH=C(Me) <sub>2</sub>	-OH	Me	1
21	S	168-170 <sup>(34)</sup>	H	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OH	Me	10
22	S	175-176	H	Me	-CH(Me)CH=CH <sub>2</sub>	-OH	Me	14



Cpd	Source	Melting point (°C)	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	IC <sub>50</sub> (μM)
23	S	158-160	H	Me	-CH <sub>2</sub> CH=CHMe	-OH	Me	20
24	S	149.5-150 <sup>(35)</sup>	H	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OAc	Me	21
25	C	159-161	H	Me	H	-OMe	H	60
26	S	101-102 <sup>(36)</sup>	H	Me	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	H	10
27	S	254-256 <sup>(32)</sup>	H	Me	H	-OH	I	>100
28	S	151.5-153	H	Me	H	-OCH <sub>2</sub> -CH=C(CH <sub>3</sub> ) <sub>2</sub>	I	22
29	S	129-130	H	Me	H	-O(CH <sub>2</sub> ) <sub>2</sub> Me	I	50
30	S	194-195	H	Me	H	-OMe	I	70
31	D	119-120	H	H	H	-OMe	OMe	33
32	C	118-121	H	H	H	-OMe	H	50
33	S	182-183	H	CF <sub>3</sub>	-CH <sub>2</sub> CCl=CH <sub>2</sub>	-OH	Me	20
34	S	143-144.5	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> (CO)Me	Me	40
35	S	112-113	H	CF <sub>3</sub>	H	-O(CH <sub>2</sub> ) <sub>2</sub> Me	Me	100
36	S	76-77	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	H	>200
37	S	197-198	H	CF <sub>3</sub>	H	-OH	Me	>200
38	S	160.5-161.5	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> C≡CH	Me	>300
39	S	97-98	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	H	>100
40	S	151-152	H	CF <sub>3</sub>	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OAc	Me	>100
41	S	112.5-113.5 <sup>(37)</sup>	H	CF <sub>3</sub>	H	-OMe	H	>100
42	S	110.5-112	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> C≡CH	H	>100
43	S	174-175	H	CF <sub>3</sub>	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OH	Me	>300
44	S	114.5-115.5	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	>300
45	S	109.5-111 <sup>(38)</sup>	H	CF <sub>3</sub>	H	-OMe	Me	>300
46	S	158-159	H	CF <sub>3</sub>	H	-OCH <sub>2</sub> (CO)Me	H	>300
47	S	186-188 <sup>(39)</sup>	H	CF <sub>3</sub>	H	-OH	H	>300



Cpd	Source	Melting point (°C)	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	IC <sub>50</sub> (μM)
48	S	149.5-150 <sup>(35)</sup>	H	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OAc	Me	21
49	S	198-198.5 <sup>(d)(40,41)</sup>	Br	Me	-CH <sub>2</sub> (CHBr)CH <sub>2</sub> Br	-OH	Me	1
50	S	158-160 <sup>(40,41)</sup>	Br	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OH	Me	5
51	S	269-271 <sup>(42)</sup>	Br	Me	H	-OH	Me	10
52	S	143-143.5	Br	H	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	2
53	S	131.5-133	Br	H	H	-OCH <sub>2</sub> CH=CMe <sub>2</sub>	Me	5
54	S	139.5-140	Br	H	H	OCH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Me	5
55	S	181-182	Br	H	H	-OCH <sub>2</sub> C≡CH	Me	10
56	S	154-155	Br	H	H	-OMe	Me	13
57	S	209-211 <sup>(42)</sup>	Br	H	H	-OH	Me	20
58	S	147-148	Br	Me	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	5
59	S	231-232	Br	Me		-OAc	Me	20
60	S	158-160 <sup>(41)</sup>	Br	Me	-CH <sub>2</sub> (CHBr)CH <sub>2</sub> Br	-OAc	Me	48
61	S	160-161 <sup>(41)</sup>	Br	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OAc	Me	140
62	S	221-223 <sup>(40,41)</sup>	CN	Me	-CH <sub>2</sub> (CHBr)CH <sub>2</sub> Br	-OH	Me	3
63	S	170-171 <sup>(41)</sup>	CN	Me	-CH <sub>2</sub> (CHBr)CH <sub>2</sub> Br	-OAc	Me	48
64	S	234-235 <sup>(40,41)</sup>	CN	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OH	Me	>300
65	S	120-124 <sup>(d)(40,41)</sup>	CN	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OAc	Me	>300
66	S	172-173 <sup>(40,41)</sup>	CN	Me	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	>300
67	S	175-176 <sup>(40,41)</sup>	F	Me	-CH <sub>2</sub> (CHBr)CH <sub>2</sub> Br	-OH	Me	120
68	S	192-193 <sup>(43)</sup>	F	Me	H	-OH	Me	>200



Cpd	Source	Melting point (°C)	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	IC <sub>50</sub> (μM)
69	S	117-118 <sup>(40,41)</sup>	F	Me	H	-OCH <sub>2</sub> CH=CH <sub>2</sub>	Me	>300
70	S	163-164 <sup>(40,41)</sup>	F	Me	-CH <sub>2</sub> CH=CH <sub>2</sub>	-OH	Me	>300
71	S	228-230 <sup>(d)</sup>	I	Me	H	-OH	Me	61

<sup>(\*)</sup>: Compound 1 is umbelliferone also known as 7-hydroxycoumarin. An IC<sub>50</sub> expressed as > (number) means no activity was observed up to that value, usually the solubility limit of the compound for the assay's conditions. A = Sigma-Aldrich; B = Regis Chemical; C = TCI America Inc.; D = Elder (Valeant) Pharmaceuticals. S means that the compound was synthesized for these studies either following an existing procedure or a new one developed for that purpose, which is described in detail in the experimental part. Published methods for specific coumarins are indicated by references provided in the table and in the experimental part. Obvious decomposition upon melting is noted by <sup>(d)</sup>.



**Table 2**

Percent linearization of plasmid DNA by coumarins plus UVA light

Concentration ( $\mu$ M)	Percentage of nicked plasmid				
	Compd 14	Compd 48	Compd 58	Compd 62	Compd 71
0	0.2	0.2	1.0	1.2	2.2
1	1.0	4.0	3.2	3.2	6.8
10	8.0	60.5	12.0	55.4	80.2
100	20.5	90.8	20.2	96.6	98.6

Table 2. DNA nicking data from plasmids analyzed in neutral agarose gels (Figure 2, panel B). In these experiments, plasmids were treated with increasing concentrations of the coumarins followed by UVA light. Image analysis was used to calculate the percent of plasmid unwinding. Note that compounds **48**, **62** and **71** were more effective in nicking DNA.