



Published in final edited form as:

Angew Chem Int Ed Engl. 2019 August 05; 58(32): 10956–10960. doi:10.1002/anie.201904861.

Nickel-Catalyzed Arylboration of Alkenylarenes: Synthesis of Boron-Substituted Quaternary Carbons and Regiodivergent Reactions

Liang-An Chen¹, Alan R. Lear¹, Pin Gao², Prof. M. Kevin Brown¹


¹Department of Chemistry, Indiana University, 800 E. Kirkwood Ave. Bloomington, IN 47405

²Department of Chemistry, School of Science, Xi'an Key Laboratory of Sustainable Energy Material Chemistry, MOE Key Laboratory for Nonequilibrium Synthesis and Modulation of Condensed Matter, Xi'an Jiaotong University, Xi'an, Shaanxi, 710049, China.

Abstract

A method for the construction of boron-substituted quaternary carbons or diarylquaternary carbons by arylboration of highly substituted alkenylarenes is presented. A wide range of alkenes and arylbromides can participate in this reaction thus allowing for a diverse assortment of products to be prepared. In addition, a solvent dependent regiodivergent arylboration of 1,2-disubstituted alkenylarenes is presented, thus greatly increasing the scope of products that can be accessed.

Graphical Abstract



Keywords

Nickel; Boron; Arylboration; Alkene; Cross Coupling

Alkene difunctionalization is a significant synthetic strategy as molecular complexity can be rapidly generated.¹ Within this broad area, alkene carboboration has emerged as an effective strategy that allows for a diverse range of groups to be installed across the alkene by taking advantage of the synthetic versatility of the C-B bond.^{2,3} Over the last five years, alkene arylboration methods have emerged with contributions from our lab⁴ and others.⁵ The arylboration of alkenylarenes is well developed with mono-, 1,2-di-, and 1,1-disubstituted alkenes (Scheme 1A).^{4,5} Notably, functionalization of trisubstituted alkenylarenes remains a

challenge. We sought to develop an arylboration of these classes of substrates because the products would contain boron-substituted quaternary carbons or diarylquaternary carbons, motifs that are currently difficult to access (Scheme 1B). In particular, synthesis of boron-substituted quaternary carbons from widely available alkenes was seen as a key motivation for this study. Notable advances in the synthesis of boron substituted quaternary carbons from alkenes include conjugate addition/allylic substitution,⁶ hydroboration/diboration,⁷ conjunctive cross coupling,⁸ and aminoboration^{9,10} The majority of the known approaches utilize alkenes with strong activating group (e.g. substituted acrylates) or directing groups. The approach outlined herein is appealing as simple alkenylarenes are the starting materials and two bonds and two stereogenic centers are generated.

Early attempts from our group to develop arylboration reactions of trisubstituted alkenylarenes with Cu/Pd-catalysis were unsuccessful.^{4a-e} However, our recent discovery of a Ni-catalyzed arylboration of unactivated alkenes,^{4f} prompted us to explore Ni-catalysis in the context of highly substituted alkenylarenes. Herein, we disclose a method for the arylboration of a wide range of trisubstituted alkenylarenes that leads to the formation of challenging boron-substituted quaternary carbons or diarylquaternary carbons (Scheme 1B). In addition, we have uncovered a solvent dependent regiodivergent arylboration of 1,2-disubstituted alkenylarenes that further increases the utility of this method (Scheme 1C).

Arylboration of trisubstituted alkenylarenes under previously reported Ni-catalysis conditions resulted in low to moderate yield of the desired products due to incomplete conversion.^{4f} Evaluation of reaction conditions ultimately led to the finding that use of DMA as solvent led to high yield of product (compare Table 1, entries 1 and 2). Unexpectedly, it was also found that premixing KOEt and (Bpin)₂ prior to the addition of substrates and catalyst led to higher yields (compare Table 1, entries 1 and 3). Presumably premixing of KOEt and (Bpin)₂ serves to sequester KOEt. It has been reported that alkoxide bases and Ni(II) salts can result in the formation of nanoparticles, which may deactivate the Ni-catalyst in this case.¹¹ While 5 mol % NiCl₂(DME) and 3.0 equiv aryl bromide were found to be optimal, lower catalyst loading or fewer equivalents of aryl bromide still resulted in useful yields (Table 1, entries 4 and 5). In addition, the reaction could be setup without the aid of glovebox techniques with little change in efficiency (Table 1, entries 6). Finally, it should be noted that the optimal conditions were particularly crucial for arylbromides bearing electron-withdrawing groups (compare Table 1, entries 7–9).

Under the optimized conditions arylbromides bearing electron donating, electron withdrawing and sterically demanding substituents were well tolerated (Scheme 2). Various functional groups such as amines, alcohols, esters, and amides did not greatly impede the reaction. Finally, heterocycles could also be employed, albeit with slightly reduced yields. The primary limitation with respect to arylbromide scope is that strongly electron-withdrawing substituents such as CF₃ resulted in low yields of product and formation of significant quantities of ArBpin.

Regarding the scope of alkenes that can be used, several points are noteworthy (Scheme 3): 1) For the synthesis of boron-substituted quaternary carbons, the substitution pattern of the aryl group could be altered with various electron withdrawing, electron donating, sterically

demanding substituents. 2) Both cyclic and acyclic substrates are suitable for this process. 3) The formation of **30** and **37** occurred with high levels of diastereoselectivity. In each case, the arylboration occurred from the least hindered face of the alkene. 3) For substrates with acyclic substituents, yield decreased with more sterically demanding substituents (compare **2** and **33**). 4) The synthesis of diarylquaternary carbons can be achieved with both acyclic and cyclic substrates (products **38-44**).

The arylboration could also be performed on gram scale (Scheme 4). Furthermore, the C-B bond could be elaborated through oxidation (product **47**) and Matteson homologation/oxidation (product **48**). Alcohol **48** could be advanced to Glucocorticoid receptor modulators through standard sequences.¹² Previously, these molecules were prepared by a non-stereoselective 6-step sequence. The route illustrated here required 3–4 operations, is stereoselective, and represents an orthogonal approach to the construction of biologically active molecules from alkenes.

In the course of these investigations, 1,2-disubstituted alkenylarenes were also probed. Under the standard conditions, **51** was generated in excellent selectivity (Table 2, entry 1). Interestingly, the observation was made that if toluene was used as a co-solvent, the formation of regioisomer **52** (>20:1 dr) was observed (Table 2, entry 2). Increasing the amount of toluene revealed that regioisomer **52** could be generated as the major product, albeit in low yield (Table 2, entry 7). Continued optimization led to the finding that higher reaction temperatures (60 °C vs. rt), use of KOMe (vs. KOEt), and excess alkene (relative to arylbromide) resulted in exclusive formation of **52** (Table 2, entry 12). Thus, through a simple change in solvent and minor modification to reaction conditions, the arylboration could be tuned to favor either regioisomer **51** or **52**. Prior studies have demonstrated that product **51** can be prepared by a Cu/Pd-catalyzed arylboration.^{4,5} The Ni-catalyzed reaction reported herein offers an alternative that utilizes a simple catalyst. Methods for the synthesis of the regioisomeric product **52** do not exist. However, a recent report from Song, has demonstrated that styrene derivatives, but not 1,2-disubstituted alkenylarenes, can undergo Pd-catalyzed arylboration with aryldiazonium salts and (Bpin)₂.^{5c}

Under conditions outlined in Table 2 entries 1 and 12 both acyclic and cyclic 1,2-disubstituted alkenylarenes participated in the reactions with excellent regioselectivity and diastereoselectivity (products **51-60**). While arylboration in DMA worked well with trisubstituted alkenes (Schemes 2–3), the conditions with toluene did not proceed, and starting material was recovered.

To account for the different regioselectivities observed in different solvents, two catalytic cycles are proposed (Scheme 6). In the case of reactions performed in DMA, and on the basis of prior studies, we propose that [Ni]-Bpin complex **62** is generated, which undergoes reaction with an alkene.^{4f,13} The regioselectivity of the migratory insertion occurs to generate benzyl-[Ni]-complex **63** regardless of alkene substitution pattern. The formation of product may occur through an oxidative addition with ArBr followed by reductive elimination via **64**. In the case of reactions in toluene, it is proposed that migratory insertion occurs from Ni(II)ArBr complex **66**.¹⁴ The regioselectivity is determined through formation

of benzyl-[Ni]-complex **67**. Transmetalation with (Bpin)₂ and reductive elimination then occurs to generate product.

Support for the proposed catalytic cycles is illustrated in Scheme 7. It was found that addition of MeOH to the reaction mixture resulted in formation of protoboration adduct **69**, thus suggesting that Bpin is incorporated prior to the aryl group.^{4f} In addition, reaction of vinyl fluoride **70** led to the formation of vinyl boronic ester **71**. Since direct cross coupling with vinyl fluorides is slow at ambient temperatures,^{15,16} it is suggested that the product formation occurs via 1,2-elimination of the corresponding alkyl-[Ni]-complex.¹⁷ With respect to the mechanistic proposal for the reaction in toluene, it was observed that reaction of α -methyl styrene (**72**) led to products resulting from a Heck-type reaction (**74**) and hydroarylation (**75**). The formation of these adducts can be rationalized with competitive reactions of the corresponding benzyl-[Ni]-complex via β -hydride elimination or protonation, respectively.¹⁸ At this stage, the role of solvent in determining the regioselectivity is not clear and is under investigation.

In summary, the synthesis of challenging boron-substituted quaternary carbons and diarylquaternary carbons has been achieved through an arylboration reaction. Furthermore, the arylboration regioselectivity of 1,2-disubstituted alkenylarenes can be tuned by altering the solvent, revealing a practical system in which diverse and versatile products can be accessed.¹⁹

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements




We thank Indiana University and the National Institutes of Health (R01GM114443 and R35GM131755) for generous financial support. This project was partially funded by the Vice Provost for Research through the Research Equipment Fund.

References


1. a) Saini V, Stokes BJ, Sigman MS, *Angew. Chem. Int. Ed* 2013, 52, 11206 *Angew. Chem.* 2013, 125, 11414–11428 b) Coombs JR, Morken JP, *Angew. Chem. Int. Ed. Engl* 2016, 55, 2636–2649. *Angew. Chem.* 2016, 128, 2682. [PubMed: 26764019]
2. For reviews regarding carboboration, see: (a) Shimizu Y, Kanai M, *Tetrahedron Lett* 2014, 55, 3727. (b) Semba K, Fujihara T, Terao J, Tsuji Y, *Tetrahedron* 2015, 71, 2183. (c) Lazreg F, Nagra F, Cazin CSJ, *Coord. Chem. Rev* 2015, 293–294, 48. (d) Neeve EC, Geier SJ, Mkhaliid IAI, Westcott SA, Marder TB, *Chem. Rev* 2016, 116, 9091. [PubMed: 27434758] (e) Hemming D, Fritzemeier R, Westcott SA, Santos WL, Steel PG, *Chem. Soc. Rev* 2018, 47, 7477. [PubMed: 30206614]
3. Sandford C, Aggarwal VK, *Chem. Commun* 2017, 53, 5481.
4. a) Smith KB, Logan KM, You W, Brown MK, *Chem. Eur. J* 2014, 20, 12032; [PubMed: 25113669] b) Logan KM, Smith KB, Brown MK, *Angew. Chem. Int. Ed* 2015, 54, 5228; *Angew. Chem.* 2015, 127, 5317; c) Logan KM, Brown MK, *Angew. Chem. Int. Ed* 2017, 56, 851; *Angew. Chem.* 2017, 129, 869; d) Smith KB, Brown MK, *J. Am. Chem. Soc* 2017, 139, 7721; [PubMed: 28581713] e) Sardini SR, Brown MK, *J. Am. Chem. Soc* 2017, 139, 9823. [PubMed: 28699339] f) Bergman AM, Dorn SK, Smith KB, Logan KM, *Angew. Chem. Int. Ed* 2019, 58, 1719. f) Logan KM, Sardini SR, White SD, Brown MK, *J. Am. Chem. Soc* 2017, 140, 159. [PubMed: 29271650]

5. a) Semba K, Nakao Y, *J. Am. Chem. Soc* 2014, 136, 7567. [PubMed: 24810227] b) Semba K, Ohtagaki Y, Nakao Y, *Org. Lett* 2016, 18, 3956. [PubMed: 27490821] c) Yang K, Song Q, *Org. Lett* 2016, 18, 5460. [PubMed: 27786482] d) Chen B, Cao P, Yin X, Liao Y, Jiang L, Ye J, Wang M, Liao J *ACS Catal* 2017, 4, 2425. e) Wang W, Ding C, Li Y, Li Y, Peng L, Yin G *Angew. Chem. Int. Ed* 2019, 58, 4612. f) Liu Z, Li X, Zeng T, Engle KM, *ACS Catal* 2019, 9, 3260. f) For a 1,1-arylboration, see: Nelson HM, Williams BD, Miro J, Toste FD *J. Am. Chem. Soc* 2015, 137, 3213. [PubMed: 25723255]
6. a) Chen IH, Yin L, Itano W, Kanai M, Shibasaki M, *J. Am. Chem. Soc* 2009, 131, 11664; [PubMed: 19653692] b) Chen IH, Yin L, Kanai M, Shibasaki M, *Org. Lett* 2010, 12, 4098; [PubMed: 20722382] c) O'Brien JM, Lee K, Hoveyda AH, *J. Am. Chem. Soc* 2010, 132, 10630–10633; [PubMed: 20681680] d) Radomkit S, Hoveyda AH, *Angew. Chem. Int. Ed* 2014, 53, 3387; *Angew. Chem.* 2014, 126, 3455; e) Feng X, Yun J, *Chem. Eur. J* 2010, 16, 13609. [PubMed: 21053219] f) Guzman-Martinez AH Hoveyda, *J. Am. Chem. Soc* 2010, 132, 10634. [PubMed: 20681681]
7. a) Blaisdell TP, Caya TC, Zhang L, Sanz-Marco A, Morken JP, *J. Am. Chem. Soc* 2014, 136, 9264. [PubMed: 24941137] b) Kerchner HA, Montgomery J, *Org. Lett* 2016, 18, 5760; [PubMed: 27786484] c) Shoba VM, Thacker NC, Bochat AJ, Takacs JM, *Angew. Chem. Int. Ed* 2016, 55, 1465; *Angew. Chem.* 2016, 128, 1487; d) Chakrabarty S, Takacs JM, *J. Am. Chem. Soc* 2017, 139, 6066. [PubMed: 28414243]
8. a) Silvi M, Sandford C, Aggarwal VK, *J. Am. Chem. Soc* 2017, 139, 5736; [PubMed: 28402109] b) Kischkewitz M, Okamoto K, Muck-Lichtenfeld C, Studer A, *Science* 2017, 355, 936; [PubMed: 28254937] c) Myhill JA, Zhang L, Lovinger GJ, Morken JP, *Angew. Chem., Int. Ed* 2018, 57, 12799; *Angew. Chem.* 2018, 130, 12981; d) Tao Z, Robb KA, Panger JL, Denmark SE, *J. Am. Chem. Soc* 2018, 140, 15621 [PubMed: 30411878]
9. Kato K, Hirano K, Miura M, *J. Org. Chem* 2017, 82, 10418. [PubMed: 28847150]
10. For selected examples of alternative approaches, see: a) Kato K, Hirano K, Miura M, *J. Org. Chem* 2017, 82, 10418. [PubMed: 28847150] b) Atack TC, Cook SP, *J. Am. Chem. Soc* 2016, 138, 6139. [PubMed: 27158838]
11. Buslov I, Song F, Hu X *Angew. Chem. Int. Ed* 2016, 55, 12295–12299. *Angew. Chem.* 2016, 128, 12483.
12. Duan J, Jiang B *PCT Int. Appl* 2007073503
13. Ni(I) may form through a comproportion pathway, as illustrated in Scheme 6. For a review regarding Ni(I), see: Lin C-Y, Power PP, *Chem. Soc. Rev* 2017, 46, 5347. [PubMed: 28675200]
14. Gao P, Chen L-A, Brown MK, *J. Am. Chem. Soc* 2018, 140, 10653. [PubMed: 30125098]
15. Liu X-W, Echavarren J, Zarate C, Martin R, *J. Am. Chem. Soc* 2015, 137, 12470. [PubMed: 26397717]
16. For β -fluoro elimination of an alkyl-Ni-complex, see: Zhou L, Zhu C, Bi P, Feng C, *Chem. Sci* 2019, 10, 1144. [PubMed: 30774912]
17. For a Cu-catalyzed variant, see: Sakaguchi H, Uetake Y, Ohashi M, Niwa T, Ogoshi S, Hosoya T, *J. Am. Chem. Soc* 2017, 139, 12855. [PubMed: 28849929]
18. The proton source may arise from the byproduct of β -hydride elimination.
19. During the preparation of this manuscript a Ni-catalyzed aryloboration of vinylarenes was reported, see: Wang W, Ding C, Pang H, Yin G *Org. Lett* 10.1021/acs.orglett.9b01120.


A) Arylboration of Alkenylarenes





B) This Work: Functionalization of Trisubstituted Alkenylarenes




C) This Work: Solvent Dependent Regioselective Arylboration




Scheme 1.
Arylboration of Alkenylarenes.

**Scheme 2.**


Reaction with Various Arylbromides. Yield of product after isolation. NMR yield determined by ¹H NMR analysis of the unpurified reaction mixture with an internal standard. ^[a] Reaction with the 4-OMeAr analog of **1** used. ^[b] Products **7** and **16** were oxidized to the alcohol before isolation. Reactions performed on 0.4 mmol scale.

**Scheme 3.**


Arylboration of Trisubstituted Alkenes. See Scheme 2 for details.




Scheme 4.
Gram Scale Reaction and Further Transformations.

**Scheme 5.**

Regiodivergent Arylboration. See Table 2, entries 1 and 12 for conditions (0.4 mmol scale). Yield of product after isolation. NMR yield determined by ^1H NMR analysis of the unpurified reaction mixture with an internal standard. ^[a] Reaction with 1.0 equiv alkene and 1.2 equiv ArBr.




Scheme 6.
Putative Catalytic Cycles.



Scheme 7.
Mechanistic Experiments.

Table 1.

Change from Standard Conditions




entry	change from standard conditions	product	yield (%) ^[a]
1	no change	2	98
2	THF instead of DMA	2	21
3	no premixing of (Bpin) ₂ and KOEt	2	79
4	1.5 equiv ArBr instead of 3.0 equiv ArBr	2	65
5	2 mol % NiCl ₂ (DME) instead of 5 mol % NiCl ₂ (DME)	2	75
6	glovebox free reaction setup	2	94
7	no change	3	65
8	THF instead of DMA	3	20
9	no premixing of (Bpin) ₂ and KOEt	3	29

Reactions performed on 0.2 mmol scale.

^[a]Yield determined by ¹H NMR analysis of the unpurified reaction mixture with an internal standard.

Table 2.

Regiodivergent Arylboration of 1,2-Disubstituted Alkenylarenes



entry	solvent	temp (°C)	base	yield (%) ^[a]	51	:	52
1	DMA	rt	KOEt	98	>99	:	1
2	toluene:DMA (1:1)	rt	KOEt	98	93	:	1
3	toluene:DMA (5:1)	rt	KOEt	42	34	:	1
4	toluene:DMA (10:1)	rt	KOEt	28	14	:	1
5	toluene:DMA (50:1)	rt	KOEt	15	4.6	:	1
6	toluene:DMA (160:1)	rt	KOEt	17	2.3	:	1
7	toluene	rt	KOEt	15	1	:	9
8	toluene	60	KOEt	55	1	:	2.2
9	toluene	60	KOMe	42	1	:	5
10 ^[b]	toluene	60	KOMe	42	1	:	36
11 ^{[b],[c]}	toluene	60	KOMe	45	1	:	39
12 ^{[c],[d]}	toluene	60	KOMe	58	1	:	>99

Reactions performed on 0.2 mmol scale.

^[a] Combined yield of **51** and **52** determined by GC using a calibrated internal standard.^[b] ArBr (1.2 equiv).^[c] NiCl₂(DME) (15 mol%).^[d] Alkene (2.0 equiv), ArBr, (1.0 equiv).