

CORRECTION

Open Access



Correction to: Neutrophils are a main source of circulating suPAR predicting outcome in critical illness

Hendrik Gussen¹, Philipp Hohlstein¹, Matthias Bartneck¹, Klaudia Theresa Warzecha¹, Lukas Buendgens¹, Tom Luedde¹, Christian Trautwein¹, Alexander Koch¹ and Frank Tacke^{1,2*}

Correction to: J Intensive Care (2019) 7:26

<https://10.1186/s40560-019-0381-5>

In the original publication of this article [1], the content of Authors' contributions needs to be revised as below:

HG and PH collected blood samples, performed FACS analyses, and conducted data analyses. MB and KTW supported the in vitro experiments. LB, TL, and CT contributed to data collection and analysis. AK and FT designed and supervised the study. HG drafted the manuscript and FT made the final corrections. All authors corrected and approved the manuscript.

Published online: 02 August 2019

Reference

1. Gussen, et al. *J Intensive Care*. 2019;7:26. <https://doi.org/10.1186/s40560-019-0381-5>.

* Correspondence: frank.tacke@gmx.net

¹Department of Medicine III, RWTH-University Hospital Aachen, 52074 Aachen, Germany

²Department of Hepatology/Gastroenterology, Charité University Medical Center, Augustenburger, Platz 1, 13353 Berlin, Germany

