



HHS Public Access

Author manuscript

Reprod Toxicol. Author manuscript; available in PMC 2019 August 11.

Published in final edited form as:

Reprod Toxicol. 2018 August ; 79: 96–123. doi:10.1016/j.reprotox.2018.06.005.

Effects of Bisphenol A and its Analogs on Reproductive Health: A Mini Review

Jacob Steven Siracusa¹, Lei Yin^{1,2}, Emily Measel¹, Shenuxan Liang¹, Xiaozhong Yu^{1,*}

¹Department of Environmental Health Science, College of Public Health, University of Georgia, Athens, Georgia 30602

²ReproTox Biotech LLC, Athens 30602, Georgia

Abstract

Known endocrine disruptor bisphenol A (BPA) has been shown to be a reproductive toxicant in animal models. Its structural analogs: bisphenol S (BPS), bisphenol F (BPF), bisphenol AF (BPAF), and tetrabromobisphenol A (TBBPA) are increasingly being used in consumer products. However, these analogs may exert similar adverse effects on the reproductive system, and their toxicological data are still limited. This mini-review examined studies on both BPA and BPA analog exposure and reproductive toxicity. It outlines the current state of knowledge on human exposure, toxicokinetics, endocrine activities, and reproductive toxicities of BPA and its analogs. BPA analogs showed similar endocrine potencies when compared to BPA, and emerging data suggest they may pose threats as reproductive hazards in animal models. While evidence based on epidemiological studies is still weak, we have utilized current studies to highlight knowledge gaps and research needs for future risk assessments.

Keywords

Bisphenol A; Bisphenol F; Bisphenol S; Bisphenol AF; Tetrabromobisphenol A; Reproductive toxicity

1. Introduction

Bisphenol A (BPA) is a high production volume (HPV) chemical, commonly used in food packaging materials, dental sealants, medical devices and thermal receipts [1]. Exposure to BPA is ubiquitous via ingestion, inhalation, and dermal contact [2, 3]. The Centers for Disease Control and Prevention (CDC) has reported measurable levels of BPA in urine samples in over 90% of the United States population [4, 5]. BPA has demonstrated endocrine disrupting effects by interacting with various physiological receptors, such as estrogen receptor α/β (ER α/β), estrogen-related receptor γ , androgen receptor (AR), and thyroid hormone receptor [6-13]. Numerous studies have investigated the reproductive toxicity of BPA, and extensive reviews were conducted to address the strength of the

*Correspondence should be addressed to Department of Environmental Health Science, College of Public Health, University of Georgia, Athens, Georgia, United States. 30602 Fax: 706.542.7472; yuxz@uga.edu.

The authors declare that there are no conflicts of interest.

evidence regarding BPA toxicity [14]. In 2006, an expert panel composed of the National Institute of Environmental Health Sciences (NIEHS), the National Institute of Dental Craniofacial Research (NIDCR), the U.S. Environmental Protection Agency (EPA), and Commonweal, reviewed human exposure to BPA *in vivo* and *in vitro* [15]. The subpanel of experts that focused on *in vivo* animal studies found contradictory results among the studies. However, they were confident that BPA impacted the male and female reproductive system [14]. Peretz et al. summarized studies published from 2007 to 2013 to examine the associations between BPA and adverse reproductive outcomes [14]. Based on the evidence from experimental animals and human exposure from 2007–2013, the authors concluded that BPA at doses below the LOAEL (50 mg/kg/day) impacted female reproduction and had potential adverse effects on the male reproductive system [14]. Furthermore, recent epidemiological studies have indicated that BPA exposure may potentially be associated with alterations in hormone levels, impairment of ovary and uterine function, and reduction of sperm quality [16-20]. Current data from experimental studies have suggested that BPA exposure adversely affected oocyte quality and maturation, decreased sperm production and quality, damaged testicular cells, perturbed hormone levels, and disrupted ovary function and uterine morphology in animal models [21-28].

Due to widespread exposure and concerns that BPA is a reproductive toxicant, the public drove manufacturers to abandon the use of BPA and introduce analogous chemicals in baby bottles, sippy cups, and infant formula packaging [29, 30]. The U.S. Food and Drug Administration then ruled that BPA would no longer be used in the products mentioned above [29, 30]. BPA analogs are being used as crosslinking reagents and flame retardants in the plastics industry to produce “BPA-free” products. However, the usage of these chemicals is expected to rise globally despite a lack of production data for these analogs. Recently, the prevalence of BPA analogs in the environment, foods, consumer products, and human urine samples have been reported [31-35]. With high degrees of structural similarities to BPA, these analogs may potentially have a similar endocrine disrupting capacity and the potential to exert adverse effects on the reproductive system. Emerging evidence suggests that BPA analogs interact with various physiological receptors, such as estrogen receptors α and β , androgenic receptors, and aryl hydrocarbons receptors [36, 37]. Compared to BPA, little is known about the reproductive toxicity of these analogs. Here, we have reviewed the current literature on BPA, its analogs, and male/female reproduction to summarize the current state of knowledge and the gaps in that knowledge, and to highlight future research directions that could provide valuable information for toxicity evaluation and risk assessment. In addition, the environmental occurrences, human biomonitoring data, toxicokinetics, and the endocrine disrupting capacity of BPA analogs have also been included. This review is structured into 7 topics: (1) BPA and female reproductive health; (2) BPA and male reproductive health; (3) BPS and reproductive health; (4) BPF and reproductive health; (5) BPAF and reproductive health; (6) TBBPA and reproductive health; (7) conclusion and research needs. This review of recent literature focuses on the effects of BPA and its analogs on the male and female reproductive system. We hope the information and conclusions in this review could direct future studies and be useful in risk assessment and in the formation of regulations regarding BPA and its analogs.

2. Search Strategy

We performed a literature search to identify journal articles related to BPA, BPS, BPF, BPAF, and TBBPA exposure and reproduction. The research included articles published between 2014 and 2017 for BPA and all years up to 2017 for BPA analogs. An electronic search was performed in Pubmed (<http://www.ncbi.nlm.nih.gov/pubmed>) and Google Scholar (<https://scholar.google.com/>). Pubmed was selected to identify journal articles as it is considered a main and reliable literature source. Google Scholar was used as a much broader search engine to collect and analyze any literature that may not have been included in Pubmed. Search terms included:

{ 'Bisphenol A' OR 'BPA' OR 'bisphenol S' OR 'BPS' OR '4,4'-sulfonyldiphenol' OR 'bisphenol F' OR 'BPF' OR '4,4'-dihydroxydiphenyl-methane' OR 'bisphenol AF' OR 'BPAF' OR 'hexafluorobisphenol A' OR 'tetrabromobisphenol A' OR 'TBBPA' OR '2,2', 6,6'-Tetrabromo-4,4'-isopropylidenediphenol' } AND { 'reproductive' OR 'oocyte' OR 'ovary' OR 'uterus' OR 'testes' OR 'sperm' OR 'Leydig cell' OR 'Sertoli cell' OR 'steroidogenesis' OR 'estradiol' OR 'testosterone' OR 'follicle-stimulating hormone' OR 'luteinizing hormone' OR 'thyroid' OR 'pregnenolone' }.

3. BPA and female reproductive health

3.1 Oocyte production and quality

Experimental studies conducted prior to 2014 provided compelling evidence that BPA had the potential to affect two stages of oogenesis. The onset of meiosis in fetal ovaries, germ cell nest breakdown, and follicle formation were concluded to be the main causes of BPA adversely affecting maturing oocytes. Three studies found that gestational exposure to low-dose BPA in mice induced increased expression of *Stra8* and a variety of meiotic genes in C57BL/6 mice [38], where longer gestational exposure down-regulated the expression of *Stra8*, *Dazl*, and *Nobox* in CD-1 mice [39]. These results suggested that BPA exposure can cause alterations of gene expression in germ cells and early meiocytes. In neonatally exposed lambs, low-dose BPA was reported to increase the number of multi-oocyte follicles [40], where another study using macaques with low-dose dietary BPA exposure also showed increased numbers of oocytes present in secondary and antral follicles at birth, and continuous exposure (<1 ng/mL in maternal serum) led to the increased incidence of unenclosed oocytes [41]. Prior to 2014, there was some evidence in an *in vitro* study that BPA altered spindle formation at a dose of 43.8 μ M in MF-1 mouse oocytes [42], and in a study of follicle-enclosed oocytes, BPA at a dose of 30 μ M impaired spindle alignment and caused meiotic arrest [43]. Experimental animal studies conducted after 2014 reported that BPA exposure affected oocyte quality, fertilization, and maturation (overview in Table 1), which is consistent with previous studies [14]. In a more recent study, low-dose BPA (50 μ g/kg) was given orally to adult C57BL/6J mice and significantly decreased the percentage of fertilized oocytes without any ovulation changes [25]. In an *in vitro* study, it was reported that BPA treatment decreased meiosis progression and increased spindle abnormalities, including abnormal spindle morphology and chromosome alignment at doses of 15 and 30 ng/ml [44]. Additionally, bovine oocytes exposed to BPA at a dose of 130 nM showed significant increases in DNA damage and apoptosis, while gene expression in the blastocysts

was not altered after oocyte fertilization [24]. Wang et al. reported that porcine oocytes exposed to BPA at a dose of 250 μ M showed a decrease in maturation rate, increases in reactive oxygen species (ROS), abnormal cytoskeletons, increases in apoptosis/autophagy rates, and alteration of the epigenetic pattern [28]. Similarly, Nakano et al. reported that BPA (2 μ g/ml) decreased the maturation rate and induced cell cycle delay and spindle abnormalities in ICR mouse oocytes [45]. The data from the most recent experimental studies further strengthen the evidence that BPA affects oocyte quality, number, maturation and induced meiotic arrest of oocytes.

3.2 Steroidogenesis

Prior to 2014, multiple epidemiological studies analyzed associations between BPA exposure and ovarian steroid hormone production in women. Three studies conducted examined women undergoing *in vitro* fertilization (IVF) and concluded that prior to oocyte retrieval, there was a decrease in peak serum estradiol levels [46-48]. However, in a study conducted by Galloway et al. on adults living in Chianti, Italy, BPA was not associated with changes in estradiol or testosterone levels in women [49]. Therefore, in 2014, the authors concluded that further studies are needed to examine the association between exposure to BPA and steroidogenesis in women [14]. Since 2014, several epidemiological studies have investigated the association between BPA exposure and hormone production in women (overview in Table 2). More recently, a study examined Korean girls with precocious puberty, and found urinary BPA levels were associated with higher levels of testosterone (T), estradiol (E2), progesterone (P4), and the alterations of steroid metabolism among individuals with high levels of BPA. However, the authors concluded that the BPA levels were not directly associated with the onset of precocious puberty [19]. However, Akin et al. reported that BPA levels were associated with a high level of serum T in polycystic ovary syndrome (PCOS) patients in Turkey [50]. Miao et al. found that urinary BPA concentrations were positively associated with higher prolactin, E2, and P4 levels in Chinese female workers with occupational BPA exposure [51]. Additionally, in the National Health and Nutrition Examination Survey (NHANES) conducted between 2011–2012, BPA was associated with higher total T levels in female adolescents [20]. However, a prospective cohort study conducted by Minguez-Alarcon et al. showed that BPA was not associated with peak E2 levels in women with at least one *in vitro* fertilization (IVF) cycle [52]. Zhou et al. also reported that no association between BPA levels and serum follicle-stimulating hormone (FSH) levels was observed in infertile women with PCOS [53]. While there are inconsistencies across studies, multiple epidemiological studies have indicated that urinary or serum BPA levels were associated with alterations of hormone levels in females. It must be noted that many of these studies examined women with impaired reproductive functions and used single-spot urine analysis. Therefore it is important for future studies to examine the associations between long-term BPA exposure and hormone levels of the general population.

Prior to 2014, experimental studies examined the effects of BPA exposure on female steroidogenesis in laboratory animals. In two studies utilizing Sprague-Dawley rats and ICR mice, low-dose exposure to BPA induced increased levels of testosterone and progesterone [54, 55]. While studies used a variety of strains of rats, mice, and lambs exposed

gestationally or gestationally and neonatally, no effect on steroidogenesis was observed [40, 56-58]. *In vitro* studies demonstrated that exposure to 44 and 440 μM BPA inhibited estradiol, testosterone, androstenedione, estrone, dehydroepiandrosterone, and progesterone production, and decreased *StAR* and *Cyp11a1* expression in cultured intact murine antral follicles [59, 60]. Another *in vitro* study examined the effects of BPA at doses 0.1, 1 and 10 μM on porcine granulosa cells; in this study, the lowest dose increased estradiol levels, whereas the higher doses decreased them [61]. At this point in time, the studies indicated BPA exhibited dose-dependent adverse effects on steroidogenesis. Since 2014, experimental studies have examined the effects of BPA exposure on female steroidogenesis using the rodent model (overview in Table 3). In one study of Wistar rats, low-dose neonatal BPA exposure via intraperitoneal injection decreased serum P4 levels without any changes in E2 levels [62]. On the other hand, BPA increased serum P4 levels coupled with high mRNA expression of 3β -hydroxysteroid dehydrogenase (3β -HSD) in Wistar rats with prenatal and neonatal exposure via drinking water [63]. Furthermore, low-dose BPA exposure via drinking water increased serum luteinizing hormone (LH), E2 levels, and follicle numbers, while there were no changes in serum FSH levels observed in Wistar rats [64]. In other strains of rats, Delclos et al. reported that SD rats with prenatal and neonatal oral exposure to BPA showed increased E2 and prolactin levels while decreased P4 levels were observed only at the highest dose group (300000 $\mu\text{g}/\text{kg}$) [23]. Sadowski et al. reported that Long-Evans rats exposed to low-dose BPA during the prenatal and neonatal period showed a decrease in FSH levels but no changes in other hormone levels or maze behavior [65]. Low-dose BPA (50 $\mu\text{g}/\text{kg}$) given orally to adult C57BL/6J mice during 3 estrous cycles did not affect serum E2, FSH, and LH levels [25]. Only one *in vitro* study reported that BPA exposure (2 or 20 mg/ml) decreased P4 synthesis and the protein expression levels of steroidogenesis enzymes (3β -HSD, CYP11A1, and CYP19A1) in human granulosa cells [66]. Collectively, the current data indicate that BPA adversely affects female steroidogenesis, but its effects vary by animal species, strain, exposure route, and exposure window *in vivo*, while *in vitro* studies have indicated that BPA has adverse effects on steroidogenesis depending on the concentration tested.

3.3 Ovary and Uterine

In 2014, there was a shortage of epidemiological studies examining the associations between PCOS and BPA exposure. In a case-control study of 71 women with PCOS and 100 without, an association was observed between serum BPA levels and increased testosterone and androstenedione levels in PCOS patients [67]. Prenatal and neonatal low-dose exposure to BPA in rodents led to the disruption of estrous cyclicity, increased testosterone production and ovarian cysts [54, 68, 69]. High-dose exposure in rodents, on the other hand, led to increased ovarian cysts and accumulation of large antral follicles [54, 68]. In the BPA-treated rodents, cystic appearing follicles differ from the ovarian phenotype in women with PCOS, while the ovarian phenotype of BPA-treated rodents differed from the ovarian phenotype in women with PCOS [54, 68]. Due to the differences in ovarian phenotypes between women with PCOS and the rodent models with PCOS, the report concluded that additional studies were needed to determine why the outcomes differ in animal models and women when exposed to BPA. There have been a limited number of studies conducted investigating the effects of BPA on the ovaries in women (overview in Table 2). Since 2014,

one case-control study of 112 women between the ages of 13 and 19 suggested that urinary BPA levels were associated with a higher risk of PCOS [50]. Additionally, in 268 infertile women diagnosed with PCOS, urinary BPA levels were associated with a significant decrease in antral follicle count (AFC) [53]. In a study conducted on 62 market cashiers diagnosed with PCOS, the serum levels of BPA and the LH/FSH ratio were significantly increased, whereas TSH levels were decreased compared to healthy women with similar jobs [70]. On the other hand, a case-control study of 52 women found no association between urinary BPA concentrations and risk of PCOS [71]. Given the limited number of studies and participants in each study, it appears there is an association between BPA exposure and PCOS symptoms; however, further studies are required to validate these current findings.

Since then, experimental studies in rodents (summarized in Table 4.1) have indicated that prenatal and postnatal exposure to low-dose BPA resulted in decreased ovarian weights, follicle numbers, and primordial follicle recruitment in Wistar rats, as well as increasing the number of corpora lutea and causing a delay in vaginal opening [62, 63, 72]. Additionally, very high-dose BPA exposure (300 mg/kg) in SD rats led to an increase in cystic follicles and decreases in corpora lutea and antral follicles [23]. Zhou et al. reported that BPA exposure decreased germ cell nest breakdown and altered the expression of key genes involved in apoptosis and anti-oxidation in CD-1 mice ovaries *in vitro*, though the effects varied by ovary collection times and exposure doses [73]. Furthermore, BPA has been reported to reduce small and large primary and secondary follicle numbers, and to increase DNA damage markers (γ -H2AX, ATM,) and DNA repair genes in the F344 rat ovary *in vitro* [74]. These data suggested that DNA damage may be one of the potential mechanisms of ovarian toxicity induced by BPA [74]. A study regarding BPA exposure on the ovaries of F1 generation mice resulted in a non-monotonic dose-relationship with several transgenerational effects found in the F3 generation [22]. These data indicated the possibility that BPA is capable of causing epigenetic alterations, which are passed transgenerationally, and can impact fertility, spermatogenesis and social and behavioral activity [75].

Prior to 2014, there were limited data regarding BPA exposure and uterine endometrium in women. In a case-control study on a population of 69 women, it was suggested that BPA concentration may have an association with the occurrence of endometriosis [76]. Buck Louis et al., in their study of 495 individuals as an operative cohort and 131 women in a population cohort, reported no association between exposure and endometriosis; however, the authors point out that the study was not designed to investigate BPA exposure nor powered to assess endometriosis [77]. Therefore, the panel in 2014 saw fit to declare more epidemiological studies were required before concluding whether BPA had adverse impacts on the uterine endometrium. Since 2014, more epidemiological data has been released. Three studies have reported the impact of BPA exposure on uterine morphology in women (overview in Table 2). In a case-control study of women with endometriosis, Upson et al. found that urinary BPA levels were associated with non-ovarian pelvic endometriosis, but had no relation to ovarian endometriosis [78]. Conversely, Minguez-Alarcon et al. reported that there was no association between urinary BPA levels and endometrial wall thickness in 256 women undergoing IVF. Younger women (<37 years old) were found to have a thicker endometrial wall as urinary BPA concentrations increased, while older women (> 37 years old) had a thinner endometrial wall [52]. Additionally, urinary BPA levels were significantly

higher in 68 patients with endometriosis compared to the BPA levels in the control group [79]. Given the limited number of studies and lack of BPA exposure data during endometriosis development, future studies are needed to elucidate the association between chronic BPA exposure and uterine morphology alterations in the general population.

Prior to 2014, experimental animal studies supported the premise that BPA exposure had adverse effects on the uterus. Adult female Balb-c mice prenatally and neonatally exposed to low-dose BPA developed endometrial-like structures which expressed *Hoxa10*, a transcription factor responsible for the mediation of the proliferation of stromal tissue [80]. Increased expression of *Hox10* was further supported by studies that examined low- and high-dose prenatal exposure in adult CD-1 and ICR mice [81, 82]. This finding was further supported by studies utilizing low- and high-dose exposure on prenatally exposed CD-1 and ICR mice, wherein prenatal low-dose BPA exposure to adult mice had been reported to induce benign and malignant lesions and endometrial polyps in the uterus, as well as perturb the Wölfian duct [69, 80]. Since 2014, multiple experimental studies have added to our understanding of the effects of BPA exposure on the uterus in rodent models (overview in Table 4.2). In two studies conducted on CD-1 mice, low-dose BPA exposure increased gland nest density, periglandular collagen accumulation, and abnormal endometrial epithelial and stromal functions [83, 84]. Additionally, Calhoun et al. reported that postnatal exposure to BPA via food consumption (400 µg/kg) altered three genes (HOXA13, WNT4, and WNT5A) involved in reproductive organ development in rhesus macaque uterus, while fetal uteri did not show any notable changes in uterus histology, proliferation, or expression of hormone receptors [85]. In one study on Wistar rats, prenatal and postnatal exposure to low-dose BPA resulted in decreased glandular proliferation rates and α -actin expression at PND90, along with increased abnormal luminal and glandular epithelium at PND360 [27]. In addition, low-dose uterine BPA exposure in SD rats *ex vivo* decreased the force of uterine contractions [86]. Kim et al. reported that short-time exposure to high-dose BPA led to a rapid and transient increase in *Egr1* via the ER-ERK1/2 pathway in ICR mice [87]. Prior *in vitro* studies supported the hypothesis that BPA exposure had adverse effects on the uterus. Exposure to 50 µM BPA was found to decrease the proliferation of cultured human endometrial endothelial cells after 48 hours [88]. In another study examining culture primary heterogeneous populations of uterine cells, a dose of 10 µM BPA inhibited uterine cell contractions, increased oxytocin-related pathways and decreased prostaglandin-related signaling after 48 hours [89]. Since 2014, more *in vitro* studies have been conducted. One study found that BPA (2.28 ng/mL) upregulated genes involved in cell differentiation and proliferation and downregulated genes involved in mitosis and adhesion in uterine smooth muscle cells [90]. It is important to note that in human endometrial endothelial cells and uterine leiomyoma, BPA exposure increased expression of vascular endothelial growth factor (IVGF) at the micromolar level [91, 92]. Wang et al. reported BPA exposure increased growth rate and colony-forming efficiency via CoX-2 in an endometrial carcinoma cell line [93]. Collectively, these studies have provided compelling evidence suggesting that BPA adversely affected the uterine both *in vivo* and *in vitro*, though the assessment endpoints varied across all studies.

4. BPA male reproductive toxicities

4.1 Sperm production and quality

Epidemiological data conducted prior to 2014 examined the associations between BPA exposure and sperm quality and production. While these studies provided consistency among results, the studies were limited by lack of evidence associating exposure and quality of semen. Occupationally exposed men [94] and patients recruited from infertility clinics [95] had higher urinary BPA levels, which were associated with decreased sperm count and motility. Another study examined fertile men and found no association between urinary BPA concentrations and changes in semen parameters [96]. Since 2014, many epidemiological studies have built on our understanding of how BPA affects sperm production and quality, but many limitations are still prevalent (overview in Table 2). In a prospective cohort study of male partners undergoing IVF treatments, the natural log-transformed urinary BPA concentrations were associated with a lower natural log-transformed sperm count, concentration, and motility [97]. Additionally, a study of young men from Denmark reported that men in the highest quartile of urinary BPA levels also had significantly lower sperm progressive motility when compared to the men in the lowest quartile [18]. On the other hand, in a study of male partners trying to conceive without any intervention methods, urinary BPA levels were associated with lower sperm DNA fragmentation [17]. This lower fragmentation rate may be explained by the low BPA levels detected in the study population when compared to NHANES data or the data obtained from sperm analysis conducted the day after sample collection [17]. Given the limited information and discrepancies among results, additional studies using more sensitive and reliable measurements to capture continuous BPA exposure and conduct semen analysis are required to determine if there is any association between BPA exposure and sperm quality in the general population. One potential reason why this data may not be currently available is that the cost per sample can be quite expensive. While this may work for small sample size, the data may not be indicative of the general population.

The earliest animal experimental studies on the effects of BPA on sperm suggested adverse consequences on spermatogenesis in adulthood following either prenatal or early postnatal exposure. Multiple studies analyzed the effects of gestational exposure to low-dose BPA and found decreased numbers of elongated spermatids in the seminiferous tubules of pubertal ICR mice and reduced sperm counts in Holtzman rats [98, 99]. In 2010, Minamiyama reported that BPA-induced decreases in rat sperm motility had been prevented by co-administering antioxidant *n*-acetylcysteine [100]. The lab suggested that the impaired sperm motility may be related to the increased levels of reactive oxygen species. In more recent animal studies examining BPA exposure and male reproduction (overview in Table 5), prenatal and postnatal exposure to low-dose BPA adversely affected sperm production and quality. Despite the variations in exposure windows, duration, species or strains, and endpoints examined, several findings in the previous literature were commonly observed: decreased sperm number, induction of sperm apoptosis and oxidative stress, and the alteration of seminiferous tubule morphology. Many studies reported that both prenatal and postnatal exposure to low-dose BPA decreased sperm numbers in mice and rats [101-108]. Additionally, induction of testicular oxidative stress or apoptosis was observed in mice and

rats exposed to low-dose BPA during the prenatal or postnatal periods [106, 108-112]. Low-dose BPA exposure was reported to delay meiosis, as well as inducing the accumulation of chromosomal abnormalities and meiotic DNA double-strand breaks (DSBs) in the late meiotic stage [113]. Few studies have indicated that low-dose BPA exposure potentially altered the epigenetic pattern in testes, including increased rates of methylation of *Igf2* and decreased protein lysine acetylation levels in rats [114, 115]. In multiple strains of mice, however, exposure to BPA did not affect chromosome pairing and synapsis [115, 116]. Overall, the strength regarding current evidence for risk assessment regarding low-dose BPA exposure is still limited. The relevance of the doses in the experimental animal studies to human exposure is still unclear since the internal (urinary or serum) BPA levels in tested animals were not examined, therefore making extrapolation of these findings to humans difficult.

Several *in vitro* studies have further investigated the modes of action that BPA utilizes for testicular toxicity (overview in Table 5). In two studies using the mouse spermatocyte GC-2 cell line, BPA exposure (20 μM) induced germ cell apoptosis via the Ca^{2+} /CaM/CaMKII signaling pathway and the PERK/EIF2 α /chop pathway [111, 117]. Increases in tyrosine phosphorylation via PKA were observed in the primary spermatozoa isolated from mice and rats [118, 119]. Additionally, BPA exposure (50 μM) induced early DNA damage responses and perturbed cytoskeleton in the C18-4 spermatogonial cell line [120]. BPA exposure at a very high concentration (300 μM) to human spermatozoa decreased the mitochondrial membrane potential and increased both apoptosis and DNA oxidative damage marker (8-hydroxy-2'-deoxyguanosine) [121]. Another study found that short-time BPA exposure at an extremely high-dose (1 mM) increased velocity straight linear, homogeneity of progressive movement velocity and intracellular free Ca^{2+} concentration in human spermatozoa [122]. Collectively, this *in vitro* data suggests that BPA exposure may impact male germ cells through perturbation of Ca^{2+} homeostasis and cytoskeleton structure, induction of apoptosis, and DNA damage.

4.2 Sertoli and Leydig Cells

Sertoli and Leydig cells play crucial roles in maintaining the normal functions of spermatogenesis. The functions of the Sertoli cells include the formation of the blood-testis barrier, secretion proteins, and growth factors that nurture and regulate apoptosis and mitosis of germ cells [123, 124]. Leydig cells are responsible for synthesizing and secreting T to support spermiogenesis [125]. Prior to 2014, few studies had analyzed the effects of BPA exposure on Leydig and Sertoli cells to determine whether BPA had a direct effect on these cells and to see if there was an effect on steroid hormone production. Continuous high-dose pubertal exposure to male Wistar/ST rats was reported to decrease the cell number and steroidogenic enzyme expression in Leydig cells [126]. However, gestational plus neonatal exposure to low dose BPA increased Leydig cell numbers in Long-Evans rats in adulthood via the up-regulation of mitogenic factors [127]. Nanjappa et al. also reported that adult-exposed Leydig isolates to low-dose BPA decreased testosterone production and the expression of steroidogenic enzymes [127]. Since 2014, multiple *in vitro* studies have examined the effects of BPA on Sertoli and Leydig cells (overview in Table 6). In three studies using the Sertoli cell line TM4, nanomolar BPA increased proliferation via

promoting energy metabolism, GPR 30, and ER α/β , while micromolar BPA decreased cellular proliferation via inducing oxidative stress and activating CaM-CaMKII-ERK and mitochondrial apoptotic pathways [128-130]. In studies conducted on primary rat Sertoli cells, BPA (~50 μM) induced cellular apoptosis via ROS, mitochondrial dysfunction, and activated HNKs/p38 MAPK, NF- κB , and Pten/Akt pathways [26, 93, 131]. An important finding was that in human primary Sertoli cells, BPA exposure significantly decreased cell viability, expression levels of occludin, ZO-1, β -catenin, and AR at a dose of 20 μM without changes in F-actin expression or localization [132]. Another study concluded that BPA treatment perturbed actin filaments starting at 0.4 μM and induced improper localization of actin regulatory proteins Arp3 and Eps8 at a dose of 200 μM [133]. An *in vivo* study demonstrated that high-dose BPA exposure decreased Leydig cell numbers and StAR protein levels in the adult rat testis [127, 134]. On the other hand, low-dose BPA exposure induced Leydig cell proliferation with increased protein expression of proliferating cell nuclear antigen, MAPK, AR, and ER α/β . Low-dose BPA was found to increase secretion of the anti-Mullerian hormone in the adult rat testis [127, 134]. In another *in vitro* study, BPA exposure decreased T biosynthesis and E2 levels, mRNA expression of aromatase, and the steroidogenic enzyme 17 α -hydroxylase/17–20 lyases in rat primary Leydig cells [135]. In Leydig cell line TM3, micromolar doses of BPA decreased cellular proliferation rates and increased cell migration and invasion [136]. Given the current *in vitro* and *in vivo* studies, BPA exposure affected the proliferation and steroidogenesis of rodent Sertoli and Leydig cells. Furthermore, the results of the experimental studies indicated that BPA exposure may be associated with decreased hormone levels in male animals and suggested direct effects of BPA on Sertoli and Leydig cells [14].

4.3 Steroidogenesis

Before 2014, the epidemiological studies conducted regarding BPA exposure and male steroidogenesis had conflicting reports. The INChianti study noticed an association between increased urinary concentrations of BPA and increased serum testosterone levels but noticed no association with estradiol levels in males [49]. However, cross-sectional studies of 167 and 302 fertile men reported no associations between BPA exposure and testosterone levels [96, 137]. One of these studies conducted by Mediola et al. found that urinary BPA concentration was associated with decreased free androgen index (the ratio of free androgen index to luteinizing hormone) and a decrease in the ratio of free testosterone to luteinizing hormones in the male partners of pregnant women [96]. In contrast, Meeker et al. found associations between high concentrations of BPA and increased serum levels of follicle-stimulating hormone, and decreased levels of inhibin B and the ratio of estradiol to testosterone [137]. Since 2014, five more epidemiological studies have examined the association between urinary BPA levels and hormone levels in men (overview in Table 2). A study conducted on young men from Denmark reported that those with BPA levels above the lowest quartile had higher levels of T, LH, and E2 as compared to the men in the lowest quartile [18]. In two cross-sectional studies of male workers occupationally exposed to BPA, urinary BPA concentrations were associated with higher sex hormone-binding globulin (SHBG) and lower androstenedione (AD) [138, 139]. In male children and adolescents in the NHANES 2011–2012 study, BPA was associated with lower total T levels [20]. However, a retrospective cohort study determined that there was no association between

maternal (3rd trimester) or childhood BPA exposure on hormone levels in boys ages 8–14 [140]. These data indicated that BPA exposure have the capability of perturbing the hormone levels in men. However, the strength of the study was limited by the study population with occupational exposure, the use of single spot urine samples, and the lack of a co-exposure assessment. Many occupational exposure studies on BPA have already been conducted. For instance, Hines et al. found that 1.4% of workers involved with industries that manufacture and use BPA in the United States exceeded the US EPA oral reference dose for BPA (50 µg/kg-day) [141]. Another study conducted in Finland showed that the highest occupational exposure levels were found in the manufacturing of thermal paper, and that increased urinary levels were noticed in liquid paint factories [142]. These studies provide insight into occupational exposure to BPA. However, these studies do not provide any health analysis of the participants, which weakens their strength. For future studies, it is imperative to analyze both occupational exposure and subsequent health of the participants to determine whether BPA exposure has any effects on male steroidogenesis.

Data prior to 2014 regarding experimental studies suggested that BPA exposure decreased hormone levels in male animals. However, this conclusion was considered contradictory. One study examined gestational through neo-natal exposure to low-dose BPA in CD-1 mice, and reported that there was a decrease in testosterone levels [143]; but this was not reported in a study examining the effects of *in utero* exposure on adult C57BL/6 mice [144]. Studies examined low-dose exposure to gestationally or neonatally exposed Holtzman rats [99, 145], adult-exposed albino rats [146], and adult-exposed Wistar rats [147] and concluded that these species experienced decreased testosterone levels. However, when examining gestational and neonatal exposure of low-dose BPA in Long Evans rats [148] or SD rats [56, 149], testosterone levels were not affected. Therefore, there was conflicting evidence in 2014 that associated BPA with decreased expression of steroidogenic enzymes and even affected the levels of follicle-stimulating hormone, which are required to start steroidogenesis. Recent experimental studies have assessed prenatal exposure to low-dose BPA, and it has been reported to alter the hormone levels in mice and rats, but the data are not consistent (overview in Table 7). Gamez et al. reported that low-dose BPA exposure led to increased serum LH and FSH levels in young Wistar rats [150]. On the contrary, a study conducted on adult Wistar rats found that BPA exposure decreased serum T, LH, and FSH levels and increased the E2 level [104]. In two studies using SD rats, findings reported decreased serum T and E2 levels with postnatal low-dose BPA exposure [101]. BPA at an extremely high dose (300 mg/kg) significantly increased serum thyroxine (T4) levels without changes in serum triiodothyronine (T3) and FSH levels [23]. BPA exposure was reported to decrease serum T levels in Swiss albino and C57BL/6 mice, but these doses varied between 0.5 µg/kg to 100 mg/kg [151, 152]. Additionally, Sadowski et al. reported that Long-Evans rats exposed to low-dose BPA showed a decrease in FSH levels at weaning [65]. An *in vitro* study examined fetal testis explants from rats, mice, and humans, and concluded that BPA exposure reduced basal T secretion at a concentration of 10 µM in rats, 1 µM in mice, and 10 nM in human explants, respectively [153]. These data suggested that BPA exposure could impair male steroidogenesis in rodents, but the effects vary by species, strain, exposure window, and duration.

5. BPA analogs and reproductive health

5.1 BPS and reproductive health

BPS is structurally similar to BPA and is now used in a variety of common consumer products as a BPA alternative. BPS has been detected in food, indoor dust, personal care products, sediment, and paper products such as currency and cashier's receipt [32, 154-156]. BPS exposure frequently occurs through ingestion, inhalation, and dermal contact. BPS was detected in 81% of the human urine samples in the United States and in seven Asian countries with a mean concentration of 0.654 ng/mL, which was comparable to BPA [33]. Neither the metabolic nor biological fate of BPS has been fully examined. An *in vitro* study indicated that glucuronidation was the major metabolic pathway for BPS [157]. As a potential EDC, BPS has been examined in the National Toxicology Program (NTP) Tox21 High Throughput Screening (HTS) Program and was classified as an estrogen agonist with a weak affinity for the ER [158, 159]. In addition, Rochester and Bolden reviewed the endocrine activity of BPS by comparing BPS's estrogenic binding potency against estradiol (EE₂), a positive control, on a plethora of estrogenic receptors. They then ran an experiment to examine BPA's estrogenic potency and compared this value to EE₂'s potency. The average estrogenic potency for BPS was calculated to be 0.32 ± 0.28 by dividing the estrogenic potency of BPA from the estrogenic binding potency of BPS, indicating BPS had a similar affinity to ER receptors as BPA [160]. This study also indicated that BPS was in the same order of magnitude as BPA regarding its androgenic, antiandrogenic, antiestrogenic and aryl hydrocarbon binding affinity, and in inhibitory hormone signaling of adipocytes [160]. A study conducted by Rosenmai et al. found that BPS bound to estrogen receptors and affected estrogenic and antiandrogenic activity in a similar manner to BPA [161].

However, there have only been a few *in vivo* studies that have examined the effects of BPS exposure on female and male reproductive systems (overview in Table 1). Yamasaki et al. reported that in the immature rat utero-trophic assay, BPS exposure significantly increased absolute and relative uterine wet weights and blotted weights at doses of 20 and 500 mg/kg, but not at a dose of 200 mg/kg [162]. In SD adult rats, BPS exposure induced testicular reactive oxygen species (ROS) and lipid peroxidation, altered seminiferous epithelium morphology, and decreased antioxidant enzyme activity and plasma T levels in the highest dose group (50 µg/kg) in spermatogonia, spermatocytes, and spermatids [163]. However, there were no significant changes in the populations of these testicular cells [163]. As vertebrates, zebrafish have conserved pharmacological targets and nervous system structures, which is comparable with mammals. This therefore makes zebrafish an ideal specimen to study, as they have a large genetic database, short lifespan, and high fecundity. Zebrafish have been widely used as a model to identify targets as well as modes of the action of EDCs [12]. In two studies that examined zebrafish, both adult and developmental exposure to BPS resulted in increased plasma estradiol (E₂) levels in both sexes, while adult exposure altered the E₂ levels at lower doses when compared to developmental exposure. Additionally, male fish exhibited more sensitive responses when compared to the female fish [164, 165]. BPS exposure in male zebrafish at different points in their lifespan had decreased T levels [164, 165]. Chen et al. reported that *Caenorhabditis elegans* exposed to BPS showed

increased germline apoptosis and activation of DNA damage checkpoint kinase CHK-1. BPS is also known to cause distinct alterations of gene expression at the whole transcriptome level compared to BPA [166]. Consequentially, these current studies have provided compelling evidence that BPS exposure altered hormone homeostasis and impaired reproductive organs in different sexes, exposure windows, and species. More *in vivo* studies are needed to validate these current findings and examine the effects of BPS exposure on multiple rodent strains.

In an *in vitro* study, Eladak et al. reported that BPS exposure decreased basal testosterone (T) secretion in mouse or human fetal testis explants, starting at doses of 100 nM and 1000 nM, respectively [153]. When compared to BPA, BPS treatment induced more significant changes in T secretion in the mouse explants, but fewer changes in human explants [153]. In an *in vitro* test on the Leydig cell line, MA-10, a dose of 10 μ M BPS significantly increased P4 and P5 levels and increased gene expression levels of CYP51 and 5 α -Red1 [167]. In addition, BPS exposure also decreased cell viability and increased early DNA damage responses and abnormal cytoskeleton structure at a dose of 50 μ M in a mouse spermatogonial cell line C18-4 [120]. The dose responses obtained of cell viability, cell cycle alteration, DNA damage, and cytoskeleton were comparable to BPA, but less drastic when compared to BPAF or TBBPA [120].

5.2 BPF and reproductive health

BPF is a component of epoxy resins widely used in tank and pipe linings, industrial floors, coatings, dental sealants, and food packaging materials [160]. BPF has been found in indoor dust and various food items such as beverages, dairy products, meats, seafood, cereals, fruits, and vegetables [32, 155]. It has had been reported that BPF was metabolized and transformed to nonactive sulfates, rather than glucuronides, and excreted through urine in pregnant rats. A study found that active BPF was distributed to multiple tissues, including the liver, uterus, placenta and fetus [168]. In the United States population, BPF was found in approximately 60% of urine samples, and the concentrations ranged from 0.15 to 0.54 μ g/L [169]. The average estrogenic hormonal potency for BPF, as compared to BPA, was reported to be 1.07 ± 1.20 , calculated in the same manner as BPS [160]. Therefore, BPF may have an effect on the endocrine system equal to or possibly more potent than that of BPA.

Limited data are available for BPF exposure and reproductive outcomes in experimental animals (overview in Table 2). Two studies conducted on female Wistar rats suggested that short-time postnatal exposure to BPF at doses over 100 mg/kg increased uterine relative weight [162, 170]. However, another study did not find any changes in reproductive organ weights in young adult female SD rats exposed to BPF at concentrations reaching 500 mg/kg [171]. Meanwhile, in male SD rats, BPF exposure increased testis weight without significant spermatological changes at a dose of 500 mg/kg. In the same study, rats in both sexes showed decreased serum triiodothyronine (T3) levels and increased thyroxine (T4) levels. The opposite trends in T3 and T4 levels and no changes to serum thyroid-stimulating hormone (TSH) levels suggested that these observed effects were not related to the endocrine-mediated effects of BPF [171].

According to the results of recent *in vitro* studies, BPF's effects on steroidogenesis are equivocal to BPA. BPF has been reported to increase T secretion and expression of steroidogenic genes (Cyp51 and 5 α Red1) at a dose of 100 μ M in MA-10 Leydig cells [167]. Eladak et al. reported that BPF exposure decreased basal T secretion in mouse fetal testis explants starting at a dose of 1000 nM, which followed a similar dose-response curve as BPA. In human fetal testis explants, BPF treatment reduced T levels at a dose of 10 nM with a non-monotonic dose-response curve (10 nM) [153]. The discrepancy in the current data could be the result of different biotransformations of BPF into its metabolites the *in vitro* models and the different species and strains tested. Collectively, the data suggest that BPF exposure may alter steroidogenesis both *in vitro* and *in vivo*, but its effects on the reproductive organs, oogenesis, spermatogenesis, and embryonic development remain inconclusive. Future studies are necessary to fully validate the current findings in multiple rodent models and determine the consequences of BPF-induced hormone alterations on reproductive functions.

5.3 BPAF and reproductive health

BPAF is used widely as a crosslinking agent and a monomer in the plastics industry. There is limited information available on the occurrence of BPAF in consumer products and human urine or blood samples. In the United States, BPAF has been found in various food items with lower detection rates (6% - 30%) compared to BPA [155]. BPAF was found to be metabolized primarily through glucuronidation and excreted through feces and urine in SD rats [172, 173]. In humans, BPAF was detected in urine samples at concentrations ranging from below the detection level to 3.93 μ g/L in China and Saudi Arabia [174, 175]. It is important to note that BPAF exhibited estrogenic binding potencies greater than those of BPA [36, 37]. Additionally, BPAF exhibited a preferential affinity (three times stronger) for ER β compared to ER α [176].

Relatively few studies have examined the effects of BPAF exposure on steroidogenesis *in vivo* (overview in Table 3). Two studies performed on zebrafish demonstrated that exposure to BPAF during development led to increases in E2 levels in the female fish [177, 178]. The gonadal examination carried out on the zebrafish indicated that exposure to 1 mg/L BPAF induced acellular areas in the testis and retarded oocyte development [178]. In those same studies, BPAF exposure decreased T levels in male fish and also altered testicular morphology [177, 178]. In contrast, no decreases in T production were observed in SD rat dams prenatally exposed to BPAF at a dose of 750 mg/kg [179]. Furthermore, Li et al. reported that BPAF could be transferred via cord blood and lactation, finally accumulating in the offspring's testes [137]. Offspring exposed to BPAF prenatally and postnatally showed a significant increase in testicular T levels and alterations of genes involved in cell differentiation and meiosis [180]. Feng et al. reported that BPAF exposure decreased serum T level and T biosynthesis (200 mg/kg), and increased serum luteinizing hormone (LH) (50 mg/kg) and follicle-stimulating hormone (FSH) (10 mg/kg) levels in adult male SD rats [181]. Recently, a study revealed that BPAF exposure uniquely impaired pregnancies and sexual development in rats at doses of ~80 and ~280 mg/kg, whereas BPA exposure did not alter these reproductive endpoints at similar concentrations [182]. These data suggest that BPAF might be a more potent endocrine disruptor than BPA.

In *in vitro* studies, exposure to BPAF inhibited mouse oocyte maturation at concentrations of 50 and 100 µg/mL and induced cell cycle arrest by the activation of spindle assembly checkpoints [45]. In a recent multi-parametric high-content analysis (HCA) of mouse spermatogonial cells, BPAF exhibited the highest spermatogonial toxicity when compared to TBBPA, BPA, and BPF [86]. Exposure to 1 µM BPAF altered the nuclear morphology and induced cell cycle arrest, while a concentration of 10 µM caused cytoskeleton perturbation and multinucleation of cells, with DNA damage recognized at a concentration of 25 µM [120]. This *in vitro* study indicated that BPAF exposure may target the spermatogonia cells as observed *in vivo* in a fertility study [182]. The effects of BPAF on steroidogenesis are still unclear and likely differ depending on different exposure windows. Additionally, further studies should be conducted to examine the effects of BPAF exposure on hormone levels and multiple reproductive endpoints in various experimental strains, species, and exposure windows.

5.4 TBBPA and reproductive health

TBBPA's primary use is as a reactive flame retardant in plastics, paper, textiles, and circuit boards. The general population can be exposed to TBBPA through inhalation, dermal contact, and ingestion of fish and shellfish [183]. TBBPA is absorbed by the gastrointestinal tract, then metabolized to glucuronides and sulfates, which in turn are excreted through feces in animal models [184]. Several studies have examined TBBPA exposure levels on the general population. Serum TBBPA levels were below the level of detection (0.03 ng/L) in pregnant Canadian women, but detectable in 5% of Inuit adults with concentrations up to 480 ng/L [185]. In the United States, TBBPA was found in approximately 35% of human breast milk samples at levels between 50–350 pg/kg/day [186]. However, the current data demonstrated that TBBPA did not interact with ER α/β or AR in a panel of *in vitro* bioassays [187].

Van der Ven et al. established a one-generation reproduction study and a subacute toxicity study to examine dose-response relationship of TBBPA and its effects on the endocrine system [188]. The subacute toxicity test was conducted as a 28-day repeat dose study according to OECD407 guidelines, including doses of 0, 30, 100, 300 mg TBBPA/kg body weight. In the two-generation study, exposure started 10 weeks or 2 weeks before mating and was continued during mating, pregnancy and lactation. No exposure-related histopathological changes were observed in any of the assessed organs in either experiment. There were also no effects on endpoints of reproduction -- i.e., mating success, number of implantation sites, and litter size. The cauda epididymis sperm count and morphology were not affected in either experiment. However, a significant dose-dependent increase of the weight of the testis and pituitary weight were observed in F1 male animals. Decreased circulating thyroxine (T4) levels were observed in both the subacute and one-generational study [188].

Similarly, another two-generation reproductive study was performed on SD rats exposed to TBBPA at doses of 0, 10, 100 or 1,000 mg/kg/day in corn oil by oral gavage [189]. The F0 generation was treated during a pre-mating period of 10 weeks and the 2-week mating period. The only significant decrease occurred in serum T4 levels in both the F0 and F1 generations

at a dose of 1000 mg/kg [189]. Cope et al. reported a two-generation reproduction study on reproductive, developmental and neurobehavioral effects of TBBPA at oral doses of 10, 100, and 1000 mg/kg BW/day in SD rats [190]. These findings are consistent with other multi-generational studies, as there were no marked effects on various reproductive parameters including time to the vaginal opening and anogenital distance, sperm motility, concentration, and morphology in F1 or F2 generations. The decrease in T4 levels in SD rats exposed to a dose of 1000 mg/kg of TBBA was the only change observed [190].

Zatecka et al. conducted a two-generational study focused on the trans-generational effects of TBBPA in CD1 outbred mice [191]. Gestational exposure to TBBPA was through drinking water (35 µg/kg) to generate the F1 offspring. Experimental and control animals of the F1 generation were bred in various conditions to evaluate the trans-generational effects on the reproductive system. Significantly reduced testicular weight, increased prostate weight, and increased seminal vesical weights were observed in the F2 generation when both parents (F1) were treated with TBBPA. However, no changes in sperm parameters were observed, and decreased the thickness of the seminiferous epithelium and increased apoptotic cells in the testes by TUNEL staining were observed in both F1 and F2 males [191]. Utilizing C57BL/6J mice exposed to the same level of TBBPA during the gestation, lactation, pre-pubertal, and pubertal periods up to the age of 70 days, Zatecka, et al. found TBBPA treatment did not induce significant changes in any of the general reproductive system parameters such as the anogenital distance, reproductive organ weight, and sperm count and morphology. However, significant reductions in T and T3 levels, and ratios of protamine 1/protamine 2, and increased total protamine/DNA ratio in sperm and apoptotic spermatozoa were observed [192].

A study of B6C3F1/N mice exposed to TBBPA for two years (500 mg/kg) showed an increased incidence of uterine epithelial tumors including adenomas, adenocarcinomas, and malignant mixed Mullerian tumors [193]. In an experiment utilizing zebrafish, TBBPA exposure decreased egg production (0.047 µM) and increased premature egg production (1.5 µM) [194]. In contrast, there were no changes in the ovary or uterine weight observed in ICR mice exposed to 1.0% TBBPA in their diet [195].

In two *in vitro* studies, TBBPA exposure was reported to increase T secretion in Leydig cell line MA-10 at micromolar levels [167, 196]. Ogunbayo et al. reported that TBBPA induced Sertoli cell death via disruption of Ca²⁺ signaling and through affecting Ca²⁺ transport proteins [197]. In a recent study with the mouse spermatogonia cell line C18-4, TBBPA exposure exhibited a dose-dependent induction of nuclear morphological changes from the high content analysis (HCA), including nuclear area and nuclear shape (LWR and P2A). TBBPA also induced significant increases of LWR at concentrations of 5, 10 and 25 µM and a significant increase of P2A at a concentration of 25 µM for 72 h. TBBPA treatments of 25 µM for 48 h, and of 10, 25 µM for 72 h led to an increase of cells in the G2/M phase accompanied by an increase of apoptotic cells. Additionally, TBBPA exposure (25 µM) induced DNA damage responses and perturbed cytoskeleton structure [120]. Compared to BPA and BPS, TBBPA exhibited higher spermatogonial toxicity, including dose- and time-dependent alterations impacting nuclear morphology, cell cycle progression, DNA damage responses, and perturbation of the cytoskeleton [120]. Taken collectively, current studies

indicated that the effects of TBBPA on reproduction varied in a limited number of studies. No severe reproductive dysfunctions were consistently observed, even at the higher doses. Some studies of TBBPA revealed alterations in T3 or T4 levels, though those changes did not accompany any histological changes in the thyroid gland or other reproductive organs. In *in vitro* studies, TBBPA exposure impacted testicular cells at relatively low doses compared to BPA. Future *in vivo* animal studies are needed to determine TBBPA's reproductive toxicity and molecular mechanism.

6. Discussion and Conclusion

This review performed a literature search of epidemiological and experimental animal studies, which examined the reproductive toxicities of BPA analogs. While there were several studies that monitored urinary or serum levels of BPA analogs, the association between BPA analog exposure and reproductive dysfunction have not been explored. Emerging evidence from animal models suggests that exposure to these chemicals could adversely affect reproductive functions, including oocyte and sperm quality, steroidogenesis, and ovary and testis functions. These adverse effects might vary based on different testing species, exposure periods, and duration.

It is also worth noting that our lab's previous study using mouse spermatogonial cells combined with high-content analysis (HCA) revealed BPAF exhibited the highest testicular toxicity (20% maximal effect concentration $EC_{20} \approx 10 \mu\text{M}$), followed by TBBPA, BPA, and BPS [120]. BPAF induced a dose-dependent increase of multinucleated germ cells (MNGs). Induction of MNGs has been reported following gestational exposure to di-(n-butyl) phthalate (DBP) [198-201]. The accumulation of MNGs in the testis has been associated with the formation of carcinoma *in situ* (CIS) cells, the known precursor to testicular germ cell cancer (TGCC) in humans [202]. Additionally, a recently published *in vivo* study showed consistent data that BPAF exposure uniquely impaired pregnancies and sexual development in rats compared to BPA at similar doses [182]. BPS uniquely reduced the reproductive lifespan of, and induced distinct transcriptome changes in, *C. elegans* [166]. Therefore, given the similarities of their estrogenic potencies to BPA, current data indicates that BPA analogs, especially BPAF and TBBPA, may disrupt reproductive functions in an ER-independent manner.

The studies reviewed here provide insufficient toxicological and epidemiological data to characterize and determine the reproductive effects of BPA analogs. In animal models, absorption, distribution, metabolism, and excretion of BPA analogs, as well as their estrogenic potencies, should be determined and evaluated. The effects of developmental and adult exposure should be characterized in multiple species. Currently, the NTP is in the process of carrying out toxicological studies of BPS and BPAF on rodents, and toxicokinetic studies of BPS following oral and intravenous exposure. In *in vitro* studies, adverse signaling pathways need to be elucidated. As endocrine disruptors, the potential interactions of bisphenols and hormone receptors on different testicular cell lines and their toxic consequences need to be determined, using gain- and loss-of-function approaches. Human studies analyzing the associations between analog exposure levels and hormone levels, oocyte and sperm quality, and pregnancy outcomes should be examined. The single urine

spot sample is usually used in BPA studies, but it does not reflect the long-term chemical exposure or determine the real exposure level the progression of a disease's development. Future epidemiological studies need to employ an advanced exposure assessment to monitor long-term BPA analog exposure on individuals. These multifaceted data will potentially allow for comparing information across BPA analogs and a better risk assessment of bisphenols in general.

Given the current data gap, the following types of research studies are needed in the future:

- Environmental studies are needed to better elucidate the environmental occurrence of BPA analogs and determine sources and pathways of human exposure.
- Epidemiological studies should be conducted in the general population with precise measurement of chronic exposure level. Co-exposure should also be considered.
- *In vivo* studies are needed to better elucidate the metabolic pathways of BPA analogs and target the internal doses and exposure times that are relevant to human exposure.
- *In vitro* studies are needed to elucidate the unique mechanisms and the mode of actions of BPA analogs on the reproductive systems, especially the molecular mechanism of the formation of MNGs by BPAF.

Acknowledgments,

We thank Jake Maas for proofreading the final version. This work was supported by the Centers for Disease Control and Prevention, The National Institute for Occupational Safety and Health (NIOSH) under award number R21 OH 010473; the National Institute of Environmental Health Sciences of the National Institutes of Health under award number R43ES027374; Alternatives Research & Development Foundation (ARDF).

Abbreviations:

3β-HSD	3 β -hydroxysteroid dehydrogenase
AR	androgen receptor
BPA	bisphenol A
BPAF	bisphenol AF
BPF	bisphenol F
BPS	bisphenol S
CDC	the US Centers for Disease Control and Prevention
E2	estradiol
EC20	20% maximal effect concentration
ER α/β	estrogen receptor α/β

ER	endoplasmic reticulum
EPA	the United States Environmental Protection Agency
FDA	the U.S. Food and Drug Administration
FSH	follicle-stimulating hormone
GD	gestational day
HTS	high throughput screening
LH	luteinizing hormone
LOAEL	the lowest adverse effect level
NHANES	the National Health and Nutrition Examination Survey
NTP	the National Toxicology Program
P4	progesterone
PF	post-fertilization
PND	postnatal day
ROS	reactive oxygen species
T	testosterone
T3	triiodothyronine
T4	thyroxine
TBBPA	tetrabromobisphenol A
TSH	thyroid-stimulating hormone

References

- [1]. Rochester JR, Bisphenol A and human health: A review of the literature, *Reprod Toxicol* 42 (2013) 132–155. [PubMed: 23994667]
- [2]. Kang JH, Kondo F, Katayama Y, Human exposure to bisphenol A, *Toxicology* 226(2–3) (2006) 79–89. [PubMed: 16860916]
- [3]. Vandenberg LN, Hauser R, Marcus M, Olea N, Welshons WV, Human exposure to bisphenol A (BPA), *Reprod Toxicol* 24(2) (2007) 139–177. [PubMed: 17825522]
- [4]. Calafat AM, Ye XY, Wong LY, Reidy JA, Needham LL, Exposure of the US population to bisphenol A and 4-tertiary-octylphenol: 2003–2004, *Environ Health Persp* 116(1) (2008) 39–44.
- [5]. Lakind JS, Naiman DQ, Daily intake of bisphenol A and potential sources of exposure: 2005–2006 National Health and Nutrition Examination Survey, *J Expo Sci Env Epid* 21(3) (2011) 272–279.
- [6]. Dong S, Terasaka S, Kiyama R, Bisphenol A induces a rapid activation of Erk1/2 through GPR30 in human breast cancer cells, *Environ Pollut* 159(1) (2011) 212–218. [PubMed: 20875696]
- [7]. Gould JC, Leonard LS, Maness SC, Wagner BL, Conner K, Zacharewski T, Safe S, McDonnell DP, Gaido KW, Bisphenol A interacts with the estrogen receptor alpha in a distinct manner from estradiol, *Mol Cell Endocrinol* 142(1–2) (1998) 203–214. [PubMed: 9783916]

- [8]. Kuiper GGJM, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, van der Burg P, Gustafsson JA, Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta, *Endocrinology* 139(10) (1998) 4252–4263. [PubMed: 9751507]
- [9]. Matsushima A, Kakuta Y, Teramoto T, Koshihara T, Liu XH, Okada H, Tokunaga T, Kawabata S, Kimura M, Shimohigashi Y, Structural evidence for endocrine disruptor bisphenol a binding to human nuclear receptor ERR gamma, *J Biochem* 142(4) (2007) 517–524. [PubMed: 17761695]
- [10]. Okada H, Tokunaga T, Liu XH, Takayanagi S, Matsushima A, Shimohigashi Y, Direct evidence revealing structural elements essential for the high binding ability of bisphenol A to human estrogen-related receptor-gamma, *Environ Health Persp* 116(1) (2008) 32–38.
- [11]. Richter CA, Birnbaum LS, Farabollini F, Newbold RR, Rubin BS, Talsness CE, Vandenberg JG, Walser-Kuntz DR, vom Saal FSV, In vivo effects of bisphenol A in laboratory rodent studies, *Reprod Toxicol* 24(2) (2007) 199–224. [PubMed: 17683900]
- [12]. Tokarz J, Moller G, de Angelis MH, Adamski J, Zebrafish and steroids: What do we know and what do we need to know?, *J Steroid Biochem* 137 (2013) 165–173.
- [13]. Wozniak AL, Bulayeva NN, Watson CS, Xenoestrogens at picomolar to nanomolar concentrations trigger membrane estrogen receptor-alpha-mediated Ca²⁺ fluxes and prolactin release in GH3/B6 pituitary tumor cells, *Environ Health Persp* 113(4) (2005) 431–439.
- [14]. Peretz J, Vrooman L, Rieke WA, Hunt PA, Ehrlich S, Hauser R, Padmanabhan V, Taylor HS, Swan SH, VandeVoort CA, Flaws JA, Bisphenol A and Reproductive Health: Update of Experimental and Human Evidence, 2007–2013, *Environ Health Persp* 122(8) (2014) 775–786.
- [15]. vom Saal FS, Akingbemi BT, Belcher SM, Birnbaum LS, Crain DA, Eriksen M, Farabollini F, Guillette LJ, Hauser R, Heindel JJ, Ho S-M, Hunt PA, Iguchi T, Jobling S, Kanno J, Keri RA, Knudsen KE, Laufer H, LeBlanc GA, Marcus M, McLachlan JA, Myers JP, Nadal A, Newbold RR, Olea N, Prins GS, Richter CA, Rubin BS, Sonnenschein C, Soto AM, Talsness CE, Vandenberg JG, Vandenberg LN, Walser-Kuntz DR, Watson CS, Welshons WV, Wetherill Y, Zoeller RT, Chapel Hill bisphenol A expert panel consensus statement: Integration of mechanisms, effects in animals and potential to impact human health at current levels of exposure, *Reprod Toxicol* 24(2) (2007) 131–138. [PubMed: 17768031]
- [16]. Dodge LE, Williams PL, Williams MA, Missmer SA, Toth TL, Calafat AM, Hauser R, Paternal Urinary Concentrations of Parabens and Other Phenols in Relation to Reproductive Outcomes among Couples from a Fertility Clinic, *Environ Health Persp* 123(7) (2015) 665–671.
- [17]. Goldstone AE, Chen Z, Perry MJ, Kannan K, Louis GMB, Urinary bisphenol A and semen quality, the LIFE Study, *Reprod Toxicol* 51 (2015) 7–13. [PubMed: 25462789]
- [18]. Lassen TH, Frederiksen H, Jensen TK, Petersen JH, Joensen UN, Main KM, Skakkebaek NE, Juul A, Jorgensen N, Andersson AM, Urinary Bisphenol A Levels in Young Men: Association with Reproductive Hormones and Semen Quality, *Environ Health Persp* 122(5) (2014) 478–484.
- [19]. Lee SH, Kang SM, Choi MH, Lee J, Park MJ, Kim SH, Lee WY, Hong J, Chung BC, Changes in steroid metabolism among girls with precocious puberty may not be associated with urinary levels of bisphenol A, *Reprod Toxicol* 44 (2014) 1–6. [PubMed: 23557689]
- [20]. Scinicariello F, Buser MC, Serum Testosterone Concentrations and Urinary Bisphenol A, Benzophenone-3, Triclosan, and Paraben Levels in Male and Female Children and Adolescents: NHANES 2011–2012, *Environ Health Perspect* 124(12) (2016) 1898–1904. [PubMed: 27383665]
- [21]. Barbonetti A, Castellini C, Di Giammarco N, Santilli G, Francavilla S, Francavilla F, In vitro exposure of human spermatozoa to bisphenol A induces pro-oxidative/apoptotic mitochondrial dysfunction, *Reprod Toxicol* 66 (2016) 61–67. [PubMed: 27686954]
- [22]. Berger A, Ziv-Gal A, Cudiamat J, Wang W, Zhou CQ, Flaws JA, The effects of in utero bisphenol A exposure on the ovaries in multiple generations of mice, *Reprod Toxicol* 60 (2016) 39–52. [PubMed: 26746108]
- [23]. Delclos KB, Camacho L, Lewis SM, Vanlandingham MM, Latendresse JR, Olson GR, Davis KJ, Patton RE, da Costa GG, Woodling KA, Bryant MS, Chidambaram M, Trbojevich R, Juliar BE, Felton RP, Thorn BT, Toxicity Evaluation of Bisphenol A Administered by Gavage to Sprague Dawley Rats From Gestation Day 6 Through Postnatal Day 90, *Toxicol Sci* 139(1) (2014) 174–197. [PubMed: 24496637]

- [24]. Ferris J, Mahboubi K, MacLusky N, King WA, Favetta LA, BPA exposure during in vitro oocyte maturation results in dose-dependent alterations to embryo development rates, apoptosis rate, sex ratio and gene expression, *Reprod Toxicol* 59 (2016) 128–38. [PubMed: 26686065]
- [25]. Moore-Ambriz TR, Acuna-Hernandez DG, Ramos-Robles B, Sanchez-Gutierrez M, Santacruz-Marquez R, Sierra-Santoyo A, Pina-Guzman B, Shibayama M, Hernandez-Ochoa I, Exposure to bisphenol A in young adult mice does not alter ovulation but does alter the fertilization ability of oocytes, *Toxicol Appl Pharm* 289(3) (2015) 507–514.
- [26]. Qi SQ, Fu WJ, Wang CM, Liu CJ, Quan C, Kourouma A, Yan MS, Yu TT, Duan P, Yang KD, BPA-induced apoptosis of rat Sertoli cells through Fas/FasL and JNKs/p38 MAPK pathways, *Reprod Toxicol* 50 (2014) 108–116. [PubMed: 25461909]
- [27]. Vigezzi L, Bosquiazio VL, Kass L, Ramos JG, Munoz-De-Toro M, Luque EH, Developmental exposure to bisphenol A alters the differentiation and functional response of the adult rat uterus to estrogen treatment, *Reprod Toxicol* 52 (2015) 83–92. [PubMed: 25666754]
- [28]. Wang T, Han J, Duan X, Xiong B, Cui XS, Kim NH, Liu HL, Sun SC, The toxic effects and possible mechanisms of Bisphenol A on oocyte maturation of porcine in vitro, *Oncotarget* 7(22) (2016) 32554–32565. [PubMed: 27086915]
- [29]. FDA., Indirect Food Additives: Adhesives and Components of Coatings, 2013, pp. 41840–41843.
- [30]. FDA., Indirect Food Additives: Polymers, 2012, pp. 41899–41902.
- [31]. Geens T, Roosens L, Neels H, Covaci A, Assessment of human exposure to Bisphenol-A, Triclosan and Tetrabromobisphenol-A through indoor dust intake in Belgium, *Chemosphere* 76(6) (2009) 755–760. [PubMed: 19535125]
- [32]. Liao CY, Liu F, Guo Y, Moon HB, Nakata H, Wu Q, Kannan K, Occurrence of Eight Bisphenol Analogues in Indoor Dust from the United States and Several Asian Countries: Implications for Human Exposure, *Environ Sci Technol* 46(16) (2012) 9138–9145. [PubMed: 22784190]
- [33]. Liao CY, Liu F, Alomirah H, Loi VD, Mohd MA, Moon HB, Nakata H, Kannan K, Bisphenol S in Urine from the United States and Seven Asian Countries: Occurrence and Human Exposures, *Environ Sci Technol* 46(12) (2012) 6860–6866. [PubMed: 22620267]
- [34]. Shi ZX, Jiao Y, Hu Y, Sun ZW, Zhou XQ, Feng JF, Li JG, Wu YN, Levels of tetrabromobisphenol A, hexabromocyclododecanes and polybrominated diphenyl ethers in human milk from the general population in Beijing, China, *Sci Total Environ* 452 (2013) 10–18. [PubMed: 23500394]
- [35]. Ye X, Wong LY, Kramer J, Zhou X, Jia T, Calafat AM, Urinary Concentrations of Bisphenol A and Three Other Bisphenols in Convenience Samples of U.S. Adults during 2000–2014, *Environ Sci Technol* 49(19) (2015) 11834–9. [PubMed: 26360019]
- [36]. Kitamura S, Suzuki T, Sanoh S, Kohta R, Jinno N, Sugihara K, Yoshihara S, Fujimoto N, Watanabe H, Ohta S, Comparative study of the endocrine-disrupting activity of bisphenol A and 19 related compounds, *Toxicol Sci* 84(2) (2005) 249–259. [PubMed: 15635150]
- [37]. Stossi F, Bolt MJ, Ashcroft FJ, Lamerdin JE, Melnick JS, Powell RT, Dandekar RD, Mancini MG, Walker CL, Westwick JK, Mancini MA, Defining Estrogenic Mechanisms of Bisphenol A Analogs through High Throughput Microscopy-Based Contextual Assays, *Chem Biol* 21(6) (2014) 743–753. [PubMed: 24856822]
- [38]. Lawson C, Gieske M, Murdoch B, Ye P, Li Y, Hassold T, Hunt PA, Gene Expression in the Fetal Mouse Ovary Is Altered by Exposure to Low Doses of Bisphenol A, *SOCIETY FOR THE STUDY OF REPRODUCTION*, United States, 2011, p. 79.
- [39]. Zhang X-F, Zhang L-J, Feng Y-N, Chen B, Feng Y-M, Liang G-J, Li L, Shen W, Bisphenol A exposure modifies DNA methylation of imprint genes in mouse fetal germ cells, *Molecular Biology Reports* (9) (2012) 8621. [PubMed: 22699882]
- [40]. Rivera OE, Varayoud J, Rodríguez HA, Muñoz-de-Toro M, Luque EH, Neonatal exposure to bisphenol A or diethylstilbestrol alters the ovarian follicular dynamics in the lamb, *Reprod Toxicol* 32 (2011) 304–312. [PubMed: 21722727]
- [41]. Patricia AH, Crystal L, Mary G, Brenda M, Helen S, Alyssa M, Terry H, Catherine AV, Bisphenol A alters early oogenesis and follicle formation in the fetal ovary of the rhesus monkey, *Proceedings of the National Academy of Sciences of the United States of America* (43) (2012) 17525. [PubMed: 23012422]

- [42]. Eichenlaub-Ritter U, Vogt E, Cukurcam S, Sun F, Pacchierotti F, Parry J, Exposure of mouse oocytes to bisphenol A causes meiotic arrest but not aneuploidy, *Mut.Res.-Genetic Toxicology and Environmental Mutagenesis* 651 (2008) 82–92.
- [43]. Lenie S, Cortvrindt R, Eichenlaub-Ritter U, Smits J, Continuous exposure to bisphenol A during in vitro follicular development induces meiotic abnormalities, *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 651(1) (2008) 71–81.
- [44]. Ferris J, Favetta LA, King WA, Bisphenol A Exposure during Oocyte Maturation in vitro Results in Spindle Abnormalities and Chromosome Misalignment in *Bos taurus*, *Cytogenet Genome Res* 145(1) (2015) 50–58. [PubMed: 25871885]
- [45]. Nakano K, Nishio M, Kobayashi N, Hiradate Y, Hoshino Y, Sato E, Tanemura K, Comparison of the effects of BPA and BPAF on oocyte spindle assembly and polar body release in mice, *Zygote* 24(2) (2016) 172–180. [PubMed: 25925194]
- [46]. Bloom MS, Kim D, vom Saal FS, Taylor JA, Cheng G, Lamb JD, Fujimoto VY, Bisphenol A exposure reduces the estradiol response to gonadotropin stimulation during in vitro fertilization, *Fertility and Sterility* 96(3) (2011) 672–677.e2. [PubMed: 21813122]
- [47]. Shelley E, Paige LW, Stacey AM, Jodi AF, Katharine FB, Antonia MC, Xiaoyun Y, John CP, Diane W, Russ H, Urinary Bisphenol A Concentrations and Implantation Failure among Women Undergoing in Vitro Fertilization, *Environ Health Persp* (7) (2012) 978.
- [48]. Mok-Lin E, Ehrlich S, Williams PL, Petrozza J, Wright DL, Calafat AM, Ye X, Hauser R, Urinary bisphenol A concentrations and ovarian response among women undergoing IVF, *International Journal Of Andrology* 33(2) (2010) 385–393. [PubMed: 20002217]
- [49]. Tamara G, Riccardo C, Jack G, Luigi F, Stefania B, Anna Maria C, Cathryn M, Paul M, David M, Daily Bisphenol A Excretion and Associations with Sex Hormone Concentrations: Results from the InCHIANTI Adult Population Study, *Environ Health Persp* (11) (2010) 1603.
- [50]. Akin L, Kendirci M, Narin F, Kurtoglu S, Saraymen R, Kondolot M, Kocak S, Elmali F, The endocrine disruptor bisphenol A may play a role in the aetiopathogenesis of polycystic ovary syndrome in adolescent girls, *Acta Paediatr* 104(4) (2015) E171–E177. [PubMed: 25469562]
- [51]. Miao MH, Yuan W, Yang F, Liang H, Zhou ZJ, Li RS, Gao ES, Li DK, Associations between Bisphenol A Exposure and Reproductive Hormones among Female Workers, *Int J Env Res Pub He* 12(10) (2015) 13240–13250.
- [52]. Minguez-Alarcon L, Gaskins AJ, Chiu YH, Williams PL, Ehrlich S, Chavarro JE, Petrozza JC, Ford JB, Calafat AM, Hauser R, Team ES, Urinary bisphenol A concentrations and association with in vitro fertilization outcomes among women from a fertility clinic, *Hum Reprod* 30(9) (2015) 2120–2128. [PubMed: 26209788]
- [53]. Zhou W, Fang F, Zhu W, Chen Z-J, Du Y, Zhang J, Bisphenol A and Ovarian Reserve among Infertile Women with Polycystic Ovarian Syndrome, *Int J Env Res Pub He* 14(1) (2016) 18.
- [54]. Marina F, Nadia B, Victoria L-L, Carlos L, Neonatal Exposure to Bisphenol A and Reproductive and Endocrine Alterations Resembling the Polycystic Ovarian Syndrome in Adult Rats, *Environ Health Persp* (9) (2010) 1217.
- [55]. Tan W, Huang H, Wang Y, Wong TY, Wang CC, Leung LK, Bisphenol A differentially activates protein kinase C isoforms in murine placental tissue, *Toxicol Appl Pharm* 269 (2013) 163–168.
- [56]. Kobayashi K, Kubota H, Ohtani K, Hojo R, Miyagawa M, Lack of effects for dietary exposure of bisphenol A during in utero and lactational periods on reproductive development in rat offspring, *Journal of Toxicological Sciences* 37(3) (2012) 565–573. [PubMed: 22687996]
- [57]. Mendoza-Rodríguez CA, García-Guzmán M, Baranda-Avila N, Morimoto S, Perrot-Appianat M, Cerbón M, Administration of bisphenol A to dams during perinatal period modifies molecular and morphological reproductive parameters of the offspring, *Reprod Toxicol* 31 (2011) 177–183. [PubMed: 21055461]
- [58]. Varayoud J, Ramos JG, Bosquiazzo VL, Lower M, Munoz-de-Toro M, Luque EH, Neonatal Exposure to Bisphenol A Alters Rat Uterine Implantation-Associated Gene Expression and Reduces the Number of Implantation Sites, *ENDOCRINE SOCIETY, United States*, 2011, p. 1101.

- [59]. Peretz J, Gupta RK, Singh J, Hernández-Ochoa I, Flaws JA, Bisphenol A Impairs Follicle Growth, Inhibits Steroidogenesis, and Downregulates Rate-Limiting Enzymes in the Estradiol Biosynthesis Pathway, *Toxicol Sci* 119(1) (2011) 209–217. [PubMed: 20956811]
- [60]. Ziv-Gal A, Craig ZR, Wang W, Flaws JA, Bisphenol A inhibits cultured mouse ovarian follicle growth partially via the aryl hydrocarbon receptor signaling pathway, *Reprod Toxicol* 42 (2013) 58–67. [PubMed: 23928317]
- [61]. Grasselli F, Baratta L, Baioni L, Bussolati S, Ramoni R, Grolli S, Basini G, Bisphenol A disrupts granulosa cell function, Elsevier Science B.V., Amsterdam., United States, 2010, p. 34.
- [62]. Li YC, Zhang WC, Liu J, Wang WX, Li H, Zhu JL, Weng SZ, Xiao SH, Wu TT, Prepubertal bisphenol A exposure interferes with ovarian follicle development and its relevant gene expression, *Reprod Toxicol* 44 (2014) 33–40. [PubMed: 24051130]
- [63]. Santamaria C, Durando M, de Toro MM, Luque EH, Rodriguez HA, Ovarian dysfunctions in adult female rat offspring born to mothers perinatally exposed to low doses of bisphenol A, *J Steroid Biochem* 158 (2016) 220–230.
- [64]. Gamez JM, Penalba R, Cardoso N, Bernasconi PS, Carbone S, Ponzo O, Pandolfi M, Scacchi P, Reynoso R, Exposure to a low dose of bisphenol A impairs pituitary-ovarian axis in prepubertal rats Effects on early folliculogenesis, *Environ Toxicol Phar* 39(1) (2015) 9–15.
- [65]. Sadowski RN, Park P, Neese SL, Ferguson DC, Schantz SL, Juraska JM, Effects of perinatal bisphenol A exposure during early development on radial arm maze behavior in adult male and female rats, *Neurotoxicol Teratol* 42 (2014) 17–24. [PubMed: 24440629]
- [66]. Mansur A, Adir M, Yerushalmi G, Hourvitz A, Gitman H, Yung Y, Orvieto R, Machtinger R, Does BPA alter steroid hormone synthesis in human granulosa cells in vitro?, *Hum Reprod* 31(7) (2016) 1562–1569. [PubMed: 27112698]
- [67]. Kandaraki E, Chatzigeorgiou A, Livadas S, Palioura E, Economou F, Koutsilieris M, Palimeri S, Panidis D, Diamanti-Kandarakis E, Endocrine disruptors and polycystic ovary syndrome (PCOS): elevated serum levels of bisphenol A in women with PCOS, *The Journal Of Clinical Endocrinology And Metabolism* 96(3) (2011) E480–E484. [PubMed: 21193545]
- [68]. Adewale HB, Jefferson WN, Newbold RR, Patisaul HB, Neonatal Bisphenol-A Exposure Alters Rat Reproductive Development and Ovarian Morphology Without Impairing Activation of Gonadotropin-Releasing Hormone Neurons1, *Biology of Reproduction* 81(4) (2009) 690–699. [PubMed: 19535786]
- [69]. Retha RN, Wendy NJ, Elizabeth P-B, Prenatal Exposure to Bisphenol A at Environmentally Relevant Doses Adversely Affects the Murine Female Reproductive Tract Later in Life, *Environ Health Persp* (6) (2009) 879.
- [70]. Vahedi M, Saeedi A, Poorbaghi SL, Sepehrimanesh M, Fattahi M, Metabolic and endocrine effects of bisphenol A exposure in market seller women with polycystic ovary syndrome, *Environ Sci Pollut Res Int* 23(23) (2016) 23546–23550. [PubMed: 27614642]
- [71]. Vagi SJ, Azziz-Baumgartner E, Sjodin A, Calafat AM, Dumesic D, Gonzalez L, Kato K, Silva MJ, Ye XY, Azziz R, Exploring the potential association between brominated diphenyl ethers, polychlorinated biphenyls, organochlorine pesticides, perfluorinated compounds, phthalates, and bisphenol a in polycystic ovary syndrome: a case-control study, *Bmc Endocr Disord* 14 (2014).
- [72]. Patisaul HB, Mabrey N, Adewale HB, Sullivan AW, Soy but not bisphenol A (BPA) induces hallmarks of polycystic ovary syndrome (PCOS) and related metabolic co-morbidities in rats, *Reprod Toxicol* 49 (2014) 209–218. [PubMed: 25242113]
- [73]. Zhou CQ, Wang W, Peretz J, Flaws JA, Bisphenol A exposure inhibits germ cell nest breakdown by reducing apoptosis in cultured neonatal mouse ovaries, *Reprod Toxicol* 57 (2015) 87–99. [PubMed: 26049153]
- [74]. Ganesan S, Keating AF, Bisphenol A-Induced Ovotoxicity Involves DNA Damage Induction to Which the Ovary Mounts a Protective Response Indicated by Increased Expression of Proteins Involved in DNA Repair and Xenobiotic Biotransformation, *Toxicol Sci* 152(1) (2016) 169–180. [PubMed: 27208089]
- [75]. Wolstenholme JT, Goldsby JA, Rissman EF, Transgenerational effects of prenatal bisphenol A on social recognition, *Horm Behav* 64(5) (2013) 833–9. [PubMed: 24100195]

- [76]. Cobellis L, Colacurci N, Trabucco E, Carpentiero C, Grumetto L, Measurement of bisphenol A and bisphenol B levels in human blood sera from healthy and endometriotic women, John Wiley & Sons, Ltd, Great Britain, 2009, p. 1186.
- [77]. Buck Louis GM, Peterson CM, Chen Z, Croughan M, Sundaram R, Stanford J, Varner MW, Kennedy A, Giudice L, Fujimoto VY, Bisphenol A and phthalates and endometriosis: the Endometriosis: Natural History, Diagnosis and Outcomes Study, Elsevier Science B.V., Amsterdam., United States, 2013, p. 162.
- [78]. Upton K, Sathyanarayana S, De Roos AJ, Koch HM, Scholes D, Holt VL, A population-based case-control study of urinary bisphenol A concentrations and risk of endometriosis, *Hum Reprod* 29(11) (2014) 2457–2464. [PubMed: 25205760]
- [79]. Simonelli A, Guadagni R, De Franciscis P, Colacurci N, Pieri M, Basilicata P, Pedata P, Lamberti M, Sannolo N, Miraglia N, Environmental and occupational exposure to bisphenol A and endometriosis: urinary and peritoneal fluid concentration levels, *Int Arch Occup Environ Health* 90(1) (2017) 49–61. [PubMed: 27718009]
- [80]. Signorile PG, Spugnini EP, Mita L, Mellone P, D'Avino A, Bianco M, Diano N, Caputo L, Rea F, Viceconte R, Portaccio M, Viggiano E, Citro G, Pierantoni R, Sica V, Vincenzi B, Mita DG, Baldi F, Baldi A, Pre-natal exposure of mice to bisphenol A elicits an endometriosis-like phenotype in female offspring, *General and Comparative Endocrinology* 168 (2010) 318–325. [PubMed: 20350546]
- [81]. Bromer JG, Zhou Y, Taylor MB, Doherty L, Taylor HS, Bisphenol-A exposure in utero leads to epigenetic alterations in the developmental programming of uterine estrogen response, *The FASEB Journal* 24(7) (2010) 2273–2280. [PubMed: 20181937]
- [82]. Hiyama M, Choi EK, Wakitani S, Tachibana T, Khan H, Kusakabe KT, Kiso Y, Bisphenol-A (BPA) Affects Reproductive Formation across Generations in Mice (NOTE), JAPANESE SOCIETY OF VETERINARY SCIENCE, Japan, 2011, p. 1211.
- [83]. Li Q, Davila J, Kannan A, Flaws JA, Bagchi MK, Bagchi IC, Chronic Exposure to Bisphenol A Affects Uterine Function During Early Pregnancy in Mice, *Endocrinology* 157(5) (2016) 1764–74. [PubMed: 27022677]
- [84]. Kendzierski JA, Belcher SM, Strain-specific induction of endometrial periglandular fibrosis in mice exposed during adulthood to the endocrine disrupting chemical bisphenol A, *Reprod Toxicol* 58 (2015) 119–130. [PubMed: 26307436]
- [85]. Calhoun KC, Padilla-Banks E, Jefferson WN, Liu LW, Gerrish KE, Young SL, Wood CE, Hunt PA, VandeVoort CA, Williams CJ, Bisphenol A Exposure Alters Developmental Gene Expression in the Fetal Rhesus Macaque Uterus, *Plos One* 9(1) (2014).
- [86]. Salleh N, Giribabu N, Feng AOM, Myint K, Bisphenol A, Dichlorodiphenyltrichloroethane (DDT) and Vinclozolin Affect ex-vivo Uterine Contraction in Rats via Uterotonin (Prostaglandin F2 alpha, Acetylcholine and Oxytocin) Related Pathways, *Int J Med Sci* 12(11) (2015) 914–925. [PubMed: 26640411]
- [87]. Kim HR, Kim YS, Yoon JA, Lyu SW, Shin H, Lim HJ, Hong SH, Lee DR, Song H, *Egr1* is rapidly and transiently induced by estrogen and bisphenol A via activation of nuclear estrogen receptor-dependent ERK1/2 pathway in the uterus, *Reprod Toxicol* 50 (2014) 60–67. [PubMed: 25461906]
- [88]. Aghajanova L, Giudice LC, Effect of bisphenol A on human endometrial stromal fibroblasts in vitro, *Reproductive BioMedicine Online* 22(3) (2011) 249–256. [PubMed: 21273127]
- [89]. An BS, Ahn HJ, Kang HS, Jung EM, Yang H, Hong EJ, Jeung EB, Effects of estrogen and estrogenic compounds, 4-tert-octylphenol, and bisphenol A on the uterine contraction and contraction-associated proteins in rats, Elsevier Science B.V., Amsterdam., Netherlands, 2013, p. 27.
- [90]. Kang SY, Song JY, Cho HH, Gene expression analysis of uterine smooth muscle cells exposed to bisphenol A, *Toxicology and Environmental Health Sciences* 6(4) (2014) 261–267.
- [91]. Helgestam M, Davey E, Stavreus-Evers AL, Olovsson M, Bisphenol A affects human endometrial endothelial cell angiogenic activity in vitro, *Reprod Toxicol* 46 (2014) 69–76. [PubMed: 24632125]

- [92]. Shen Y, Ren ML, Feng X, Cai YL, Gao YX, Xu Q, An evidence in vitro for the influence of bisphenol A on uterine leiomyoma, *Eur J Obstet Gyn R B* 178 (2014) 80–83.
- [93]. Wang KH, Kao AP, Chang CC, Lin TC, Kuo TC, Bisphenol A-induced epithelial to mesenchymal transition is mediated by cyclooxygenase-2 up-regulation in human endometrial carcinoma cells, *Reprod Toxicol* 58 (2015) 229–233. [PubMed: 26546977]
- [94]. Li D-K, Zhou Z, Miao M, He Y, Wang J, Ferber J, Herrinton LJ, Gao E, Yuan W, Urine bisphenol-A (BPA) level in relation to semen quality, *Fertility and Sterility* 95(2) 625–630.e4. [PubMed: 21035116]
- [95]. Meeker JD, Calafat AM, Hauser R, Urinary Bisphenol A Concentrations in Relation to Serum Thyroid and Reproductive Hormone Levels in Men from an Infertility Clinic, *ACS AMERICAN CHEMICAL SOCIETY, United States*, 2010, p. 1458.
- [96]. Jaime M, Niels J, Anna-Maria A, Antonia MC, Xiaoyun Y, Redmon JB, Erma ZD, Christina W, Amy S, Sally WT, Fan L, Shanna HS, Are Environmental Levels of Bisphenol A Associated with Reproductive Function in Fertile Men?, *Environ Health Persp* (9) (2010) 1286.
- [97]. Knez J, Kranvogel R, Breznic BP, Voncina E, Vlasisavljevic V, Are urinary bisphenol A levels in men related to semen quality and embryo development after medically assisted reproduction?, *Fertility and Sterility* 101(1) (2014) 215–22. [PubMed: 24182411]
- [98]. Okada A, Kai O, Effects of estradiol-17beta and bisphenol A administered chronically to mice throughout pregnancy and lactation on the male pups' reproductive system, *Asian Journal Of Andrology* 10(2) (2008) 271–276. [PubMed: 18286211]
- [99]. Salian S, Doshi T, Vanage G, Perinatal exposure of rats to Bisphenol A affects the fertility of male offspring, *Life Sciences* 85 (2009) 742–752. [PubMed: 19837096]
- [100]. Minamiyama Y, Ichikawa H, Takemura S, Kusunoki H, Naito Y, Yoshikawa T, Generation of reactive oxygen species in sperms of rats as an earlier marker for evaluating the toxicity of endocrine-disrupting chemicals, *Free Radical Research* 44(12) (2010) 1398–1406. [PubMed: 20815788]
- [101]. Gurmeet KSS, Rosnah I, Normadiah MK, Das S, Mustafa AM, Detrimental Effects of Bisphenol a on Development and Functions of the Male Reproductive System in Experimental Rats, *Excli J* 13 (2014) 151–160. [PubMed: 26417249]
- [102]. Vilela J, Hartmann A, Silva EF, Cardoso T, Corcini CD, Varela AS, Martinez PE, Colares EP, Sperm impairments in adult vesper mice (*Calomys laucha*) caused by in utero exposure to bisphenol A, *Andrologia* 46(9) (2014) 971–978. [PubMed: 24147964]
- [103]. LI Y.-h., Fei D, Fen Y, ZHOU X.-y., PAN H.-j., Yang L, Runsheng L, Pubertal exposure to bisphenol A affects the reproduction of male mice and sex ratio of offspring, *Journal of Reproduction and Contraception* 26(1) (2015) 14–21.
- [104]. Wisniewski P, Romano RM, Kizys MML, Oliveira KC, Kasamatsu T, Giannocco G, Chiamolera MI, Dias-Da-Silva MR, Romano MA, Adult exposure to bisphenol A (BPA) in Wistar rats reduces sperm quality with disruption of the hypothalamic-pituitary-testicular axis, *Toxicology* 329 (2015) 1–9. [PubMed: 25575453]
- [105]. Hass U, Christiansen S, Boberg J, Rasmussen MG, Mandrup K, Axelstad M, Low-dose effect of developmental bisphenol A exposure on sperm count and behaviour in rats, *Andrology-Us* 4(4) (2016) 594–607.
- [106]. Kalb AC, Kalb AL, Cardoso TF, Fernandes CG, Corcini CD, Varela AS, Martinez PE, Maternal Transfer of Bisphenol A During Nursing Causes Sperm Impairment in Male Offspring, *Arch Environ Con Tox* 70(4) (2016) 793–801.
- [107]. Rahman MS, Kwon WS, Karmakar PC, Yoon SJ, Ryu BY, Pang MG, Gestational Exposure to Bisphenol-A Affects the Function and Proteome Profile of F1 Spermatozoa in Adult Mice, *Environ Health Perspect* (2016).
- [108]. Wang HF, Liu M, Li N, Luo T, Zheng LP, Zeng XH, Bisphenol A Impairs Mature Sperm Functions by a CatSper-Relevant Mechanism, *Toxicol Sci* 152(1) (2016) 145–54. [PubMed: 27125968]
- [109]. Quan C, Wang C, Duan P, Huang W, Chen W, Tang S, Yang K, Bisphenol a induces autophagy and apoptosis concurrently involving the Akt/mTOR pathway in testes of pubertal SD rats, *Environ Toxicol* (2016).

- [110]. Quan C, Wang C, Duan P, Huang W, Yang K, Prenatal bisphenol a exposure leads to reproductive hazards on male offspring via the Akt/mTOR and mitochondrial apoptosis pathways, *Environ Toxicol* (2016).
- [111]. Yin L, Dai Y, Cui Z, Jiang X, Liu W, Han F, Lin A, Cao J, Liu J, The regulation of cellular apoptosis by the ROS-triggered PERK/EIF2 α /chop pathway plays a vital role in bisphenol A-induced male reproductive toxicity, *Toxicol Appl Pharmacol* 314 (2016) 98–108. [PubMed: 27894913]
- [112]. Wang P, Luo CH, Li QY, Chen S, Hue Y, Mitochondrion-mediated apoptosis is involved in reproductive damage caused by BPA in male rats, *Environ Toxicol Phar* 38(3) (2014) 1025–1033.
- [113]. Liu C, Duan WX, Zhang L, Xu SC, Li RY, Chen CH, He MD, Lu YH, Wu HJ, Yu ZP, Zhou Z, Bisphenol A exposure at an environmentally relevant dose induces meiotic abnormalities in adult male rats, *Cell Tissue Res* 355(1) (2014) 223–232. [PubMed: 24085620]
- [114]. Chen Z, Zuo X, He D, Ding S, Xu F, Yang H, Jin X, Fan Y, Ying L, Tian C, Ying C, Long-term exposure to a ‘safe’ dose of bisphenol A reduced protein acetylation in adult rat testes, *Sci Rep* 7 (2017) 40337. [PubMed: 28067316]
- [115]. Vrooman LA, Oatley JM, Griswold JE, Hassold TJ, Hunt PA, Estrogenic Exposure Alters the Spermatogonial Stem Cells in the Developing Testis, Permanently Reducing Crossover Levels in the Adult, *Plos Genet* 11(1) (2015).
- [116]. Mao ZX, Xia W, Chang HL, Huo WQ, Li YY, Xu SQ, Paternal BPA exposure in early life alters Igf2 epigenetic status in sperm and induces pancreatic impairment in rat offspring, *Toxicol Lett* 238(3) (2015) 30–38. [PubMed: 26276081]
- [117]. Qian WY, Wang YX, Zhu JY, Mao CF, Wang Q, Huan F, Cheng J, Liu YQ, Wang J, Xiao H, The toxic effects of Bisphenol A on the mouse spermatocyte GC-2 cell line: the role of the Ca²⁺-calmodulin-Ca²⁺/calmodulin-dependent protein kinase II axis, *J Appl Toxicol* 35(11) (2015) 1271–1277. [PubMed: 26096086]
- [118]. Rahman MS, Kwon WS, Lee JS, Yoon SJ, Ryu BY, Pang MG, Bisphenol-A affects male fertility via fertility-related proteins in spermatozoa, *Sci Rep* 5 (2015) 9169. [PubMed: 25772901]
- [119]. Wan X, Ru Y, Chu C, Ni Z, Zhou Y, Wang S, Zhou Z, Zhang Y, Bisphenol A accelerates capacitation-associated protein tyrosine phosphorylation of rat sperm by activating protein kinase A, *Acta Biochim Biophys Sin (Shanghai)* 48(6) (2016) 573–80. [PubMed: 27174873]
- [120]. Liang S, Yin L, Shengyang Yu K, Hofmann MC, Yu X, High-Content Analysis Provides Mechanistic Insights into the Testicular Toxicity of Bisphenol A and Selected Analogues in Mouse Spermatogonial Cells, *Toxicol Sci* 155(1) (2017) 43–60. [PubMed: 27633978]
- [121]. Barbonetti A, Castellini C, Di Giammarco N, Santilli G, Francavilla S, Francavilla F, In vitro exposure of human spermatozoa to bisphenol A induces pro-oxidative/apoptotic mitochondrial dysfunction, *Reprod Toxicol* 66 (2016) 61–67. [PubMed: 27686954]
- [122]. Kotwicka M, Skibinska I, Piworun N, Jendraszak M, Chmielewska M, Jedrzejczak P, Bisphenol A modifies human spermatozoa motility in vitro, *Journal of Medical Science* 85(1) (2016) 39–45.
- [123]. Haywood M, Spaliviero J, Jimenez M, King NJ, Handelsman DJ, Allan CM, Sertoli and germ cell development in hypogonadal (hpg) mice expressing transgenic follicle-stimulating hormone alone or in combination with testosterone, *Endocrinology* 144(2) (2003) 509–17. [PubMed: 12538611]
- [124]. Lui WY, Mruk D, Lee WM, Cheng CY, Sertoli cell tight junction dynamics: Their regulation during spermatogenesis, *Biology of Reproduction* 68(4) (2003) 1087–1097. [PubMed: 12606453]
- [125]. Payne AH, Hormonal-Regulation of Cytochrome-P450 Enzymes, Cholesterol Side-Chain Cleavage and 17-Alpha-Hydroxylase C17–20 Lyase in Leydig-Cells, *Biology of Reproduction* 42(3) (1990) 399–404. [PubMed: 2160293]
- [126]. Nakamura D, Yanagiba Y, Duan Z, Ito Y, Okamura A, Asaeda N, Tagawa Y, Li C, Taya K, Zhang S-Y, Naito H, Ramdhan DH, Kamijima M, Nakajima T, Bisphenol A may cause testosterone reduction by adversely affecting both testis and pituitary systems similar to estradiol, *Toxicol Lett* 194 (2010) 16–25. [PubMed: 20144698]

- [127]. Nanjappa MK, Simon L, Akingbemi BT, The Industrial Chemical Bisphenol A (BPA) Interferes with Proliferative Activity and Development of Steroidogenic Capacity in Rat Leydig Cells, *Biology of Reproduction* 86(5) (2012).
- [128]. Ge LC, Chen ZJ, Liu H, Zhang KS, Su Q, Ma XY, Huang HB, Zhao ZD, Wang YY, Giesy JP, Du J, Wang HS, Signaling related with biphasic effects of bisphenol A (BPA) on Sertoli cell proliferation: A comparative proteomic analysis, *Bba-Gen Subjects* 1840(9) (2014) 2663–2673.
- [129]. Ge LC, Chen ZJ, Liu HY, Zhang KS, Liu H, Huang HB, Zhang G, Wong CKC, Giesy JP, Du J, Wang HS, Involvement of activating ERK1/2 through G protein coupled receptor 30 and estrogen receptor alpha/beta in low doses of bisphenol A promoting growth of Sertoli TM4 cells, *Toxicol Lett* 226(1) (2014) 81–89. [PubMed: 24495410]
- [130]. Qian WY, Zhu JY, Mao CF, Liu JL, Wang YX, Wang Q, Liu YQ, Gao R, Xiao H, Wang J, Involvement of CaM-CaMKII-ERK in bisphenol A-induced Sertoli cell apoptosis, *Toxicology* 324 (2014) 27–34. [PubMed: 24905940]
- [131]. Wang C, Qi S, Liu C, Yang A, Fu W, Quan C, Duan P, Yu T, Yang K, Mitochondrial Dysfunction and Ca²⁺ Overload in Injured Sertoli Cells Exposed to Bisphenol A, *Environ Toxicol* (2016).
- [132]. de Freitas ATAG, Ribeiro MA, Pinho CF, Peixoto AR, Domeniconi RF, Scarano WR, Regulatory and junctional proteins of the blood-testis barrier in human Sertoli cells are modified by monobutyl phthalate (MBP) and bisphenol A (BPA) exposure, *Toxicol in Vitro* 34 (2016) 1–7. [PubMed: 26922907]
- [133]. Xiao X, Mruk DD, Tang EI, Wong CK, Lee WM, John CM, Turek PJ, Silvestrini B, Cheng CY, Environmental toxicants perturb human Sertoli cell adhesive function via changes in F-actin organization mediated by actin regulatory proteins, *Hum Reprod* 29(6) (2014) 1279–91. [PubMed: 24532171]
- [134]. Nakamura D, Yanagiba Y, Duan ZW, Ito Y, Okamura A, Asaeda N, Tagawa Y, Li CM, Taya K, Zhang SY, Naito H, Ramdhan DH, Kamijima M, Nakajima T, Bisphenol A may cause testosterone reduction by adversely affecting both testis and pituitary systems similar to estradiol, *Toxicol Lett* 194(1–2) (2010) 16–25. [PubMed: 20144698]
- [135]. Akingbemi BT, Sottas CM, Koulova AI, Klinefelter GR, Hardy MP, Inhibition of testicular steroidogenesis by the xenoestrogen bisphenol a is associated with reduced pituitary luteinizing hormone secretion and decreased steroidogenic enzyme gene expression in rat Leydig cells, *Endocrinology* 145(2) (2004) 592–603. [PubMed: 14605012]
- [136]. Chen ZJ, Zhang KS, Ge LC, Liu H, Chen LK, Du J, Wang HS, Signals involved in the effects of bisphenol A (BPA) on proliferation and motility of Leydig cells: a comparative proteomic analysis, *Toxicol Res-Uk* 5(6) (2016) 1573–1584.
- [137]. Meeker JD, Ehrlich S, Toth TL, Wright DL, Calafat AM, Trisini AT, Ye X, Hauser R, Semen quality and sperm DNA damage in relation to urinary bisphenol A among men from an infertility clinic, *Reprod Toxicol* 30 (2010) 532–539. [PubMed: 20656017]
- [138]. Liu XQ, Miao MH, Zhou ZJ, Gao ES, Chen JP, Wang JT, Sun F, Yuan W, Li DK, Exposure to bisphenol-A and reproductive hormones among male adults, *Environ Toxicol Phar* 39(2) (2015) 934–941.
- [139]. Zhuang WL, Wu KS, Wang YK, Zhu HJ, Deng ZZ, Peng L, Zhu GH, Association of Serum Bisphenol-A Concentration and Male Reproductive Function Among Exposed Workers, *Arch Environ Con Tox* 68(1) (2015) 38–45.
- [140]. Ferguson KK, Peterson KE, Lee JM, Mercado-Garcia A, Blank-Goldenberg C, Tellez-Rojo MM, Meeker JD, Prenatal and peripubertal phthalates and bisphenol A in relation to sex hormones and puberty in boys, *Reprod Toxicol* 47 (2014) 70–76. [PubMed: 24945889]
- [141]. Hines CJ, Jackson MV, Deddens JA, Clark JC, Ye X, Christianson AL, Meadows JW, Calafat AM, Urinary Bisphenol A (BPA) Concentrations among Workers in Industries that Manufacture and Use BPA in the USA, *Annals of Work Exposures and Health* 61(2) (2017) 164–182. [PubMed: 28395354]
- [142]. Heinälä M, Ylinen K, Tuomi T, Santonen T, Porras SP, Assessment of Occupational Exposure to Bisphenol A in Five Different Production Companies in Finland, *Annals of Work Exposures and Health* 61(1) (2017) 44–55. [PubMed: 28395312]

- [143]. Xi W, Lee CKF, Yeung WSB, Giesy JP, Wong MH, Zhang X, Hecker M, Wong CKC, Effect of perinatal and postnatal bisphenol A exposure to the regulatory circuits at the hypothalamus-pituitary-gonadal axis of CD-1 mice, *Reprod Toxicol* 4 (2011) 409. [PubMed: 21182934]
- [144]. LaRocca J, Boyajian A, Brown C, Smith SD, Hixon M, Effects of In Utero Exposure to Bisphenol A or Diethylstilbestrol on the Adult Male Reproductive System, *Birth defects research. Part B, Developmental and reproductive toxicology* 92(6) (2011) 526–533. [PubMed: 21922642]
- [145]. Salian S, Doshi T, Vanage G, Neonatal exposure of male rats to Bisphenol A impairs fertility and expression of sertoli cell junctional proteins in the testis, *Toxicology* 265 (2009) 56–67. [PubMed: 19782717]
- [146]. El-Beshbishy HA, Aly HAA, El-Shafey M, Lipoic acid mitigates bisphenol A-induced testicular mitochondrial toxicity in rats, *Toxicology & Industrial Health* 29(10) (2013) 875. [PubMed: 22623521]
- [147]. D’Cruz SC, Jubendradass R, Jayakanthan M, Rani SJA, Mathur PP, Bisphenol A impairs insulin signaling and glucose homeostasis and decreases steroidogenesis in rat testis: An in vivo and in silico study, *Food and Chemical Toxicology* (3–4) (2012) 1124. [PubMed: 22142692]
- [148]. Howdeshell KL, Furr J, Lambright CR, Wilson VS, Ryan BC, Gray LE, Gestational and Lactational Exposure to Ethinyl Estradiol, but not Bisphenol A, Decreases Androgen-Dependent Reproductive Organ Weights and Epididymal Sperm Abundance in the Male Long Evans Hooded Rat, Oxford University Press, Great Britain, 2008, p. 371.
- [149]. Qiu L-L, Wang X, Zhang X.-h., Zhang Z, Gu J, Liu L, Wang Y, Wang X, Wang S-L, Decreased androgen receptor expression may contribute to spermatogenesis failure in rats exposed to low concentration of bisphenol A, *Toxicol Lett* 219 (2013) 116–124. [PubMed: 23528252]
- [150]. Gamez JM, Penalba R, Cardoso N, Ponzo O, Carbone S, Pandolfi M, Scacchi P, Reynoso R, Low dose of bisphenol A impairs the reproductive axis of prepuberal male rats, *J Physiol Biochem* 70(1) (2014) 239–246. [PubMed: 24271643]
- [151]. Chouhan S, Yadav SK, Prakash J, Westfall S, Ghosh A, Agarwal NK, Singh SP, Increase in the expression of inducible nitric oxide synthase on exposure to bisphenol A: A possible cause for decline in steroidogenesis in male mice, *Environ Toxicol Phar* 39(1) (2015) 405–416.
- [152]. Zang Z, Ji S, Xia T, Huang S, Effects of Bisphenol A on Testosterone Levels and Sexual Behaviors of Male Mice, *Advances in Sexual Medicine* 6(04) (2016) 41.
- [153]. Eladak S, Grisin T, Moison D, Guerquin MJ, N’Tumba-Byn T, Pozzi-Gaudin S, Benachi A, Livera G, Rouiller-Fabre V, Habert R, A new chapter in the bisphenol A story: bisphenol S and bisphenol F are not safe alternatives to this compound, *Fertility and Sterility* 103(1) (2015) 11–21. [PubMed: 25475787]
- [154]. Liao CY, Liu F, Kannan K, Bisphenol S, a New Bisphenol Analogue, in Paper Products and Currency Bills and Its Association with Bisphenol A Residues, *Environ Sci Technol* 46(12) (2012) 6515–6522. [PubMed: 22591511]
- [155]. Liao CY, Kannan K, Concentrations and Profiles of Bisphenol A and Other Bisphenol Analogues in Foodstuffs from the United States and Their Implications for Human Exposure, *J Agr Food Chem* 61(19) (2013) 4655–4662. [PubMed: 23614805]
- [156]. Liao CY, Kannan K, A Survey of Alkylphenols, Bisphenols, and Triclosan in Personal Care Products from China and the United States, *Arch Environ Con Tox* 67(1) (2014) 50–59.
- [157]. Skledar DG, Schmidt J, Fic A, Klop i I, Trontelj J, Dolenc MS, Finel M, Maši LP, Influence of metabolism on endocrine activities of bisphenol S, *Chemosphere* 157(Supplement C) (2016) 152–159. [PubMed: 27213244]
- [158]. Dreier DA, Connors KA, Brooks BW, Comparative endpoint sensitivity of in vitro estrogen agonist assays, *Regul Toxicol Pharmacol* 72(2) (2015) 185–93. [PubMed: 25896097]
- [159]. Huang R, Sakamuru S, Martin MT, Reif DM, Judson RS, Houck KA, Casey W, Hsieh JH, Shockley KR, Ceger P, Fostel J, Witt KL, Tong W, Rotroff DM, Zhao T, Shinn P, Simeonov A, Dix DJ, Austin CP, Kavlock RJ, Tice RR, Xia M, Profiling of the Tox21 10K compound library for agonists and antagonists of the estrogen receptor alpha signaling pathway, *Sci Rep* 4 (2014) 5664. [PubMed: 25012808]

- [160]. Rochester JR, Bolden AL, Bisphenol S and F: A Systematic Review and Comparison of the Hormonal Activity of Bisphenol A Substitutes, *Environ Health Persp* 123(7) (2015) 643–650.
- [161]. Rosenmai AK, Dybdahl M, Pedersen M, Alice van Vugt-Lussenburg BM, Wedeby EB, Taxvig C, Vinggaard AM, Are structural analogues to bisphenol a safe alternatives?, *Toxicol Sci* 139(1) (2014) 35–47. [PubMed: 24563381]
- [162]. Yamasaki K, Noda S, Imatanaka N, Yakabe Y, Comparative study of the uterotrophic potency of 14 chemicals in a uterotrophic assay and their receptor-binding affinity, *Toxicol Lett* 146(2) (2004) 111–120. [PubMed: 14643963]
- [163]. Ullah H, Jahan S, Ul Ain Q, Shaheen G, Ahsan N, Effect of bisphenol S exposure on male reproductive system of rats: A histological and biochemical study, *Chemosphere* 152 (2016) 383–391. [PubMed: 26994432]
- [164]. Ji K, Hong S, Kho Y, Choi K, Effects of Bisphenol S Exposure on Endocrine Functions and Reproduction of Zebrafish, *Environ Sci Technol* 47(15) (2013) 8793–8800. [PubMed: 23806087]
- [165]. Naderi M, Wong MYL, Gholami F, Developmental exposure of zebrafish (*Danio rerio*) to bisphenol-S impairs subsequent reproduction potential and hormonal balance in adults, *Aquat Toxicol* 148 (2014) 195–203. [PubMed: 24508763]
- [166]. Chen YC, Shu L, Qiu ZQ, Lee DY, Settle SJ, Hee SQ, Telesca D, Yang X, Allard P, Exposure to the BPA-Substitute Bisphenol S Causes Unique Alterations of Germline Function, *Plos Genet* 12(7) (2016).
- [167]. Roelofs MJE, van den Berg M, Bovee TFH, Piersma AH, van Duursen MBM, Structural bisphenol analogues differentially target steroidogenesis in murine MA-10 Leydig cells as well as the glucocorticoid receptor, *Toxicology* 329 (2015) 10–20. [PubMed: 25576683]
- [168]. Cabaton N, Chagnon M-C, Lhuguenot J-C, Cravedi J-P, Zalko D, Disposition and Metabolic Profiling of Bisphenol F in Pregnant and Nonpregnant Rats, *J Agr Food Chem* 54(26) (2006) 10307–10314. [PubMed: 17177575]
- [169]. Ye X, Wong LY, Kramer J, Zhou X, Jia T, Calafat AM, Urinary Concentrations of Bisphenol A and Three Other Bisphenols in Convenience Samples of U.S. Adults during 2000–2014, *Environ Sci Technol* 49(19) (2015) 11834–9. [PubMed: 26360019]
- [170]. Stroheker T, Chagnon MC, Pinnert MF, Berges R, Canivenc-Lavier MC, Estrogenic effects of food wrap packaging xenoestrogens and flavonoids in female Wistar rats: a comparative study, *Reprod Toxicol* 17(4) (2003) 421–432. [PubMed: 12849853]
- [171]. Higashihara N, Shiraishi K, Miyata K, Oshima Y, Minobe Y, Yamasaki K, Subacute oral toxicity study of bisphenol F based on the draft protocol for the “Enhanced OECD Test Guideline no. 407”, *Arch Toxicol* 81(12) (2007) 825–832. [PubMed: 17628788]
- [172]. Li M, Yang Y, Yang Y, Yin J, Zhang J, Feng Y, Shao B, Biotransformation of Bisphenol AF to Its Major Glucuronide Metabolite Reduces Estrogenic Activity, *Plos One* 8(12) (2013) e83170. [PubMed: 24349450]
- [173]. Yang Y, Yin J, Yang Y, Zhou N, Zhang J, Shao B, Wu Y, Determination of bisphenol AF (BPAF) in tissues, serum, urine and feces of orally dosed rats by ultra-high-pressure liquid chromatography–electrospray tandem mass spectrometry, *Journal of Chromatography B* 901 (2012) 93–97.
- [174]. Asimakopoulos AG, Xue JC, De Carvalho BP, Iyer A, Abualnaja KO, Yaghmoor SS, Kumosani TA, Kannan K, Urinary biomarkers of exposure to 57 xenobiotics and its association with oxidative stress in a population in Jeddah, Saudi Arabia, *Environ Res* 150 (2016) 573–581. [PubMed: 26654562]
- [175]. Yang YJ, Guan J, Yin J, Shao B, Li H, Urinary levels of bisphenol analogues in residents living near a manufacturing plant in south China, *Chemosphere* 112 (2014) 481–486. [PubMed: 25048943]
- [176]. Matsushima A, Liu XH, Okada H, Shimohigashi M, Shimohigashi Y, Bisphenol AF Is a Full Agonist for the Estrogen Receptor ER alpha but a Highly Specific Antagonist for ER beta, *Environ Health Persp* 118(9) (2010) 1267–1272.
- [177]. Shi JC, Jiao ZH, Zheng S, Li M, Zhang J, Feng YX, Yin J, Shao B, Long-term effects of Bisphenol AF (BPAF) on hormonal balance and genes of hypothalamus-pituitary-gonad axis and

- liver of zebrafish (*Danio rerio*), and the impact on offspring, *Chemosphere* 128 (2015) 252–257. [PubMed: 25723718]
- [178]. Yang XX, Liu YC, Li J, Chen MJ, Peng D, Liang Y, Song MY, Zhang J, Jiang GB, Exposure to Bisphenol AF disrupts sex hormone levels and vitellogenin expression in zebrafish, *Environmental Toxicology* 31(3) (2016) 285–294. [PubMed: 25213402]
- [179]. Furr JR, Lambright CS, Wilson VS, Foster PM, Gray LE, A Short-term In Vivo Screen Using Fetal Testosterone Production, a Key Event in the Phthalate Adverse Outcome Pathway, to Predict Disruption of Sexual Differentiation, *Toxicol Sci* 140(2) (2014) 403–424. [PubMed: 24798384]
- [180]. Li J, Sheng N, Cui RN, Feng YX, Shao B, Guo XJ, Zhang HX, Dai JY, Gestational and lactational exposure to bisphenol AF in maternal rats increases testosterone levels in 23-day-old male offspring, *Chemosphere* 163 (2016) 552–561. [PubMed: 27567155]
- [181]. Feng YX, Yin J, Jiao ZH, Shi JC, Li M, Shao B, Bisphenol AF may cause testosterone reduction by directly affecting testis function in adult male rats, *Toxicol Lett* 211(2) (2012) 201–209. [PubMed: 22504055]
- [182]. Sutherland V., McIntyre B., Pelch K., Waidyanatha S., Conley JM., Gray LE., Foster PM, A Comparison of In Vivo Reproductive and Developmental Toxicity (DART) Endpoints for Bisphenol AF and Bisphenol A, the Society of Toxicology 56th Annual Meeting, Baltimore, Maryland, 2017.
- [183]. Driffield M, Harmer N, Bradley E, Fernandes AR, Rose M, Mortimer D, Dicks P, Determination of brominated flame retardants in food by LC-MS/MS: diastereoisomer-specific hexabromocyclododecane and tetrabromobisphenol A, *Food Addit Contam A* 25(7) (2008) 895–903.
- [184]. Lai DY, Kacew S, Dekant W, Tetrabromobisphenol A (TBBPA): Possible modes of action of toxicity and carcinogenicity in rodents, *Food and Chemical Toxicology* 80 (2015) 206–214. [PubMed: 25818463]
- [185]. Dallaire R, Ayotte P, Pereg D, Dery S, Dumas P, Langlois E, Dewailly E, Determinants of Plasma Concentrations of Perfluorooctanesulfonate and Brominated Organic Compounds in Nunavik Inuit Adults (Canada), *Environ Sci Technol* 43(13) (2009) 5130–5136. [PubMed: 19673318]
- [186]. Carignan CC, Abdallah MAE, Wu N, Heiger-Bernays W, McClean MD, Harrad S, Webster TF, Predictors of Tetrabromobisphenol-A (TBBP-A) and Hexabromocyclododecanes (HBCD) in Milk from Boston Mothers, *Environ Sci Technol* 46(21) (2012) 12146–12153. [PubMed: 22998345]
- [187]. Molina-Molina JM, Amaya E, Grimaldi M, Saenz JM, Real M, Fernandez MF, Balaguer P, Olea N, In vitro study on the agonistic and antagonistic activities of bisphenol-S and other bisphenol-A congeners and derivatives via nuclear receptors, *Toxicol Appl Pharm* 272(1) (2013) 127–136.
- [188]. Van der Ven LTM, de Kuil TV, Verhoef A, Verwer CM, Lilienthal H, Leonards PEG, Schauer UMD, Canton RF, Litens S, De Jong FH, Visser TJ, Dekant W, Stern N, Hakansson H, Slob W, Van den Berg M, Vos JG, Piersma AH, Endocrine effects of tetrabromobisphenol-A (TBBPA) in Wistar rats as tested in a one-generation reproduction study and a subacute toxicity study, *Toxicology* 245(1–2) (2008) 76–89. [PubMed: 18255212]
- [189]. ECB., European union risk assessment report—2,2',6,6'-tetrabromo-4,4'-isopropylidenediphenol (tetrabromobisphenol-A or TBBP-A) (CAS: 79–94-7) Part II—human health, vol 63, EUR 22161 EN. Institute for Health and Consumer Protection, European Chemicals Bureau, European Commission Joint Research Centre, 4th Priority List, Luxembourg: Office for Official Publications of the European Communities, in: I.f.H.a.C. Protection (Ed.) 2012, pp. 1–157.
- [190]. Cope RB, Kacew S, Dourson M, A reproductive, developmental and neurobehavioral study following oral exposure of tetrabromobisphenol A on Sprague-Dawley rats, *Toxicology* 329 (2015) 49–59. [PubMed: 25523853]
- [191]. Zatecka E, Ded L, Elzeinova F, Kubatova A, Dorosh A, Margaryan H, Dostalova P, Peknicova J, Effect of tetrabromobisphenol A on induction of apoptosis in the testes and changes in expression of selected testicular genes in CD1 mice, *Reprod Toxicol* 35 (2013) 32–9. [PubMed: 22677475]

- [192]. Zatecka E, Castillo J, Elzeinova F, Kubatova A, Ded L, Peknicova J, Oliva R, The effect of tetrabromobisphenol A on protamine content and DNA integrity in mouse spermatozoa, *Andrology-Us* 2(6) (2014) 910–917.
- [193]. Dunnick JK, Sanders JM, Kissling GE, Johnson CL, Boyle MH, Elmore SA, Environmental Chemical Exposure May Contribute to Uterine Cancer Development: Studies with Tetrabromobisphenol A, *Toxicol Pathol* 43(4) (2015) 464–473. [PubMed: 25476797]
- [194]. Kuiper RV, van den Brandhof EJ, Leonards PEG, van der Ven LTM, Wester PW, Vos JG, Toxicity of tetrabromobisphenol A (TBBPA) in zebrafish (*Danio rerio*) in a partial life-cycle test, *Arch Toxicol* 81(1) (2007) 1–9. [PubMed: 16738895]
- [195]. Tada Y, Fujitani T, Yano N, Takahashi H, Yuzawa K, Ando H, Kubo Y, Nagasawa A, Ogata A, Kamimura H, Effects of tetrabromobisphenol A, brominated flame retardant, in ICR mice after prenatal and postnatal exposure, *Food and Chemical Toxicology* 44(8) (2006) 1408–1413. [PubMed: 16716481]
- [196]. Dankers AC, Roelofs MJ, Piersma AH, Sweep FC, Russel FG, van den Berg M, van Duursen MB, Masereeuw R, Endocrine disruptors differentially target ATP-binding cassette transporters in the blood-testis barrier and affect Leydig cell testosterone secretion in vitro, *Toxicol Sci* 136(2) (2013) 382–91. [PubMed: 24014645]
- [197]. Ogunbayo OA, Lai PF, Connolly TJ, Michelangeli F, Tetrabromobisphenol A (TBBPA), induces cell death in TM4 Sertoli cells by modulating Ca(2+) transport proteins and causing dysregulation of Ca(2+) homeostasis, *Toxicol in Vitro* 22(4) (2008) 943–952. [PubMed: 18329244]
- [198]. Saffarini CM, Heger NE, Yamasaki H, Liu T, Hall SJ, Boekelheide K, Induction and persistence of abnormal testicular germ cells following gestational exposure to di-(n-butyl) phthalate in p53-null mice, *J Androl* 33(3) (2012) 505–13. [PubMed: 21868749]
- [199]. Kleymenova E, Swanson C, Boekelheide K, Gaido KW, Exposure in utero to di(n-butyl) phthalate alters the vimentin cytoskeleton of fetal rat Sertoli cells and disrupts Sertoli cell-gonocyte contact, *Biol Reprod* 73(3) (2005) 482–90. [PubMed: 15901642]
- [200]. Mahood IK, Scott HM, Brown R, Hallmark N, Walker M, Sharpe RM, In utero exposure to di(n-butyl) phthalate and testicular dysgenesis: comparison of fetal and adult end points and their dose sensitivity, *Environ Health Perspect* 115 Suppl 1 (2007) 55–61. [PubMed: 18174951]
- [201]. Fisher JS, Macpherson S, Marchetti N, Sharpe RM, Human ‘testicular dysgenesis syndrome’: a possible model using in-utero exposure of the rat to dibutyl phthalate, *Hum Reprod* 18(7) (2003) 1383–94. [PubMed: 12832361]
- [202]. Ferrara D, Hallmark N, Scott H, Brown R, McKinnell C, Mahood IK, Sharpe RM, Acute and long-term effects of in utero exposure of rats to di(n-butyl) phthalate on testicular germ cell development and proliferation, *Endocrinology* 147(11) (2006) 5352–62. [PubMed: 16916955]
- [203]. Camacho L, Basavarajappa MS, Chang CW, Han T, Kobets T, Koturbash I, Surratt G, Lewis SM, Vanlandingham MM, Fuscoe JC, da Costa GG, Pogribny IP, Delclos KB, Effects of oral exposure to bisphenol A on gene expression and global genomic DNA methylation in the prostate, female mammary gland, and uterus of NCTR Sprague-Dawley rats, *Food and Chemical Toxicology* 81 (2015) 92–103. [PubMed: 25862956]
- [204]. Saegusa Y, Fujimoto H, Woo GH, Inoue K, Takahashi M, Mitsumori K, Hirose M, Nishikawa A, Shibutani M, Developmental toxicity of brominated flame retardants, tetrabromobisphenol A and 1,2,5,6,9,10-hexabromocyclododecane, in rat offspring after maternal exposure from mid-gestation through lactation, *Reprod Toxicol* 28(4) (2009) 456–467. [PubMed: 19577631]

Table 1.

BPA and oocyte outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Moore-Ambroz et al., 2015	Mouse	C57BL/6 J	Oral dose ^a	3 estrous cycle	50 µg/kg	After the 3 estrous cycles	Young (PND 28-32) female rats were exposed to BPA at a dose of 50 µg/kg during the first estrous cycle. In <i>in vivo</i> and <i>in vitro</i> fertilization assay, BPA exposure significantly decreased the percentage of fertilized oocytes. No changes of ovulation were observed	[25]
Ferris et al., 2015	Bovine		<i>In vitro</i> oocyte	24 h	15 and 30 ng/ml		In bovine oocyte, exposure to 30 ng/ml BPA (130 nM) decreased meiosis progression and increased spindle abnormalities, including abnormal spindle morphology and chromosome alignment	[44]
Wang et al., 2016b	Porcine		<i>In vitro</i> oocyte		250 µM		BPA exposure (250 µM) increased ROS, abnormal cytoskeletons, apoptosis/autophagy rate and altered epigenetic pattern in porcine oocytes	[28]
Ferris et al., 2016	Bovine		<i>In vitro</i> cumulus oocyte complexes	8 days	65 and 130 nM		Bovine oocytes exposed to BPA were fertilized in embryo culture and showed significant DNA damage and apoptosis at 130 nM, while gene expression in blastocysts was not altered.	[24]
Nakano et al., 2016	Mouse	ICR	<i>In vitro</i> oocyte	6, 9, 12, 15 and 18 h	2, 20, 50 and 100 µg/ml		BPA exposure inhibited oocyte maturation at concentrations of 50 and 100 µg/ml and delayed the cell cycle at a dose of 2 µg/ml. Further BPA treatment caused spindle abnormalities and activated the spindle assembly checkpoint at a dose of 50 µg/ml	[45]

^aPlacing a pipette tip with the dosing solution in the corner of the animal mouth.

Table 2.

BPA and human reproductive outcomes.

Source	Study design	Study Population	Sample Size	Measurement Time	BPA concentrations	Summary	Reference
Ferguson et al., 2014	Retrospective cohort	Boys ages 8–14 (ELEMENT) ^a	118	At the time of hormone measurement	NA	In 118 boys during the peripubertal period, parental (3rd trimester) or childhood BPA exposure was not associated with hormone levels, adrenarche or puberty.	[140]
Knez et al., 2014	Prospective cohort	Male partners undergoing IVF treatments (Slovenia)	149	Same day as follicle aspiration	Geometric mean 1.55 ng/mL	In 149 couples undergoing IVF, the natural logarithm transformed urinary BPA concentration was associated with lower natural logarithm transformed sperm count (b 1/4 0.241, 95% CI 0.470 to 0.012), natural logarithm transformed sperm concentration (b 1/4 0.219, 95% CI 0.436 to 0.003), and sperm vitality (b 1/4 2.660, 95% CI 4.991 to 0.329) in male partner, while the embryo development parameters were not associated with BPA exposure.	[97]
Lassen et al., 2014	Cross-sectional	Young men from general population (Denmark)	308	Same day as blood sample collection	Median 3.25 ng/mL	In 308 young men from Denmark's general population, men with BPA concentrations above the lowest quartile had higher serum T, LH, E2, and free T levels, as compared with the lowest quartile (p = 0.02). Men in the highest quartile of BPA had lower sperm progressive motility compared with men in the lowest quartile (−6.7 percentage points, 95% CI, −11.76, −1.63).	[18]
Lee et al., 2014	Case-control	Girls with central precocious puberty (Korea)	40 Peripheral PP/42 Central PP/32 Controls	Same day as blood sample collection	8.7 ± 7.6 µg/g creatinine (peripheral-PP), 8.0 ± 9.9 µg/g (central-PP), 6.6 ± 7.3 µg/g (control)	In 82 PP patients, BPA levels were slightly higher compared with 32 age-matched healthy girls but not associated with the onset of PP. Altered steroid metabolism was associated with urinary BPA levels, and levels of T, E2, and P4 were significantly increased in individuals with higher BPA level.	[19]
Upson et al., 2014	Case-control	Women with confirmed endometriosis (WREN) ^c	143 Cases/287 Controls	After disease diagnosis	Median 1.32 ng/mL (cases), 1.24 ng/mL (control)	In 143 endometriosis cases and 287 population-based controls, total urinary BPA concentrations was associated with non-ovarian pelvic endometriosis (second versus lowest quartile, OR 3.0; 95% CI, 1.2, 7.3; third versus lowest quartile, OR 3.0; 95% CI, 1.1, 7.6), but not associated with ovarian endometriosis.	[78]
Vagi et al., 2014	Case-control	Female PCOS patients from an urban academic medical center in Los Angeles	52 Cases/50 Controls	Same day as blood sample collection	Geometric mean 1.6 µg/L (PCOS case), 2.1 µg/L (control)	In 52 patients diagnosed with PCOS, the urinary BPA concentrations were not associated with increased risk of PCOS.	[71]
Akin et al., 2015	Case-control	Female PCOS patients from a Paediatric Endocrinology Outpatient Clinic (Turkey)	112 Cases/61 Controls	During the early follicular phase	Geometric mean 1.61 ng/mL (PCOS case), 0.8 ng/mL (control)	Among 112 girls with PCOS and 61 controls between 13 and 19 years of age, serum bisphenol A levels markedly increased in PCOS cases. Bisphenol A was significantly correlated with total T (r = 0.52), free T (r = 0.44) (p < 0.05).	[50]
Goldstone et al., 2015	Prospective cohort	Male partners of couples trying to become pregnant (LIFE)	418	Two days before semen collection	Geometric mean 0.55 ng/mL	In 418 males from Michigan and Texas, 2005–2009, BPA was negatively associated with DNA fragmentation in adjusted linear regression (̂ = −0.0544, p = 0.035). No	[17]

Source	Study design	Study Population	Sample Size	Measurement Time	BPA concentrations	Summary	Reference
Liu et al., 2015	Cross-sectional	Male workers with occupational exposure (China)	592	Pre and post-shift	Median 685.9 µg/g creatinine (exposed), 4.2 µg/g creatinine (unexposed)	associations between BPA exposure and other sperm parameters were observed. Among 592 male workers, urine BPA level was associated with increased prolactin ($p < 0.001$), E2 ($p < 0.001$), sex SHBG level ($p = 0.001$), and a reduced androstenedione ($p < 0.001$) and free androgen index level ($p = 0.021$).	[138]
Miao et al., 2015	Cross-sectional	Female workers with occupational exposure (China)	356	Pre and post-shift	Geometric mean 22 µg/g creatinine (exposed), 0.4 µg/g creatinine (unexposed)	In 106 exposed and 250 unexposed female workers, urinary BPA levels were positively associated with higher PRL and PROG levels ($p < 0.05$). The BPA levels were positively associated with E2 levels among exposed workers ($p = 0.05$) and negatively associated with FSH levels among the unexposed group ($p < 0.005$).	[51]
Minguez-Alarcon et al., 2015	Prospective cohort	Female completed at least one IVF cycle (Massachusetts General Hospital Fertility Center)	256	Same day of oocyte retrieval	Geometric mean 1.87 ng/mL	In 256 women undergoing IVF, Urinary BPA concentrations were not associated with endometrial wall thickness or peak E2 levels. Younger women (< 37 years old) had thicker endometrial thickness across increasing quartiles of urinary BPA concentrations, while older women (> 37 years old) had thinner endometrial thickness across increasing quartiles of urinary BPA concentrations.	[52]
Zhuang et al., 2015	Cross-sectional	Male workers with occupational exposure (China)	559	Pre-shift	Median 18.75 ng/mL (exposed) 3.37 ng/mL (unexposed)	In 281 exposed and 278 unexposed male workers, serum BPA levels were increased after occupational exposure. The serum BPA level was associated with lower serum AD level (0.18 ng/mL; 95 % CI -0.22 to -0.13) and higher serum SHBG level (2.79 nmol/L; 95 % CI 2.11-3.46), 104	[139]
Scinicariello et al., 2016	Cross-sectional	Male and female children (6-11 years) and adolescents (12-19 years) from general population (NHANES 2011-2012)	588	NA	Geometric mean 1.40-1.94 ng/mL	In 558 child and adolescent participants, urinary BPA concentrations were associated with significantly lower TT in male adolescents, and significantly higher TT in female adolescents (p -trend=0.01).	[20]
Vahedi et al., 2016	Case-control	Female PCOS patients who worked as marketing sellers (Iran)	62 Cases/62 Controls	NA	0.48 ng/mL (PCOS case), 0.16 ng/mL (control)	In 62 marketing professionals with PCOS and 62 healthy women with a similar job, serum BPA levels, LH-FSH ratio were significantly increased in PCOS cases, while TSH levels were significantly decreased.	[70]
Zhou et al., 2016	Cross-sectional	Infertile women with PCOS (China)	268	NA	Median 2.35 ng/mL	In 268 infertile women diagnosed with PCOS, urinary BPA level was associated with a significant decrease in AFC ($\beta = -0.34$, 95% CI = -0.60, -0.08; $p = 0.01$). But no association between BPA level and FSH was observed.	[53]
Simonelli et al., 2017	Case-control	Female endometriosis patients	68 Cases/60 Controls	During the consultation	Mean 5.31 ng/mL (cases), 1.64 ng/mL (control)	In 68 patients with endometriosis, urinary BPA level was significantly higher as compared with BPA level in the control group.	[79]

^aThe Early Life Exposure in Mexico to Environmental Toxicants project

Siracusa et al.

^qThe Longitudinal Investigation of Fertility and the Environment Study

^pThe Women's Risk of Endometriosis Study

^rThe Environment and Reproductive Health Study

^sThe Health Outcomes and Measures of the Environment Study

^tThe Maternal-Infant Research on Environmental Chemicals Study

^uThe North Carolina Early Pregnancy Study

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

BPA and female steroidogenesis in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Delclos, et al., 2014	Rat	SD	Oral gavage	GD 6-PND 90	2.5–300000 µg/kg	PND 80	SD female rats were exposed to BPA orally from GD 6 through PND 90. The chronic BPA exposure resulted in increases in serum E2 and prolactin level only at 100000 and 300000 µg/kg groups. Progesterone was significantly decreased at 300,000 µg BPA/kg BW/day groups.	[23]
Li et al., 2014	Rat	Wistar	Intraperitoneal injection	PND 28-PND 35	10, 40 and 160 mg/kg	PND 35	Pre-puberty female Wistar rats were exposed to BPA for one week. BPA exposure decreased at 40 and 160 mg BPA/kg BW/day, while serum E2 level was not altered.	[62]
Sadowski et al., 2014	Rat	Long-Evans	Oral gavage	GD 0-PND 9	4, 40 and 400 µg/kg	PND 23	Long-Evans rats received oral administration of BPA at 4 µg/kg, 40 µg/kg, or 400 µg/kg throughout pregnancy, and the pups received direct oral administration of BPA between PND 1–9. Rats in both genders had decreased levels of FSH at weaning at doses of 4 and 400 µg/kg. But no changes in other hormone levels or maze behavior were observed.	[65]
Gamez et al., 2015	Rat	Wistar	Drinking water	GD 0-PND 21	3 µg/kg	PND 30	Wistar mated rats were treated with BPA in their drinking water (estimated 3 µg/kg) until their offspring were weaned at 21 days after birth. Serum LH and E2 levels were significantly increased in the treatment group with increased follicle number, while no changes in serum FSH level were observed.	[64]
Moore-Ambriz et al., 2015	Mouse	C57BL/6 J	Oral dose	3 estrous cycle	50 µg/kg	After the 3 estrous cycles	Young (PND 28-32) female rats were exposed to BPA at a dose of 50 µg/kg during the first estrous cycle. In <i>in vivo</i> and <i>in vitro</i> fertilization assay, BPA exposure significantly decreased the percentage of fertilized oocytes. No changes of ovulation and hormone levels were observed.	[25]
Santamaria et al., 2016	Rat	Wistar	Drinking water	GD 9-PND 21	0.5 and 50 µg/kg	PND 90	In adult female rat offspring with gestational and lactational exposure, serum P4 levels were significantly increased in BPA 0.5 and 50 µg/kg groups coupled with high mRNA expression of 3b-hydroxysteroid dehydrogenase (3b-HSD).	[63]
Mansur et al., 2016	Human		<i>In vitro</i> granulosa cells	48 h	0.2, 0.02, 2.0, 20 mg/ml		Treatment of human granulosa cells after 48 h resulted in significantly lower progesterone biosynthesis at doses of 2 or 20 mg/ml and lowered E2 level at a dose of 20 mg/ml. These two concentrations significantly reduced the mRNA and protein levels of 3b-HSD, CYP11A1, and CYP19A1 without affecting STAR and 17b-hydroxysteroid dehydrogenase mRNA expression.	[66]

Table 4.1.

BPA and ovary outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Delclos, et al., 2014	Rat	SD	Oral gavage	GD 6-PND 90	2.5–3000000 µg/kg	PND 21	SD rats exposed to BPA orally from GD 6 through PND 90. Adverse effects of BPA exposure on the ovary, including increased cystic follicles, depleted corpora lutea, and antral follicles at two BPA high dose (100000 or 300000 µg/kg).	[23]
Li et al., 2014	Rat	Wistar	Intraperitoneal injection	PND 28-PND 35	10, 40 and 160 mg/kg	PND 35	Pre-puberty female Wistar rats were exposed to BPA for one week. BPA exposure significantly decreased rat ovarian weights and follicle numbers at doses of 40 and 160 mg/kg. In addition, the mRNA and protein levels of the germline alpha (FIGLA) and oocyte-specific histone H1 variant (H1FOO) genes decreased at 160 mg/kg and 10–160 mg/kg, respectively.	[62]
Patisaul et al., 2014	Rat	Wistar	Dietary exposure and Drinking water	GD 6-PND 40	0.18-0.44 mg/kg	PND 120	Wistar rats exposed to BPA during GD 6-PND 40 showed delayed vaginal opening. But no changes in corpora lutea and cystic follicles were observed.	[72]
Berger et al., 2016	Mouse	FVB	Oral dose	GD 11-PND 0	0.5, 20, and 50 µg/kg	PND 4 and 21 (F1-F3)	To examine the transgenerational effects of BPA on the ovary, FVB mice exposed to BPA in utero and ovaries of F1-3 generations were collected at PND 4 and PND 21 to examine expression levels of multiple genes in apoptosis, antioxidant, igf family, hormone receptor, and steroidogenesis. Overall, more significant changes in gene expression were observed at PND 21, while no clear dose-dependent responses showed.	[22]
Santamaria et al., 2016	Rat	Wistar	Drinking water	GD 9-PND 21	0.5 and 50 µg/kg	PND 90	In adult female rat offspring with Maternal exposure to low doses of BPA during gestation and breastfeeding, ovaries from both BPA-treated groups showed decreased ovaries weight, primordial follicle recruitment, and a greater number of corpora lutea.	[63]
Gamez et al., 2015	Rat	Wistar	Drinking water	GD 0-PND 21	3 µg/kg	PND 30	No changes in ovarian weight and its relative weight.	[64]
Zhou et al., 2015	Mouse	CD-1	<i>In vitro</i> ovary	24-192 h	0.1, 1, 5, and 10 µg/mL	PND1, 2, 4 and 8	In ovaries of newborn mice, BPA exposure increased germ cells and decreased primordial follicle in ovaries. Further, BPA exposure altered the expression of key genes involved in apoptosis and antioxidant, while the effects varied by ovary collection times and exposure doses.	[73]
Ganesan and Keating, 2016	Rat	F344	<i>In vitro</i> ovary	48-192 h	440 µM	PND 4	In F344 rat ovaries, BPA reduced small primary, large primary and secondary follicle numbers and increased DNA damage markers, γ-H2AX, ATM, and DNA repair genes.	[74]

Table 4.2.

BPA and uterine outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Calhoun et al., 2014	Monkey	Rhesus Macaque	Dietary exposure	GD 100-GD 165	400 µg/kg	GD 165 (F1)	The pregnant rhesus macaques were exposed orally to BPA. Fatal uteri did not show notable changes in uterus histology, proliferation, expression of hormone receptors. At GD 165, BPA exposure altered expression levels of three genes (HOXA13, WNT4, and WNT5A) involved in reproductive organ development.	[85]
Kim et al., 2014	Mouse	ICR	Intraperitoneal injection	0.5, 1, 2, 4, 6 and 12 h	10, 20, 50, 100, 200 and 500 mg/kg	Adult	Egr1 (early growth response 1) was rapidly and transiently induced by BPA (starting at 20 mg/kg) in stromal cells surrounding implanting blastocyst via nuclear estrogen receptor (ER)-ERK1/2 pathway.	[87]
Kendzioriski and Belcher, 2015	Mouse	C57Bl/6N and CD-1	Dietary exposure	4 weeks	0.004, 0.04, 0.4, 4, and 40 mg/kg	Adult	In adult C57Bl/6N and CD-1 mice, BPA non-monotonically increased gland nest density and periglandular collagen accumulation, collagen I and III expressions, decreased matrix metalloproteinase 2 (MMP2) and MMP14 expression, and increased immune response were observed in 4 mg/kg BPA CD-1 group only.	[84]
Camacho et al., 2015	Rat	SD	Oral gavage	GD 6-PND 90	2.5, 8, 25, 80, 260, 840, 2700, 100,000 and 300,000 µg/kg	PND 4, PND 90	In SD rats exposed to BPA orally, no changes in global genomic DNA methylations in uterus were observed. Changes in prostate and female mammary glands were only observed in BPA-high dose groups (100,000 and 300,000 µg/kg).	[203]
Salleh et al., 2015	Rat	SD	Ex-vivo	3-5 min	1×10 ⁻⁸ -1×10 ⁻⁴ M	Adult	In <i>ex-vivo</i> uterus contraction assay, uterine contractile force decreased with increasing doses of BPA.	[86]
Vigezzi et al., 2015	Rat	Wistar	Drinking water	GD 9-PND 21	0.5 and 50 µg/kg	PND 90, PND 360	Wistar rats were exposed to BPA via drinking water through GD 9 to PND 21. The female offspring showed decreased glandular proliferation (0.5 and 50 µg/kg) and alpha-actin expression (50 µg/kg) at PND 90 and increased the incidence of abnormalities in the luminal (50 µg/kg) and glandular epithelium (0.5 and 50 µg/kg) at PND 360.	[27]
Li et al., 2016	Mouse	CD-1	Oral	PND 22-GD 9	60 and 600 µg/kg	GD 9	In CD-1 mice, BPA exposure affected puberty and estrous cyclicity, decreased embryo implantation, PGR and HAND2 expression in uterine with enhanced activation of fibroblast growth factor and MAPK signaling in the epithelium, and increased improper endometrial epithelial and stromal functions.	[83]
Kang et al., 2014			<i>In vitro</i> uterine smooth muscle cells	48 h	2.28 ng/mL		In uterine smooth muscle cells, BPA treatment upregulated genes involved in cell differentiation, cellular metabolic processes, cell proliferation and smooth muscle contraction and decreased the genes involved in mitosis, lipid metabolism, regulation of muscle cell differentiation and cell adhesion.	[90]

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Helmesstam et al., 2014	Human		<i>In vitro</i> endometrial endothelial cell	24 h	0.01, 1, 100, 10000 nM		The HECC cells were co-cultured with primary endometrial stromal cells to mimic the <i>in vivo</i> situation and treated with environmentally relevant doses of BPA. BPA exposure did not alter the HECC viability and proliferation, and increased tube formation and vascular endothelial growth factor (VEGF)-D protein expression at a dose of 10000 nM.	[91]
Shen et al., 2014	Human		<i>In vitro</i> uterine leiomyoma	24, 48 and 72 h	1, 2.5, 5, 10 and 20 μ M		The BPA exposure increased the protein expression level of ER α , IGF-1, and VEGF at doses of 2.5, 5 and 10 μ M in uterine leiomyomas.	[92]
Wang et al., 2015	Human		<i>In vitro</i> endometrial carcinoma cells line (RL95-2)	7 days	1 \times 10 ⁻¹⁰ -1 \times 10 ⁻⁴ M		The BPA exposure increased gene expression of cyclooxygenase-2 (COX-2) and epithelial-mesenchymal transition (EMT), promote growth rate and colony-forming efficiency in a nonmonotonic manner in human endometrial carcinoma cells line.	[93]

Table 5.

BPA and sperm outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Gurmeet et al., 2014	Rat	SD	Oral gavage	PND 28-70	1, 5 and 100 mg/kg	PND 70	In prepubertal SD rats, 6-week BPA oral exposure increased intercellular junction disruptions, sloughing of germ cells, and immature germ cells and cellular debris. No testis weight change was observed. Further BPA exposure decreased plasma T and E2 levels in all treatment groups.	[101]
Liu et al., 2014	Rat	Wistar	Oral gavage	60 days	20 µg/kg	Adult	In Wistar rats, a 60-day exposure to BPA at an environmentally relevant dose resulted in significant increase in the proportion of stage VII seminiferous epithelium and the decrease in stage VII. Further, BPA exposure induced the expression level of γ -H2AX, ATM and SCP3, suggesting DNA damage and early meiosis inhibition.	[138] 80
Vilela et al., 2014	Mouse	Vesper	Oral gavage	GD 0-PND 0	40, 80 and 200 µg/kg	PND 70	In vesper mice received utero BPA exposure, BPA treatment significantly reduced the sperm with normal morphology at a dose of 200 µg/kg, mitochondrial integrity at doses of 80 and 200 µg/kg, and <i>in vitro</i> penetration rate starting at 40 µg/kg, whereas no changes in DNA integrity and acrosome integrity were observed.	[102]
Wang et al., 2014	Rat	SD	Oral gavage	4 weeks	50, 100, and 200 mg/kg	Adult	In SD rat, BPA exposure increased germ cell apoptosis in testes and decreased sperm concentration at doses of 100 and 200 mg/kg. Further, BPA exposure significantly increased protein and mRNA levels of cytochrome C, apoptosis-inducing factor, caspase-3/9, Bax, and decreased protein and gene levels of Bcl-2 in all treatment groups. Also, the abnormal structure of mitochondria was observed at a dose of 200 mg/kg.	[112]
LI et al., 2015	Mouse	C57BL/6J	Intraperitoneal injection	PND 21-28	50 mg/kg	PND 56	In C57BL/6J mice receiving BPA exposure, a decrease in epididymal sperm number and increase in sperm deformity rate were observed. Abnormal seminiferous tubules with sloughing of germ cells were observed, while male fertility was not affected.	[103]
Mao et al., 2015	Rat	SD	Oral gavage	GD 0-PND 21	40 µg/kg	PND 56 (F1)	The paternal BPA exposure induced Igf2 DMR2 methylation and decreased mRNA expression of IGf2 in F1 sperm, which potentially associated with hypermethylation of Igf2 observed in islets of male F2 offspring.	[116]
Vrooman et al., 2015	Mouse	Cd-1; C57BL/6J; C3H/HeJ	Oral gavage	PND 1-PND 12	20 and 500 µg/kg	PND 12/84/365	In multiple strains of mice, BPA exposure did not alter the chromosome pairing and synapsis in spermatogonial stem cells.	[115]
Wisniewski et al., 2015	Rat	Wistar	Oral gavage	PND 50-90	5 and 25mg/kg	PND 105	In Wistar rats exposed to BPA orally, serum T, LH, and FSH levels were decreased, and the E2 level was increased in both treatment groups. In addition, the gene expression levels of hormone receptors were increased in the hypothalamus. BPA exposure reduced sperm production, reserves and transit time in both treatment groups.	[104]
Hass et al., 2016	Rat	Wistar	Oral gavage	GD 7-PND 22	25, 250, 5000 and 50000 µg/kg	3 months	Prenatal and postnatal BPA exposure significantly decreased the sperm count only at a dose of 25 µg/kg in Wistar rat. No changes in reproductive organ were observed.	[105]

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Kalb et al., 2016	Mouse	Swiss Albino	Oral gavage	PND 0-PND 21	300, 900, and 3000 µg/kg	Adult	During the breastfeeding period, maternal BPA exposure altered sperm parameters (sperm motility, normal morphology, membrane integrity, acrosomal integrity, DNA integrity, and mitochondrial functionality) at all treatment groups. BPA 3000 µg/kg treatment also induced Testicular degeneration. In addition, the total antioxidant capacity was impaired at doses of 900 and 3000 µg/kg.	[106]
Quan et al., 2016a	Rat	SD	Intraperitoneal injection	20 days	2, 10, 50 mg/kg	6 months	In SD rats, BPA exposure induced oxidative stress (SOD activity, GSH-Px activity, and MDA level) at a dose of 50 mg/kg and sperm malformation rate and apoptosis in testis at all treatment groups. The mRNA expression of genes in Akt/mTOR pathway were altered in the different treatment groups. In addition, the serum T (2, 10, 50 mg/kg), FSH (10 and 50 mg/kg) and LH (50 mg/kg) levels were decreased.	[109]
Quan et al., 2016b	Rat	SD	Oral gavage	GD 14-21	1, 10, 100 mg/kg	PND 21	In SD rat male offspring, utero BPA exposure induced oxidative stress (MDA level) at doses of 10 and 100 mg/kg and apoptosis in testis, and altered seminiferous tubules morphology at all treatment groups. The mRNA and protein expression of genes in Akt/mTOR pathway were altered in the different treatment group. In addition, the serum T (100 mg/kg), FSH (100 mg/kg) and LH (1, 10, 100 mg/kg) levels were decreased.	[110]
Rahman et al., 2016	Mouse	CD-1	Oral gavage	GD 7-14	50, 5000 and 50000 µg/kg	PND 120	Gestational exposure to BPA inhibited sperm count, motility parameters, and intracellular ATP levels in a dose-dependent manner. BPA exposure reduced numbers of stage VIII seminiferous epithelial cells in testis at doses of 5000 and 50000 µg/kg and decreased PKA activity and tyrosine phosphorylation in spermatozoa at doses of 5000 and 50000 µg/kg. Further, BPA exposure altered the expression levels multiple proteins involved in ATP production, oxidative stress response, and fertility.	[107]
Wang et al., 2016	Mouse	C57BL/6	Oral gavage	8 weeks	10, 50 and 250 µg/kg	Adult	In C57BL/6J mice, BPA exposure decreased sperm viability at a dose of 250 µg/kg and motility at all treatment groups. Sperm-specific Ca ²⁺ channel currents were reduced in all treatment groups. And the mRNA and protein levels of CatSper subunits were reduced at doses of 50 and 250 µg/kg. Similar results were observed <i>in vitro</i> sperm treated with BPA.	[108]
Yin et al., 2016	Mouse	Kunming	Oral gavage	5 weeks	3, 30 and 300 mg/kg	Adult	In Kunming mice, BPA exposure induced spermatogenic cellular apoptosis, sloughing of germ cells, and seminiferous tubules vacuolation in all treatment groups.	[111]
Chen et al., 2017	Rat	Wistar	Oral gavage	35 weeks	50 µg/kg	Adult	In Wistar rats receiving 50 µg/kg BPA exposure for 35 weeks, no apoptosis or sperm deformation was observed. BPA exposure decreased protein lysine acetylation levels and increased the expression of histone deacetylase Sirt1, ERβ expression and its binding with caveolin-1 (Cav-1) in testes.	[114]
Qian et al., 2015	Mouse		<i>In vitro</i> spermatoocyte GC-2 cell line	0.5-48 h	0.02-20 µM		In mouse spermatoocyte GC-2 cell line, BPA treatment decreased cell viability, the release of mitochondrial cytochrome c and the activation of caspase-3 at doses from 0.2 to 20 µM. Further, BPA exposure impaired Ca ²⁺ homeostasis and induced Ca ²⁺ sensor proteins (CaM and CaMKII) expressions starting at 0.02 and 0.2 µM, respectively. The pretreatment with Ca chelator or sensor protein inhibitor could partially attenuate BPA-induced cellular injury.	[117]

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Rahman et al., 2015	Mouse	ICR	<i>In vitro</i> spermatozoa	6 h	0.0001, 0.01, 1, and 100 µM		BPA exposure significantly affected sperm motility, fertilization rate, and intracellular ATP levels at a dose of 100 µM. Also, BPA induced the protein kinase A (PKA) activity, tyrosine phosphorylation and fertility-related proteins (peroxiredoxin-5, glutathione peroxidase 4, glyceraldehyde-3-phosphate dehydrogenase, and succinate dehydrogenase), except β-actin in spermatozoa.	[118]
Barbonetti et al., 2016	Human		<i>In vitro</i> spermatozoa	4 and 20 h	10–800 µM		In human spermatozoa, BPA exposure decreased the sperm motility, viability and mitochondrial membrane potential, and increased apoptosis, DNA oxidative damage marker, 8-hydroxy-2'-deoxyguanosine starting at 300 µM.	[121]
Kotwicka et al., 2016	Human		<i>In vitro</i> spermatozoa	5-60 min	10–10, 10–8 and 10–6 M		In human spermatozoa, BPA treatment induced a transient increase of velocity straight linear (VSL) and homogeneity of progressive movement velocity (HPMV) at 15 min after stimulation. 1h BPA exposure did not alter vitality phosphatidylserine membrane translocation, for all doses. BPA at a dose of 10–6 mol/L induced a rapid and transient increase of intracellular free calcium ions concentration.	[122]
Wan et al., 2016	Rat	SD	<i>In vitro</i> mature sperm	10-300 min	1-600 µg/ml		In rat sperm, exposure to BPA significantly decreased sperm motility in a dose and time-dependent manner. Further, capacitation-associated protein tyrosine phosphorylation was induced by BPA treatment, while the induction could be blocked by PKA inhibitor.	[119]
Yin et al., 2016	Mouse		<i>In vitro</i> spermatocyte GC-2 cell line	24 and 48 h	20, 40 and 80 µM		In mouse spermatocyte GC-2 cell line, BPA exposure decreased cell viability starting at 50 µM for 24 h and 5 µM for 48 h and induced apoptosis, ER stress, ROS mitochondrial damage and the mitochondrial apoptotic pathway starting at 20 µM for 48 h. Knocking down the PERK/EIF2α/chop pathway (one of the ER stress pathways) partially recovered the BPA-induced cell apoptosis.	[111]
Liang et al., 2017	Mouse		<i>In vitro</i> Spermatogonial cell line C18-4	24, 48 and 72 h	0.1, 1, 10, and 50 µM		The BPS exposure significantly decreased cell viability, altered nuclear morphology, increased gamma-H2AX expression level, and perturbed cytoskeleton and cell cycle progression at a dose of 50 µM in spermatogonial cell line C18-4.	[120]

Table 6.

BPA and Sertoli/Leydig cells in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Ge et al., 2014a	Mouse		<i>In vitro</i> Sertoli cell line TM4	24 and 48 h	10–8 M and 10–5 M		Exposure to 10– 5 MBPA induced oxidative stress and inhibited cell proliferation. Exposure to 10– 8 M BPA increased intercellular ATP activities of mitochondria, and proliferation in Sertoli cell line TM4.	[128]
Ge et al., 2014b	Mouse		<i>In vitro</i> Sertoli cell line TM4	24 and 48 h	10–8 M and 10–3 M		Exposure to 1 and 10 nM BPA significantly induced the Sertoli cell proliferation via activating ERK1/2 through GPR30 and ER alpha/beta.	[129]
Qi et al., 2014	Rat		<i>In vitro</i> primary Sertoli cell culture	24 h	30, 50, 70 and 90 μ M		In primary rat Sertoli cells, BPA exposure significantly reduced cell viability and induced apoptosis starting at a dose of 50 μ M. Further, BPS exposure activated JNKs/p38 MPAK and Fas/FasL signaling pathways and induced translocation of NF- κ B at doses of 50 and 70 μ M.	[26]
Qian et al., 2014	Mouse		<i>In vitro</i> Sertoli cell line TM4	12, 24 and 48 h	0.02, 0.2, 2.0 and 20 μ M		In primary rat Sertoli cells, BPA exposure significantly reduced cell viability and induced apoptosis starting at a dose of 50 μ M. Further, mitochondrial mass loss, membrane potential decrease, cytochrome c release, Bcl-2 family members down-regulation and caspases-3 up-regulation were observed in TM4 cells treated with BPA. Additionally, the expression of CaM, ERK1/2, and phosphorylation of CaMKII significantly increased after BPA treatment. Treatments with CaM, ERK1/2 and CaMKII inhibitor attenuated BPA-induced cell damage.	[130]
Xiao et al., 2014	Human		<i>In vitro</i> primary Sertoli cell culture	24, 48 and 72 h	0.4, 4, 40 and 300 μ M		In human Sertoli cells, BPA exposure significantly decreased the cell viability, expression levels of ZO-1, N-Cadherin and beta-Catenin at a dose of 200 μ M. BPA treatment also increased truncation and depolymerization of actin microfilaments starting at 0.4 μ M and improper localization of actin regulatory proteins Arp3 and Eps8 at a dose of 200 μ M.	[133]
Wang et al., 2015	Rat		<i>In vitro</i> primary Sertoli cell culture	24 h	30, 50, and 70 μ M		In primary rat Sertoli cells, BPA exposure decreased cell viability and induced apoptosis at doses of 50 and 70 μ M. BPA exposure caused the elevation of Pten expression and the inactivation of Akt, and then triggered the caspase3, which led to apoptosis of Sertoli cells.	[132]
de Freitas et al., 2016	Human		<i>In vitro</i> primary Sertoli cell culture	6 and 24 h	20 μ M		In human Sertoli cells, BPA exposure significantly decreased the cell viability, expression levels of occludin, ZO-1, β -catenin, and AR at a dose of 20 μ M for 6 and 48 h without changes in F-actin expression or localization.	[131]
Wang et al., 2016	Rat		<i>In vitro</i> primary Sertoli cell culture	30, 50, and 70 μ M			In primary rat Sertoli cells, BPA exposure induced ROS, intracellular Ca ²⁺ release at doses of 50 and 70 μ M and cellular apoptosis at a dose of 70 μ M. Pretreatment with N-acetyl-L-cysteine attenuated the BPA-induced cellular apoptosis.	[135]
Akingbemi et al., 2004	Rat	Long-Evans	<i>In vitro</i> primary Leydig cell culture	18 h	0.01 nM		Treatment of adult Leydig cells with 0.01 nM BPA decreased T biosynthesis and E2 level, mRNA expression of aromatase and the steroidogenic enzyme 17 α -hydroxylase/17–20 lyases.	[134]
Nakamura et al., 2010	Rat	Wistar/ST	Intraperitoneal injection	6 weeks	0, 20, 100 and 200mg/kg	Adult	200 mg/kg BPA significantly decreased Leydig cell numbers with decreases in the STAR protein level in the rat testis	[134]

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Naniappa et al., 2012	Rat	Long-Evans	Oral gavage	GD 12-PND 21	2.5 and 25 µg/kg	Adult	BPA exposure (2.5 and 25 µg/kg) significantly promoted Leydig cell division during the prepubertal period and increased Leydig cell numbers at PND 90. In Leydig cells, BPA treatment induced the protein expression of proliferating cell nuclear antigen, MAPK, AR, and ER, increased secretion of the anti-Müllerian hormone. Also, BPA suppressed protein expressions of the LH receptor and the 17beta-hydroxysteroid dehydrogenase enzyme.	[127]
Chen et al., 2016	Mouse		<i>In vitro</i> Leydig cell line TM3	24, 48 and 72 h	10 ⁻⁸ to 10 ⁻³ M		In Leydig TM3 cells, BPA greater than 10 ⁻⁶ M inhibited the proliferation of Leydig TM3 in a dose-dependent manner. The proteomic study revealed BPA could modulate the expression of proteins related to cell structure and motility and cellular metabolism. Further, BPA induced cell migration at a dose of 10 ⁻⁵ M via galectin-1 and ERK1/2.	[136]

Table 7.

BPA and male steroidogenesis in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Delclos et al., 2014	Rat	SD	Oral gavage	GD 6-PND 90	2.5–300000 µg/kg	PND 15	In male offspring, BPA exposure significantly increased serum T4 level at the highest BPA dose group. No changes in serum T3 and FSH levels and testicular descent were observed.	[23]
Gamez et al., 2014	Rat	Wistar	Drinking water	GD 6-PND 21	3 µg/kg	PND 35	In Wistar rats exposed to BPA, a decrease in testicular weight was observed, while seminal vesicles weight, relative weights of testes and seminal vesicles were not changed.	[150]
Gurmeet, et al., 2014	Rat	SD	Oral gavage	PND 28-70	1, 5 and 100 mg/kg	PND 70	In prepubertal SD rats, 6-week BPA oral exposure increased intercellular junction disruptions, sloughing of germ cells, and immature germ cells and cellular debris. No testis weight change was observed. Further BPA exposure decreased plasma T and E2 levels in all treatment groups.	[101]
Sadowski et al., 2014	Rat	Long-Evans	Oral gavage	GD 0-PND 9	4, 40 and 400 µg/kg	PND 23	Long-Evans rats received oral administration of BPA at 4 µg/kg, 40 µg/kg, or 400 µg/kg throughout pregnancy, and the pups received direct oral administration of BPA between postnatal days 1–9. Rats had decreased levels of FSH at weaning at doses of 4 and 400 µg/kg but no changes in other hormone levels and maze behavior.	[65]
Chouhan et al., 2015	Mouse	Swiss albino	Intraperitoneal injection	60 days	0.5, 50 and 100 µg/kg	Adult	In Swiss albino mice, 60-day intraperitoneal BPA exposure significantly decreased sperm count, STAR expression in the testis and serum T level in all treatment groups.	[151]
Quan et al., 2016a	Rat	SD	Intraperitoneal injection	20 days	2, 10, 50 mg/kg	6 months	In SD rats, BPA exposure induced oxidative stress (SOD activity, GSH-Px activity, and MDA level) at a dose of 50 mg/kg and sperm malformation rate and apoptosis in testis at all treatment groups. The mRNA expression of genes in Akt/mTOR pathway were altered in the different treatment group. In addition, the serum T (2, 10, 50 mg/kg) FSH (10 and 50 mg/kg) and LH (50 mg/kg) levels were decreased.	[109]
Quan et al., 2016b	Rat	SD	Oral gavage	GD 14-21	1, 10, 100 mg/kg	PND 21	In SD rat male offspring, utero BPA exposure induced oxidative stress at doses of 10 and 100 mg/kg and apoptosis in testis, altered seminiferous tubules morphology at all treatment groups. The mRNA and protein expression of genes in Akt/mTOR pathway were altered in the different treatment group. In addition, the serum T (100 mg/kg), FSH (100 mg/kg) and LH (1, 10, 100 mg/kg) levels were decreased.	[110]
Wisniewski et al., 2015	Rat	Wistar	Oral gavage	PND 50-90	5 and 25mg/kg	PND 105	In Wistar rats exposed to BPA orally, serum T, LH, and FSH levels were decreased, and the E2 level was increased in both treatment groups. Also, the gene expression levels of hormone receptors were increased in the hypothalamus. BPA exposure reduced sperm production, reserves and transit time in both treatment groups.	[104]
Zang et al., 2016	Mouse	C57BL/6	Intraperitoneal injection	21 days	10, 50 and 100 mg/kg	Adult	In C57BL/6 mice, BPA exposure significantly impaired sexual behavior and decreased testis weight in all treatment groups and epididymis at 100 mg/kg group. Further BPA exposure decreased serum T levels at a dose of 100 mg/kg and intratesticular T at doses of 50 and 100 mg/kg.	[152]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	n	Summary	Reference
Eladak <i>et al.</i> , 2015	Human/ Mouse/R at		<i>In vitro</i> fetal testis explants	24-72 h	0.001-10000 nM			BPA exposure significantly reduced basal T secretion levels in rat fetal testis explants at a dose of 10000 nM, in mouse at 1000 and 10000 nM, and in human starting at 10 nM.	[153]

Table 8.

BPS and reproductive outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Yamasaki et al., 2004	Rat	Wistar	Subcutaneous injection	3 days	20, 100 and 500 mg/kg	PND 24	BPS showed weaker estrogen receptor binding capacity as compared with E2 (0.0055%). In the immature rat uterotropic assay, BPS treatment significantly increased absolute and relative uterine wet weight and blotted weight at doses of 20 and 500 mg/kg.	[162]
Ji et al., 2013	Fish	Danio ratio	Water	21 days	0.5, 5, and 50 µg/L	Adult	Adult zebrafish were exposed to 0.5, 5, and 50 µg/L of BPS for 21 days. Egg production and the gonad somatic index in female fish was significantly decreased at 0.5 µg/L BPS. Plasma concentrations of E2 were significantly increased in both genders. In male fish, significant decreases of T level were observed with up-regulation of cyp19a and down-regulation of cyp17 and 17βhsd transcripts at 50 µg/L BPS. In an F1 generation, parental BPS exposure resulted in delayed and decrease in hatching rate. Continuous BPS exposure in the F1 further increased malformation rates.	[164]
Naderi et al., 2014	Fish	Danio ratio	Water	75 days	0.1, 1, 10 and 100 µg/l	Adult	Zebrafish embryos were exposed to various concentrations of BPS for 75 days. The plasma E2 Levels were significantly increased in both genders starting at a dose of 100 µg/l. In male fish, BPS exposure decreased plasma T, T3, T4 levels and sperm count at doses of 10 and 100 µg/l. In female fish, BPS exposure decreased T3 and T4 levels at a dose of 100 µg/l and egg production at doses of 10 and 100 µg/l	[165]
Ullah et al., 2016	Rat	SD	Oral gavage	28 days	1, 5, 25 and 50 µg/kg	Adult	In adult rats exposed to various concentrations of BPS, a significant increase in the testicular ROS and lipid peroxidation was observed at a dose of 50 µg/kg with reduced antioxidant enzyme activity. Further BPS decreased plasma T level and altered the testicular morphology (thin seminiferous epithelium; reduction the tubular epithelium area; empty lumen) at a dose 50 µg/kg. No changes in spermatogenesis were observed.	[163]
Eladak et al., 2015	Human/ Mouse/Rat		<i>In vitro</i> fetal testis explants	24-72 h	10, 100, 1000 and 10000 nM		BPS exposure significantly reduced basal T secretion levels in mouse fetal testis explants starting at 100 nM, and in human starting at 1000 nM.	[153]
Chen et al., 2016	<i>Caenorhabditis Elegans</i>			4 days	125, 250 and 500 µM		BPS exposure increased embryonic lethality, germline nuclear loss and apoptosis at doses of 125, 250 and 500 µM in <i>C. Elegans</i> . 500 µM BPS exposure induced DNA damage and impaired homologous chromosome synapsis. Further BPS show distinct alterations of gene expression at whole transcriptome level as compared with BPA.	[166]
Roelofs et al., 2015	Mouse		<i>In vitro</i> Leydig cells MA-10	48 h	10 µM		BPS exposure significantly increased P4 levels sand gene (5αRed1) in the MA-10 cells at a dose of 10 µM.	[167]
Liang et al., 2017	Mouse		<i>In vitro</i> Spermatogonial cell line C18-4	24, 48 and 72 h	0.1, 1, 10, and 50 µM		The BPS exposure significantly decreased cell viability, altered nuclear morphology, increased gamma-H2AX expression level, and perturbed cytoskeleton and cell cycle progression at a dose of 50 µM in spermatogonial cell line C18-4.	[120]

Table 9.

BPF and reproductive outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Stroheker et al., 2003	Rat	Wistar	Oral gavage	4 days	25, 50, 100, and 200 mg/kg	PND 26	In immature Wistar rats, BPF exposure increased vaginal cornification and uterus relative weight at a dose of 200 mg/kg. BPF showed weaker estrogen receptor binding capacity as compared with E2 (0.0719%). In the immature rat heterotrophic assay, BPF treatment significantly increased absolute and relative uterine wet weight and blotted weight at all treatment group in a dose-dependent manner.	[170]
Yamasaki et al., 2004	Rat	Wistar	Subcutaneous injection	3 days	100, 300 and 1000 mg/kg	PND 24	In SD rats exposed to BPF orally, serum T3 levels increased, and serum T4 levels decreased in both genders at a dose of 500 mg/kg. In male rats, the testis weight significantly increased at a dose of 500 mg/kg.	[162]
Higashihara et al., 2007	Rat	SD	Oral gavage	28 days	20, 100 and 500 mg/kg	Adult	BPF exposure significantly increased T levels at a dose of 100 µM, P4 levels and genes (Cyp51 and 5aRed1) at a dose of 10 µM in the MA-10 cells.	[167]
Roelofs et al., 2015	Mouse		<i>In vitro</i> Leydig cells MA-10	48 h	0.01–100 µM		BPF exposure significantly reduced basal T secretion levels in mouse fetal testis explants starting at 1000 nM, and in human starting at 10000 nM.	[153]
Eladak et al., 2015	Human/Mouse/Rat		<i>In vitro</i> fetal testis explants	24–72 h	10, 100, 1000 and 10000 nM			

Table 10.

BPAF and reproductive outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
Feng et al., 2012	Rat	SD	Oral gavage	14 days	2, 10, 50 and 200 mg/kg	Adult	In adult rats, orally to BPAF at doses from 2 to 200 mg/kg, the concentration of BPAF in testes increased in a dose-dependent manner. In addition, 14 days of exposure decreased serum T level and the expression levels of genes and proteins in T biosynthesis pathway, and increased serum LH and FSH levels in rats given a dose of 200 mg/kg.	[181]
Furr et al., 2014	Rat	SD	Oral gavage	GD 14-GD 18	200, 300, 400, 500 and 750 mg/kg	GD 18	The prenatal exposure to BPAF (200-750 mg/kg) did not alter T secretion in dams on GD 18.	[179]
Shi et al., 2015	Fish	Danio ratio	Water	PF 4h - PF 140d	5, 25 and 125 µg/L	PF 140d	Zebrafish were exposed to BPAF at 5, 25 and 125 µg/L from 4-hour pf to 120 pf, representing the period from embryo to adult. The increases in E2 levels were observed in both genders, while decreases in T levels were only observed in the male fishes. In offspring, increase in malformation and decrease in survival rate were observed only at maternal exposure to BPAF at 125 µg/L.	[177]
Li et al., 2016	Rat	SD	Oral gavage	GD 3-GD 19 PND 3-PND 19	100 mg/kg	PND 23 PND 23	Female rats were exposed to BPAF (100 mg/kg) orally during gestation or lactation. HPLC-MS/MS analysis showed that BPAF was transferred via cord blood and lactation, finally bio-accumulating in the offspring testes. Offspring exposed to BPAF both prenatally and postnatally showed a significant increase in testis T levels and alterations in genes involved in cell differentiation and meiosis in testes.	[180]
Yang et al., 2014	Fish	Danio ratio	Water	28 days	0.05, 0.25 and 1 mg/L	~ PF 80d	Zebrafish exposed to BPAF at a concentration of 1 mg/L showed the acellular area in the testis with an increase in T levels in males and retardation of oocyte development in females.	[178]
Nakano et al., 2016	Mouse	ICR	<i>In vitro</i> oocyte	6, 9, 12, 15 and 18 h	2, 20, 50 and 100 µg/ml		BPAF exposure inhibited oocyte maturation at concentrations of 50 and 100 µg/ml and delayed the cell cycle at a dose of 2 µg/ml. Further BPAF treatment caused spindle abnormalities, activated the spindle assembly checkpoint but showed no difference in the MAD2 localization at a dose of 50 µg/ml.	[45]
Liang et al., 2017	Mouse		<i>In vitro</i> Spermatogonial cell line C18-4	24, 48 and 72 h	0.1, 1, 5, 10 and 25 µM		The BPAF exposure significantly decreased cell viability, altered nuclear morphology, increased gamma-H2AX expression level, and perturbed cytoskeleton and cell cycle progression with EC 20 ≈ 10 µM. Further, BPAF treatment at 25 µM induced the formation of multinucleated cells with active DNA synthesis and cells with dot-like structure.	[120]

Table 11.

TBBPA and reproductive outcomes in experimental studies.

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
ECB., 2006	Rat	SD	Oral gavage	10 weeks pre-mating period, 2 weeks mating period, gestation and lactation	10, 100 and 1000 mg/kg	After mating/weaning	In SD rats, TBBPA exposure showed no effects on organ weight and sperm quality. Serum T4 levels were decreased in males and females of the F0, and F1 generations and serum T3 levels were decreased in F0 males only at high dose groups. No change in FSH level was observed.	[189]
Tada et al., 2006	Mice	ICR	Dietary exposure	GD 0-PND 27	0.01%, 0.1% or 1.0% in diet	PND 27	In ICR mice receiving maternal dietary exposure through GD 0 to PND 27, no changes in testes, ovary or uterine weight were observed.	[195]
Kuiper, et al., 2007	Fish	Danio ratio	Water	30 days 47 days	0.023-1.5 µM	Adult Offspring ph.47 d	Adult zebrafish were exposed to waterborne TBBPA for 30 days, and their offspring received TBBPA exposure up to 47 days. In F0 female fish, TBBPA exposure decreased egg production starting at 0.047 µM and increased premature oocytes at a dose of 1.5 µM. In offspring, exposure to 6 µM TBBPA resulted in embryo malformation.	[194]
Van der Ven, et al., 2008	Rat	Wistar	Dietary exposure	Paternal and maternal exposure started 10 and 2 weeks before mating and ended in lactation	3, 10, 30, 100, 300, 1,000 and 3,000 mg/kg	Adult	Maternal and paternal TBBPA exposure resulted in decreases in circulating T4 level and testes weight, increase in circulating T3 level at male offspring, and a decrease in circulating T4 in female offspring. In addition, the concentrations of TBBPA metabolites in plasma increased with increasing exposure levels.	[188]
Saegusa, et al., 2009	Rat	SD	Dietary exposure	GD 10-PND 20	100, 1000 or 10,000 ppm	PND 77	In SD rats receiving the prenatal and postnatal exposure to TBBPA, no changes of testes, ovaries and uterus weights were observed. Decreases in serum T3 levels were observed only at low [high?] exposure levels.	[204]
Zatecka, et al., 2013	Mouse	CD1	Drinking water	Life time	35 µg/kg	PND 70	In CD-1 mice, two-generation exposure to TBBPA resulted in decreases in testes weight, seminal vesicles weight and expression of the gene for the androgen receptor. Further, TBBPA exposure increased in apoptotic cells and apoptotic genes expressions in seminiferous tubules. However, no changes in sperm parameters were observed.	[191]
Zatecka, et al., 2014	Mouse	C57BL/6J	Drinking water	GD 0-PND 70	35 µg/kg	PND 70	C57BL/6J mice were exposed to TBBPA during the GD 0 to PND 70. The TBBPA treatment significantly decreased protamine 1/protamine 2 the ratio increased total protamine/DNA ratio and increased number of TUNEL positive spermatozoa.	[192]
Cope, et al., 2015	Rat	SD	Oral gavage	GD 0-GD 9	100, 300 and 1000 mg/kg	PND 60	In SD rats, exposure to 100 mg/kg TBBPA resulted in a decrease in circulating T4 levels in rats. No changes in sperm motility, concentration, abnormal sperm percentage were observed in F0/1/2 generations.	[190]
Dunnick, et al., 2015	Mouse	Wistar Han rats /	Oral gavage	2 years	250, 500 and 1000 mg/kg	Adult	In 2-year TBBPA oral exposure, TBBPA increased uterine epithelial tumors including adenomas, adenocarcinomas, and	[193]

Source	Animal	Strain	Exposure route	Time of exposure	Doses	Age at collection	Summary	Reference
		B6C3F1/ N					malignant mixed Mullerian tumors in rats at doses of 500 and 1000 mg/kg. In the testes of treated male rats, atrophy of the germinal epithelium and testicular interstitial cell adenomas was observed (no significance).	
Ogunbayo et al., 2008	Mouse		<i>In vitro</i> Sertoli cell line TM4	0-18 h	0-60 μ M		In mouse Sertoli cells TM4, TBBPA treatment significantly increased cell death starting at a dose of 10 μ M and cellular Ca^{2+} levels starting at a dose of 5 μ M. Further, TBBPA treatment inhibited sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPases starting at a dose of 0.4 μ M and activated the Ryanodine receptor Ca^{2+} channel starting at a dose of 2 μ M.	[197]
Dankers et al., 2013	Mouse		<i>In vitro</i> Leydig cell line MA-10	24 h	0-30 μ M		TBBPA exposure significantly increased T levels at a dose of 10 and 30 μ M via multidrug resistance proteins. Further, 10 μ M TBBPA exposure significantly induced the expression levels of StAR, Cyp11A1, and Cyp17 in Leydig cell line MA-10.	[196]
Roelofs et al., 2015	Mouse		<i>In vitro</i> Leydig cell line MA-10	48 h	0.01–100 μ M		TBBPA exposure significantly increased T levels at a dose of 30 μ M, P4 levels and gene (SaRed1) at a dose of 10 μ M in the MA-10 cells.	[167]
Liang et al., 2017	Mouse		<i>In vitro</i> Spermatogonial cell line C18-4	24, 48 and 72 h	0.1, 1, 5, 10, and 25 μ M		The TBBPA exposure significantly decreased cell viability, altered nuclear morphology, increased gamma-H2AX expression level, and perturbed cytoskeleton and cell cycle progression at a dose of 25 μ M in spermatogonial cell line C18-4.	[120]