

## Communications

# Diversity of Bacteria Exhibiting Bile Acid-inducible 7 $\alpha$ -dehydroxylation Genes in the Human Gut

Marius Vital<sup>a,b,\*</sup>, Tatjana Rud<sup>b</sup>, Silke Rath<sup>b</sup>, Dietmar H. Pieper<sup>b,1</sup>, Dirk Schlüter<sup>a,1</sup>

<sup>a</sup> Institute for Medical Microbiology and Hospital Epidemiology, Hannover Medical School, 30625 Hannover, Germany

<sup>b</sup> Microbial Interactions and Processes Research Group, Helmholtz Centre for Infection Research, 38124 Braunschweig, Germany

## ARTICLE INFO

## Article history:

Received 24 May 2019

Received in revised form 17 July 2019

Accepted 24 July 2019

Available online 26 July 2019

## Keywords:

Bile acids

Gut microbiota

Microbiome

Metagenomics

7 $\alpha$ -dehydroxylation

Systems biology

## ABSTRACT

The secondary bile acids deoxycholic acid (DCA) and lithocholic acid (LCA), formed by gut microbiota from primary bile acids via a multi-step 7 $\alpha$ -dehydroxylation reaction, have wide-ranging effects on host metabolism and play an important role in health and disease. A few 7 $\alpha$ -dehydroxylating strains have been isolated, where bile acid-inducible (*bai*) genes were organized in a gene cluster and encoded major enzymes involved. However, only little is known on diversity and abundance of intestinal bacteria catalysing DCA/LCA formation in the human gut *in situ*. In this study, we took the opportunity to screen metagenome-assembled genomes (MAGs) from sequence data of stool samples provided by two recent studies along with newly available gut-derived isolates for the presence of the *bai* gene cluster. We revealed in total 765 and 620 MAGs encoding the potential to form DCA/LCA that grouped into 21 and 26 metagenomic species, respectively. The majority of MAGs (92.4 and 90.3%) were associated with a *Ruminococcaceae* clade that still lacks an isolate, whereas less MAGs belonged to *Lachnospiraceae* along with eight new isolates (n total = 11) that contained the *bai* genes. Only a few MAGs were linked to *Peptostreptococcaceae*. Signatures for horizontal transfer of *bai* genes were observed. This study gives a comprehensive overview of the diversity of *bai*-exhibiting bacteria in the human gut highlighting the application of metagenomics to unravel potential functions hidden from current isolates. Eventually, isolates of the identified main MAG clade are required in order to prove their capability of 7 $\alpha$ -dehydroxylating primary bile acids.

© 2019 The Authors. Published by Elsevier B.V. on behalf of Research Network of Computational and Structural Biotechnology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The primary bile acids cholic acid (CA) and chenodeoxycholic acid (CDCA) are produced from cholesterol in the liver and are subsequently conjugated to taurine or glycine residues. After their excretion into the duodenum they enable absorption of dietary lipids, cholesterol and fat-soluble vitamins that are essential for lipid metabolism and host health [1]. Additionally, bile acids act as signaling molecules regulating their own synthesis, uptake, transportation, and detoxification, and are involved in overall lipid, glucose and energy metabolisms via binding on nuclear and G-protein-coupled bile acid receptors that are expressed throughout the body [2]. The majority of secreted bile (95%) is reabsorbed along the entire gut by active transportation and passive diffusion, subsequently reconstituted in the liver and again


secreted into the duodenum, which is referred to as the enterohepatic circulation [1]. Gut microbiota directly act on bile acids substantially modifying the composition of the bile acid pool. As a first step, bacteria initiate their deconjugation via bile salt hydrolases rendering bile acids susceptible to various subsequent bacterial transformations including 7 $\alpha$ -dehydroxylation, dehydrogenation, and epimerization that lead to the generation of secondary bile acids [1,3]. Deoxycholic acid (DCA) and lithocholic acid (LCA) comprise the majority of secondary bile acids and are formed from CA and CDCA, respectively, via 7 $\alpha$ -dehydroxylation, a multi-step process that primarily occurs in the colon. Upon reabsorption DCA is reconstituted, but not rehydroxylated, which leads to its accumulation in the bile acid pool comprising a substantial part of total bile (around 25%, with large interindividual variations) [4]. In contrast, LCA is reconstituted and additionally sulfonated in the liver promoting its excretion from the body.

Secondary bile acids have wide-ranging effects on host health. On the one hand they promote disease with high levels being cytotoxic and associated with an increased risk of cholesterol gallstone disease and colon cancer [5]. Furthermore, a recent study demonstrated their role in hepatocellular carcinoma via modulating the immune system [6]. On the other hand, numerous studies described their antimicrobial

\* Corresponding author at: Institute for Medical Microbiology and Hospital Epidemiology, Hannover Medical School, OE5210, Carl-Neuberg-Str. 1, 30625 Hannover, Germany.

E-mail address: [vital.marius@mh-hannover.de](mailto:vital.marius@mh-hannover.de) (M. Vital).

<sup>1</sup> Both authors contributed equally.



**Fig. 1.** Approximately-Maximum-Likelihood tree of *bai*-gene-containing reference genomes based on 92 housekeeping genes. In blue, metagenome-derived species-level genome bins (SGBs) of study A are depicted, whereas SGBs from study B are shown in red. Abundances, i.e., number of *bai*-gene-containing metagenome-assembled genomes (MAGs) associated with individual SGBs, are shown on the right. Names of isolates are displayed in bold with those highlighted in gold derived from recent isolation efforts. Names shown in grey represent gut-derived isolates and SGBs not exhibiting the *bai* gene cluster (only high quality SGBs were considered).

effects highlighting their ability to provide colonization resistance against *Clostridioides difficile* [c.f. 7].

Enzymes involved in 7 $\alpha$ -dehydroxylation are encoded by bile acid inducible (*bai*) genes that were previously identified in a few strains of *Lachnospiraceae* and *Peptostreptococcaceae* [1,8]. However, despite their pivotal role for host physiology, only little is known on diversity of 7 $\alpha$ -dehydroxylating bacteria in the human gut *in situ*. Our recent, extensive survey on publicly available metagenomic/metatranscriptomic datasets suggested that the *bai* gene cluster is present and expressed in most individuals, yet only in a small fraction (<1%) of total intestinal bacteria [8]. Its main representative was an uncultivated member of the order *Clostridiales*, namely, *Firmicutes bacterium* CAG:103 that recruited 63.9% of all *bai*-associated reads, which displayed high amino acid sequence similarities to the reference. Only a minor portion was linked to *Lachnospiraceae* (4.7%) and *Peptostreptococcaceae* (1.9%), which include the species *Clostridium scindens* and *Clostridium hiranonis*, respectively.

## 2. Results & Discussion

Recently, vast amounts of metagenome-assembled genomes (MAGs) of gut bacteria became available from two signature studies, namely, Pasolli et al. [9] and Almeida et al. [10], which are referred to as study A and B, respectively, in the following text. Furthermore, efforts from Forster et al. [11] and Zou et al. [12] substantially increased cultured representatives of intestinal bacteria. We took the opportunity to screen for *bai*-containing genomes in those references in order to get a comprehensive overview of bacteria exhibiting this crucial function in the gut environment and to expand current reference sequences. Eleven new isolates containing the gene cluster were identified (Fig. 1). For metagenomic-based data, both studies yielded very similar results with 765 and 620 *bai*-exhibiting MAGs obtained from study A and B, respectively, which represented 0.51% (A) and 0.73% (B) of all stool-associated MAGs in these studies (excluding infants, which were devoid of *bai*-containing MAGs), and clustered into 21 (A) and 26 (B) species-level genome bins (SGBs) (Fig. 1). For detailed distribution of *bai*-containing bacteria in human gut microbiota, i.e., their total abundance and abundances of individual clades, we want to refer the reader to our previous report [8]. Most *bai*-exhibiting MAGs were associated with *Ruminococcaceae* – 92.4 and 90.3% of all *bai*-containing MAGs were associated with this family in study A and B, respectively. The vast majority of these MAGs, 97.5% (A) and 98.2% (B), were closely related to the previously identified metagenome-derived *Firmicutes bacterium* CAG:103, with the bulk harboring all *bai* genes (*baiA-I*). Despite recent isolation efforts, this clade still lacks a cultured representative; most closely related gut isolates were the *Ruminococcaceae* genera *Oscillibacter*, *Intestinimonas*, and *Pseudoflavonifractor*. BLASTing of selected housekeeping genes from main SGBs against NCBI's non-redundant protein database revealed the same genera as mentioned above as their closest relatives. It should be mentioned that although SGBs, which surround *Firmicutes bacterium* CAG:103 in the phylogenetic tree shown in Fig. 1, formed a functionally coherent group, we detected several medium quality SGBs, i.e., genome bins with completeness <90% and/or contamination >5%, that were devoid of the *bai* gene cluster and interleaved with *bai*-containing SGBs (data not shown). Furthermore, major SGBs contained many non *bai*-exhibiting MAGs, where, for instance, only 66.2 and 66.1% of total MAGs of the SGBs CosteaPI\_2017\_\_SID713B026-11-90-0\_\_bin.59 and CosteaPI\_2017\_\_SID713A023-11-0-0\_\_bin.14, respectively, harbored the target genes (no data is available for study B). It is, thus, questionable whether this clade is truly functionally consistent and care should be taken based on analyses using SGBs as representatives for *bai*-containing bacteria, since this might lead to overestimations in their abundance. Only 55 (A) and 56 (B) MAGs were associated with *Lachnospiraceae*, whereas the majority of new isolates belonged to this family. MAGs from the *Lachnospiraceae* formed two main clades intermitted by non *bai*-exhibiting *Dorea* species. A tiny fraction of MAGs was

associated with *Peptostreptococcaceae*. Phylogenetic analysis of *bai* genes showed signatures of horizontal gene transfer, where *Lachnospiraceae*-associated *bai* genes related to those identified in *Dorea* sp. AM58-8 grouped with sequences of the main *Ruminococcaceae* clade that contained the majority of MAGs (Fig. S1). *Bai* genes of the *Peptostreptococcaceae* related to those of *C. hiranonis* clustered in-between genes of the *Lachnospiraceae* species *C. scindens* and *C. hylemonae*, respectively. Genes of members associated with *C. sordelli* formed an outgroup. While *baiA-I* catalyze the oxidative arm of 7 $\alpha$ -dehydroxylation, enzymes of the reductive arm are largely unknown [1]. Recently, a flavoprotein (*baiN*) isolated from *C. scindens* was suggested to play a role in the reductive reactions [13]. Screening for *baiN* in the present study did yield hits in most genomes, however, amino acid identities were often low, even for *C. hiranonis*, a verified 7 $\alpha$ -dehydroxylating bacterium (Table S1). Furthermore, also non *baiA-I*-containing taxa exhibited similar genes, which was already shown in the original publication [13]. We could, hence, not convincingly point *baiN* homologous gene sequences outside of the main *Lachnospiraceae* clade that includes *C. scindens* and *C. hylemonae* and more detailed investigations, including biochemical testing, needs to be performed in order to unravel enzymes encoding the reductive arm in *baiA-I*-exhibiting bacteria revealed in this study.

In conclusion, this study gives an extensive overview of the diversity of *bai*-exhibiting bacteria in the human gut and highlights the application of metagenomics to unravel potential functions hidden from current isolates. Obtained reference sequences will assist guided isolation of target bacteria enabling *in vitro* experiments to validate 7 $\alpha$ -dehydroxylation of primary bile acids.

## 3. Materials and Methods

Metagenome-assembled genomes (MAGs) from study A [9] and B [10] as well as genomes of isolates from Zou et al. [12] were downloaded, subjected to prokka (v. 1.13.3, default mode) [14], and screened for individual *bai* genes (*baiA-I*) using Hidden Markov Chain Models (HMM) as described previously [8]. Protein sequence score cut-offs were set at 50% of the lowest protein reference from our previous database [8] and all genomes exhibiting  $\geq 4$  genes in synteny (defined as being separated by  $\leq 10$  genes based on locus tag) were selected as candidates. Manual inspections were performed for all genes based on phylogenetic trees. Study A provided association of each MAG with its representative species-level genome bin (SGB) representing all MAGs spanning a 95% genetic similarity. For study B, genetic distances of all *bai*-exhibiting MAGs were calculated using Mash (v. v. 2.1.1, option “-s 1e4” for sketching) [15] that were subsequently clustered into SGBs ( $\geq 95\%$  genetic similarity) using hierarchical clustering in R (v. 3.5.2, function `stats::hclust` followed by function `stats::cutree`). Raw data of new isolates from Forster et al. (2019) [11] were downloaded, quality filtered using fastp (v. 0.20.0, options “-5 20 -3 20 -1 70”) [15], and assembled via SPAdes on paired-end read mode (v. 3.13.0, option “-careful”) [16]. Contigs were then subjected to prokka before screening with HMMs. All reference gene sequences are available at <https://www.pathofunctions.com>. The tree shown in Fig. 1 was constructed from 92 housekeeping genes using UBCG (v. 3.0, default mode) [17], whereas concatenated sequences of *baiCD,E,H*, that were found in most genomes, were used to construct the tree in Fig. S1 applying FastTree (v. 2.1.10, default mode) [18]. Both trees were visualized with ggtree (v. 1.14.6) [19]. Screening for *baiN* was performed by BLASTing reference sequences (EDS08212.1, ZP\_03776912.1) against all genomes shown in Fig. 1 using DIAMOND (v. 0.9.24) [20]; only top hits were recorded.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.csbj.2019.07.012>.

## Declaration of Competing Interest

There is no conflict of interest.

## Acknowledgements

This work was funded by RESIST (Excellence Cluster), 1.7.2019–30.6.2022, and by the Helmholtz Association's Initiatives on Personalized Medicine (iMed) and Aging and Metabolic Programming (AMPro).

## References

- [1] Ridlon JM, Harris SC, Bhowmik S, Kang DJ, Hylemon PB. Consequences of bile salt biotransformations by intestinal bacteria. *Gut Microbes* 2016;7:22–39.
- [2] Li T, Chiang JYL. Bile acid signaling in metabolic disease and drug therapy. *Pharmacol Rev* 2014;66:948–83.
- [3] Foley MH, O'Flaherty S, Barrangou R, Theriot CM. Bile salt hydrolases: gatekeepers of bile acid metabolism and host-microbiome crosstalk in the gastrointestinal tract. *PLoS Pathog* 2019;15:e1007581.
- [4] Ridlon JM, Kang D-J, Hylemon PB. Bile salt biotransformations by human intestinal bacteria. *J Lipid Res* 2006;47:241–59.
- [5] Wells JE, Hylemon PB. Identification and characterization of a bile acid 7 $\alpha$ -dehydroxylation operon in *Clostridium* sp. strain TO-931, a highly active 7 $\alpha$ -dehydroxylating strain isolated from human feces. *Appl Environ Microbiol* 2000;66:1107–13.
- [6] Ma C, Han M, Heinrich B, Fu Q, Zhang Q, Sandhu M, et al. Gut microbiome – mediated bile acid metabolism regulates liver cancer via NKT cells. *Science* 2018;360:1–9.
- [7] Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gouberne A, et al. Precision microbiome reconstitution restores bile acid mediated resistance to *Clostridium difficile*. *Nature* 2014;517:205–8.
- [8] Rath S, Rud T, Karch A, Pieper DH, Vital M. Pathogenic functions of host microbiota. *Microbiome* 2018;6(174):1–13.
- [9] Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, et al. Extensive unexplored human microbiome diversity revealed by over 150,000 genomes from metagenomes spanning age, geography, and lifestyle. *Cell* 2019;176:649–62.
- [10] Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, et al. A new genomic blueprint of the human gut microbiota. *Nature* 2019;499:504.
- [11] Forster SC, Kumar N, Anonye BO, Almeida A, Viciani E, Stares MD, et al. A human gut bacterial genome and culture collection for improved metagenomic analyses. *Nat Biotechnol* 2019;37:186–92.
- [12] Zou Y, Xue W, Luo G, Deng Z, Qin P, Guo R, et al. 1,520 reference genomes from cultivated human gut bacteria enable functional microbiome analyses. *Nat Biotechnol* 2019;37:179–85.
- [13] Harris SC, Devendran S, Alves JMP, Mythen SM, Hylemon PB, Ridlon JM. Identification of a gene encoding a flavoprotein involved in bile acid metabolism by the human gut bacterium *Clostridium scindens* ATCC 35704. *Biochim Biophys Acta Mol Cell Biol Lipids* 2018;1863:276–83.
- [14] Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–9.
- [15] Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol* 2016;17:1–14.
- [16] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012;19:455–77.
- [17] Na SI, Kim YO, Yoon SH, Ha S, Baek I, Chun J. UBCG: up-to-date bacterial core gene set and pipeline for phylogenomic tree reconstruction. *J Microbiol* 2018;56:281–5.
- [18] Price MN, Dehal PS, Arkin AP. FastTree 2 - approximately maximum-likelihood trees for large alignments. *PLoS One* 2010;5:e9490.
- [19] Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. Ggtree: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol Evol* 2017;8:28–36.
- [20] Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat Methods* 2015;12:59–60.