



Published in final edited form as:

Metalomics. 2019 August 01; 11(8): 1430–1440. doi:10.1039/c9mt00042a.

Copper chaperone ATOX1 is required for MAPK signaling and growth in *BRAF* mutation-positive melanoma

Ye-Jin Kim^a, Gavin J. Bond^b, Tiffany Tsang^c, Jessica M. Posimo^a, Luca Busino^{a,d}, Donita C. Brady^{a,d,*}

^aDepartment of Cancer Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

^bBiochemistry Major Program, Department of Chemistry, College of Arts and Sciences, University of Pennsylvania, Philadelphia, PA, 19104, USA.

^cCell and Molecular Biology Graduate Group, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

^dAbramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

Abstract

Copper (Cu) is a tightly regulated micronutrient that functions as a structural or catalytic cofactor for specific proteins essential for a diverse array of biological processes. While the study of the extremely rare genetic diseases, Menkes and Wilson, has highlighted the requirement for proper Cu acquisition and elimination in biological systems for cellular growth and proliferation, the importance of dedicated Cu transport systems, like the Cu chaperones ATOX1 and CCS, in the pathophysiology of cancer is not well defined. We found that *ATOX1* was significantly overexpressed in human blood, breast, and skin cancer samples, while *CCS* was significantly altered in human brain, liver, ovarian, and prostate cancer when compared to normal tissue. Further analysis of genetic expression data in Cancer Cell Line Encyclopedia (CCLE) revealed that *ATOX1* is highly expressed in melanoma cell lines over other cancer cell lines. We previously found that Cu is required for BRAF^{V600E}-driven MAPK signaling and melanomagenesis. Here we show that genetic loss of *ATOX1* decreased BRAF^{V600E}-dependent growth and signaling in human melanoma cell lines. Pharmacological inhibition of ATOX1 with a small molecule, DCAC50, decreased the phosphorylation of ERK1/2 and reduced the growth of *BRAF* mutation-positive melanoma cell lines in a dose-dependent manner. Taken together, these results suggest that targeting the Cu chaperone ATOX1 as a novel therapeutic angle in BRAF^{V600E}-driven melanomas.

* bradyd@penmedicine.upenn.edu.

Conflicts of Interest

D.C.B holds ownership in Merlon Inc. D.C.B. is an inventor on the patent application 20150017261 entitled “Methods of treating and preventing cancer by disrupting the binding of copper in the MAP kinase pathway”. No potential conflicts of interest were disclosed by the other authors.

Introduction

While growth factors and nutrients are well-appreciated regulators of cell proliferation, the contribution of transition metals to pro-proliferative cellular processes is understudied. The transition metal Cu is an essential micronutrient for most living organisms by regulating a diverse array of biological processes, such as cellular respiration, free radical detoxification, pigmentation, kinase signal transduction, lipolysis, and neuropeptide processing¹⁻³. However, excess Cu accumulation is toxic in all organisms tested due, in large part, to free Cu ions reacting with hydrogen peroxide to form hydroxyl radicals⁴. The importance of intact Cu homeostatic mechanisms to cell growth control is underscored by the various growth phenotypes associated with aberrant Cu excretion and absorption in Menkes and Wilson disease, respectively^{1,4-9}. Therefore, biological systems depend upon dedicated machinery for the proper acquisition, distribution, and elimination of Cu¹⁰.

In mammals, the majority of Cu acquisition is achieved through the high affinity Cu¹⁺ transport function of the transmembrane protein CTR1^{11,12}. Evolutionarily conserved Cu chaperones, ATOX1, CCS, and COX17, facilitate the efficient delivery of Cu to cuproenzymes located in the trans-Golgi network, cytosol, or mitochondria¹³⁻¹⁹, while the large quantities of molecules like glutathione (GSH) and metallothionein limit cellular toxicity from free Cu ions²⁰. Specifically, the redox sensitive Cu chaperone ATOX1 delivers Cu to the P-type ATPases ATP7A or ATP7B, depending on the cell type, which facilitate the transport of Cu into the lumen of the trans-Golgi network where Cu loading of Cu-dependent enzymes, like tyrosinase, lysyl oxidase, and ceruloplasmin, occurs²¹⁻²³. Within the cytosol, CCS activates Cu,Zn-superoxide dismutase (SOD1) through direct insertion of the Cu cofactor required for catalyzing the disproportionation of superoxide to hydrogen peroxide and dioxygen^{24,25}. Finally, COX17 shuttles Cu within the inner membrane space of the mitochondria and utilizes two accessory proteins, COX11 and SCO1, to metallate the CuA and CuB sites of cytochrome c oxidase^{13,26-28}. While the physiological relevance of these Cu chaperones has been elucidated through the study of Menkes and Wilson disease, the contribution of dedicated Cu delivery systems in the context of additional pathophysiologies such as cancer, neurodegeneration, fatty liver disease, impaired wound healing, and cardiovascular disease remains to be explored.

Within the context of tumor initiation and tumor progression, some cancers may increase serum Cu levels several years prior to its diagnosis and enhanced mortality in cancer patients is associated with higher levels of Cu^{29,30}, but the biological significance of these observations is not well-established. Molecularly, the expression of the Cu chaperones ATOX1, CCS, and COX17 is elevated in non-small cell lung carcinoma (NSCLC) when compared to normal lung tissue³¹. However, ATOX1 may play a more significant role in tumorigenesis and progression than other Cu chaperones³². Specifically, functional studies revealed that genetic knockdown of *ATOX1* reduced the Cu-stimulated growth of NSCLC cell lines³¹. Further supporting a role for ATOX1 in tumorigenesis, it was recently demonstrated that ATOX is highly overexpressed in breast cancer patients at both the mRNA and protein level and is functionally required for breast cancer cell migration³³. In an effort to disrupt intracellular Cu transport to block cancer cell proliferation, Wang *et al.* developed a small molecular inhibitor, DCAC50, targeting the Cu chaperones ATOX1 and CCS³⁴.

Treatment with DCAC50 selectively reduced cell viability by promoting redox stress and reducing ATP levels of both human breast and lung cancer cell lines³⁴. Together, these studies establish ATOX1 as an integral Cu chaperone necessary for various tumorigenic properties and support further investigation of the molecular mechanisms by which ATOX1 contributes to tumorigenesis.

In the present study, we assessed the expression of the Cu chaperones *ATOX1* and *CCS* across multiple tumor subtypes and found that *ATOX1* was significantly upregulated in both melanoma patient samples and cell lines and *ATOX1* overexpression correlated with poor survival. Given that our previous work demonstrated that genetic ablation of *CTR1* or treatment with Cu chelators reduced the growth BRAF^{V600E}-driven melanoma by dampening MAPK pathway activation^{35,36}, we interrogated the importance of *ATOX1* in *BRAF* mutation-positive melanoma. Here we demonstrate that targeting *ATOX1* is a novel vulnerability in *BRAF* mutation-positive melanoma by influencing oncogene addicted kinase signaling.

Methods

Data mining.

ATOX1 or *CCS* mRNA expression in tumor and normal tissue samples was obtained from the Gene Expression across Normal and Tumor tissue (GENT) web-based genome database GENT (Gene Expression across Normal and Tumors; <http://medicalgenome.kribb.re.kr/GENT/reference.php>, Korea Research Institute of Bioscience and Biotechnology). *ATOX1* and *CCS* mRNA expression in 1036 cancer cell lines was obtained from the Cancer Cell Line Encyclopedia (CCLE) database (<https://portals.broadinstitute.org/ccle> at The Broad Institute). Cancer cell lines in which mRNA expression was greater than two standard deviations from the mean were defined as low or high expression. The Cancer Genome Atlas (TCGA) skin cutaneous melanoma dataset was queried from cBioPortal for *ATOX1* mRNA expression and patient survival data was stratified into top or bottom five percent of *ATOX1* mRNA expression.

Cell culture.

A375, HPAC, SW48, and WM88 cell lines were obtained from the American Type Culture Collection and Rockland Immunochemicals, respectively. HEK-HT cells were described previously³⁷. Parental cell lines were cultured in Dulbecco's Modified Eagle Media (DMEM, Gibco) supplemented with 10% v/v fetal bovine serum (FBS, GE Lifesciences) and 100 U/mL penicillin and streptomycin (P/S) antibiotics (Gibco). A375 and WM88 melanoma cells stably expressing pLenti-CRISPRV2puro constructs were maintained as above supplemented with 2ug/mL puromycin (Invitrogen). A375 and WM88 melanoma cell lines were stably infected with lentiviruses derived from pLentiCRISPRV2 (see plasmids below) using established protocols.

Plasmids.

pLenti-CRISPRV2puro (Addgene plasmid #52961) was previously described. pLenti-CRISPRV2puro-*Rosa26*-sgRNA was created to express the mouse *Rosa26* target sequence

5'-CCCGATCCCCTACCTAGCCG. pLentiCRISPRV2puro-*ATOX1*- sgRNA #2 was created to express the human *ATOX1* target sequence 5'- TCAGAAGCACGAGTTCTCTG. pLentiCRISPRV2puro- *ATOX1*- sgRNA #4 was created to express the human *ATOX1* target sequence 5'- TCAATGCAGACCTTCTTGTT.

Colony formation assay.

A375 and WM88 cells stably expressing pLenti-CRISPRV2puro constructs were seeded at 5.0×10^3 cells per well in 6-well plates. After seven days cells were washed once with 1X Phosphate Buffered Saline (PBS) and stained with 1mL of crystal violet staining solution (0.5% w/v crystal violet in 25% v/v methanol) for 15 minutes. After 15 minutes all wells were washed in distilled water until background staining was minimal. Crystal violet stained colonies were imaged on the ChemiDoc Touch Imaging System (Bio-Rad) imager prior to dissolving in 10% acetic acid for 15 minutes and subsequent measurement of absorbance at 590nm (Synergy, BioTek). For DCAC50 treatments, A375 (5.0×10^3), HEK-HT (1.0×10^3), HPAC (5.0×10^3), SW48 (2.0×10^4), and WM88 (5.0×10^3) cells were seeded at the indicated densities per well in 6-well plates and treated 24 hours after with vehicle or a final concentration of 1, 2.5, 5, 10, or 25 μ M DCAC50 (Sigma) for seven days. After 7 days, colonies were processed as above.

Immunoblot analysis.

A375 and WM88 parental cells or those stably expressing pLentiCRISPRv2 constructs were treated as indicated and washed with cold PBS and lysed with cold RIPA buffer containing 1X EDTA-free Halt™ protease and phosphatase inhibitor cocktail halt protease and phosphatase inhibitors (Thermo Scientific). Protein concentration was determined by BCA Protein Assay (Pierce) using BSA as a standard. Equal amount of lysates were resolved by SDS-PAGE using standard techniques, and protein was detected with the following primary antibodies: mouse anti-ATOX (1:1000; WH0000475M1, Sigma Aldrich), mouse anti- β -ACTIN (1:10000; 3700, Cell Signaling), rabbit anti-phospho(Thr202/Tyr204)-ERK1/2 (1:1000; 9101, Cell Signaling), mouse anti-ERK1/2 (1:1000; 9107, Cell Signaling), rabbit anti-phospho(Ser217/221)-MEK1/2 (1:1000; 9154S, Cell Signaling), anti-mouse MEK1/2 (1:1000; 4694S, Cell Signaling), rabbit anti-phospho(Thr180/Tyr182)-p38 (1:1000; 9211S, Cell Signaling), rabbit anti-p38 (1:1000; 9212S, Cell Signaling), rabbit anti-phospho(Thr183/Tyr185)-JNK (1:1000; 9251S, Cell Signaling), rabbit anti-JNK (1:1000; 9252S, Cell Signaling), rabbit anti-LAMIN A/C (1:1000; 2032S, Cell Signaling), or mouse anti- α -TUBULIN (1:1000; 3873S, Cell Signaling) followed by detection with one of the horseradish peroxidase conjugated secondary antibodies: goat anti-rabbit IgG (1:2000; 7074, Cell Signaling) or goat anti-mouse (1:2000; 7076, Cell Signaling) using SignalFire (Cell Signaling) or SignalFire Elite ECL (Cell Signaling) detection reagents. The fold change in the ratio of phosphorylated protein to total protein was measured in Image Studio Lite (LI-COR Biosciences) software by boxing each band using the rectangular selection tool and calculating the signal of the band in pixels. The signal of the phosphorylated protein band in pixels was normalized to the signal of the total protein band in pixels. The average fold change is shown in figures. For nuclear and cytosolic fractionations, cells were plated in 10cm dish (Genesee Scientific, 25–202), washed with cold 1X PBS, and nuclear and

cytosolic fractions were separated using nuclear/cytosol fractionation kit (BioVision K266) according to manufacturer's protocols.

Statistical analyses.

Data are represented as the mean \pm SEM. The sample size (n) indicates the experimental replicates or number of samples analyzed. For all experiments, each data point analyzed was from an independent biological sample. Statistical significance was typically determined using a one-way ANOVA followed by a Tukey's multi-comparisons test or Mantel-Cox test, in which significance was determined as $P < 0.05$. All statistical analyses were performed using GraphPad Prism 7 software.

Results

***ATOX1* overexpression in melanoma correlates with poor survival**

Although the varied expression of the human Cu proteome across cancer subtypes was recently defined³², the characterization did not include all human cancers, such as melanoma. These gaps leave the involvement of the Cu transport system in various other cancers incompletely annotated and support a need for future investigation. To fill this gap, we examined the expression of Cu transport systems across multiple human cancer and normal tissues utilizing the Gene Expression across Normal and Tumor tissue (GENT) web-accessible database. We found that *ATOX1* mRNA expression was significantly higher in blood, breast, and skin cancer when compared to the expression in their respective normal tissues (Fig. 1a). Conversely, *CCS* mRNA expression was significantly increased in brain and ovarian cancers and reduced in liver and prostate cancers when compared to their respective normal tissue samples (Fig. 1b). To substantiate these findings in human cancer cell lines, we surveyed *ATOX1* and *CCS* transcript levels in the Cancer Cell Line Encyclopedia (CCLE) database, which provides detailed gene expression analysis for over 1,000 cancer cell lines. Interestingly, *ATOX1* mRNA levels were highest amongst melanoma cell lines when compared to all other cancer cell lines and tended to be several orders of magnitude higher than *CCS* mRNA levels (Fig. 1c). Since there was significant overexpression of *ATOX1* in melanoma within both datasets, we utilized The Cancer Genome Atlas (TCGA) skin cutaneous melanoma patient data to query if altered *ATOX1* mRNA expression correlated with survival outcome³⁸. Patient samples displaying elevated *ATOX1* mRNA expression significantly correlated with a shorter median survival (Fig. 1d). These data suggest a genetic link between *ATOX1* expression and the development or progression of melanoma.

Genetic deletion of *ATOX1* reduced melanoma growth

To investigate the functional importance of *ATOX1* in melanoma, we deleted *ATOX1* via CRISPR/Cas9 in human melanoma cell lines, A375 and WM88, which overexpress *ATOX1* based on CCLE data analysis (Fig. 2a and b). We choose to investigate A375 and WM88 as they harbor oncogenic mutations in the serine/threonine kinase *BRAF*, which is the most commonly mutated gene in cutaneous melanoma and underlies therapeutic targeting via the FDA-approved small molecule kinase inhibitors dabrafenib and vemurafenib³⁸⁻⁴⁴. Targeted disruption of *ATOX1* with two different sgRNAs significantly reduced the colony formation

of A375 and WM88 cells when plated at low density when compared to control (Fig. 2c–f). Thus, our results demonstrate that *BRAF* mutation-positive melanoma cell lines depend on *ATOX1* for growth, suggesting that its overexpression contributes to melanomagenesis.

Genetic deletion of *ATOX1* decreased MAPK pathway activation in *BRAF*^{V600E} mutant melanoma

In the context of melanoma, oncogenic *BRAF*^{V600E} constitutively phosphorylates and activates the MEK1 and MEK2 kinases, which in turn phosphorylate and activate the ERK1 and ERK2 kinases, hyperactivating the canonical Mitogen Activated Protein Kinase Pathway (MAPK) to promote uncontrolled melanocytes growth and cancer initiation and progression³⁹. Thus, targeting either oncogenic *BRAF*^{V600E} and the MEK1/2 kinases has proven to be a clinically effective strategy to prolong survival in metastatic melanoma^{42,45–48}. Our lab previously established Cu as an essential micronutrient for *BRAF*^{V600E}-driven MAPK signaling, tumor growth of human melanoma cell lines, and melanomagenesis in a genetically engineered mouse model^{35,36}. Capitalizing on the dependence of oncogenic *BRAF*^{V600E} for MEK1/2-mediated signaling for growth and survival⁴⁹, we investigated whether canonical MAPK pathway activation was altered in response to genetic knockout of *ATOX1*. We tested and found that stable knockout of endogenous *ATOX1* by CRISPR/Cas9 in the human melanoma cell lines A375 and WM888 specifically dampened phosphorylation of ERK1/2 (Fig. 3a–d). Despite hyperactivation of additional MAPK pathway downstream of oncogene induced stress in melanoma⁵⁰, phosphorylation of p38 and JNK were unaltered in melanoma cell lines without *ATOX1* expression (Fig. 3a–d). These data suggest that canonical MAPK pathway activation is specifically diminished in the absence of *ATOX1* and may contribute to the reduced growth of *BRAF* mutation-positive melanoma cells in the absence of *ATOX1*. Cu is required for *BRAF*^{V600E}-driven MEK1/2 kinase activity through a direct Cu-MEK1/2 interaction³⁶, but it remains to be determined whether well-established Cu transport systems participate in Cu loading of these protein kinases. To determine whether *ATOX1* supplies Cu to MEK1/2, we interrogated whether phosphorylation of ERK1/2 could be restored by treating *ATOX1* knockout melanoma cell lines with exogenous Cu. Interestingly, acute, excess Cu treatment was sufficient to stimulate the MAPK pathway above baseline, suggesting that MEK1/2 may not be fully occupied with Cu in A375 and WM88 cells (Fig. 3e–f). In the absence of *ATOX1*, we found that the reduced phosphorylation of ERK1/2 was rescued by exogenous Cu to basal levels. Thus, *ATOX1* is required for *BRAF*^{V600E}-mutant melanoma MAPK pathway activation indirectly of a supporting Cu binding of MEK1/2.

Acute inhibition of *ATOX1* reduced melanoma growth

Previous studies have suggested that Cu-chelators used in the treatment of Wilson disease could be exploited in *BRAF*^{V600E}-driven melanoma as a unique vulnerability by forestalling MAPK signaling^{35,36}. In agreement, we hypothesized that treatment with the small molecule inhibitor DCAC50, which was previously shown to inhibit *ATOX1* and CCS by blocking Cu transfer between these chaperones and downstream interacting target proteins³⁴, would acutely inhibit *ATOX1* and reduce the growth of melanoma cell lines dependent on *BRAF*^{V600E}-mediated activation of the MAPK pathway. In both A375 and WM88 melanoma cell lines, DCAC50 treatment attenuated colony formation in a dose-dependent

fashion (Fig. 4a–d). Thus, inhibition of ATOX1 with concentrations of the small molecule inhibitor DCAC50 that are known to be selective and effective in reducing the proliferation of NSCLC are equally potent in dampening the growth of BRAF mutant melanoma cell lines³⁴.

To address whether DCAC50 mechanistically reduced BRAF^{V600E}-driven melanoma cell line colony formation by solely targeting overexpressed ATOX1, A375 and WM88 melanoma cell lines stably expressing a control sgRNA or two different sgRNAs targeting *ATOX1* were treated with increasing concentrations of DCAC50 and growth at low density was evaluated (Fig. 4e–h). While DCAC50 reduced the colony formation of A375 and WM88 cells in a dose-dependent manner when plated at low density when compared to vehicle control, stable knockout of *ATOX1* with two different sgRNAs failed to further decrease the antiproliferative activity of DCAC50 at concentrations in which significant reductions in colony growth were observed in the parental lines (Fig. 4e–h). Thus, inhibition of BRAF^{V600E}-mutant melanoma cell growth by the small molecule DCAC50 is due in large part to targeting only one of its molecular targets, ATOX1.

Given these results, we hypothesized that human cancer cell lines with lower ATOX1 expression will be less sensitive to DCAC50. To test this hypothesis, we further analyzed the genetic expression data in CCLE and found that *ATOX1* expression is lower in the pancreatic cancer cell line HPAC and the colon cancer cell line SW48 when compared to the melanoma cell lines A375 and WM88 (Supplemental Fig. 1a). In contrast to the hypothesis, HPAC and SW48 colony formation was reduced following treatment with DCAC50 in a dose-dependent manner (Supplemental Fig. 1b–e). However, DCAC50 displayed negligible effects on primary human embryonic kidney cells immortalized with SV40 T-antigen and telomerase (HEK-HT) and is therefore not broadly toxic to normal cell growth (Supplemental Fig. 1f,g). Since ATOX1 is not responsible for directly delivering Cu to MEK1/2 in cells, the mechanism by which DCAC50 reduced the growth of these cancer cells remained to be determined. Several studies have reported that ATOX1 not only functions in the cytosol but also translocates to the nucleus to facilitate the transcription of genes like *CCND1* to drive cell growth and proliferation^{51–53}. We tested and found that ATOX1 is expressed at undetectable levels in several human cancer cell lines (HPAC and SW48) and an immortalized human cell line (HEK-HT) and is exclusively localized to the cytosolic fraction of A375 and WM88 melanoma cells (Supplemental Fig. 2a). Taken together, these findings suggest that *ATOX1* expression levels may not be a predictive marker of sensitivity to the small molecule DCAC50, but indicate that the growth of cancer cells is typically selectively inhibited due in part to targeting either Cu chaperone.

DCAC50 treatment attenuated MAPK pathway activation

Finally, to investigate whether acute inhibition of ATOX1 would decrease MAPK pathway activation in an analogous fashion to genetic deletion of *ATOX1*, A375 and WM88 melanoma cells were treated with increasing concentrations of DCAC50. Similar to loss of *ATOX1*, phosphorylated ERK1/2 was reduced in response to DCAC50 treatment when compared to vehicle control, while the activation of other MAPKs remained elevated (Fig. 5a–d). These results indicate that ATOX1 is required for MAPK signaling whether directly

through a mechanism by which ATOX1 interacts with MAPK proteins themselves or indirectly by disabling Cu-requiring proteins.

Discussion

We define here for the first time the landscape of differential expression for the Cu chaperones ATOX1 and CCS in human cancer, which revealed selective upregulation of *ATOX1* in the context of skin cutaneous melanoma that correlated with poorer survival in patients. These findings agree with a study from Blockhuys *et al.* that defined the human Cu proteome and uncovered ATOX1 overexpression in all molecular subtypes of breast cancer³². However, the genetic mechanism(s) driving the elevated expression of ATOX1 in breast cancer or melanoma remains to be elucidated. One intriguing hypothesis is that transcription factor dysregulation in melanoma is causing high expression of ATOX1, which is predicted to be regulated by MTF1 and cMYC-MAX both of which have been implicated in melanoma initiation and progression. Cutaneous melanoma can be classified genetically based on the frequency of mutations in *BRAF*, *RAS*, *NFI*, and triple-WT established and these four subtypes provided a framework for clinical management of the disease³⁸. While *BRAF*^{V600E}-driven melanoma is the most prevalent alteration⁴³, *ATOX1* overexpression was detected in all four genomic subtypes of cutaneous melanoma, suggesting a potential unifying mechanism for the observed upregulation and dependence across the cancer. Our findings highlight a need for additional exploration of Cu transport systems in cancer by further investigating the molecular players governing their dysregulation in hopes of uncovering new facets of the system that can be targeted therapeutically.

Functionally, we established that ATOX1 is required for the growth of *BRAF* mutation-positive melanoma and its genetic deletion or pharmacologic inhibition is associated with a dampening of oncogenic MAPK pathway activation. However, the extent to which the reduction in oncogenic kinase signaling underlies the requirement for ATOX1 in melanoma cell growth remains to be defined. Based on previous studies, several molecular mechanisms could collectively contribute to our findings. First, our recent work demonstrated that Cu directly binds to MEK1/2 to influence the amplitude of the RAF-MEK-ERK cascade in response to proliferative signals⁵⁴. Unlike Mg²⁺ that is required for ATP coordination and in turn kinase activity, Cu allosterically potentiates the ability of MEK1/2 to phosphorylate ERK1/2 in a dose-dependent manner. Further studies showed either genetically (ie. *CTRI* knockdown) or pharmacologically (ie. Cu chelators) decreasing Cu levels in the setting of MEK1/2 dependent *BRAF* mutation-positive melanomas reduced tumor growth^{35,36}. Thus, an as yet unknown, dedicated cellular delivery mechanism of Cu to MEK1/2 is likely to exist that could be mediated by ATOX1. However, we established here that reduced MAPK pathway activation in the absence of *ATOX1* could be rescued by exogenous Cu treatment (Fig. 3e–h), suggesting that ATOX1 is not directly required for Cu-mediated activation of the MAPK pathway. Second, ATOX1 traditionally functions as the Cu chaperone for delivery of Cu to the trans-Golgi network via the P-type ATPase ATP7A in melanocytes to support the Cu-loading into Cu-dependent enzymes like LOX and tyrosinase. Both Zhu *et al.* and Matsui *et al.* have provided convincing evidence that loss of ATP7A function either by genetic deletion or pharmacologic inhibition with the gastric proton pump inhibitor omeprazole can reduce oncogenic RAS-mediated transformed cell growth due in large part to elevated ROS

levels or block melanomagenesis potentially through tyrosinase degradation^{55,56}. Therefore, overexpression of ATOX1 in melanoma and other cancers may simply support the metalation of Cu-dependent enzymes localized to the secretory compartment that are essential for tumorigenesis. In support for this hypothesis, a phase II clinical trial in triple negative breast cancer patients treated with the Cu chelator tetrathiomolybdate revealed that tumor microenvironment remodeling mediated through the Cu-dependent activity of LOX was ablated^{57,58}. Finally, while several reports have suggested that ATOX1 regulates the expression of genes required for cell growth and proliferation by translocating to the nucleus^{51–53}, ATOX1 is predominately localized to the cytosolic fraction of human melanoma cell lines (Supplemental Fig. 2a). In conclusion, this study highlights the importance of exploring additional Cu homeostasis machinery in the context of cancer and more specifically, suggests that inhibiting ATOX1 with the small molecule DCAC50 may be a promising strategy for the treatment of *BRAF* mutation-positive cancer in combination with other MAPK pathway inhibitors.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank F. Daldal, A. Dancis, J.M. Davis, and E.S. Witze (University of Pennsylvania) for technical support, discussions, and/or review of the manuscript. This research was funded by the National Institutes of Health (NIH) grants GM124749 (D.C.B.) and CA207513 (L.B.) and the Pew Scholars Program in Biomedical Science Award #50359 (D.C.B.). T.T. is supported by NCI NRSA Fellowship (F31) F31CA243294. J.M.P. is supported by American Cancer Society Postdoctoral Fellowship 131203PF1714701CCG.

References

1. Kim B-E, Nevitt T and Thiele DJ, *Nat Chem Biol*, 2008, 4, 176–185. [PubMed: 18277979]
2. Thiele DJ and Gitlin JD, *Nat Chem Biol*, 2008, 4, 145–147. [PubMed: 18277968]
3. Festa RA and Thiele DJ, *Curr Biol*, 2011, 21, R877–83. [PubMed: 22075424]
4. Ala A, Walker AP, Ashkan K and Dooley JS, *Lancet*, 2007, 369, 397–408. [PubMed: 17276780]
5. Kaler SG, *Ann N Y Acad Sci*, 2014, 1314, 64–68. [PubMed: 24735419]
6. Graper ML, Huster D, Kaler SG, Lutsenko S and Thiele DJ, *Ann N Y Acad Sci*, 2014, 1314, v–vi.
7. Vulpe C, Levinson B, Whitney S, Packman S and Gitschier J, *Nature Genetics*, 1993, 3, 7–13. [PubMed: 8490659]
8. Bull PC, Thomas GR, Rommens JM, Forbes JR and Cox DW, *Nature Genetics*, 1993, 5, 327–337. [PubMed: 8298639]
9. Tanzi RE, Petrukhin K, Chernov I, Pellequer JL, Wasco W, Ross B, Romano DM, Parano E, Pavone L and Brzustowicz LM, *Nature Genetics*, 1993, 5, 344–350. [PubMed: 8298641]
10. Kaplan JH and Lutsenko S, *J Biol Chem*, 2009, 284, 25461–25465. [PubMed: 19602511]
11. Ohrvik H and Thiele DJ, *Ann N Y Acad Sci*, 2014, 1314, 32–41. [PubMed: 24697869]
12. Lee J, Prohaska JR and Thiele DJ, *Proc Natl Acad Sci USA*, 2001, 98, 6842–6847. [PubMed: 11391005]
13. Glerum DM, Shtanko A and Tzagoloff A, *J Biol Chem*, 1996, 271, 14504–14509. [PubMed: 8662933]
14. Wernimont AK, Huffman DL, Lamb AL, O'Halloran TV and Rosenzweig AC, *Nat Struct Mol Biol*, 2000, 7, 766–771.

15. Pufahl RA, Singer CP, Peariso KL, Lin SJ, Schmidt PJ, Fahrni CJ, Culotta VC, Penner-Hahn JE and O'Halloran TV, *Science*, 1997, 278, 853–856. [PubMed: 9346482]
16. Anastassopoulou Ioanna, Banci Lucia, Bertini Ivano, Cantini Francesca, Katsari A, Efthalia and Rosato A, *Solution Structure of the Apo and Copper(I)-Loaded Human Metallochaperone HAH1†*, American Chemical Society, 2004.
17. Banci Lucia, Bertini Ivano, Chasapis CT, Rosato A and Tenori L, *Biochem Biophys Res Commun*, 2007, 364, 645–649. [PubMed: 17961510]
18. Robinson NJ and Winge DR, *Annu Rev Biochem*, 2010, 79, 537–562. [PubMed: 20205585]
19. Cobine PA, Pierrel F and Winge DR, *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 2006, 1763, 759–772. [PubMed: 16631971]
20. Banci L, Bertini I, Ciofi-Baffoni S, Kozyreva T, Zovo K and Palumaa P, *Nature*, 2010, 465, 645–648. [PubMed: 20463663]
21. Hatori Y and Lutsenko S, *Antioxid Redox Signal*, 2013, 19, 945–957. [PubMed: 23249252]
22. Lutsenko S, *Metallomics*, 2016, 8, 840–852. [PubMed: 27603756]
23. Hatori Y, Clasen S, Hasan NM, Barry AN and Lutsenko S, *J Biol Chem*, 2012, 287, 26678–26687. [PubMed: 22648419]
24. Culotta VC, Klomp LWJ, Strain J, Casareno RLB, Krems B and Gitlin JD, *J Biol Chem*, 1997, 272, 23469–23472. [PubMed: 9295278]
25. Wong PC, Waggoner D, Subramaniam JR, Tessarollo L, Bartnikas TB, Culotta VC, Price DL, Rothstein J and Gitlin JD, *Proc Natl Acad Sci USA*, 2000, 97, 2886–2891. [PubMed: 10694572]
26. Horng Y-C, Leary SC, Cobine PA, Young FBJ, George GN, Shoubridge EA and Winge DR, *J Biol Chem*, 2005, 280, 34113–34122. [PubMed: 16091356]
27. Horng Y-C, Cobine PA, Maxfield AB, Carr HS and Winge DR, *J Biol Chem*, 2004, 279, 35334–35340. [PubMed: 15199057]
28. Leary SC, Kaufman BA, Pellicchia G, Guercin G-H, Mattman A, Jaksch M and Shoubridge EA, *Hum. Mol. Genet*, 2004, 13, 1839–1848. [PubMed: 15229189]
29. Coates RJ, Weiss NS, Daling JR, Rettmer RL and Warnick GR, *Cancer Res*, 1989, 49, 4353–4356. [PubMed: 2743325]
30. Brewer GJ and Merajver SD, *Integr Cancer Ther*, 2002, 1, 327–337. [PubMed: 14664727]
31. Cai H and Peng F, *Oncology Reports*, 2013, 30, 269–275. [PubMed: 23624903]
32. Blockhuys S, Celauro E, Hildesjö C, Feizi A, Stål O, Fierro-González JC and Wittung-Stafshede P, *Metallomics*, 2017, 9, 112–123. [PubMed: 27942658]
33. Blockhuys S and Wittung-Stafshede P, *Biochem Biophys Res Commun*, 2017, 483, 301–304. [PubMed: 28027931]
34. Wang J, Luo C, Shan C, You Q, Lu J, Elf S, Zhou Y, Wen Y, Vinkenborg JL, Fan J, Kang H, Lin R, Han D, Xie Y, Karpus J, Chen S, Ouyang S, Luan C, Zhang N, Ding H, Merckx M, Liu H, Chen J, Jiang H and He C, *Nat Chem*, 2015, 7, 968–979. [PubMed: 26587712]
35. Brady DC, Crowe MS, Greenberg DN and Counter CM, *Cancer research*, 2017, 77, 6240–6252. [PubMed: 28986383]
36. Brady DC, Crowe MS, Turski ML, Hobbs GA, Yao X, Chaikuad A, Knapp S, Xiao K, Campbell SL, Thiele DJ and Counter CM, *Nature*, 2014, 509, 492–496. [PubMed: 24717435]
37. Hahn WC, Counter CM, Lundberg AS, Beijersbergen RL, Brooks MW and Weinberg RA, *Nature*, 1999, 400, 464–468. [PubMed: 10440377]
38. The Cancer Genome Atlas Network, *Cell*, 2015, 161, 1681–1696. [PubMed: 26091043]
39. Holderfield M, Deuker MM, McCormick F and McMahon M, *Nat Rev Cancer*, 2014, 14, 455–467. [PubMed: 24957944]
40. Flaherty KT, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, O'Dwyer PJ, Lee RJ, Grippo JF, Nolop K and Chapman PB, *N Engl J Med*, 2010, 363, 809–819. [PubMed: 20818844]
41. Falchook GS, Long GV, Kurzrock R, Kim KB, Arkenau TH, Brown MP, Hamid O, Infante JR, Millward M, Pavlick AC, O'Day SJ, Blackman SC, Curtis CM, Lebowitz P, Ma B, Ouellet D and Kefford RF, *Lancet*, 2012, 379, 1893–1901. [PubMed: 22608338]

42. Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, Dummer R, Garbe C, Testori A, Maio M, Hogg D, Lorigan P, Lebbe C, Jouary T, Schadendorf D, Ribas A, O'Day SJ, Sosman JA, Kirkwood JM, Eggermont AMM, Dreno B, Nolop K, Li J, Nelson B, Hou J, Lee RJ, Flaherty KT, G. A. McArthurBRIM-3 Study Group, *N Engl J Med*, 2011, 364, 2507–2516. [PubMed: 21639808]
43. Pollock PM, Harper UL, Hansen KS, Yudt LM, Stark M, Robbins CM, Moses TY, Hostetter G, Wagner U, Kakareka J, Salem G, Pohida T, Heenan P, Duray P, Kallioniemi O, Hayward NK, Trent JM and Meltzer PS, *Nat Genet*, 2002, 33, 19–20. [PubMed: 12447372]
44. Bollag G, Tsai J, Zhang J, Zhang C, Ibrahim P, Nolop K and Hirth P, *Nat Rev Drug Discov*, 2012, 11, 873–886. [PubMed: 23060265]
45. Flaherty KT, Infante JR, Daud A, Gonzalez R, Kefford RF, Sosman J, Hamid O, Schuchter L, Cebon J, Ibrahim N, Kudchadkar R, Burris HA, Falchook G, Algazi A, Lewis K, Long GV, Puzanov I, Lebowitz P, Singh A, Little S, Sun P, Allred A, Ouellet D, Kim KB, Patel K and Weber J, *N Engl J Med*, 2012, 367, 1694–1703. [PubMed: 23020132]
46. Long GV, Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, Garbe C, Jouary T, Hauschild A, Grob J-J, Chiarion-Sileni V, Lebbe C, Mandalà M, Millward M, Arance A, Bondarenko I, Haanen JBAG, Hansson J, Utikal J, Ferraresi V, Kovalenko N, Mohr P, Probachai V, Schadendorf D, Nathan P, Robert C, Ribas A, Demarini DJ, Irani JG, Swann S, Legos JJ, Jin F, Mookerjee B and Flaherty K, *Lancet*, 2015, 386, 444–451. [PubMed: 26037941]
47. Infante JR, Fecher LA, Falchook GS, Nallapareddy S, Gordon MS, Becerra C, Demarini DJ, Cox DS, Xu Y, Morris SR, Peddareddigari VGR, Le NT, Hart L, Bendell JC, Eckhardt G, Kurzrock R, Flaherty K, Burris HA and Messersmith WA, *Lancet Oncol*, 2012, 13, 773–781. [PubMed: 22805291]
48. Falchook GS, Lewis KD, Infante JR, Gordon MS, Vogelzang NJ, Demarini DJ, Sun P, Moy C, Szabo SA, Roadcap LT, Peddareddigari VGR, Lebowitz PF, Le NT, Burris HA, Messersmith WA, O'Dwyer PJ, Kim KB, Flaherty K, Bendell JC, Gonzalez R, Kurzrock R and Fecher LA, *Lancet Oncol*, 2012, 13, 782–789. [PubMed: 22805292]
49. Solit DB, Garraway LA, Pratilas CA, Sawai A, Getz G, Basso A, Ye Q, Lobo JM, She Y, Osman I, Golub TR, Sebolt-Leopold J, Sellers WR and Rosen N, *Nature*, 2006, 439, 358–362. [PubMed: 16273091]
50. Corazzari M, Rapino F, Ciccocanti F, Giglio P, Antonioli M, Conti B, Fimia GM, Lovat PE and Piacentini M, *Cell Death Differ*, 2015, 22, 946–958. [PubMed: 25361077]
51. Itoh S, Kim HW, Nakagawa O, Ozumi K, Lessner SM, Aoki H, Akram K, McKinney RD, Ushio-Fukai M and Fukai T, *J Biol Chem*, 2008, 283, 9157–9167. [PubMed: 18245776]
52. Ohrvik H and Wittung-Stafshede P, *Int J Mol Sci*, 2015, 16, 16728–16739. [PubMed: 26213915]
53. Matson Dzebo M, Blockhuys S, Valenzuela S, Celauro E, Esbjörner EK and Wittung-Stafshede P, *Comput Struct Biotechnol J*, 2018, 16, 443–449. [PubMed: 30455854]
54. Turski ML, Brady DC, Kim HJ, Kim B-E, Nose Y, Counter CM, Winge DR and Thiele DJ, *Mol Cell Biol*, 2012, 32, 1284–1295. [PubMed: 22290441]
55. Zhu S, Shanbhag V, Wang Y, Lee J and Petris M, *J Cancer*, 2017, 8, 1952–1958. [PubMed: 28819394]
56. Matsui MS, Petris MJ, Niki Y, Karaman-Jurukovska N, Muizzuddin N, Ichihashi M and Yarosh DB, *Journal of Investigative Dermatology*, 2015, 135, 834–841. [PubMed: 25337692]
57. Chan N, Willis A, Kornhauser N, Ward MM, Lee SB, Nackos E, Seo BR, Chuang E, Cigler T, Moore A, Donovan D, Cobham MV, Fitzpatrick V, Schneider S, Wiener A, Guillaume-Abraham J, Aljom E, Zelkowitz R, Warren JD, Lane ME, Fischbach C, Mittal V and Vahdat L, *Clin Cancer Res*, 2016, 23, 666–676. [PubMed: 27769988]
58. Jain S, Cohen J, Ward MM, Kornhauser N, Chuang E, Cigler T, Moore A, Donovan D, Lam C, Cobham MV, Schneider S, Hurtado Rúa SM, Benkert S, Mathijssen Greenwood C, Zelkowitz R, Warren JD, Lane ME, Mittal V, Rafii S and Vahdat LT, *Ann Oncol*, 2013, 24, 1491–1498. [PubMed: 23406736]

Significance to metallomics

Maintaining Cu homeostasis is critical for the function of Cu-dependent enzymes that are required for normal physiology and development. While aberrant Cu accumulation is associated with cancer initiation and progression, the Cu homeostasis machinery that contributes to cell intrinsic and extrinsic mechanisms necessary for tumorigenesis remains largely undefined. We present here an analysis of Cu chaperone expression across human cancers that illuminated a novel dependence on *ATOX1* for cancer cell growth in the context of *BRAF* mutation-positive melanoma. Our findings connect intracellular Cu transport via *ATOX1* to oncogenic kinase signaling that could be leveraged as a novel therapeutic opportunity in melanoma.




Figure 1. *ATOX1* overexpression in melanoma correlates with poor survival.

a,b, mRNA expression of *ATOX1* (**a**) or *CCS* (**b**) from tumor or normal tissue samples in the GENT web-based database. Statistical analysis was performed using a one-way ANOVA followed by Tukey's multiple-comparison test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

c, log₂ mRNA expression of *ATOX1* versus *CCS* in 1036 cancer cell lines from the CCLE database. Non-melanoma cell lines (n=975), melanoma cell lines (n=61), or A375 and WM88 are indicated with gray, red, or black dots, respectively. Dashed lines indicate two standard deviations from mean mRNA expression and were used as cut-off values to designate low (left, bottom), normal (middle), or high expression (right, top). **d**, Kaplan-Meier survival analysis of TCGA skin cutaneous melanoma patients with high *ATOX1* (red line, n= 23) or low *ATOX1* (black line, n=23) mRNA expression. Results were compared using a Mantel-Cox test. ** $P < 0.01$.




Figure 2. Genetic deletion of *ATOX1* reduces colony formation of *BRAF* mutation-positive melanoma cell lines.

a,b, Immunoblot detection of *ATOX1* or β -ACTIN from A375 or WM88 melanoma cells stably expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4). **c,d**, Representative images of crystal violet stained colonies from A375 or WM88 melanoma cells stably expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4). **e,f**, Quantification of crystal violet staining from three independent experiments plated in triplicate represented as normalized percent growth \pm SEM of A375 or WM88 melanoma cells expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4). Statistical analysis was performed using a one-way ANOVA followed by Tukey's multiple-comparisons test. ** $P < 0.01$, **** $P < 0.0001$.




Figure 3. Genetic deletion of *ATOX1* decreases MAPK pathway activation.

a,b, Immunoblot detection of P-ERK1/2, T-ERK1/2, P-MEK1/2, T-MEK1/2, P-p38, T-p38, P-JNK, T-JNK, or β-ACTIN from A375 or WM88 melanoma cells stably expressing sgRNA against *ROSA* (–) or *ATOX1* (#2 or #4). Quantification: P-ERK1/2/T-ERK1/2 normalized to *ROSA* control. **c,d**, Quantification of the ratio of P-ERK1/2 to T-ERK1/2 normalized to *ROSA* control ± SEM in A375 and WM88 melanoma cells stably expressing sgRNA against *ROSA* (–) or *ATOX1* (#2 or #4) from three independent biological replicates. Statistical analysis was performed using a one-way ANOVA followed by Tukey’s multiple-

comparisons test. * $P < 0.05$, *** $P < 0.001$. **e,f**, Immunoblot detection of P-ERK1/2, T-ERK1/2, or β -ACTIN from A375 or WM88 melanoma cells stably expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4) treated with vehicle or 10 μ M CuCl₂ for 20 minutes. Quantification: P-ERK1/2/T-ERK1/2 normalized to *ROSA* control. **g,h**, Quantification of the ratio of P-ERK1/2 to T-ERK1/2 normalized to *ROSA* control \pm SEM in A375 and WM88 melanoma cells stably expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4) treated with vehicle or 10 μ M CuCl₂ for 20 minutes from three independent biological replicates. Statistical analysis was performed using a one-way ANOVA followed by Tukey's multiple-comparisons test. ns = non-significant, * $P < 0.05$.




Figure 4. Acute inhibition of ATOX1 reduces melanoma growth.

a,b, Representative images of crystal violet stained colonies from A375 or WM88 melanoma cells treated with vehicle or increasing concentrations of DCAC50. **c,d**, Quantification of crystal violet staining from three independent experiments plated in triplicate represented as normalized percent growth \pm SEM of A375 or WM88 melanoma cells treated with vehicle or increasing concentrations of DCAC50. Statistical analysis was performed using a one-way ANOVA followed by Tukey's multiple-comparisons test. * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$. **e,f**, Representative images of crystal violet stained colonies from A375 or WM88 melanoma cells stably expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4) treated with vehicle or increasing concentrations of DCAC50. **g,h**, Quantification of crystal violet staining from three independent experiments plated in triplicate represented as normalized percent growth \pm SEM of A375 or WM88 melanoma cells stably expressing sgRNA against *ROSA* (-) or *ATOX1* (#2 or #4) treated with vehicle

or increasing concentrations of DCAC50. Statistical analysis was performed using a one-way ANOVA followed by Tukey's multiple-comparisons test. ns = non-significant, **P<0.01, ****P < 0.0001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 5. DCAC50 treatment attenuates MAPK pathway activation.
a,b, Immunoblot detection of P-ERK1/2, T-ERK1/2, P-p38, T-p38, P-JNK, T-JNK, or β -ACTIN from A375 or WM88 melanoma cells treated with vehicle or increasing concentrations of DCAC50 (μ M) for 24 hours. Quantification: P-ERK1/2/T-ERK1/2 normalized to *ROSA* control. **c,d**, Quantification of the ratio of P-ERK1/2 to T-ERK1/2 normalized to vehicle control \pm SEM in A375 and WM88 melanoma cells treated with vehicle or increasing concentration of DCAC50 for 24 hours from three independent biological replicates.