



Published in final edited form as:

Dev Psychopathol. 2019 October ; 31(4): 1423–1437. doi:10.1017/S0954579418001013.

Effects of Victimization on Mental Health and Substance Use Trajectories in Young Sexual Minority Men

Gregory Swann, Emily Forscher, Emily Bettin, Michael E. Newcomb, Brian Mustanski

Department of Medical Social Sciences, Northwestern University Feinberg School of Medicine, Chicago, IL; Northwestern University Institute for Sexual and Gender Minority Health and Wellbeing, Chicago, IL

Abstract

Objective: Young sexual minority men (YSMM) experience more victimization and are at higher risk for mental health and substance use problems compared to heterosexual youth. We attempt to understand change over time in the experience of these constructs among YSMM.

Method: Data were taken from a diverse community-based sample of YSMM (N=450, baseline mean age 18.93) surveyed every six months for 2.5 years. Multilevel modeling was used to model within-person change in victimization, internalizing symptoms, externalizing symptoms, alcohol frequency, marijuana use, and illicit drug use. We tested the indirect effect of concurrent and time-lagged victimization on the association between age and mental health and substance use.

Results: Victimization, internalizing symptoms, and externalizing symptoms decreased over time. Concurrent victimization was associated with higher internalizing symptoms, externalizing symptoms, alcohol use, marijuana use, and illicit drug use. Analysis of indirect effects suggested that the association between victimization and mental health and substance use outcomes decreased as participants transitioned from adolescence into adulthood.

Conclusions: This study found that the reduction in victimization that YSMM experience as they grow older is associated with a reduction in negative mental health and substance use outcomes. Prevention efforts to limit victimization exposure may reduce health disparities for YSMM.

Keywords

Victimization; Gay; Substance Use; Mental Health; Internalized Stigma

INTRODUCTION

Young sexual minority men (YSMM) are at higher risk for mental health disorders (Bostwick et al., 2014; Fergusson, Horwood, & Beautrais, 1999; Hershberger & D’Augelli, 1995; King et al., 2008; McLaughlin, Hatzenbuehler, Xuan, & Conron, 2012; Remafedi, French, Story, Resnick, & Blum, 1998) and engage in substance use more frequently

compared to heterosexual male youth (Corliss, Rosario, Wypij, Fisher, & Austin, 2008; Hughes & Eliason, 2002; Marshal et al., 2008; Newcomb, Birkett, Corliss, & Mustanski, 2014; Talley, Hughes, Aranda, Birkett, & Marshal, 2014). Previous research has demonstrated an association between experiences of victimization (which include bullying, being threatened or attacked with a weapon, and physical and sexual assault) for YSM and higher symptoms of both mental health problems (Birkett, Newcomb, & Mustanski, 2015; Burton, Marshal, Chisolm, Sucato, & Friedman, 2013; Poteat & Espelage, 2007; Russell, Ryan, Toomey, Diaz, & Sanchez, 2011; Swann, Minshew, Newcomb, & Mustanski, 2016) and increased substance use (Bontempo & D'Augelli, 2002; Huebner, Thoma, & Neilands, 2015). Mental health symptoms and substance use problems have different patterns of change over the transition from adolescence to young adulthood. Mental health problems decrease over the transition for YSM (Birkett et al., 2015) and substance use issues increase (Halkitis et al., 2014; Marshal, Friedman, Stall, & Thompson, 2009; Swann, Bettin, Clifford, Newcomb, & Mustanski, 2017). However, the association that both share with victimization suggests that victimization may be one driver of mental health and substance use disparities between YSM and heterosexual young men.

Minority stress theory describes a model through which victimization and stigma-related stress results in more negative outcomes for members of minority groups, such as YSM (Hatzenbuehler, 2009; Meyer, 2003). The theory posits that sexual minorities face higher levels of bullying and victimization due to their stigmatized minority status. In turn, this minority-group-specific victimization and stigma results in more negative health outcomes, such as substance use problems and symptoms of mental health issues. Recent work has provided empirical support for this framework in LGBTQ samples, including finding that minority stress increases mental health problems via increases in rumination (Liao, Kashubeck-West, Weng, & Deitz, 2015; Schwartz, Stratton, & Hart, 2016), deficits in emotion regulation and higher general life stress (Burton, Wang, & Pachankis, 2018), and lowered self-compassion (Liao et al., 2015). The effects of minority stress are especially important to understand for YSM over the developmental period from adolescence into young adulthood. Youth are making important transitions such as the move from high school into either college or the work force over this period (Arnett & Hughes, 2012). Identity development is still occurring and for some sexual minority youth the process of incorporating their sexual identity is still ongoing (Morgan, 2013). Brain development is also continuing to occur, including development of the prefrontal cortex that is essential to impulse control and executive functioning, into a person's 20's (Lebel, Walker, Leemans, Phillips, & Beaulieu, 2008). This period of continued transition and development across multiple arenas make it a critical time for victimization and stigma-related stress to have a negative and potentially long-lasting impact.

Past research has shown that YSM are particularly vulnerable to victimization. Generally, sexual and gender minority (SGM) youth, including YSM, report high rates of victimization compared to their heterosexual peers (Bontempo & D'Augelli, 2002; Pilkington & D'Augelli, 1995; Robin et al., 2002; Russell & Joyner, 2001; Shields, Whitaker, Glassman, Franks, & Howard, 2012). The rates of victimization differ depending on the exact question asked. Bontempo and D'Augelli (2002), using data from the Youth Risk Behavior Survey (YRBS), found that 24% of gay and bisexual identifying males reported

experiencing more than 10 incidents of victimization at school in the previous year compared to 2.7% for heterosexual males. Robin and colleagues (2002), also reporting on YRBS data, found that for their two samples, percentage of youth who reported being threatened or injured with a weapon at school in the previous year was 13.4% for youth who reported having only same-sex relationships, ranged between 38.6-45.3% for youth reporting both-sex relationships, and ranged between 8.3-9.1% for youth who reported only opposite sex relationships. More recently, Shields and researchers (2012) found that 62% of LGB identifying youth reported at least one victimization indicator (threatened or injured with a weapon, bullied at school, or in a physical fight) compared to 31% of heterosexual youth.

For YSMM, victimization is strongly associated with increased endorsement of mental health problems, such as depression, internalizing symptoms, and post-traumatic stress (Birkett et al., 2015; Whitbeck, Chen, Hoyt, Tyler, & Johnson, 2004). There is also evidence that experiences of victimization can be more damaging for YSMM. Compared to heterosexual peers, YSMM with comparable levels of victimization were more likely to attempt suicide (Bontempo & D'Augelli, 2002). The association between victimization and mental health problems, coupled with their higher risk for victimization, make YSMM a particularly vulnerable group for negative mental health outcomes.

Among YSMM, victimization related to their minority identity is not only concurrently observed with higher rates of substance use, specifically, binge drinking, marijuana use, and cocaine use (Bontempo & D'Augelli, 2002; Mustanski, Andrews, Herrick, Stall, & Schnarrs, 2014; Robin et al., 2002), but victimization also serves as a risk factor for increased substance use. A meta-analysis found that victimization was one of the strongest risk factors for substance use, defined as use of alcohol, marijuana, cocaine, and ecstasy use, among YSMM (Goldbach, Tanner-Smith, Bagwell, & Dunlap, 2014). In a recent analysis of the effect of victimization on alcohol use, greater experience of victimization specific to their minority identity was significantly associated with heavy alcohol use in YSMM, even after controlling for baseline alcohol use and concurrent affiliation with substance-using peers (Dermody, Marshal, Burton, & Chisolm, 2016). This suggests that victimization is a key predictor of substance use for YSMM.

Past research has found associations between victimization, mental health problems, and substance abuse for YSMM, but what is less clear is how these relationships change over time. Birkett and colleagues (2015) followed SGM youth over the course of three and a half years. They found that victimization was strongly associated with psychological distress and that victimization mediated the developmental decline in psychological distress. In other words, the decrease in victimization over time partly explained SGM youth's similar decrease in symptoms of psychological distress. Building off this work, it is unknown if victimization will also mediate additional outcomes that have been shown to change significantly over the course of adolescence, such as the externalizing spectrum of mental health issues (Hicks et al., 2007) and substance use problems, including escalating rates of alcohol, marijuana, and hard drug use (Swann et al., 2017). Furthermore, it is also not clear how additional risk and protective factors for minority stress, such as internalized stigma and social support, might impact these associations, despite evidence linking mental health and

substance use (i.e., alcohol, marijuana, ecstasy, cocaine, opiates, and sedatives) with both internalized stigma (Hequembourg & Dearing, 2013; Livingston, Oost, Heck, & Cochran, 2015; Newcomb & Mustanski, 2010) and (lack of) social support (McConnell, Birkett, & Mustanski, 2016; Ryan, Russell, Huebner, Diaz, & Sanchez, 2010).

Internalized stigma refers to the negative impact on a person's self-concept that occurs from exposure to societal bias (Shidlo, 1994). In the case of YSMM, it is the internalization of negative attitudes and biases toward same-sex sexuality. Internalized stigma has been found to be a predictor of higher levels of internalizing mental health problems, including symptoms of depression and anxiety disorders, for sexual minorities (Feinstein, Davila, & Dyar, 2017; Kaysen et al., 2014; Newcomb & Mustanski, 2010; Pachankis, Sullivan, Feinstein, & Newcomb, 2018; Puckett, Levitt, Horne, & Hayes-Skelton, 2015; Puckett, Mereish, Levitt, Horne, & Hayes-Skelton, 2018). Hatzenbuehler's mediation framework suggests that internalized stigma also leads to higher levels of externalizing problems, including conduct problems and substance abuse, as stigma-related stress increases vulnerability in SGM to psychological processes that are generally predictive of psychopathology (Hatzenbuehler, 2009). Specifically, stigma leads to an increase in emotion dysregulation and general negative affect, which increases engagement in maladaptive coping behaviors (e.g., substance use).

Experiences of victimization and discrimination have been found to not only predict internalized stigma, but internalized stigma has also been shown to mediate the association between experiences of discrimination and mental health problems (Puckett, Newcomb, Garofalo, & Mustanski, 2016; Szymanski & Ikizler, 2013). The association between internalized stigma and alcohol and drug use is less clear, with some research studies finding a positive association (Hequembourg & Dearing, 2013; Livingston et al., 2015; Moody, Starks, Grov, & Parsons, 2018), others finding none (Amadio, 2006; Amadio & Chung, 2004; Ross et al., 2001), and at least one finding a negative association but only for LGBT individuals who were low on depressive symptoms (Span & Derby, 2009).

Whereas experiences of victimization and the internalization of stigma have been documented as risk factors for higher rates of mental health and substance use problems, social support is a potential protective factor (Davidson & Demaray, 2007). Higher perceived social support has been associated with fewer symptoms of depression and lower rates of suicidal ideation (Liu & Mustanski, 2012; McConnell et al., 2016; Ryan et al., 2010; Safren & Heimberg, 1999; Teasdale & Bradley-Engen, 2010) for LGBT youth including YSMM. There is also evidence of a similar effect of perceived social support as a protective factor against alcohol, marijuana, and illicit drug (i.e., cocaine, ecstasy, opiate) problems in more general samples of young adults (Newcomb & Bentler, 1988) and sexual minority women (Lehavot & Simoni, 2011).

For the present study, we plan to expand upon the previous research that has found an indirect effect of within-person age on psychological distress through victimization by testing whether this effect is also true for symptoms of externalizing disorders and levels of substance use. We hypothesize that victimization will mediate the association between age and mental health (both internalizing and externalizing symptoms) and substance use

outcomes both concurrently and over time. In addition, we hypothesize that gay-related stigma will be associated with higher mental health symptoms and substance use, and that it will mediate the association between victimization and our outcomes, such that as participants get older they will experience less stigma from victimization resulting in less mental health and substance use problems. Finally, we hypothesize that social support will moderate the effects of victimization on our outcomes, so that YSMM who report higher support experience fewer negative outcomes.

METHOD

Participants and Procedures

Data were collected as part of Crew 450: a longitudinal Chicago-based study examining a syndemic of psychosocial stressors associated with HIV in a cohort of 450 YSMM. To be eligible for the study, individuals were required to be between 16 and 20 years old at baseline, assigned male at birth, speak English, have reported a sexual encounter with a man or identified as gay or bisexual, and available for two years of follow up. Participants were recruited through a modified form of respondent-driven sampling (RDS) (Heckathorn, 1997) that allowed for a greater number of seeds than standard RDS (Kuhns et al., 2015; Newcomb, Ryan, Garofalo, & Mustanski, 2014). The initial convenience sample (i.e., “seeds”; $N = 172$; 38.2%) was recruited from YSMM-frequented venues, school/organizational outreach, flyers posted in the community, and geo-social network applications.

Six waves of data were included in these analyses, collected every six months over approximately 2.5 years. Retention at follow-up waves was high: 85.8%, 80.7%, 75.6%, 75.4%, and 75.6% respectively. Participants completed surveys using computer-assisted self-interview (CASI) technology during in-person visits. Compensation was provided at each wave: \$70 for baseline and \$45 for each follow-up. Individuals provided consent/assent prior to participation, and all procedures for the study were approved by the Institutional Review Boards (IRBs) of the primary investigators’ institutions with a waiver of parental permission under 45 CFR 46.408(c) (Mustanski, 2011).

The average age of the sample was 18.93 ($SD = 1.29$) at baseline. The majority of the sample identified their primary race/ethnicity as African American/Black ($N = 240$, 53.3%), followed by Hispanic/Latino ($N = 90$, 20.0%), non-Hispanic White ($N = 81$, 18.0%), participants who identified their race as “other” ($N = 24$, 5.3%), Asian ($N = 8$, 1.8%), and American Indian or Alaska native ($N = 7$, 1.6%). Most of the sample described their sexual orientation as “only gay/homosexual” ($N = 226$, 50.2%), followed by “mostly gay/homosexual” ($N = 103$, 22.9%), “bisexual” ($N = 96$, 21.3%), “mostly heterosexual” ($N = 11$, 2.4%), “other” ($N = 11$, 2.4%), and “only heterosexual” ($N = 3$, 0.7%). When asked at baseline to report their highest level of education, the largest group was “some high school” ($N = 172$, 38.2%), followed by “some college” ($N = 152$, 33.8%), “high school diploma” ($N = 104$, 23.1%), and 4.8% ($N = 22$) that fell into all other answer options (8th grade/GED/trade school certificate/undergraduate degree). Most of the sample identified as current students (75.1%) and were not currently working at baseline (68.9%).

Measures

Demographics.—Questions about demographic characteristics, such as age, race/ethnicity, and sexual identity were assessed at baseline. Race/ethnicity response options of American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and Other were collapsed in to a single “other” category for use as a covariate.

Internalizing/Externalizing Problems.—Participants completed the Adult Self-Report (ASR) developed by Achenbach System of Empirically Based Assessment (ASEBA) (Achenbach, 2009). All questions utilized a 3-point response scale: not true (0), somewhat or sometimes true (1), or very true or often true (2). Total scores for eight syndrome scales were calculated by taking the sum of ASEBA-identified items. Second-order factor analysis has shown that the syndrome scales Anxious/Depressed, Withdrawn, Somatic Complaints create one broad-band group (labeled Internalizing), while syndrome scales Aggressive Behavior, Rule-Breaking Behavior, and Intrusive scales form a second (labeled Externalizing). Reliability at baseline was excellent for both ASR Internalizing ($\alpha = .93$) and ASR Externalizing ($\alpha = .91$).

Alcohol Use.—Items assessing alcohol use were taken from the Task Force on Recommended Alcohol Questions (National Council on Alcohol Abuse and Alcoholism Recommended Sets of Alcohol Consumption Questions - October 15-16, 2003). The recall period was changed from 12 months to 6 months. Total alcohol consumed was scored by multiplying typical quantity (“how many alcoholic drinks did you have on a typical day when you drank alcohol?”) with frequency (“during the past 6 months, how often did you usually have any kind of drink containing alcohol?”). Quantity was scored on an 11-point scale (0 = 0 drinks, 5 = 7-8 drinks, 10 = 25 or more drinks) and frequency was scored on a 10-point scale (0 = never, 9 = every day). Total scores had a possible range of 0-90.

Marijuana and Illicit Drug Use.—Participants reported whether they had used any of the following drugs in the past six months: marijuana, cocaine, methamphetamines, prescription stimulants, prescription depressants, heroin, other opiates (e.g., morphine, codeine, Demerol), MDMA, psychedelics, gamma hydroxybutyrate (GHB), ketamine, and other inhalants. For each drug endorsed, participants answered “During the past 6 months, how many times did you use [insert drug]?” on a 7-point scale from 0 (0 times) to 6 (every day or almost every day). This frequency score was used for marijuana use. Due to low endorsement of other illicit drugs, frequency was dichotomized and a sum score of total unique substances used in the previous six months was computed.

Victimization.—Sexual orientation-based victimization was assessed through 24 items adapted from previous research with gay and bisexual men (Kuhns, Vazquez, & Ramirez-Valles, 2008; Ramirez-Valles, Kuhns, Campbell, & Diaz, 2010). Themes included mockery, rejection, harassment, refusal of services, and experiences of threats and physical violence. Responses fell on a 4-point scale (1 = never, 2 = once or twice, 3 = a few times, 4 = many times). The total victimization score was calculated by taking the mean of all 24 items ($\alpha = .86$ at baseline).

Social Support.—The 12-item Multidimensional Scale of Perceived Social Support (MSPSS) (Zimet, Dahlem, Zimet, & Farley, 1988) was used to assess social support. Previous research has shown the MSPSS to have high reliability and consistent factor structure (Canty-Mitchell & Zimet, 2000; Dahlem, Zimet, & Walker, 1991; Zimet, Powell, Farley, Werkman, & Berkoff, 1990). A Total Support score was calculated by taking the mean of all items (rated on a 7-point scale: 1 = very strongly disagree, 7 = very strongly agree), with higher scores indicating higher support ($\alpha = .89$ at baseline).

Internalized Homophobia.—The 8-item Desire to be Straight (DS) Subscale comes from The Internalized Gay-Related Stigma (ISM) measure created for this study. All items in this subscale were originally used in the Internalized Homosexual Stigma scale found in Ramirez-Valles et al. (2010) and recently validated for this sample by Puckett, Newcomb, Ryan, et al. (2016). Items are scored on a 4-point scale (1 = strongly disagree, 4 = strongly agree), with higher scores indicating more internalized homophobia. The subscale was computed by taking the mean of all items, and reliability at baseline was good ($\alpha = .88$).

Statistical Analyses

Multilevel models were conducted in MPlus in order to test for the within-person effects of age, victimization, social support, and internalized gay-related stigma on mental health and substance use outcomes. Between-subject baseline age, sexual orientation (coded as bisexual vs. other), and dummy-coded race (with White identification as the reference group) were included as covariates. The baseline age of the sample ranged from ages 16 to 20 which meant that participants began Crew 450 at different points in their own development. For this reason, we took an age-based approach to modeling change over time and treated it as our marker of change instead of a wave-based approach. Recruitment at different ages was also a factor in using a multilevel approach for analysis instead of autoregressive cross-lagged models that would have been wave-focused. Latent growth curve models with individually-varying assessment schedules would have been another alternative but interpreting time-varying covariates that vary by wave within that framework can be a challenge compared to performing similar analyses within the multilevel framework.

The developmental trajectories of the within-person variables, as well as the mental health and substance use outcomes, were first modeled by testing the effects of age on each variable individually. The substance use outcomes were treated as count data and estimated using a Poisson distribution. We initially tested for quadratic effects within these models but, because of the complexity, we did not carry over the quadratic effects into the subsequent multivariate analyses even when significant. We followed up the developmental trajectory models by analyzing the within-person effects as concurrent predictors of internalizing and externalizing symptoms. The model tested is shown in Figure 1. Social support was tested as a moderator of the association between victimization and mental health. We also tested for two indirect pathways using the Preacher, Zyphur, and Zhang (2010) framework for testing multilevel mediation: 1) age on mental health mediated by victimization, and 2) age on mental health mediated by victimization and internalized stigma. We only included the second indirect pathway in models where the association between victimization and internalized stigma was significant and the association between internalized stigma and the

outcome was significant. MPlus does not allow for bootstrapping in multilevel models but recommends instead using the Bayes estimator for models that include the estimation of indirect effects (Muthén, 2010). The Bayesian approach is better for estimating indirect effects because it does not assume a normal distribution. When using the Bayes estimator, MPlus reports one-tailed significance tests ($p < .025$ is considered significant instead of $p < .05$) and 95% credibility intervals. The Bayes estimator has been used for all models that included indirect effects.

We also took a time-lagged approach by testing whether previous wave victimization, social support, and internalized gay-related stigma that occurred six months prior predicted internalizing and externalizing after controlling for mental health at the previous wave. We repeated the concurrent and time-lagged models with alcohol, marijuana, and drug use as the outcomes in the place of mental health.

RESULTS

Differences in Retention

YSMM in the sample participated in an average of 4.80 (SD = 1.64) waves of data collection in the first two and half years of the study. Bivariate linear regression models were run to identify any differences in retention based on demographic covariates or any of the primary study variables. Participants who reported using marijuana more often at baseline ($B = -0.09$, $p = .010$), who reported using more illicit drugs at baseline ($B = -0.14$, $p = .001$), or who reported higher internalized stigma at baseline ($B = -0.30$, $p = .007$) participated in significantly fewer waves. Bisexual-identifying participants participated in significantly fewer waves compared to participants who reported any other sexual orientation ($B = -0.55$, $p = .004$). There were no significant differences in retention based on race/ethnicity, age, baseline internalizing symptoms, externalizing symptoms, alcohol use, victimization, or social support.

Developmental Trajectories

Intraclass correlations (ICC) and developmental trajectories for each variable are presented in Table 1. The ICC is a measure of consistency between time points. ICC values are inversely related to within-person variance and values closer to one indicate less within-person variance (i.e., less change within individuals across measurement periods). Based on the ICC, 45.8% of variance in internalizing symptoms and 45.6% of variance in externalizing symptoms was within-person. Both internalizing and externalizing symptoms had a significant within-person effect of age such that for each year of age, participants decreased in their number of internalizing symptoms by 12.99 and externalizing symptoms by 11.35 on average. There was also a significant quadratic trend for both that suggested that this pattern of decreasing symptoms slowed as participants aged. In terms of between-person differences, participants who were older at baseline had higher internalizing and externalizing symptoms, and Black participants had fewer symptoms compared to White participants. There was no significant difference for bisexual participants in comparison to participants with other identities.

Within-person variance was 44.1% for alcohol use and 42.3% for marijuana use. Drug use had the lowest amount of within-person variance at 38.0%, indicating that illicit drug use was more consistent within-persons compared to alcohol and marijuana use. There was no significant effect of age for alcohol, marijuana, or drug use. Black and Latino/Hispanic participants were significantly lower on all three substance use outcomes compared to White participants. Participants in the “other” race/ethnicity category had significantly lower rates of alcohol and drug use compared to White participants. There was no difference for bisexual-identifying participants compared to participants who identified with a different sexual orientation.

Within-person variance for victimization was 49.7%. There was a within-person effect of age on victimization such that for each additional year of age participants’ victimization scores lowered by 0.71. There was also a small but significant quadratic effect that suggested that this decrease begins to slow at older ages. Participants who identified as bisexual reported higher levels of victimization compared to other sexual orientation groups. Participants who were older at baseline also reported significantly higher victimization. Within-person variance was the lowest for internalized stigma at 37.5%. The within-person effect of age showed that internalized stigma decreased as participants got older by 0.10 for each additional year. Similar to victimization, participants who were older at baseline or who identified as bisexual reported higher internalized stigma. Participants who identified as Black or who fell in the “other” racial category also reported significantly higher internalized stigma compared to White participants. For social support, 67.1% of variance was within-person. There was no effect of age on social support and no between-person demographic differences.

Concurrent Predictors of Internalizing and Externalizing Symptoms

The concurrent within-person effects of age, victimization, internalized stigma, and social support were included in a single model in order to assess the effect of these variables on internalizing and externalizing symptoms at the same wave (see Table 2). Higher levels of victimization were a significant predictor of higher levels of both internalizing ($B = 6.78, p < .001$) and externalizing symptoms ($B = 6.57, p < .001$). Victimization was also significantly associated with internalized stigma such that participants who reported higher victimization also had higher stigma ($B = 0.41, p < .001$). Higher internalized stigma was also significantly associated with higher internalizing ($B = 2.55, p < .001$) and externalizing symptoms ($B = 1.35, p < .001$). High social support was associated with lower internalizing symptoms ($B = -1.22, p < .001$) and externalizing symptoms ($B = -0.41, p < .001$). Social support did not significantly moderate the pathway between victimization and internalizing ($B = 0.41, p = .130$) or externalizing ($B = 0.49, p = .100$).

There was a significant indirect effect of age on internalizing symptoms through victimization (95% CI: $-0.45, -0.23$; $B = -0.35, p < .001$) and through both victimization and internalized stigma (95% CI: $-0.19, -0.09$; $B = -0.14, p < .001$). The decrease in victimization as participants got older was associated with a decrease in the effect of victimization on internalizing symptoms both directly and through a decrease in internalized stigma. For each additional year of age, the direct effect of victimization on internalizing

decreased by 0.35 and the indirect effect of internalized stigma through victimization decreased by 0.14. These same indirect effects was also present for externalizing symptoms (age through victimization: 95% CI: $-0.43, -0.25$; $B = -0.33, p < .001$; age through victimization and internalized stigma: 95% CI: $-0.18, -0.10$; $B = -0.14, p < .001$). For each additional year older, the direct effect of victimization on externalizing decreased by 0.33 and the indirect effect of internalized stigma through victimization decreased by 0.14.

There were between-person differences based on age, such that participants who were older at baseline reported higher internalizing symptoms ($B = 0.92, p < .001$). Black participants reported fewer internalizing ($B = -3.85, p < .001$) and externalizing ($B = -2.96, p < .001$) symptoms compared to White participants. There were no differences for participants who identified as Latino/Hispanic or for participants in the “other” race/ethnicity category. There also was no difference based on sexual orientation.

Time-lagged Predictors of Internalizing and Externalizing

The time-lagged effects of victimization, internalized stigma, social support, and past wave internalizing and externalizing symptoms on current wave internalizing and externalizing symptoms are presented in Table 2. Higher past wave victimization was associated with higher current internalizing ($B = 1.67, p = 0.18$) and externalizing ($B = 1.74, p = .002$) symptoms. Past wave internalized stigma and social support were not associated with current internalizing or externalizing symptoms, including no significant effects of social support as a moderator. Past wave internalizing symptoms were positively associated with current internalizing symptoms ($B = 0.32, p < .001$), and were negatively associated with current externalizing symptoms ($B = -0.23, p < .001$). Higher previous wave externalizing symptoms were associated with higher current wave externalizing symptoms ($B = 0.58, p < .001$) but were not significantly associated with current wave internalizing symptoms ($B = -0.08, p = .115$).

There was a significant indirect effect of age on current wave internalizing symptoms through previous wave victimization such that as participants got older, victimization decreased and the effect of victimization on internalizing symptoms decreased (95% CI: $-0.16, -0.01$; $B = -0.07, p = .018$). For each additional year of age, the effect of previous wave victimization on internalizing decreased by 0.07. The indirect effect of age on current wave externalizing symptoms through previous wave victimization was also significant (95% CI: $-0.14, -0.02$; $B = -0.08, p = .002$). Decreases in past wave victimization as participants got older were associated with decreases in externalizing symptoms. For each year older, the effect of previous wave victimization on participants’ externalizing symptoms decreased by 0.08.

In the time-lagged model, participants who were older at baseline reported higher symptoms of externalizing ($B = 0.89, p < .05$) but there was no difference in internalizing symptoms based on baseline age. Similar to the concurrent model, Black participants reported fewer internalizing ($B = -3.47, p < .01$) and externalizing symptoms ($B = -2.82, p < .01$) compared to White participants. There were no differences for participants who identified as Latino/Hispanic, who fell in the “other” race/ethnicity category, or who identified as bisexual.

Concurrent Predictors of Substance Use Behaviors

The within-person effects of age, victimization, internalized stigma, and social support on substance use outcomes are presented in Table 3. Age was significantly associated with drug use ($B = 0.18, p < .001$) and marijuana use ($B = 0.12, p < .001$) such that as participants got older, their use increased. The association between age and alcohol use was not significant ($B = 0.46, p = .079$). Higher victimization was significantly associated with higher alcohol frequency ($B = 2.83, p < .001$), higher marijuana use ($B = 0.30, p = .010$), and higher drug use ($B = 0.44, p < .001$). There were no significant associations between internalized stigma and social support on substance use outcomes, including moderating effects of social support.

The indirect effects of age on alcohol frequency through victimization was significant (95% CI: $-0.23, -0.07$; $B = -0.14, p < .001$). Those who experienced larger declines in victimization also experienced larger declines in alcohol use. For each additional year of age, the effects of victimization on alcohol use decreased by 0.14. There were similar indirect effects for marijuana use through victimization (95% CI: $-0.03, 0.00$; $B = -0.02, p = .010$). The pattern was the same for indirect effects on drug use through victimization (95% CI: $-0.04, -0.01$; $B = -0.02, p < .001$). For each additional year older, the effects of victimization on marijuana use decreased by 0.02 and the effects of victimization on drug use decreased by 0.02.

There was a between-person difference for baseline age such that participants who were older at baseline reported lower marijuana use ($B = -0.16, p = .023$). There were no differences for alcohol or drug use. Black and Latino/Hispanic participants reported lower alcohol (Black: $B = -9.42, p < .001$; Latino/Hispanic: $B = -4.36, p = .001$), marijuana (Black: $B = -0.81, p = .001$; Latino/Hispanic: $B = -0.93, p = .001$), and drug use (Black: $B = -1.77, p < .001$; Latino/Hispanic: $\beta = -1.43, p < .001$) compared to White participants. Participants in the “other” race/ethnicity category were lower on alcohol ($B = -5.71, p < .001$) and drug use ($B = -1.59, p < .001$) compared to White participants, but not marijuana use.

Time-lagged Predictors of Substance Use Behaviors

The time-lagged effects of victimization, internalized stigma, social support, and previous wave substance use on current substance use are presented in Table 4. There were no significant effects of previous wave victimization, internalized stigma, or social support on current wave substance use outcomes. Higher past wave alcohol frequency ($B = 0.30, p < .001$) and marijuana use ($B = 0.80, p < .001$) was predictive of higher current wave alcohol frequency. There was no significant effect of past wave drug use on alcohol frequency. Previous wave marijuana use ($B = .68, p < .001$) and drug use ($B = 0.10, p = .003$) were predictive of current wave marijuana use such that higher past use was associated with higher current use. There was no significant effect of past wave alcohol use on marijuana use. Previous wave drug use ($B = 0.53, p < .001$) and marijuana use ($B = 0.13, p < .001$) was predictive of current wave drug use, but not alcohol use.

There were no significant indirect effects on substance use behaviors in the time-lagged models. In terms of between-person differences, Black participants reported lower alcohol use ($B = -7.05, p < .001$) and drug use ($B = -0.91, p < .001$) compared to White participants. Latino/Hispanic participants reported lower alcohol use ($B = -3.28, p = .001$) and drug use ($B = -0.73, p < .001$) compared to White participants. Participants in the “other” race/ethnicity category also reported lower alcohol ($B = -5.04, p < .001$) and drug use ($B = -0.89, p < .001$) compared to White participants. There were no differences based on baseline age or sexual orientation for substance use outcomes.

DISCUSSION

The purpose of the present study was to understand how experiences of victimization impact mental health and substance use among YSM, and how these associations change as YSM age from adolescence to young adulthood. In the concurrent predictor models, more experiences of victimization were associated with higher internalizing and externalizing symptoms, and higher alcohol and marijuana use. These findings are in concordance with past research that has found a positive association between victimization and mental health problems (Birkett et al., 2015; Swann et al., 2016) and between victimization and alcohol, marijuana, and illicit drug use for YSM (Bontempo & D’Augelli, 2002; Huebner et al., 2015). Our findings also support our hypothesis that victimization would mediate the association between age and both mental health and substance use outcomes, such that as participants transitioned from late adolescence into young adulthood, they would experience less victimization and as a result have fewer symptoms of mental health problems and substance use behaviors associated with victimization.

Our findings indicated that the decrease in both internalizing and externalizing symptoms as YSM moved into young adulthood was partially explained by the decrease in experiences of victimization over time. For internalizing symptoms, this result was true for both concurrent and time-lagged victimization and replicated previous findings with a sample of sexual minority young men and women (Birkett et al., 2015). This is the first study to extend those findings to externalizing symptoms. The results suggest that for YSM, the decline over time in symptoms of mental health problems is not just a decontextualized developmental process. Instead it is, in part, a function of the decrease in exposure to the bullying and victimization that increase rates of mental health issues in sexual minority youth both directly and via increases in internalized stigma. The results of the time-lagged model also suggested that for mental health, the effects of victimization can linger and still impact participants’ wellbeing six months later.

The current study also extends previous research analyzing the effect of victimization as a mediator of the association between age and substance use, though the effect was somewhat different than its mediating effect on developmental change in mental health symptoms. Victimization showed a similar mediating effect between age and alcohol, marijuana, and drug use, but this effect must be interpreted in the context of an overall increase in these substance use behaviors. In other words, there was an overall increase in illicit drug and marijuana use over time similar to what has been found in previous research (Halkitis et al., 2014; Swann et al., 2017), but there were substantial individual differences in these rates of

change (i.e., some decreased their use over time). Those YSMM who experienced large declines in victimization also tended to experience less substance use over time. However, given that substance use behaviors tended to increase for the sample as a whole, it is likely that other developmental factors contributed to the rate of change (e.g., increased access to substances, greater independence from family). More work is clearly needed in order to disentangle the multiple influences on substance use. We should also note that the indirect effect for substance use outcomes was not significant in the time-lagged models. This might suggest that for substance use, the effect of victimization is more proximal and did not linger into the subsequent assessment period. It is possible that YSMM turn to alcohol, marijuana, and illicit drugs to cope when victimization occurs, but they do not continue to use substances to cope unless their experiences of victimization are sustained.

In the concurrent model, we found that internalized gay-related stigma was positively associated with both internalizing and externalizing mental health symptoms. There was no significant association for stigma in the time-lagged model for mental health outcomes, or for stigma and the substance use outcomes in either the concurrent or time-lagged models. However, higher victimization was associated with higher internalized stigma in all of our models and there was evidence of decreased stigma via decreased victimization as participants got older resulting indirectly in fewer mental health symptoms in the cross-sectional model. Our findings concur with previous research that has shown associations between internalized stigma and the internalizing spectrum of mental health issues and disorders (Kaysen et al., 2014; Newcomb & Mustanski, 2010; Puckett, Levitt, et al., 2015; Puckett, Woodward, Mereish, & Pantalone, 2015). It is also one of the first to suggest that experiences with internalized stigma are not just associated with feelings of depression and anxiety, but also symptoms on the externalizing spectrum such as aggression and rule-breaking behavior. The lack of association between internalized stigma and substance use was in concordance with previous research that has failed to find a significant link between the two (Amadio, 2006; Amadio & Chung, 2004; Ross et al., 2001).

Participant reports of social support were not a significant factor in the majority of our models. The sole exception was the negative association social support had with internalizing and externalizing symptoms in the concurrent model. This finding was similar to past research that found social support to be associated with better mental health outcomes (Liu & Mustanski, 2012; McConnell et al., 2016). The current findings do not support past research that has linked social support and substance use issues (Lehavot & Simoni, 2011; Newcomb & Bentler, 1988). There was also no evidence that social support acted as a mediator between experiences of victimization and any of our outcomes. The results of our models suggest that the effect of victimization on mental health and substance use is significant even after accounting for the possible mitigating influence of social support.

Implications

The present study expanded upon previous research that showed victimization mediating the association between age and mental health problems into the areas of externalizing and substance use. It also lent further support to minority stress theory by demonstrating that

higher levels of victimization that YSMM experience are associated with higher rates of negative health outcomes (Hatzenbuehler, 2009; Meyer, 2003). The current research highlights the importance of addressing the bullying and victimization that YSMM and other SGM youth face during adolescents. Previous research has already shown that YSMM are at higher risk of being victimized (Bontempo & D'Augelli, 2002; Robin et al., 2002; Russell & Joyner, 2001; Shields et al., 2012) and being victimized multiple times (Bontempo & D'Augelli, 2002). The current study illustrates that these experiences of victimization have a negative impact on YSMM across a range of health indicators and that negative impact begins to reduce as YSMM become less exposed to victimization as they become older. However, this is not to say that we should simply allow victimization to decrease naturally over time. Efforts to reduce victimization experienced by sexual minorities, particularly at younger ages, are an important step in reducing the lifespan mental health and substance use disparities between YSMM and their heterosexual counterparts. This is especially important within the context of adolescence and young adulthood as a period of ongoing development. Negative experiences can have long-lasting psychosocial and physiological consequences. In particular, this is a crucial period for brain development where chronic stress (Eiland & Romeo, 2013), substance use (Squeglia, Jacobus, & Tapert, 2009), and mental health problems (Whittle et al., 2014) can stunt development in regions governing executive functioning and emotion regulation with long term consequences into adulthood. It is critical that we develop interventions that seek to safeguard YSMM from experiencing victimization, as well as programs to help YSMM who experience victimization cope more adaptively with these detrimental experiences.

Limitations

The multilevel framework used in the current study presented several important limitations. In the concurrent models, the order of effects cannot be confirmed. Minority stress theory suggests that YSMM experience negative health outcomes at disproportionate rates because they are exposed to more discrimination and victimization. However, within the concurrent framework, we cannot confirm that experiences of victimization preceded higher mental health problems or substance use. The time-lagged model allowed us to test for longitudinal effects, but the effects on substance use may be too proximal and not long-lasting enough to be detected six months later. The more complex models we fit were only designed to account for linear effects. We know from our initial trajectory models that internalizing symptoms, externalizing symptoms, and experiences of victimization had significant quadratic effects in their change over time. Excluding those quadratic effects from the cross-sectional and time-lagged models may have introduced additional bias into our results. We also found differences in retention based on marijuana use, illicit drug use, and internalized stigma. Participants who were higher on those variables at baseline participated in significantly fewer waves, so our models may be less accurate for YSMM who report the highest levels of substance use or experience the most internalized stigma.

Suggestions for Future Research

Additional research should expand to other SGM youth beyond just YSMM, including women who have sex with women and transgender and other gender minority youth. Researchers should also explore what factors put YSMM and other SGM groups at higher

risk for experiencing victimization. More research should also be done to further explore the associations between victimization experiences and internalized stigma and to understand more about the mechanisms by which internalized stigma mediates the association between victimization and health outcomes. Finally, researchers should follow YSMM further into adulthood to see the extent to which victimization experiences continue to decline over time and how that impacts their mental health and substance use.

Conclusions

The current study found that declines in victimization across development are an important explanatory factor in understanding changes in mental health and substance use among YSMM across development. These results illustrate a vital need to address the higher levels of victimization that YSMM are exposed to compared to their heterosexual peers. This study shows that the reduction in victimization that YSMM experience as they grow older and become better able to select their own environments is associated with a reduction in negative mental health and substance use outcomes. Prevention efforts to limit victimization exposure may be a necessary step to reducing the health disparities that YSMM and other SGM youth face.

REFERENCES

- Achenbach TM (2009). *The Achenbach System of Empirically Based Assessment (ASEBA): Development, Findings, Theory, and Applications*. Burlington, VT: University of Vermont Research Center for Children, Youth and Families.
- Amadio DM (2006). Internalized heterosexism, alcohol use, and alcohol-related problems among lesbians and gay men. *Addictive Behaviors*, 31(7), 1153–1162. doi:10.1016/j.addbeh.2005.08.013 [PubMed: 16183207]
- Amadio DM, & Chung YB (2004). Internalized Homophobia and Substance Use Among Lesbian, Gay, and Bisexual Persons. *Journal of Gay & Lesbian Social Services: Issues in Practice, Policy & Research*, 17(1). doi:10.1300/J041v17n01_06
- Arnett JJ, & Hughes M (2012). *Adolescence and emerging adulthood : a cultural approach*. Harlow, England ; New York: Pearson.
- Birkett M, Newcomb ME, & Mustanski B (2015). Does it get better? A longitudinal analysis of psychological distress and victimization in LGBTQ youth. *Journal of Adolescent Health*, 56(3), 280–285. doi:10.1016/j.jadohealth.2014.10.275 [PubMed: 25586226]
- Bontempo DE, & D’Augelli AR (2002). Effects of at-school victimization and sexual orientation on lesbian, gay, or bisexual youths’ health risk behavior. *Journal of Adolescent Health*, 30(5), 364–374. [PubMed: 11996785]
- Bostwick WB, Meyer I, Aranda F, Russell S, Hughes T, Birkett M, & Mustanski B (2014). Mental health and suicidality among racially/ethnically diverse sexual minority youths. *American Journal of Public Health*, 104(6), 1129–1136. doi:10.2105/AJPH.2013.301749 [PubMed: 24825217]
- Burton CL, Wang K, & Pachankis JE (2018). Does Getting Stigma Under the Skin Make It Thinner? Emotion Regulation as a Stress-Contingent Mediator of Stigma and Mental Health. *Clinical Psychological Science*, 2167702618755321.
- Burton CM, Marshal MP, Chisolm DJ, Sucato GS, & Friedman MS (2013). Sexual minority-related victimization as a mediator of mental health disparities in sexual minority youth: a longitudinal analysis. *Journal of Youth and Adolescence*, 42(3), 394–402. doi:10.1007/s10964-012-9901-5 [PubMed: 23292751]
- Canty-Mitchell J, & Zimet GD (2000). Psychometric properties of the Multidimensional Scale of Perceived Social Support in urban adolescents. *American Journal of Community Psychology*, 28(3), 391–400. [PubMed: 10945123]

- Corliss HL, Rosario M, Wypij D, Fisher LB, & Austin SB (2008). Sexual Orientation Disparities in Longitudinal Alcohol Use Patterns Among Adolescents Findings From the Growing Up Today Study. *Archives of Pediatrics & Adolescent Medicine*, 162(11), 1071–1078. [PubMed: 18981356]
- Dahlem NW, Zimet GD, & Walker RR (1991). The Multidimensional Scale of Perceived Social Support: a confirmation study. *Journal of Clinical Psychology*, 47(6), 756–761. [PubMed: 1757578]
- Davidson LM, & Demaray MK (2007). Social support as a moderator between victimization and internalizing-externalizing distress from bullying. *School Psychology Review*, 36(3).
- Dermody SS, Marshal MP, Burton CM, & Chisolm DJ (2016). Risk of heavy drinking among sexual minority adolescents: indirect pathways through sexual orientation-related victimization and affiliation with substance-using peers. *Addiction*, 111(9), 1599–1606. doi:10.1111/add.13409 [PubMed: 27018582]
- Eiland L, & Romeo RD (2013). Stress and the developing adolescent brain. *Neuroscience*, 249, 162–171. doi:10.1016/j.neuroscience.2012.10.048 [PubMed: 23123920]
- Feinstein BA, Davila J, & Dyar C (2017). A weekly diary study of minority stress, coping, and internalizing symptoms among gay men. *Journal of Consulting and Clinical Psychology*, 85(12), 1144. [PubMed: 29189030]
- Fergusson DM, Horwood LJ, & Beautrais AL (1999). Is sexual orientation related to mental health problems and suicidality in young people? *Archives of General Psychiatry*, 56(10), 876–880. [PubMed: 10530626]
- Goldbach JT, Tanner-Smith EE, Bagwell M, & Dunlap S (2014). Minority stress and substance use in sexual minority adolescents: a meta-analysis. *Prevention Science*, 15(3), 350–363. doi:10.1007/s11121-013-0393-7 [PubMed: 23605479]
- Halkitis PN, Siconolfi DE, Stults CB, Barton S, Bub K, & Kapadia F (2014). Modeling substance use in emerging adult gay, bisexual, and other YMSM across time: the P18 cohort study. *Drug and Alcohol Dependence*, 145, 209–216. doi:10.1016/j.drugalcdep.2014.10.016 [PubMed: 25456332]
- Hatzenbuehler ML (2009). How does sexual minority stigma “get under the skin”? A psychological mediation framework. *Psychological Bulletin*, 135(5), 707–730. [PubMed: 19702379]
- Heckathorn DD (1997). Respondent-driven sampling: A new approach to the study of hidden populations. *Social Problems*, 44, 174–200.
- Hequembourg AL, & Dearing RL (2013). Exploring shame, guilt, and risky substance use among sexual minority men and women. *Journal of Homosexuality*, 60(4), 615–638. doi: 10.1080/00918369.2013.760365 [PubMed: 23469820]
- Hershberger SL, & D’Augelli AR (1995). The impact of victimization on the mental health and suicidality of lesbian, gay, and bisexual youths. *Developmental Psychology*, 31(1), 65–74.
- Hicks BM, Blonigen DM, Kramer MD, Krueger RF, Patrick CJ, Iacono WG, & McGue M (2007). Gender differences and developmental change in externalizing disorders from late adolescence to early adulthood: A longitudinal twin study. *Journal of Abnormal Psychology*, 116(3), 433–447. doi:10.1037/0021-843X.116.3.433 [PubMed: 17696699]
- Huebner DM, Thoma BC, & Neilands TB (2015). School Victimization and Substance Use Among Lesbian, Gay, Bisexual, and Transgender Adolescents. *Prevention Science*, 16(5), 734–743. doi: 10.1007/s11121-014-0507-x [PubMed: 25529390]
- Hughes TL, & Eliason M (2002). Substance Use and Abuse in Lesbian, Gay, Bisexual and Transgender Populations. *The Journal of Primary Prevention*, 22, 263–298.
- Kaysen DL, Kulesza M, Balsam KF, Rhew IC, Blayney JA, Lehavot K, & Hughes TL (2014). Coping as a mediator of internalized homophobia and psychological distress among young adult sexual minority women. *Psychology of Sexual Orientation and Gender Diversity*, 1(3). doi:10.1037/sgd0000045
- King M, Semlyen J, Tai SS, Killaspy H, Osborn D, Popelyuk D, & Nazareth I (2008). A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. *BMC Psychiatry*, 8, 70. doi:10.1186/1471-244X-8-70 [PubMed: 18706118]
- Kuhns LM, Kwon S, Ryan DT, Garofalo R, Phillips G 2nd, & Mustanski BS (2015). Evaluation of respondent-driven sampling in a study of urban young men who have sex with men. *Journal of Urban Health*, 92(1), 151–167. doi:10.1007/s11524-014-9897-0 [PubMed: 25128301]

- Kuhns LM, Vazquez R, & Ramirez-Valles J (2008). Researching special populations: retention of Latino gay and bisexual men and transgender persons in longitudinal health research. *Health Education Research*, 23(5), 814–825. doi:10.1093/her/cym066 [PubMed: 17974545]
- Lebel C, Walker L, Leemans A, Phillips L, & Beaulieu C (2008). Microstructural maturation of the human brain from childhood to adulthood. *Neuroimage*, 40(3), 1044–1055. doi:10.1016/j.neuroimage.2007.12.053 [PubMed: 18295509]
- Lehavot K, & Simoni JM (2011). The impact of minority stress on mental health and substance use among sexual minority women. *Journal of Consulting and Clinical Psychology*, 79(2). doi: 10.1037/a0022839
- Liao KY-H, Kashubeck-West S, Weng C-Y, & Deitz C (2015). Testing a mediation framework for the link between perceived discrimination and psychological distress among sexual minority individuals. *Journal of Counseling Psychology*, 62(2), 226. [PubMed: 25867695]
- Liu RT, & Mustanski B (2012). Suicidal ideation and self-harm in lesbian, gay, bisexual, and transgender youth. *American Journal of Preventive Medicine*, 42(3), 221–228. doi:10.1016/j.amepre.2011.10.023 [PubMed: 22341158]
- Livingston NA, Oost KM, Heck NC, & Cochran BN (2015). The role of personality in predicting drug and alcohol use among sexual minorities. *Psychology of Addictive Behaviors*, 29(2), 414–419. doi:10.1037/adb0000034 [PubMed: 25347022]
- Marshal MP, Friedman MS, Stall R, King KM, Miles J, Gold MA, ... Morse JQ (2008). Sexual orientation and adolescent substance use: A meta-analysis and methodological review. *Addiction*, 103(4), 546–556. [PubMed: 18339100]
- Marshal MP, Friedman MS, Stall R, & Thompson AL (2009). Individual trajectories of substance use in lesbian, gay and bisexual youth and heterosexual youth. *Addiction*, 104(6), 974–981. [PubMed: 19344440]
- McConnell EA, Birkett M, & Mustanski B (2016). Families Matter: Social Support and Mental Health Trajectories Among Lesbian, Gay, Bisexual, and Transgender Youth. *Journal of Adolescent Health*, 59(6), 674–680. doi:10.1016/j.jadohealth.2016.07.026 [PubMed: 27707515]
- McLaughlin KA, Hatzenbuehler ML, Xuan Z, & Conron KJ (2012). Disproportionate exposure to early-life adversity and sexual orientation disparities in psychiatric morbidity. *Child Abuse & Neglect*, 36(9), 645–655. doi:10.1016/j.chiabu.2012.07.004
- Meyer IH (2003). Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychological Bulletin*, 129(5), 674–697. [PubMed: 12956539]
- Moody RL, Starks TJ, Grov C, & Parsons JT (2018). Internalized Homophobia and Drug Use in a National Cohort of Gay and Bisexual Men: Examining Depression, Sexual Anxiety, and Gay Community Attachment as Mediating Factors. *Archives of Sexual Behavior*, 47(4), 1133–1144. doi:10.1007/s10508-017-1009-2 [PubMed: 28608294]
- Morgan EM (2013). Contemporary issues in sexual orientation and identity development in emerging adulthood. *Emerging Adulthood*, 1(1), 52–66.
- Mustanski B (2011). Ethical and regulatory issues with conducting sexuality research with LGBT adolescents: a call to action for a scientifically informed approach. *Archives of Sexual Behavior*, 40(4), 673–686. doi:10.1007/s10508-011-9745-1 [PubMed: 21528402]
- Mustanski B, Andrews R, Herrick A, Stall R, & Schnarrs PW (2014). A syndemic of psychosocial health disparities and associations with risk for attempting suicide among young sexual minority men. *American Journal of Public Health*, 104(2), 287–294. doi:10.2105/AJPH.2013.301744 [PubMed: 24328641]
- Muthén B (2010). Bayesian analysis in Mplus: A brief introduction. Unpublished manuscript. www.statmodel.com/download/IntroBayesVersion, 203.
- Newcomb MD, & Bentler PM (1988). Impact of adolescent drug use and social support on problems of young adults: A longitudinal study. *Journal of Abnormal Psychology*, 97(1). doi: 10.1037/0021-843x.97.1.64
- Newcomb ME, Birkett M, Corliss HL, & Mustanski B (2014). Sexual Orientation, Gender, and Racial Differences in Illicit Drug Use in a Sample of US High School Students. *American Journal of Public Health*, 104(2), 304–310. doi:10.2105/AJPH.2013.301702 [PubMed: 24328653]

- Newcomb ME, & Mustanski B (2010). Internalized homophobia and internalizing mental health problems: a meta-analytic review. *Clinical Psychology Review*, 30(8), 1019–1029. doi:10.1016/j.cpr.2010.07.003 [PubMed: 20708315]
- Newcomb ME, Ryan DT, Garofalo R, & Mustanski B (2014). The effects of sexual partnership and relationship characteristics on three sexual risk variables in young men who have sex with men. *Archives of Sexual Behavior*, 43(1), 61–72. doi:10.1007/s10508-013-0207-9 [PubMed: 24217953]
- Pachankis J, Sullivan T, Feinstein B, & Newcomb M (2018). Young adult gay and bisexual men's stigma experiences and mental health: An 8-year longitudinal study. *Developmental Psychology*.
- Pilkington NW, & Daugelli AR (1995). Victimization of Lesbian, Gay, and Bisexual Youth in Community Settings. *Journal of Community Psychology*, 23(1), 34–56.
- Poteat VP, & Espelage DL (2007). Predicting psychosocial consequences of homophobic victimization in middle school students. *The Journal of Early Adolescence*, 27(2). doi:10.1177/0272431606294839
- Preacher KJ, Zyphur MJ, & Zhang Z (2010). A general multilevel SEM framework for assessing multilevel mediation. *Psychological Methods*, 15(3), 209–233. doi:10.1037/a0020141 [PubMed: 20822249]
- Puckett JA, Levitt HM, Horne SG, & Hayes-Skelton SA (2015). Internalized heterosexism and psychological distress: The mediating roles of self-criticism and community connectedness. *Psychology of Sexual Orientation and Gender Diversity*, 2(4). doi:10.1037/sgd0000123
- Puckett JA, Mereish EH, Levitt HM, Horne SG, & Hayes-Skelton SA (2018). Internalized heterosexism and psychological distress: The moderating effects of decentering. *Stigma and Health*, 3(1), 9.
- Puckett JA, Newcomb ME, Garofalo R, & Mustanski B (2016). The Impact of Victimization and Neuroticism on Mental Health in Young Men Who Have Sex with Men: Internalized Homophobia as an Underlying Mechanism. *Sexuality Research and Social Policy*, 13(3), 193–201. doi:10.1007/s13178-016-0239-8 [PubMed: 28138349]
- Puckett JA, Newcomb ME, Ryan DT, Swann G, Garofalo R, & Mustanski B (2016). Internalized homophobia and perceived stigma: A validation study of stigma measures in a sample of young men who have sex with men. *Sexuality Research and Social Policy*. doi:10.1007/s13178-016-0258-5
- Puckett JA, Woodward EN, Mereish EH, & Pantalone DW (2015). Parental rejection following sexual orientation disclosure: Impact on internalized homophobia, social support, and mental health. *LGBT Health*, 2(3). doi:10.1089/lgbt.2013.0024
- Ramirez-Valles J, Kuhns LM, Campbell RT, & Diaz RM (2010). Social Integration and Health Community Involvement, Stigmatized Identities, and Sexual Risk in Latino Sexual Minorities. *Journal of Health and Social Behavior*, 51(1), 30–47. [PubMed: 20420293]
- Remafedi G, French S, Story M, Resnick MD, & Blum R (1998). The relationship between suicide risk and sexual orientation: results of a population-based study. *American Journal of Public Health*, 88(1), 57–60. [PubMed: 9584034]
- Robin L, Brener ND, Donahue SF, Hack T, Hale K, & Goodenow C (2002). Associations between health risk behaviors and opposite-, same-, and both-sex sexual partners in representative samples of Vermont and Massachusetts high school students. *Archives of Pediatrics & Adolescent Medicine*, 156(4), 349–355. doi:10.1001/archpedi.156.4.349 [PubMed: 11929369]
- Ross MW, Rosser BRS, Bauer GR, Bocking WO, Robinson BBE, Rugg DL, & Coleman E (2001). Drug use, unsafe sexual behavior, and internalized homonegativity in men who have sex with men. *AIDS and Behavior*, 5(1). doi:10.1023/a:1009567707294
- Russell ST, & Joyner K (2001). Adolescent sexual orientation and suicide risk: Evidence from a national study. *American Journal of Public Health*, 91(8), 1276–1281. [PubMed: 11499118]
- Russell ST, Ryan C, Toomey RB, Diaz RM, & Sanchez J (2011). Lesbian, gay, bisexual, and transgender adolescent school victimization: implications for young adult health and adjustment. *Journal of School Health*, 81(5), 223–230. doi:10.1111/j.1746-1561.2011.00583.x [PubMed: 21517860]

- Ryan C, Russell S, Huebner D, Diaz R, & Sanchez J (2010). Family acceptance in adolescents and the health of LGBT young adults. *Journal of Child and Adolescent Psychiatric Nursing*, 23, 205–213. [PubMed: 21073595]
- Safren SA, & Heimberg RG (1999). Depression, hopelessness, suicidality, and related factors in sexual minority and heterosexual adolescents. *Journal of Consulting and Clinical Psychology*, 67(6), 859–866. [PubMed: 10596508]
- Schwartz DR, Stratton N, & Hart TA (2016). Minority stress and mental and sexual health: Examining the psychological mediation framework among gay and bisexual men. *Psychology of Sexual Orientation and Gender Diversity*, 3(3), 313–324. doi:10.1037/sgd0000180
- Shidlo A (1994). Internalized homophobia: Conceptual and empirical issues in measurement In Greene B & Herek GM (Eds.), *Psychological perspectives on lesbian and gay issues*, Vol. 1 Thousand Oaks, CA: Sage Publications Inc.
- Shields JP, Whitaker K, Glassman J, Franks HM, & Howard K (2012). Impact of Victimization on Risk of Suicide Among Lesbian, Gay, and Bisexual High School Students in San Francisco. *Journal of Adolescent Health*, 50(4), 418–420. doi:10.1016/j.jadohealth.2011.07.009 [PubMed: 22443849]
- Span SA, & Derby PL (2009). Depressive symptoms moderate the relation between internalized homophobia and drinking habits. *Journal of Gay & Lesbian Social Services: Issues in Practice, Policy & Research*, 21(1). doi:10.1080/10538720802497688
- Squeglia LM, Jacobus J, & Tapert SF (2009). The influence of substance use on adolescent brain development. *Clinical EEG and Neuroscience*, 40(1), 31–38. doi:10.1177/155005940904000110 [PubMed: 19278130]
- Swann G, Bettin E, Clifford A, Newcomb ME, & Mustanski B (2017). Trajectories of alcohol, marijuana, and illicit drug use in a diverse sample of young men who have sex with men. *Drug and Alcohol Dependence*.
- Swann G, Minshew R, Newcomb ME, & Mustanski B (2016). Validation of the Sexual Orientation Microaggression Inventory in Two Diverse Samples of LGBTQ Youth. *Archives of Sexual Behavior*, 45(6), 1289–1298. doi:10.1007/s10508-016-0718-2 [PubMed: 27067241]
- Szymanski DM, & Ikizler AS (2013). Internalized heterosexism as a mediator in the relationship between gender role conflict, heterosexist discrimination, and depression among sexual minority men. *Psychology of Men & Masculinity*, 14(2). doi:10.1037/a0027787
- Talley AE, Hughes TL, Aranda F, Birkett M, & Marshal MP (2014). Exploring alcohol-use behaviors among heterosexual and sexual minority adolescents: intersections with sex, age, and race/ethnicity. *American Journal of Public Health*, 104(2), 295–303. doi:10.2105/AJPH.2013.301627 [PubMed: 24328614]
- Teasdale B, & Bradley-Engen MS (2010). Adolescent same-sex attraction and mental health: the role of stress and support. *Journal of Homosexuality*, 57(2), 287–309. doi: 10.1080/00918360903489127 [PubMed: 20390994]
- Whitbeck LB, Chen X, Hoyt DR, Tyler KA, & Johnson KD (2004). Mental disorder, subsistence strategies, and victimization among gay, lesbian, and bisexual homeless and runaway adolescents. *Journal of Sex Research*, 41(4), 329–342. doi:10.1080/00224490409552240 [PubMed: 15765273]
- Whittle S, Lichter R, Dennison M, Vijayakumar N, Schwartz O, Byrne ML, ... Allen NB (2014). Structural brain development and depression onset during adolescence: a prospective longitudinal study. *American Journal of Psychiatry*, 171(5), 564–571. doi:10.1176/appi.ajp.2013.13070920 [PubMed: 24577365]
- Zimet GD, Dahlem NW, Zimet SG, & Farley GK (1988). The Multidimensional Scale of Perceived Social Support. *Journal of Personality Assessment*, 52(1), 30–41.
- Zimet GD, Powell SS, Farley GK, Werkman S, & Berkoff KA (1990). Psychometric characteristics of the Multidimensional Scale of Perceived Social Support. *Journal of Personality Assessment*, 55(3-4), 610–617. [PubMed: 2280326]




Figure 1.
Multilevel Model of Indirect Pathways on Study Outcomes

Developmental Trajectories

Table 1.

	Intraclass Correlation															
	0.54	0.54	0.56	0.58	0.62	0.50	0.63	0.33								
	Internalizing	Externalizing	Alcohol Q-F*	Marijuana Use*	Drug Use*	Victimization	Gay-Related Stigma	Social Support								
	Beta (SE)	p-value	Beta (SE)	p-value	Beta (SE)	p-value	Beta (SE)	p-value	Beta (SE)	p-value						
Between-person																
Intercept	114.23 (44.02)	0.009	104.54 (35.30)	0.003	1.45 (.69)	0.036	1.20 (.72)	0.099	-1.30 (1.90)	0.494	7.28 (1.37)	<.001	1.39 (.37)	<.001	5.87 (.73)	<.001
Variance	72.86 (7.58)	<.001	53.00 (4.76)	<.001	-	-	-	-	-	-	.08 (.01)	<.001	.23 (.02)	<.001	.57 (.07)	<.001
Baseline Age	2.46 (.52)	<.001	2.14 (.42)	<.001	.07 (.05)	0.134	-0.4 (.05)	0.455	.02 (.17)	0.890	.14 (.02)	<.001	.11 (.02)	<.001	-.01 (.05)	0.862
Race																
Black	-3.30 (1.21)	0.006	-2.86 (1.03)	0.005	-.88 (.11)	<.001	-.39 (.12)	0.001	-2.39 (.27)	<.001	.01 (.03)	0.667	.15 (.06)	0.018	.01 (.12)	0.910
Latino/Hispanic	-.98 (1.59)	0.537	-2.01 (1.25)	0.107	-.33 (.11)	0.003	-.48 (.16)	0.002	-1.30 (.29)	<.001	.00 (.04)	0.923	.01 (.07)	0.849	-.09 (.14)	0.550
Other	1.25 (1.95)	0.520	.24 (1.71)	0.890	-.45 (.17)	0.007	-.21 (.18)	0.248	-1.70 (.43)	<.001	.10 (.06)	0.097	.23 (.11)	0.034	.10 (.17)	0.559
White (referent)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bisexual	1.35 (1.09)	0.218	.80 (.96)	0.402	-.02 (.13)	0.856	.18 (.13)	0.155	.36 (.31)	0.248	.12 (.05)	0.011	.42 (.07)	<.001	-.05 (.12)	0.653
Within-person																
Variance	64.68 (5.67)	<.001	43.81 (4.13)	<.001	-	-	-	-	-	-	.06 (.01)	<.001	.14 (.01)	<.001	1.17 (.09)	<.001
Age at wave	-12.99 (4.71)	0.006	-11.35 (3.68)	0.002	.00 (.03)	0.949	.02 (.03)	0.499	.07 (.10)	0.464	-.71 (.14)	<.001	-.10 (.01)	<.001	-.04 (.03)	0.223
Quadratic Age at wave	.28 (.12)	0.015	.24 (.09)	0.009	-	-	-	-	-	-	.01 (.00)	<.001	-	-	-	-

Notes:

* Poisson distribution was used and MPlus does not compute variances. Q-F = quantity-frequency. Quadratic effect of age was only included in models where it was significant.

Table II.
Multilevel Cross-sectional and Time-lagged Models for Mental Health Outcomes

	Cross-sectional				Time-lagged							
	Internalizing Symptoms		Externalizing Symptoms		Internalizing Symptoms		Externalizing Symptoms					
	Estimate	p-value	95% CI	Estimate	p-value	95% CI	Estimate	p-value	95% CI			
<i>Between-person</i>												
Intercept	0.82	0.470	-13.06; 10.97	7.07	0.090	-8.6; 18.68	-1.59	0.399	-13.69; 10.50	2.49	0.297	-6.58; 11.60
Variance	62.44	<.001	54.67; 73.19	50.60	<.001	44.01; 61.06						
Baseline Age	0.92	<.001	.34; 1.58	0.57	0.040	-.01; 1.06	0.49	0.132	-.35; 1.34	0.66	0.008	.13; 1.21
Race												
Black	-3.85	<.001	-6.28; -1.70	-2.96	<.001	-5.25; -.55	-3.44	0.001	-5.60; -1.35	-2.81	<.001	-4.43; -1.23
Latino/Hispanic	-1.14	0.250	-3.80; 1.25	-1.87	0.060	-4.21; .40	-1.60	0.096	-4.09; .83	-1.30	0.079	-3.15; .54
Other	-0.16	0.490	-4.04; 3.23	-0.81	0.370	-4.03; 2.46	-0.53	0.374	-3.61; 2.69	-0.75	0.265	-3.09; 1.65
White (referent)	-	-	-	-	-	-	-	-	-	-	-	-
Sexual Orientation												
Bisexual	-0.54	0.320	-2.98; 1.55	-0.49	0.320	-2.14; 1.51	0.27	0.395	-1.79; 2.31	0.22	0.387	-1.35; 1.72
Gay/Other Orientation (referent)	-	-	-	-	-	-	-	-	-	-	-	-
Internalizing-Externalizing Correlation	37.46	<.001	29.51; 46.23	37.46	<.001	29.51; 46.23	27.20	<.001	17.00; 38.57	27.20	<.001	17.00; 38.57
<i>Within-person</i>												
Variance	61.08	<.001	55.82; 65.44	41.89	<.001	55.82; 65.44	62.79	<.001	55.89; 71.12	46.15	<.001	41.56; 51.53
Age at wave	-0.30	<.001	-.40; -.23	-0.29	<.001	-.35; -.22	-0.01	0.484	-.57; .60	-0.40	0.002	-.69; -.11
LGBT Victimization at wave	6.78	<.001	5.35; 8.03	6.57	<.001	5.74; 7.47	-	-	-	-	-	-
Internalized Stigma at wave	2.55	<.001	1.75; 3.45	1.35	<.001	.63; 1.96	-	-	-	-	-	-
Social Support at wave	-1.22	<.001	-1.59; -.91	-0.41	<.001	-.66; -.14	-	-	-	-	-	-
Internalizing at previous wave	-	-	-	-	-	-	0.32	<.001	.19; .45	-0.23	<.001	-.31; -.15
Externalizing at previous wave	-	-	-	-	-	-	-0.08	0.115	-.21; .05	0.58	<.001	.47; .68
LGBT Victimization at previous wave	-	-	-	-	-	-	1.67	0.018	.11; 3.21	1.74	0.002	.52; 2.94
Internalized Stigma at previous wave	-	-	-	-	-	-	0.81	0.063	-.22; 1.83	0.24	0.278	-.56; 1.05
Social Support at previous wave	-	-	-	-	-	-	0.01	0.492	-.43; .44	-0.09	0.298	-.44; .25

	Cross-sectional				Time-lagged				
	Internalizing Symptoms		Externalizing Symptoms		Internalizing Symptoms		Externalizing Symptoms		
	Estimate	p-value	95% CI	Estimate	p-value	95% CI	Estimate	p-value	95% CI
Age on LGBT Victimization	-0.05	<.001	-.06;-.04	-0.05	<.001	-.06;-.04	-0.05	<.001	-.06;-.03
LGBT Victimization on Stigma	0.41	<.001	.33; .48	0.41	<.001	.33; .48	0.38	<.001	.30; .46
Internalizing-Externalizing Correlation	38.56	<.001	34.60; 42.01	38.56	<.001	34.60; 42.01	37.47	<.001	32.72; 43.41
Social Support x LGBT Victimization	0.41	0.130	-.25; 1.23	0.49	0.100	-.15; 1.03	0.40	0.217	-.59; 1.38
<i>Indirect Effects</i>									
Age through LGBT Victimization	-0.35	<.001	-.45; -.23	-0.33	<.001	-.43; -.25	-0.07	0.018	-.16; -.01
Age through LGBT Victimization & Stigma	-0.14	<.001	-.19; -.09	-0.14	<.001	-.18; -.10	-	-	-

Notes: Models used Bayes estimator. P-values based on one-tail test and must be below .025 to reach significance. CI = Credibility Interval.

Table III.

Multilevel Cross-sectional Model for Substance Use Outcomes

	Alcohol Quantity-Frequency			Marijuana Use			Illicit Drug Use		
	Estimate	p-value	95% CI	Estimate	p-value	95% CI	Estimate	p-value	95% CI
Between-person									
Intercept	0.33	0.480	-12.97: 13.23	3.14	0.015	.33: 5.95	0.63	0.297	-1.67: 2.93
Variance	67.06	<.001	57.00: 78.98	3.35	<.001	2.88: 3.90	2.25	<.001	1.91: 2.65
Baseline Age	0.33	0.244	-.60: 1.24	-0.16	0.023	-.33: .00	-0.12	0.048	-.26: .02
Race									
Black	-9.42	<.001	-11.78:-7.10	-0.81	0.001	-1.32:-.30	-1.77	<.001	-2.19:-1.34
Latino/Hispanic	-4.36	0.001	-7.14:-1.65	-0.93	0.001	-1.52:-.34	-1.43	<.001	-1.93:-.93
Other	-5.71	<.001	-9.22:-2.28	-0.42	0.136	-1.18: .34	-1.59	<.001	-2.22:-.96
White (referent)	-	-	-	-	-	-	-	-	-
Sexual Orientation									
Bisexual	-0.15	0.444	-2.32: 2.03	0.38	0.055	-.09: .85	0.23	0.127	-.16: .62
Gay/Other Orientation (referent)	-	-	-	-	-	-	-	-	-
Alcohol & Marijuana Correlation	6.23	<.001	4.55: 8.09	6.23	<.001	4.55: 8.09	-	-	-
Alcohol & Illicit Drug Correlation	6.01	<.001	4.56: 7.66	-	-	-	6.01	<.001	4.56: 7.66
Marijuana & Illicit Drug Correlation	-	-	-	1.21	<.001	.90: 1.56	1.21	<.001	.90: 1.56
Within-person									
Variance	56.12	<.001	52.20: 60.46	1.79	<.001	1.66: 1.92	1.56	<.001	1.45: 1.68
Age at wave	0.46	0.079	-.13: 1.15	0.12	<.001	.05: .19	0.18	<.001	.10: .26
LGBT Victimization at wave	2.83	<.001	1.42: 4.26	0.30	0.010	.05: .55	0.44	<.001	.21: .67
Internalized Stigma at wave	0.03	0.479	-.92: .96	0.04	0.308	-.13: .22	-0.01	0.445	-.17: .14
Social Support at wave	0.03	0.429	-.31: .39	0.01	0.387	-.06: .08	-0.04	0.108	-.10: .02
Age on LGBT Victimization	-0.05	<.001	-.06:-.04	-0.05	<.001	-.06:-.04	-0.05	<.001	-.06:-.04
LGBT Victimization on Stigma	0.41	<.001	.34: .48	0.41	<.001	.34: .48	0.41	<.001	.34: .48
Alcohol & Marijuana Correlation	1.72	<.001	1.19: 2.26	1.72	<.001	1.19: 2.26	-	-	-
Alcohol & Illicit Drug Correlation	1.13	<.001	.64: 1.63	-	-	-	1.13	<.001	.64: 1.63
Marijuana & Illicit Drug Correlation	-	-	-	0.17	<.001	.08: .26	0.17	<.001	.08: .26
Social Support x LGBT Victimization	0.21	0.289	-.55: .98	-0.02	0.382	-.16: .12	0.01	0.455	-.12: .13

	Alcohol Quantity-Frequency			Marijuana Use			Illicit Drug Use		
	Estimate	p-value	95% CI	Estimate	p-value	95% CI	Estimate	p-value	95% CI
Indirect Effects									
Age through LGBT Victimization	-0.14	<.001	-.23;-.07	-0.02	0.010	-.03;.00	-0.02	<.001	-.04;-.01

Notes: Models used Bayes estimator. P-values based on one-tail test and must be below .025 to reach significance. CI = Credibility Interval.

Table IV.

Multilevel Time-lagged Model for Substance Use Outcomes

	Alcohol Quantity-Frequency			Marijuana Use			Illicit Drug Use		
	Estimate	p-value	95% CI	Estimate	p-value	95% CI	Estimate	p-value	95% CI
Between-person									
Intercept	4.66	0.171	-4.90: 14.28	1.80	0.005	.43: 3.16	1.51	0.018	.10: 2.92
Variance	20.66	<0.001	11.68: 31.56	0.14	<0.001	.04: .32	0.29	<0.001	.12: .53
Baseline Age	0.40	0.185	-.41: 1.35	-0.04	0.249	-.20: .07	-0.01	0.431	-.13: .11
Race									
Black	-7.05	<0.001	-9.07:-5.16	-0.20	0.071	-.48: .07	-0.91	<0.001	-1.21:-.63
Latino/Hispanic	-3.28	0.001	-5.36:-1.27	-0.25	0.044	-.55: .04	-0.73	<0.001	-1.04:-.44
Other	-5.04	<0.001	-7.64:-2.50	-0.13	0.248	-.49: .24	-0.89	<0.001	-1.28:-.51
White (referent)	-	-	-	-	-	-	-	-	-
Sexual Orientation									
Bisexual	-0.10	0.454	-1.75: 1.57	0.14	0.249	-.20: .07	0.02	0.429	-.22: .27
Gay/Other Orientation (referent)	-	-	-	-	-	-	-	-	-
Alcohol & Marijuana Correlation	-0.82	0.040	-1.71: .11	-0.82	0.040	-1.71: .11	-	-	-
Alcohol & Illicit Drug Correlation	0.97	0.031	-.04: 2.18	-	-	-	0.97	0.031	-.04: 2.18
Marijuana & Illicit Drug Correlation	-	-	-	-0.12	0.021	-.25: -.01	-0.12	0.021	-.25: -.01
Within-person									
Variance	57.87	<0.001	51.78: 64.82	2.15	<0.001	1.97: 2.34	1.79	<0.001	1.61: 1.98
Age at wave	-0.05	0.453	-.86: .58	-0.01	0.388	-.10: .13	-0.03	0.293	-.13: .07
Alcohol at previous wave	0.30	<0.001	.22: .39	0.01	0.066	.00: .02	0.00	0.434	-.01: .01
Marijuana at previous wave	0.80	<0.001	.46: 1.13	0.68	<0.001	.62: .74	0.13	<0.001	.08: .19
Illicit Drug at previous wave	0.10	0.319	-.33: .53	0.10	0.003	.03: .16	0.53	<0.001	.45: .61
LGBT Victimization at previous wave	-0.50	0.234	-1.86: .85	-0.03	0.397	-.25: .19	-0.07	0.277	-.28: .15
Internalized Stigma at previous wave	-0.57	0.107	-1.49: .33	-0.10	0.077	-.25: .04	0.02	0.405	-.13: .16
Social Support at previous wave	0.02	0.452	-.36: .41	0.04	0.114	-.03: .11	-0.04	0.131	-.10: .03
Age on LGBT Victimization	-0.05	<0.001	-.06:-.03	-0.05	<0.001	-.06:-.03	-0.05	<0.001	-.06:-.03
LGBT Victimization on Stigma	0.38	<0.001	.30: .46	0.38	<0.001	.30: .46	0.38	<0.001	.30: .46
Alcohol & Marijuana Correlation	2.79	<0.001	1.97: 3.61	2.79	<0.001	1.97: 3.61	-	-	-

	Alcohol Quantity-Frequency			Marijuana Use			Illicit Drug Use		
	Estimate	p-value	95% CI	Estimate	p-value	95% CI	Estimate	p-value	95% CI
Alcohol & Illicit Drug Correlation	1.34	0.001	.53: 2.15	-	-	-	1.34	0.001	.53: 2.15
Marijuana & Illicit Drug Correlation	-	-	-	2.79	<.001	1.97: 3.61	2.79	<.001	1.97: 3.61
Social Support x LGBT Victimization Indirect Effects	-0.50	0.234	-1.86: .85	0.01	0.434	-.14: .17	-0.08	0.153	-.23: .07
Age through LGBT Victimization	0.02	0.234	-.04: .09	0.00	0.397	-.01: .01	0.00	0.277	-.01: .01

Notes: Models used Bayes estimator. P-values based on one-tail test and must be below .025 to reach significance. CI = Credibility Interval.