



Published in final edited form as:

Optom Vis Sci. 2019 July ; 96(7): 470–476. doi:10.1097/OPX.0000000000001400.

Hypercapnia Impairs Vasoreactivity To Changes In Blood Pressure And Intraocular Pressure In Rat Retina

Guodong Liu, PhD^{1,2}, Grant Cull, B.Sci², Lin Wang, PhD², Bang V. Bui, PhD³

¹Department of Ophthalmology, The Tenth People's Hospital, Shanghai, Tongji University School of Medicine, Shanghai, China

²Devers Eye Institute, Legacy Research Institute, Portland, Oregon, USA

³Department of Optometry and Vision Sciences, University of Melbourne, Parkville, Victoria, Australia

Abstract

Significance: Retinal vascular tone is set to a level that facilitates the vasodilation and vasoconstriction. Hypercapnia reduces their “vasodilatory reserve” and impairs their reactivity to changes in intraocular or blood pressure. Therefore, eyes with systemic or local hypercapnia may have greater risk of developing ischemia in face of perfusion pressure perturbation.

Purpose: Retinal vessels are regulated by both myogenic and metabolic mechanisms. We considered whether alteration of metabolic status would modify the vascular response to ocular perfusion pressure (OPP) lowering in rat retina.

Methods: In pentobarbital anesthetized adult Brown Norway rats, normocapnia or hypercapnia was achieved by artificially ventilating animals with air or 5% carbon dioxide in ~30% oxygen, respectively. OPP was gradually reduced to ~20 mmHg by either lowering blood pressure (slowly drawing blood from a femoral artery/vein) or manometrically increasing intraocular pressure (IOP) under normocapnic or hypercapnic conditions. In all four groups (n = 7 eyes for each), a confocal scanning laser ophthalmoscope was used to acquire image sequences centered on the optic nerve throughout pressure modification. The diameter of arterioles and venules at various OPP levels was measured and expressed as percentage relative to their own baseline. The response of arterioles and venules to OPP lowering was compared between normocapnic and hypercapnic groups.

Results: Average arterial carbon dioxide partial pressure was 36.9 ± 2.6 mmHg in normocapnic and 64.1 ± 5.9 mmHg in hypercapnic ($P < 0.001$) animals. In the normocapnic groups, blood pressure lowering and IOP elevation resulted in significant vasodilation of both arterioles and venules ($P < 0.0001$). In the hypercapnic groups, OPP lowering induced vasodilation was

Correspondence: Bang V. Bui, Department of Optometry and Vision Sciences, the University of Melbourne, Parkville, 3010, Victoria, Australia, bvb@unimelb.edu.au.

Author Contributions:

Study design: L.W., B.V.B. and G.C; Resources: L.W., B.V.B; Writing the original manuscript: L.W. and G.D.L; Figures: G.D.L. and L.W; Project administration: G.D.L. and G.C; Data collection: G.C. and G.D.L; Analysis: G.D.L. and L.W; Review & Editing: B.V.B., L.W., G.C. and G.D.L.

Conflict of Interest: The authors declare that they have no conflict of interest.

significantly attenuated compared to the corresponding normocapnic groups ($P < 0.0001$ for both, 2-way ANOVA).

Conclusions: Hypercapnia significantly modified myogenic vascular autoregulation in response to OPP reduction.

Keywords

Metabolic regulation; Myogenic regulation; Hypercapnia; Ocular perfusion pressure

Introduction

The retina is one of the most metabolically active tissues in the body. To sustain this demand in the absence of stored energy, blood supply is tightly autoregulated by local myogenic and metabolic mechanisms.^{1, 2} Myogenic regulation refers to the adjustment of vascular resistance in order to maintain stable blood flow in face of fluctuating ocular perfusion pressure (OPP). Myogenic regulation is triggered by activation of stretch sensitive channels on the vascular endothelium³ and vascular smooth muscle cells,^{4, 5} which stimulate the releases of vasoactive peptides to either constrict or relax smooth muscle cells.^{4, 5} Metabolic regulation refers to the capacity for the retinal vascular bed to adjust blood supply in response to changes in local metabolic status, including oxygen (O₂) and carbon dioxide (CO₂) tensions, pH and the local concentrations of a range of other metabolic products.^{1, 2, 6, 7}

As CO₂ is a major metabolic product, its local tension is known to impact local vessel tone, with hypercapnia reducing (relative vasodilatation) and hypocapnia increasing (relative vasoconstriction) vessel tone.⁸⁻¹¹ Changes in local CO₂ tension modify the balance of a range of vasoactive substances including nitric oxide,^{12, 13} prostaglandin E₂¹⁴ and adenosine.¹⁵ Given that local CO₂ tension impacts vessel tone, it is likely that the capacity for the vasculature in response to OPP lowering will be affected during hypercapnia.

Studies in the cerebral circulation show that metabolic perturbation affects vasoreactivity to perfusion pressure changes. *Ex vivo* experiments of isolated rat diaphragmatic arterioles show that alteration of CO₂ levels in the perfusate significantly modify myogenic responses to changes in intraluminal pressure.¹⁶ *In vivo*, hypercapnia shortens the range of the cerebral autoregulatory plateau in response to changes in blood pressure.¹⁷ Furthermore, in human middle cerebral arteries, hypercapnia slows their responses to a rapid change in blood pressure.¹⁸ How retinal myogenic autoregulation is modified by metabolic status has yet to be investigated.

In the present study, we hypothesized that hypercapnia would impair the vasoreactivity of retinal blood vessels to OPP lowering induced by blood pressure (BP) and IOP. We tested this hypothesis in rats, where a robust myogenic autoregulatory response has been documented¹⁹ and a species increasingly used in studies of vascular physiology and ocular diseases.

Materials and methods

Animals

Experiments were conducted in young adult Brown-Norway rats between 131 ± 21 days. Rats were maintained in 22°C , 12-h light/12-h dark environment. Normal rat chow and water were available *ad libitum*. The experiments were regularly operated between 8:00AM to 12:00AM in a dedicated small quiet animal surgery suit at constant room temperature and adequate air ventilation. All experimental methods and animal care procedures adhered to Association for Research in Vision and Ophthalmology's Statement for the Use of Animals in Ophthalmic and Vision Research and were approved by Institutional Animal Care and Use Committee (Legacy Health System, Portland, Oregon).

Anesthesia and surgical preparation

General anesthesia was induced by inhalation of 2% isoflurane followed by maintenance at 1 – 2% isoflurane. The femoral arteries and veins were cannulated with polyethylene tubing (PE10 or 50, BD Intramedic, Franklin Lakes, NJ). One femoral artery was connected to a pressure transducer (BLPR2; World Precision Instruments, Sarasota, FL) and a four-channel amplifier system (Lab-78 Trax-4/24T; World Precision Instruments) for continuous BP monitoring. A femoral vein was used for administration of drugs. The other artery or vein was used for blood drawing to induce systemic hypotension as detailed below. An orotracheal intubation was performed and the anesthesia was switched to continuous intravenous infusion of sodium pentobarbital (2–5 mg/kg/hr, Nembutal, Oak Pharmaceuticals, Lake Forest, IL) via an infusion pump (Aladdin, World Precision Instruments). The reservoir was set initially at a height to maintain a baseline level of 10 mmHg. Animals were ventilated with a respirator (RSP 1002, Kent Scientific Co., Torrington, CT) with 30% oxygen at a rate between 60 and 90 breaths/min. The arterial oxygen partial pressure (PaO_2) was maintained at average of 110.6 ± 14.0 mmHg and a pH of 7.44 ± 0.10 , as measured using a blood gas analyzer (i-STAT[®], Abbott Inc. Princeton, NJ). A heating pad, set at 37°C , was used to maintain body temperature throughout the experiments. Both eyes were dilated with tropicamide (0.5%; Alcon Laboratories, Inc., Fort Worth, TX), and topical anesthetized with 0.5% proparacaine hydrochloride (Bausch & Lomb, Inc., Bridgewater, NJ).

Monitoring of EtCO_2 and PaCO_2

In order to avoid the need for repeated blood sampling, which makes stable continuous imaging difficult, arterial carbon dioxide partial pressure (PaCO_2) was calibrated to the end tidal carbon dioxide level (EtCO_2). In 12 rats, PaCO_2 ranging from 30 mmHg to 60 mmHg was measured ($n = 27$ samples) using the blood gas analyzer. The corresponding EtCO_2 was recorded simultaneously using a capnometer (RSP-300, Kent Scientific Co., Torrington, CT). The relationship between EtCO_2 and PaCO_2 was determined using linear regression ($\text{EtCO}_2 = 0.76 \text{ mmHg} \cdot \text{PaCO}_2 - 3.5309$, $R^2 = 0.79$). This function was used to convert EtCO_2 to PaCO_2 .

In the current study, we set PaCO_2 in normocapnic groups between 30 to 40 mmHg (or EtCO_2 between 20 and 27 mmHg), which was maintained by adjusting the respiratory rate

(between 60 and 90 breaths/min) and the inspiration/expiration ratio (between 15% and 45%). The PaCO₂ in the hypercapnic groups was set between 60 and 70 mmHg (or EtCO₂ between 45 and 50 mmHg) and achieved by adding 5% CO₂ to the inspired air.

Gradual OPP reduction by BP lowering and IOP elevation

OPP reduction was induced by BP lowering and IOP elevation. For BP lowering, a syringe pump was used to slowly draw blood from either a femoral artery or a vein at a rate of 0.84 ± 0.24 ml/min. Drawing started after the mean BP had stabilized between 95 and 100 mmHg. The blood draw ended when BP fell to between 20 and 25 mmHg. This rate of blood draw resulted in an average rate of blood pressure lowering of 10.7 ± 3.1 mmHg/min and the average blood volume drawn was 6.0 ± 1.8 ml.

In a randomly chosen eye, a gradual IOP elevation was achieved by cannulating the anterior chamber with a 33-G needle connected to a reservoir filled with balanced salt solution (Baxter, Deerfield, IL). To induce a gradual increase in IOP at a rate that matched the BP lowering, the reservoir was raised from 10 to 70 mmHg at a rate of 9.6 ± 0.3 mmHg/min using a modified peristaltic pump (Longer Precision Pump Co., Ltd, Hebei, China). The final target IOP (70 mmHg) was confirmed using a rebound tonometer (TONOLAB, Vantaa, Finland). The OPP was calculated as the difference between mean femoral arterial BP and IOP. The former is an estimate of ophthalmic arterial pressure, which requires no hydrostatic correction in rats; the latter is the estimate of retinal venous pressure.

Changes in vessel diameter in response to two methods of gradual OPP reduction (BP lowering and IOP elevation) were assessed in both normocapnia (as control) and hypercapnia, which were randomized assigned into 4 groups of rats: normocapnia with BP lowering; normocapnia with IOP elevation; hypercapnia with BP lowering; hypercapnia with IOP elevation. Each group includes $n = 7$ eyes. The sample sizes were determined by using

Vessel responses to BP lowering and IOP elevation were also compared with two groups of spontaneously hypercapnic rats ($n = 7$ eyes for BP group and $n = 5$ eyes for IOP group), in which animals were anesthetized in the same way but breathed normal room air, rather than being intubated and artificially ventilated (i.e., spontaneous hypercapnia versus controlled hypercapnia). In these spontaneously hypercapnic animals average PaCO₂ was 63.4 ± 11.6 mmHg, PaO₂ was 117.5 ± 32.58 mmHg and pH was 7.33 ± 0.07 .

Vessel diameter imaging and quantification

Following general preparation, rats were placed on a stage in front of a spectral-domain optical coherence tomography/confocal scanning laser ophthalmoscope (SD-OCT/cSLO, Spectralis; Heidelberg Engineering, Heidelberg, Germany). Customized rigid gas-permeable contact lenses (3.5-mm posterior radius of curvature, 5.0-mm optical zone diameter, and +5.0-diopter back vertex power) were placed on both eyes. Changes in retinal vessel diameter across a 30° field centered on the optic nerve head were recorded in infrared reflection mode (cSLO-IR) at baseline and throughout the course of OPP lowering (BP lowering or IOP elevation). More specifically, baseline vessel diameter was recorded over the first 20-seconds, after which OPP modification was initiated and continuous recording ensued until BP was decreased to 20–25 mmHg or IOP was increased to 70 mmHg.

Changes in vessel diameter during OPP lowering were quantified using Fiji (ImageJ v2.0.0; National Institutes of Health, Bethesda, MD) as described previously.²⁰ Image sequences were combined into a single stack and registered with custom-software to allow the same regions to be analyzed across time. In each eye, 3 to 6 arterioles and 3 to 6 venules were assessed. ImageJ was used to extract the light intensity profile perpendicularly across the vessel at one-disc diameter from the optic disc margin. Using custom written software, vessel diameter was quantified as the distance between the two intensity troughs, which correspond to the two edges of the vessel. Changes in diameter for each vessel were expressed as a percentage relative to its own baseline. Group data were averaged into bins spanning 10 mmHg of OPP.

Data analysis and statistics

The percentage of vessel diameter responses to OPP lowering (binned every 10 mmHg) in each experiment group were analyzed using one-way repeated measures ANOVA with Dunnett's post hoc test. Differences in vessel reactivity under normocapnia and hypercapnia were compared using 2-way ANOVA (OPP and PaCO₂) for BP lowering and IOP elevation challenges separately. The OPP at which vessel diameter began to increase or decrease with OPP lowering was determined by segmental linear regression analysis using individual diameter measurements at corresponding OPP.

The critical probability (*P*) to reject our null hypothesis was set at 5%. Data were presented as mean ± SEM (standard error of the mean) in the figures and given as mean ± SD (standard deviation) within the text. All analyses were performed using Prism 7 (GraphPad Software, Inc., La Jolla, CA).

Results

Table 1 shows baseline parameters (prior to OPP manipulation) for the four experimental groups. The values including PaCO₂ and vessel diameters in controlled hypercapnic groups were significantly higher than the normocapnic rats (*P* < 0.05), while baseline OPP in each group was similar.

Effects of controlled hypercapnia on baseline vessel diameters (prior to OPP manipulation)

Representative baseline images in Figure 1 shows that compared with a normocapnic animal (Fig. 1a), vessels were more dilated in a hypercapnic animal (Fig. 1b). The average baseline arteriolar diameter was larger in the two hypercapnic groups (41.43 ± 5.2 μm) than the two normocapnic groups (35.8 ± 4.5 μm). This represents a 19.6% hypercapnia induced increase in basal arteriolar diameter (*P* < 0.001). Similarly, hypercapnia induced a 27.1% increase in venular diameter (normocapnia: 39.5 ± 4.2 μm vs hypercapnia: 49.8 ± 5.7 μm, *P* < 0.0001). These are summarized in Table 1.

Effects of hypercapnia on vascular responses to BP lowering

The baseline OPP prior to blood drawing was 82.2 ± 1.4 and 83.0 ± 4.4 mmHg in the normocapnic and controlled hypercapnic groups (*P* = 0.07), respectively. Blood drawing

induced similar BP reductions of 70.2 ± 3.4 mmHg (from 94.1 ± 2.8 to 23.9 ± 2.3 mmHg) in normocapnic and 69.2 ± 5.5 mmHg (from 94.6 ± 3.0 to 22.8 ± 4.1 mmHg) in hypercapnic groups, respectively ($P = 0.48$). Under normocapnic conditions, retinal arteriolar and venular diameters remained stable with moderate OPP lowering. As OPP fell to 52.2 ± 1.2 mmHg for arterioles and 67.7 ± 1.7 mmHg for venules, vessel diameters started to increase steadily and reached peaks of 15.4% and 12.1% above baseline, respectively ($P < 0.0001$, 1-way ANOVA for both, Fig. 2).

Under hypercapnic conditions, arteriolar and venular responses were significantly attenuated compared with normocapnia ($P < 0.001$, 2-way ANOVA for both, Fig. 2). Specifically, arteriolar dilation only reached a maximum of 5.2% compared to the 15.4% seen in the normocapnic group (Fig. 2a). During hypercapnia, arteriolar diameter started to decline when OPP dropped to very low levels at 23.9 ± 0.9 mmHg, this is in contrast to the dilation seen in the normocapnic group at similar OPPs (Fig. 2a). For venules, hypercapnia completely abolished the vasodilation seen in the normocapnic group (Fig. 2b). At a very low OPP of 23.3 ± 1.5 mmHg retinal venules showed significant constriction ($P < 0.01$, Fig. 2b).

Normocapnic and controlled hypercapnic animals were also compared against those that were naturally ventilated (spontaneously hypercapnic groups). Figure 2a and 2b show that vessel responses to BP lowering in spontaneously hypercapnic groups were not significantly different to those under controlled hypercapnia ($P = 0.47$ and $P = 0.08$ for arterioles and venules, respectively).

Effects of hypercapnia on vascular response to IOP elevation

The baseline OPP prior to IOP elevation was similar in normocapnic (86.3 ± 2.2 mmHg) and controlled hypercapnic groups (85.6 ± 3.8 mmHg, $P = 0.13$). In the normocapnic group, gradual IOP elevation from 10 to 70 mmHg induced vasodilation in both arterioles and venules. For arterioles and venules, significant dilation started when OPP was lowered to 59.6 ± 2.1 mmHg and 74.6 ± 2.0 mmHg, respectively. Peak vasodilations of 11.9% and 4.5% was reached by arterioles and venules, respectively ($P < 0.0001$, Fig. 3).

Hypercapnia completely blunted both arteriolar and venular vasodilatation in response to IOP elevation ($P = 0.97$ and $P = 0.12$, respectively, Fig. 3). Compared to the normocapnic groups, both arteriolar and venular responses were significantly attenuated in controlled hypercapnia groups ($P < 0.0001$, 2-way ANOVA for both, Fig. 3). At very low OPPs (24.3 mmHg, Fig. 3b), venules were significantly constricted ($P = 0.0002$), whereas arterioles did not ($P = 0.26$). These hypercapnia data were adequately described using a single line regression model, rather than the two-line regression model needed to describe data from normocapnic groups. Vessel response to IOP elevation in the spontaneously hypercapnic group was blunted compared with normocapnia; an effect that was similar to the controlled hypercapnic group (Fig. 3).

Discussion

We evaluated the impact of altered metabolic status induced by hypercapnia on myogenic regulation in the rat retinal circulation. We found that hypercapnia significantly increased basal vessel diameter and blunted the capacity for vessels to dilate in react to OPP lowering.

Previous studies have shown that increasing PaCO₂ increases basal arteriolar diameter. The magnitude of such increases have been found to range between 1.4%²¹ to 8.5%¹⁴ in brain arterioles and from 3%²² to 14%²³ in retinal arterioles. In the current study, arteriolar diameter was increased by 19.6%. A likely reason for the difference between ours and previous reports may lie in differences in induced magnitude of PaCO₂ changes.^{24–26} In previous studies on human subjects, PaCO₂ levels are most often mildly elevated by between 12.0% to 25.4%.^{22, 23} In our study, direct delivery of a 5% CO₂ mixture by artificial ventilation raised PaCO₂ by ~70%, which would account for the larger basal dilation observed in our study.

In blood, PaCO₂ reacts with water in the red blood cells catalyzed by carbonic anhydrase. The dissolved carbon dioxide in the blood reacts with water to form carbonic acid, which is further disassociated into HCO³⁻ and H⁺. [Guyton, 2006 #1647] The latter is believe to be a major mediator leading to the change in vascular tone [Brian, 1998 #3388] by stimulate the release of vasoactive substances from vascular and/or glial cells.^{27, 28} In particular nitric oxide, which is synthesized by astrocytes,²⁹ Müller cells³⁰ and endothelial cells,³¹ relaxes smooth muscle cells via a cyclic guanosine monophosphate signaling pathway.³² Hypercapnia may also enhance the synthesis of prostaglandin E₂ through cyclooxygenase-1 and glutathione pathways,³³ as well it reduces calcium levels within and thus relaxes smooth muscle cells.^{34–36} Inhibition of prostaglandin E₂ reverses hypercapnia-induced vasodilation.
14

Whilst relaxation of smooth muscle cells is relevant to increased arteriolar diameter, it cannot explain our observation that venules dilated by 27.1% under hypercapnia as rat venules have virtually no smooth muscle cells. Since the inner retinal circulation is a closed system, venular dilation could arise from passive distention by increased blood volume in upstream arterioles. However, we found that retinal venules dilated significantly more than arterioles (27.1% vs. 19.6%, *P* = 0.003). Differences in the way that arterioles and venules respond to stimuli have been reported in previous studies. For example, while flickering light induced dilation of retinal arteries, veins showed little change.³⁷ In the cerebral circulation, arterioles have been shown to dilate (10%) more than venules (2%) in response to xenon gas inhalation.³⁸ In response to hypercapnia in awake mice, cerebral arterioles dilated by 15.8%, whereas venules dilated by only 2.3%.²⁶ These studies suggest that changes in venular diameter do not passively follow arterial changes²⁰ until the blood volume was reduced too low or the mechanical pressure (IOP) was too high (OPP was between 20 and 30 mmHg) that the venular wall could no longer withhold the tension.”

Myogenic mechanisms maintain stable blood flow over a wide range of OPPs; often referred to as the autoregulation plateau along the pressure-flow curve.⁶ In the cerebral circulation, hypercapnia impairs vascular responses to change in perfusion pressure,^{39, 40} as evidenced

by narrowing of the autoregulation plateau.¹⁷ Dineen et al.,¹⁸ showed that hypercapnia delays the initial transient hemodynamic responses to a rapid change in perfusion pressure in human middle cerebral arteries. Consistent with these studies, we showed for the first time that hypercapnia blunts the capacity of the retinal vasculature to compensate for OPP lowering (Figs. 2 and 3). These data support the idea that metabolic perturbation is detrimental to pressure-induced myogenic autoregulation in the retina.

Since basal vessel diameter was already increased by hypercapnia (Table 1, Fig. 1), further vasodilation in response to OPP lowering might be limited by an already low myogenic tone; or a reduced “vasodilatory reserve”.⁴¹ A previous study in isolated arterioles revealed that severe hypercapnia inhibit myogenic response by suppression of smooth muscle cells and activation of endothelium dependent mechanisms.¹⁶ Hypercapnia induced reductions in pH are also known to significantly attenuate vasodilation.⁴² In addition to the local CO₂ levels, mechanical factors also likely influence the retinal vasodilatory reserve,⁴³ when the OPP was reduced to below 20 mmHg in control eyes and by both BP and IOP.

The results from the study highlight the importance to maintain normal PaCO₂ levels when studying vascular physiology. As indicated by Moulton et al.,⁴⁴ depending on the type of anesthetic used, the respiratory system of experimental animals can be depressed to different degrees resulting in accumulation of CO₂. We show here that in pentobarbital anesthetized rats spontaneously breathing room air, PaCO₂ was as high as those rats breathing 5% CO₂ (63.4 vs. 64.2 mmHg). Thus, without knowing or controlling PaCO₂, studies employing spontaneous room air breath will underestimate vasoreactivity.

Several limitations of our study are worth noting. First in the current study we elected to use changes in retinal vessel diameter as an index of vascular responses to OPP challenge. However, the actual blood flow resulting from changes in diameter is not known. Second, although PaCO₂ was constantly adjusted to maintain levels within our target range by monitoring EtCO₂, PaO₂ was only sampled once. Although PaO₂ variation can influence vasoreactivity,¹⁰ by ventilating animals with 30% O₂ we believe that small variations in PaO₂ would not substantially influence our results. Finally, as IOP was continuously changing during the experiment, it was challenging to maintain a consistent focus during imaging. As a result, the image quality in experiments with IOP elevation was more variable than BP lowering manipulations.

Our current study provides evidence that alteration of metabolic status induced by hypercapnia profoundly impacts pressure-initiated myogenic autoregulation in rat retinal vessels. These findings have implication for understanding how the retinal vasculature copes with OPP fluctuation in conditions such as glaucoma. Tissues that already have low oxygen or high carbon dioxide tension (which can also occur with low pH arising from increased anaerobic glycolysis) would tend to have lower vessel tone, and thus reduce “vasodilatory reserve”. In those whose vasculature is already impaired (e.g. diabetes induced vascular damage, arteriosclerosis with long term systemic hypertension), a narrower autoregulatory plateau means that only a small reduction in OPP would exceed the lower limit of an already compromised “vasodilatory reserve”.

Acknowledgement:

This work was supported by the National Eye Institute Grant R01-EY019939, Legacy Good Samaritan Foundation, Portland, Oregon, USA and an Australian Research Council Future Fellowship.

Supported by the National Eye Institute Grant R01-EY019939, Legacy Good Samaritan Foundation, Portland, Oregon, USA and an Australian Research Council Future Fellowship.

References

1. Anderson DR. Introductory Comments on Blood Flow Autoregulation in the Optic Nerve Head and Vascular Risk Factors in Glaucoma. *Survey of ophthalmology* 1999;43 Suppl 1:S5–9.
2. Kiel JW. *The Ocular Circulation*. San Rafael (CA): Morgan & Claypool Life Sciences; 2010.
3. Chistiakov DA, Orekhov AN, Bobryshev YV. Effects of Shear Stress on Endothelial Cells: Go with the Flow. *Acta Physiol (Oxf)* 2017;219:382–408. [PubMed: 27246807]
4. Baek EB, Kim SJ. Mechanisms of Myogenic Response: Ca(2+)-Dependent and -Independent Signaling. *J Smooth Muscle Res* 2011;47:55–65. [PubMed: 21757855]
5. Bevan JA, Hwa JJ. Myogenic Tone and Cerebral Vascular Autoregulation: The Role of a Stretch-Dependent Mechanism. *Ann Biomed Eng* 1985;13:281–6. [PubMed: 4037457]
6. Cipolla MJ. Control of Cerebral Blood Flow. In: *The Cerebral Circulation*. San Rafael (CA); 2009.
7. Jacob M, Chappell D, Becker BF. Regulation of Blood Flow and Volume Exchange across the Microcirculation. *Crit Care* 2016;20:319. [PubMed: 27765054]
8. Geiser MH, Riva CE, Dorner GT, et al. Response of Choroidal Blood Flow in the Foveal Region to Hyperoxia and Hyperoxia-Hypercapnia. *Curr Eye Res* 2000;21:669–76. [PubMed: 11148604]
9. Hosking SL, Harris A, Chung HS, et al. Ocular Haemodynamic Responses to Induced Hypercapnia and Hyperoxia in Glaucoma. *Br J Ophthalmol* 2004;88:406–11. [PubMed: 14977778]
10. Venkataraman ST, Hudson C, Fisher JA, et al. Retinal Arteriolar and Capillary Vascular Reactivity in Response to Isoxic Hypercapnia. *Exp Eye Res* 2008;87:535–42. [PubMed: 18840429]
11. Wang L, Grant C, Fortune B, Cioffi GA. Retinal and Choroidal Vasoreactivity to Altered Paco₂ in Rat Measured with a Modified Microsphere Technique. *Exp Eye Res* 2008;86:908–13. [PubMed: 18420196]
12. Sato E, Sakamoto T, Nagaoka T, et al. Role of Nitric Oxide in Regulation of Retinal Blood Flow During Hypercapnia in Cats. *Invest Ophthalmol Vis Sci* 2003;44:4947–53. [PubMed: 14578421]
13. Smith JJ, Lee JG, Hudetz AG, et al. The Role of Nitric Oxide in the Cerebrovascular Response to Hypercapnia. *Anesth Analg* 1997;84:363–9. [PubMed: 9024030]
14. Hoiland RL, Tymko MM, Bain AR, et al. Carbon Dioxide-Mediated Vasomotion of Extra-Cranial Cerebral Arteries in Humans: A Role for Prostaglandins? *J Physiol* 2016;594:3463–81. [PubMed: 26880615]
15. Fenton RA, Rubio R, Berne RM. Adenosine and the Acid-Base State of Vascular Smooth Muscle. *J Appl Physiol Respir Environ Exerc Physiol* 1981;51:179–84. [PubMed: 6790498]
16. Nagi MM, Ward ME. Modulation of Myogenic Responsiveness by Co₂ in Rat Diaphragmatic Arterioles: Role of the Endothelium. *Am J Physiol* 1997;272:H1419–25.
17. Meng L, Gelb AW. Regulation of Cerebral Autoregulation by Carbon Dioxide. *Anesthesiology* 2015;122:196–205. [PubMed: 25401418]
18. Dineen NE, Brodie FG, Robinson TG, Panerai RB. Continuous Estimates of Dynamic Cerebral Autoregulation During Transient Hypocapnia and Hypercapnia. *J Appl Physiol* 2010;108:604–13. [PubMed: 20035062]
19. Wong VH, Armitage JA, He Z, et al. Chronic Intraocular Pressure Elevation Impairs Autoregulatory Capacity in Streptozotocin-Induced Diabetic Rat Retina. *Ophthalmic Physiol Opt* 2015;35:125–34. [PubMed: 25529024]
20. Li H, Bui BV, Cull G, et al. Glial Cell Contribution to Basal Vessel Diameter and Pressure-Initiated Vascular Responses in Rat Retina. *Invest Ophthalmol Vis Sci* 2017;58:1–8. [PubMed: 28055098]

21. Verbree J, Bronzwaer AS, Ghariq E, et al. Assessment of Middle Cerebral Artery Diameter During Hypocapnia and Hypercapnia in Humans Using Ultra-High-Field Mri. *J Appl Physiol* (1985) 2014;117:1084–9. [PubMed: 25190741]
22. Venkataraman ST, Hudson C, Fisher JA, Flanagan JG. Novel Methodology to Comprehensively Assess Retinal Arteriolar Vascular Reactivity to Hypercapnia. *Microvasc Res* 2006;72:101–7. [PubMed: 16926032]
23. Duan A, Bedggood PA, Metha AB, Bui BV. Reactivity in the Human Retinal Microvasculature Measured During Acute Gas Breathing Provocations. *Scientific reports* 2017;7:2113. [PubMed: 28522835]
24. Ide K, Eliasziw M, Poulin MJ. The Relationship between Middle Cerebral Artery Blood Velocity and End-Tidal Pco₂ in the Hypocapnic-Hypercapnic Range in Humans. *J Appl Physiol* 2003;95:129–37 [PubMed: 19278048]
25. Willie CK, Macleod DB, Shaw AD, et al. Regional Brain Blood Flow in Man During Acute Changes in Arterial Blood Gases. *J Physiol* 2012;590:3261–75. [PubMed: 22495584]
26. Nishino A, Takuwa H, Urushihata T, et al. Vasodilation Mechanism of Cerebral Microvessels Induced by Neural Activation under High Baseline Cerebral Blood Flow Level Results from Hypercapnia in Awake Mice. *Microcirculation* 2015;22:744–52. [PubMed: 26454149]
27. Fleming I, Hecker M, Busse R. Intracellular Alkalinization Induced by Bradykinin Sustains Activation of the Constitutive Nitric Oxide Synthase in Endothelial Cells. *Circ Res* 1994;74:1220–6. [PubMed: 7514511]
28. Petropoulos IK, Pournaras JA, Stangos AN, Pournaras CJ. Effect of Systemic Nitric Oxide Synthase Inhibition on Optic Disc Oxygen Partial Pressure in Normoxia and in Hypercapnia. *Invest Ophthalmol Vis Sci* 2009;50:378–84. [PubMed: 18676634]
29. Neufeld AH, Hernandez MR, Gonzalez M. Nitric Oxide Synthase in the Human Glaucomatous Optic Nerve Head. *Arch Ophthalmol* 1997;115:497–503. [PubMed: 9109759]
30. Donati G, Pournaras CJ, Munoz JL, et al. Nitric Oxide Controls Arteriolar Tone in the Retina of the Miniature Pig. *Invest Ophthalmol Vis Sci* 1995;36:2228–37. [PubMed: 7558716]
31. Ignarro LJ. Nitric Oxide as a Unique Signaling Molecule in the Vascular System: A Historical Overview. *Journal of physiology and pharmacology : an official journal of the Polish Physiological Society* 2002;53:503–14. [PubMed: 12512688]
32. Denninger JW, Marletta MA. Guanylate Cyclase and The .No/Cgmp Signaling Pathway. *Biochim Biophys Acta* 1999;1411:334–50. [PubMed: 10320667]
33. Howarth C, Sutherland B, Choi HB, et al. A Critical Role for Astrocytes in Hypercapnic Vasodilation in Brain. *J Neurosci* 2017;37:2403–14. [PubMed: 28137973]
34. Niwa K, Haensel C, Ross ME, Iadecola C. Cyclooxygenase-1 Participates in Selected Vasodilator Responses of the Cerebral Circulation. *Circ Res* 2001;88:600–8. [PubMed: 11282894]
35. Davidge ST, Baker PN, Laughlin MK, Roberts JM. Nitric Oxide Produced by Endothelial Cells Increases Production of Eicosanoids through Activation of Prostaglandin H Synthase. *Circ Res* 1995;77:274–83. [PubMed: 7614714]
36. Attwell D, Buchan AM, Charpak S, et al. Glial and Neuronal Control of Brain Blood Flow. *Nature* 2010;468:232–43. [PubMed: 21068832]
37. Polak K, Schmetterer L, Riva CE. Influence of Flicker Frequency on Flicker-Induced Changes of Retinal Vessel Diameter. *Invest Ophthalmol Vis Sci* 2002;43:2721–6. [PubMed: 12147608]
38. Fukuda T, Nakayama H, Yanagi K, et al. The Effects of 30% and 60% Xenon Inhalation on Pial Vessel Diameter and Intracranial Pressure in Rabbits. *Anesth Analg* 2001;92:1245–50. [PubMed: 11323354]
39. McCulloch TJ, Visco E, Lam AM. Graded Hypercapnia and Cerebral Autoregulation During Sevoflurane or Propofol Anesthesia. *Anesthesiology* 2000;93:1205–9. [PubMed: 11046207]
40. Haggendal E, Johansson B. Effects of Arterial Carbon Dioxide Tension and Oxygen Saturation on Cerebral Blood Flow Autoregulation in Dogs. *Acta Physiol Scand Suppl* 1965;258:27–53. [PubMed: 5869903]
41. Henrion D. Pressure and Flow-Dependent Tone in Resistance Arteries. Role of Myogenic Tone. *Arch Mal Coeur Vaiss* 2005;98:913–21. [PubMed: 16231579]

42. Aalkjaer C, Poston L. Effects of Ph on Vascular Tension: Which Are the Important Mechanisms? *J Vasc Res* 1996;33:347–59. [PubMed: 8862140]
43. Celotto AC, Capellini VK, Baldo CF, et al. Effects of Acid-Base Imbalance on Vascular Reactivity. *Braz J Med Biol Res* 2008;41:439–45. [PubMed: 18592120]
44. Moulton EM, Choi W, Boas DA, et al. Evaluating Anesthetic Protocols for Functional Blood Flow Imaging in the Rat Eye. *J Biomed Opt* 2017;22:16005. [PubMed: 28056146]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig.1. Representative baseline cSLO images from one eye in normocapnic group (a) and another in hypercapnic group (b). Vessel diameters were measured at 1-optic disc (dotted circle) distance from the optic disc margin (solid circle). a (red): arterioles; v (white): venules.




Fig. 2. Effect of hypercapnia on retinal arteriolar and venular diameters (percentage change, %) in response to BP modification. Mean (\pm SD) diameter change (%) summarized into 10 mmHg OPP bins for arterioles (a) and venules (b) under normocapnic (blue, $n = 7$), controlled hypercapnic (red, $n = 7$) and spontaneously hypercapnic (black, $n = 7$) conditions. Unfilled circles indicate significant difference from baseline (first symbols). Asterisks (*) indicate significant difference between normocapnic and controlled hypercapnic groups. The actual diameter of arterioles at baseline were $35.5 \pm 3.7 \mu\text{m}$, $39.3 \pm 2.7 \mu\text{m}$ and $40.9 \pm 5.2 \mu\text{m}$ for normocapnia, controlled hypercapnia and spontaneously hypercapnia, respectively. The actual diameter of venules at baseline were $39.4 \pm 2.9 \mu\text{m}$, $47.3 \pm 3.0 \mu\text{m}$ and $48.7 \pm 5.2 \mu\text{m}$ for normocapnia, controlled hypercapnia and spontaneously hypercapnia, respectively.




Fig. 3. Effect of hypercapnia on retinal arteriolar and venular diameters in response to IOP elevation. Mean (\pm SD) diameter change (%) summarized into 10 mmHg OPP bins for arterioles (a) and venules (b) under normocapnic (blue, $n = 7$), controlled hypercapnic (red, $n = 7$) and spontaneously hypercapnic (black, $n = 5$) conditions. Unfilled circles indicate significant difference from baseline (first symbols). Asterisks (*) indicate significant difference between normocapnic and controlled hypercapnic groups. The actual diameter of arterioles at baseline were $40.5 \pm 5.2 \mu\text{m}$, $43.6 \pm 7.1 \mu\text{m}$ and $42.8 \pm 5.3 \mu\text{m}$ for normocapnia, controlled hypercapnia and spontaneously hypercapnia, respectively. The actual diameter of venules at baseline were $39.4 \pm 2.9 \mu\text{m}$, $47.3 \pm 3.0 \mu\text{m}$ and $54.3 \pm 2.8 \mu\text{m}$ for normocapnia, controlled hypercapnia and spontaneously hypercapnia, respectively.

Table 1.

Baseline values for all experimental groups

Groups	NC_BP	HC_BP	NC_IOP	HC_IOP
OPP (mmHg)	82.2 ± 1.4	83.0 ± 4.4	86.3 ± 2.2	85.6 ± 3.8
PaCO ₂ (mmHg)	36.6 ± 2.9	64.2 ± 5.9	37.7 ± 2.9	64.0 ± 6.5
Arteriolar diameter (µm)	35.5 ± 3.7	39.3 ± 2.7	40.5 ± 5.2	43.6 ± 7.1
Venular diameter (µm)	39.4 ± 2.9	47.3 ± 3.0	36.4 ± 5.0	51.4 ± 8.2

NC: normocapnia; HC: hypercapnia; BP: blood pressure; IOP: intraocular pressure; OPP: ocular perfusion pressure; PaCO₂: arterial carbon dioxide partial pressure;