



# HHS Public Access

Author manuscript

*Neurobiol Aging*. Author manuscript; available in PMC 2020 September 01.

Published in final edited form as:

*Neurobiol Aging*. 2019 September ; 81: 22–29. doi:10.1016/j.neurobiolaging.2019.05.011.

## The relationship between domain-specific subjective cognitive decline and Alzheimer's pathology in normal elderly adults

Sepideh Shokouhi<sup>1</sup>, Alexander C Conley<sup>1</sup>, Suzanne L Baker<sup>2</sup>, Kimberly Albert<sup>1</sup>, Hakmook Kang<sup>3</sup>, Harry E Gwirtsman<sup>1,4</sup>, Paul A Newhouse<sup>1,4</sup>, The Alzheimer's Disease Neuroimaging Initiative

<sup>1</sup>Center for Cognitive Medicine, Department of Psychiatry and Behavioral Sciences, Vanderbilt University Medical Center, Nashville, TN, USA

<sup>2</sup>Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

<sup>3</sup>Department of Biostatistics, Vanderbilt University Medical Center, Nashville, TN, USA

<sup>4</sup>Geriatric Research, Education, and Clinical Center, Tennessee Valley Veterans Affairs Medical Center, Nashville, TN, USA

### Abstract

We evaluated the associations of subjective (self-reported Everyday Cognition, ECog) and objective cognitive measures with regional amyloid- $\beta$  ( $A\beta$ ) and tau accumulation in 86 clinically normal (CN) elderly subjects from the Alzheimer's Disease Neuroimaging Initiative (ADNI). Regression analyses were conducted to identify whether individual ECog domains (Memory, Language, Organization, Planning, Visuospatial, and Divided Attention) were equally or differentially associated with regional [<sup>18</sup>F]florbetapir and [<sup>18</sup>F]flortaucipir uptake and how these associations compared to those obtained with objective cognitive measures. A texture analysis, the weighted two-point correlation, was used as an additional approach for estimating the whole-brain tau burden without PET intensity normalization. While the strongest models for ECog domains included either tau (Planning and Visuospatial) or  $A\beta$  (Memory and Organization), the strongest models for all objective measures included  $A\beta$ . In  $A\beta$ -negative participants, the strongest models for all ECog domains of executive functioning included tau. Our results indicate differential associations of individual subjective cognitive domains with  $A\beta$  and tau in CN. Detailed characterization of ECog may render a valuable pre-screening tool for pathological prediction.

---

Correspondence: Sepideh Shokouhi, Ph.D., [sepideh.shokouhi@vanderbilt.edu](mailto:sepideh.shokouhi@vanderbilt.edu).

Disclosure statement

The authors report no actual or potential conflicts of interest relevant to this article.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Keywords

[<sup>18</sup>F]flortaucipir; neurofibrillary tau tangles; subjective cognitive decline; amyloid beta plaques; Aging; Alzheimer's disease; [<sup>18</sup>F]florbetapir

---

## 1- Introduction

Subjective cognitive decline (SCD) is increasingly recognized as a precursor to mild cognitive impairment (MCI). Older individuals with SCD are at higher likelihood of converting to MCI and dementia than those without SCD (Dufouil et al., 2005; Glodzik-Sobanska et al., 2007; Mitchell et al., 2014; Reisberg et al., 2010). However, SCD is etiologically heterogeneous and complex. Biomarker studies may help clarify the biological basis of SCD and pinpoint the most AD-predictive SCD characteristics as early as possible and help rule out other conditions that can lead to self-perceptions of cognitive alterations. Identifying the pathological pathways that influence subjective cognitive decline in the clinically normal elderly population is also critical for the identification and recruitment of at-risk participants in prevention trials before the onset of dementia. Self-recognized deficiencies can pertain to one or more cognitive domains, including memory, attention/processing speed, language, executive function, orientation, and visuospatial skills. Understanding how biological factors influence individual SCD domains would provide valuable early diagnosis and prognosis information.

Amyloid- $\beta$  ( $A\beta$ ) plaques and Neurofibrillary tangles (NFTs) of misfolded hyperphosphorylated tau proteins begin to accumulate decades before the onset of clinical dementia, which raises the question of how they influence the manifestation of SCD in non-demented subjects. Previous research has established an association between SCD and preclinical AD pathologies (Amariglio et al., 2012; Amariglio et al., 2018; Amariglio et al., 2015; Barnes et al., 2006; Buckley et al., 2017; Jorm et al., 2004; La Joie et al., 2016; Perrotin et al., 2012; Snitz et al., 2015; Vogel et al., 2017). Most of these studies have focused on associations between subjective deficits within the memory domain and  $A\beta$  pathology. A recent study by Buckley and colleagues (Buckley et al., 2017) reported that preclinical subjective cognitive decline is associated with both increasing entorhinal tau burden and (to a lesser extent) global  $A\beta$  burden. However, they found that tau and  $A\beta$  do not necessarily interact to influence preclinical subjective cognitive decline.

In this study, we have conducted a comprehensive examination of the associations of two major AD pathologies ( $A\beta$ , tau) with objective and subjective cognitive measures to gain a more in-depth understanding of the pathological basis of SCD, measured with the Everyday Cognition (ECog) instrument (Farias et al., 2008) in clinically normal subjects. Specifically, we have addressed the central question of whether individual ECog domains (memory, language, executive functioning, and visuospatial processing) are equally or differentially linked with regional [<sup>18</sup>F]flortaucipir and [<sup>18</sup>F] florbetapir standardized uptake value ratios (SUVRs). We have utilized multiple regression models to (1) determine associations between individual pathologies (predictors) and each ECog domain (outcomes), (2)

construct the strongest prediction model (best combination of predictors) for each ECog domain, and (3) identify interactions between key predictors in the strongest models.

In addition to the regional tau SUVR values, we have conducted a post hoc whole brain texture analysis approach, the weighted two-point correlation function ( $wS_2$ ), as an additional approach for estimating the severity of tau burden. One of the main advantages of this technique is the elimination of the reference region normalization of PET image intensity, which may be relevant to early tau characterization in normal subjects when the [ $^{18}\text{F}$ ]flortaucipir SUVR signal is low and likely to be affected by the background off-target binding (Baker SL, 2019).

## 2- Materials & Methods

### 2.1- Alzheimer's Disease Neuroimaging Initiative

Data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database ([adni.loni.usc.edu](http://adni.loni.usc.edu)). This study was approved by the Institutional Review Board of Vanderbilt University Medical Center (IRB#181429). The ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner MD, with the primary goal to test whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). Cognitive, demographic, and imaging data for this study were collected from 86 clinically normal (CN) ADNI subjects. We did not include any EMCI, MCI or AD subjects because this study focuses on associations between AD pathologies ( $A\beta$  and tau burden) and the earliest self-reported cognitive symptoms. The ADNI CN group has a Mini-Mental State Examination, MMSE (Folstein et al., 1983) memory scores of 24–30 (inclusive), a CDR of 0, and education-adjusted score on delayed recall of paragraph A from Wechsler Memory Scale Logical Memory II (E-WMS) 9 for 16 or more years of education, 5 for 8–15 years of education, 3 for 0–7 years of education. CN subjects are not diagnosed with MCI or dementia (based on the absence of significant impairment in cognitive functions and daily activity).

### 2.2- Cognitive and neuropsychological measures

Participants' global cognitive measures were obtained from their MMSE, Alzheimer's Disease Assessment Scale-cognitive subscale (ADAS-Cog), composite executive function (ADNI-EF), and composite memory (ADNI-MEM) scores (Crane et al., 2012; Gibbons et al., 2012). The ADNI-EF is a composite executive function score derived from the baseline scores of a number of tasks including Category Fluency, Clock Drawing, Digit Span Backwards, Digit Symbol Substitution, as well as Trails A and B. The ADNI-MEM is a composite score combining recognition and recall scores from a number of tasks including the ADAS-Cog, Rey Auditory Verbal Learning Test (RAVLT), MMSE and Logical Memory Task. Self-reported ECog (Farias et al., 2008) scores were used as a measure of subjective cognitive decline. The self-rated ECog questionnaire comprised six subscales, including Everyday Memory, Everyday Language, Everyday Visuospatial abilities, Everyday Planning, Everyday Organization, and Everyday Divided Attention.

### 2.3- Pre-processing of imaging data

**Tau PET**—All ADNI [<sup>18</sup>F]Flortaucipir scans were acquired at participating sites following the standardized ADNI protocol for [<sup>18</sup>F]Flortaucipir ([adni.loni.usc.edu](http://adni.loni.usc.edu)). In brief, the injection of 370 MBq ± 10% of [<sup>18</sup>F]Flortaucipir was followed by a 30 min (6X5min frames) PET scan starting at 75–105 minutes post-injection. The temporal frames were uploaded to USC's Laboratory of Neuroimaging ADNI database (LONI) where they were coregistered, averaged, and smoothed with a scanner-specific filter derived from each site's Hoffman brain phantom to obtain a common isotropic resolution of 8 mm full width at half maximum (FWHM) resolution. [<sup>18</sup>F]Flortaucipir SUVR values from six regions (Braak I-IV) (Schöll et al., 2016) were directly downloaded from the ADNI PET imaging archive. The regional SUVR values were normalized by the inferior cerebellar gray matter uptake (Baker et al., 2017) and re-grouped into Braak I (entorhinal), BraakIII-IV (limbic), and Braak V-VI (isocortical) regions of interest (ROIs).

**A $\beta$  PET**—All ADNI [<sup>18</sup>F]Florbetapir scans were acquired at participating sites following the standardized ADNI protocol for [<sup>18</sup>F]Florbetapir (Jagust et al., 2015). Regional [<sup>18</sup>F]Florbetapir SUVR values (normalized by the whole cerebellum) were directly downloaded from the ADNI PET imaging archive. We used SUVR values from four ROIs, the anterior-posterior cingulate, lateral parietal (including precuneus), frontal, and lateral temporal cortices (Farrell et al., 2018; Landau et al., 2014; Palmqvist et al., 2017; Perrotin et al., 2012; Villeneuve et al., 2015) for the subsequent statistical analyses. In addition to individual regional SUVR values, a mean composite ROI SUVR was calculated, and a threshold of 1.11 was applied to identify the A $\beta$ -positive individuals (Landau et al., 2012).

### 2.4- Statistical analysis of the associations between tau-PET measures and cognitive and behavioral measures

We utilized three types of regression models in R ([www.R-project.org](http://www.R-project.org)) to determine individual and combined associations between A $\beta$  and tau pathologies and cognitive measures. Multiple simple linear regression models (model 1) were used as the first step to determine the associations of individual cognitive measures (outcomes) with individual regional pathologies (predictors). The regional [<sup>18</sup>F] florbetapir and [<sup>18</sup>F] flortaucipir values were included separately as predictors. All models included age, gender, and APOE- $\epsilon$ 4 status (dichotomous variable) as covariates. We performed the Bonferroni correction whenever we needed to adjust for multiple comparisons. Backward-elimination regression analyses (model 2) were run for each objective and subjective (self-reported ECog) cognitive measure as outcome but included all regional [<sup>18</sup>F]florbetapir and [<sup>18</sup>F]flortaucipir SUVR values, age, sex, and APOE- $\epsilon$ 4 status as initial predictors. The backward-elimination procedure starts with an initial model that includes all predictors, followed by testing the deletion of each predictor using a chosen model fit criterion, and step-wise deleting predictors with the most statistically insignificant deterioration of the model fit. This process was used to determine which combination of predictors built the strongest model to fit a specific cognitive measure. Step-down method, i.e., backward-elimination, is usually better than the forward method (Mantel, 1970) and is also recommended over stepwise method (Harrell, 2015). The Akaike Information Criterion was used as a model selection criterion for a given set of data. The backward-elimination regression analyses were conducted first

for all CN subjects and then repeated for the A $\beta$ -negative subjects of the CN group. Interactions between predictors (model 3): We tested multiple two-way interactions between individual regional tau, regional A $\beta$ , age, sex, and APOE- $\epsilon$ 4 status. Due to a large number of predictors (10 variables) and outcomes (10 cognitive measures), we primarily focused on pairwise interactions between variables that were included in the strongest models (from step 2, backward elimination) for each subjective cognitive test.

## 2.5- Post hoc wS<sub>2</sub> Analysis

Applied to a flortaucipir image, the wS<sub>2</sub> analysis provides a quantitative characterization of spatial clustering of image intensity values in individual subjects' brain. We used this metric as an alternative approach for estimating the severity of global tau burden. We downloaded the ADNI SUV (standardized uptake value) images that were not intensity-normalized by any reference region, meaning that the uptake of the individual voxels had not been divided by the mean uptake of a reference region. Using SPM 12 (Ashburner, 2012), each tau PET SUV image was aligned with the subject's T1-weighted magnetic resonance imaging (MRI) volume and the cortical gray matter (GM) mask was segmented and applied to flortaucipir SUV image for the subsequent wS<sub>2</sub> analysis (MATLAB 2018, MathWorks, Inc., Natick, Massachusetts, United States). The wS<sub>2</sub> approach captures the image texture (spatial clustering) by pairwise comparison of the SUV signal intensities between different locations in gray matter. The outcome of this analysis is a single measure per subject that quantifies the (global) severity of spatial clustering of image intensity values as an estimate of the tau burden in the individual subject's brain. Detailed steps of this analysis are provided in the supplement section (supplement A). Applications of two-point correlation functions are described in previous literature (Blair et al., 1996; Jiao et al., 2008; Tewari et al., 2004). The three-step regression analyses (see 2.4) were repeated with the wS<sub>2</sub>-based global measure of tau burden. The wS<sub>2</sub>-based tau burden was combined with the global A $\beta$  SUVR measures in the backward-elimination regression analysis (model 2) to find the strongest model for predicting each cognitive measure. Pairwise interactions between wS<sub>2</sub>-based tau and global A $\beta$  SUVR were tested in model 3.

## 3- Results

Table 1 provides a summary of subject-specific information, including age, sex, number of APOE- $\epsilon$ 4 carriers, objective/subjective cognitive measures, global flortaucipir SUVR values, and A $\beta$ -positivity status. Out of 86 subjects, 39 were women. The average age at the time of the tau-PET scans was 78 $\pm$ 8. Out of 86 subjects, 24 subjects were APOE- $\epsilon$ 4 carriers, and 38 subjects were A $\beta$ -positive.

### 3.1- Associations of individual regional pathologies with cognitive measures while including age, sex, and APOE- $\epsilon$ 4 status as covariates

Tables 2 summarizes the results of the simple regression analyses for finding associations of each objective cognitive measure and each domain of the self-reported ECog with regional tau SUVR measures while including age, sex, and APOE- $\epsilon$ 4 as covariates. The Bonferroni-corrected P-value of 0.016 (0.05/3) was used to adjust for multiple comparisons across regional tau measures (Braak I, Braak III-IV, Braak V-VI). Associations with self-reported

**ECog measures:** The Braak SUVR values significantly predicted Everyday Planning (Braak I). Other, marginally significant ( $p < 0.09$ ) associations were found in models for predicting Everyday Visuospatial and Everyday Organization with Braak I SUVR as the predictor. **Associations with objective cognitive measures:** The Braak SUVR values significantly predicted the MMSE scores. Other, marginally significant associations were found for predicting ADNI-EF ( $p < 0.1$ ).

Table 3 summarizes the results of the simple regression analyses for associations of each objective cognitive measure and each domain of the self-reported ECog with the regional A $\beta$  SUVR measures while including age, sex, and APOE- $\epsilon 4$  as covariates. The Bonferroni-corrected P-value of 0.0125 (0.5/4) was used to adjust for multiple comparisons across regional A $\beta$  measures (parietal, frontal, temporal, cingulate). **Associations with self-reported ECog measures:** Significant associations were found in models for predicting Everyday Organization (all regions) and Everyday Memory (frontal lobe). Other, marginally significant associations ( $p < 0.098$ ) were found in Everyday Visuospatial (frontal lobe), and Everyday Language (parietal lobe). **Associations with objective cognitive measures:** Significant associations were found in models for predicting MMSE and ADAS\_cog. Other, marginally significant associations ( $p < 0.07$ ) were found in models for predicting ADNI-MEM and ADNI-EF.

### 3.2- Backward-elimination regression analysis

Table 4 summarizes the results of the backward-elimination regression analyses for each objective cognitive performance test (MMSE, ADAS\_cog, ADNI-EF, ADNI-MEM) and each subjective cognitive measure (self-reported ECog domains: Everyday Memory, Everyday Language, Everyday Visuospatial, Everyday Planning, Everyday Organization, Everyday Divided Attention) as outcomes. **Strongest models for self-reported ECog measures:** The backward-elimination regression analysis indicated that the strongest models to predict ECog measures included either only A $\beta$  (Everyday Memory, Everyday Language, Everyday Organization), only tau (Everyday Planning, Everyday Visuospatial), or none (Everyday Divided Attention). When applied to A $\beta$ -negative CN subjects, the strongest models for Everyday Visuospatial and Everyday Planning continued to include tau only (Braak I) whereas the model for Everyday Memory continued to include A $\beta$  only (cingulate cortex with marginal significance). In A $\beta$ -negative subjects, both tau and A $\beta$  were included in the strongest model for predicting Everyday Organization. The strongest model for predicting Everyday Divided Attention in A $\beta$ -negative CN subjects included tau. **Strongest models for objective cognitive measures:** The strongest models for predicting the objective cognitive measures included A $\beta$ , mostly without tau (MMSE, ADAS\_cog, ADNI-EF). The best model for predicting ADNI-MEM included both tau and A $\beta$ . When excluding A $\beta$ -positive CN subjects, the strongest models for ADNI-EF and ADNI-MEM included both pathologies.

### 3.3- Interactions between predictors

**Interactions with age—**We found significant interactions between tau and age when predicting Everyday Visuospatial ( $p = 0.027$  for Braak I SUVR), Everyday Planning ( $p = 0.002$  for Braak I SUVR and  $p = 0.04$  for Braak III-IV SUVR) and ADNI-MEM ( $p = 0.034$



for Braak III-IV SUVR). We found significant interactions between A $\beta$  (temporal lobe) and age when predicting Everyday Organization ( $p = 0.03$ ). Interactions between AD pathologies: When including all CN subjects, no interactions were found between the pathologies to predict the ECog domains. However, we found a significant interaction between A $\beta$  (cingulate cortex) and tau (Braak III-IV) when predicting Everyday Organization ( $p = 0.017$ ) only in A $\beta$ -negative subjects.

### 3.4- Analysis with wS<sub>2</sub>-based tau measures

With the simple regression model, we found significant associations between wS<sub>2</sub>-based tau and several objective cognitive measures, including MMSE ( $p < 0.0001$ ,  $R^2 = 0.30$ ), ADAS\_cog ( $p = 0.006$ ,  $R^2 = 0.28$ ), and ADNI-EF ( $p = 0.005$ ,  $R^2 = 0.24$ ). Also, we found significant associations between wS<sub>2</sub>-based tau and individual self-reported ECog measures, including Everyday Planning ( $p = 0.04$ ,  $R^2 = 0.16$ ), Everyday Organization ( $p = 0.01$ ,  $R^2 = 0.11$ ), and Everyday Memory ( $p = 0.006$ ,  $R^2 = 0.22$ ). The associations between wS<sub>2</sub>-based tau and Everyday Visuospatial ( $p = 0.06$ ) and Everyday Language ( $p = 0.07$ ) were marginal. Backward elimination regression analysis of self-reported ECog measures and interactions between predictors: Similar to the SUVR-based results, the backward elimination regression analysis confirmed that the strongest models for Everyday Visuospatial and Everyday Planning included wS<sub>2</sub>-based tau (but not A $\beta$ ). Also, the strongest model for predicting Everyday Divided Attention did not include tau or A $\beta$ . The strongest models for Everyday Organization and Everyday Memory included both wS<sub>2</sub>-based tau and A $\beta$ . We found significant interactions between wS<sub>2</sub>-based tau and age when predicting Everyday Visuospatial ( $p = 0.0001$ ), Everyday Planning ( $p = 0.0008$ ), and Everyday Organization ( $p = 0.03$ ). We found no interactions between wS<sub>2</sub>-based tau and global A $\beta$ . However, when including regional A $\beta$  SUVR values, we found a significant interaction between wS<sub>2</sub>-based tau and cingulate cortex A $\beta$  for predicting Everyday Language ( $p = 0.03$ ). Backward elimination regression analysis of objective cognitive measures and interactions between predictors: The strongest models for ADAS\_cog and ADNI-MEM included A $\beta$  only. The strongest model for MMSE and ADNI-EF included wS<sub>2</sub>-based tau and A $\beta$ . We found significant interaction between tau and age in predicting MMSE.

## 4- Discussion

Developing accessible screening tools that can predict the Alzheimer's-related pathological profile before a person begins to show more severe signs of cognitive and/or functional decline would provide an early opportunity to set apart a 'vulnerable' population for which preventive measures could be instituted and monitored. A growing body of research suggests that the presence of SCD in older normal adults predicts future decline and conversion to MCI and AD (Mitchell et al., 2014; Reisberg et al., 2010). However, SCD is etiologically heterogeneous and complex. Biomarker studies may help disentangle the biological basis of SCD and pinpoint the most AD-predictive characteristics in clinically normal subjects. Previous AD biomarker studies of SCD have primarily focused on the memory complaints (Amariglio et al., 2012; Amariglio et al., 2015; Barnes et al., 2006; Jorm et al., 2004; Kryscio et al., 2014; Snitz et al., 2015) by using either simple memory questions or constructing composite memory scores from different instruments, such as Everyday

Memory or Memory Functioning Questionnaire (Gilewski et al., 1990). Although recent evidence suggests that the associations between SCD and AD pathology (La Joie et al., 2016) are not restricted to the memory, it is not fully understood how individual SCD domains relate to A $\beta$  and tau.

The objective of this study was to gain an in-depth understanding of the early pathological bases of objective and subjective cognitive alterations with specific focus on individual domains of the self-reported ECog test. We found preliminary evidence in support of differential associations of ECog domains with A $\beta$  and tau pathologies. Our findings suggest that a detailed characterization of self-reported cognitive complaints may provide useful diagnosis and prognosis information. For instance, difficulties in thinking ahead or in developing schedules in advance of anticipated events (Everyday Planning) may be an important sign of progressing tauopathy. Regarding the associations of SCD with A $\beta$ , our findings are consistent with those from a study by Amariglio and colleagues reporting that higher [<sup>11</sup>C]PiB-PET binding in normal individuals was primarily associated with both subjective complaints related to memory and Everyday Organization (Amariglio et al., 2012). Given that the episodic memory is among the earliest cognitive domains reported in preclinical AD, it is plausible to assume that initial self-reported memory concerns are influenced by A $\beta$ , which is the earliest known AD biomarker (Jack et al., 2013). However, these relationships may change across the disease severity (Barnes et al., 2006), possibly due to the diminished self-perception of cognitive decline in MCI and AD patients with advanced neurodegeneration (Carr et al., 2000), or the plateauing of A $\beta$  accumulation at advanced disease stages.

We found that the most robust models for Everyday Planning and Everyday Visuospatial included tau only. When excluding A $\beta$ -positive participants, the strongest model for all ECog domains related to executive functioning (Everyday Planning, Everyday Organization, and Everyday Divided Attention) and visuospatial skills included tau. It is possible that subjective decline of the executive function and visuospatial skills in the absence of substantial A $\beta$  load may be an early sign of primary age-related tauopathy (PART). Josephs and colleagues (Josephs et al., 2017) analyzed 52 pathologically definite PART cases and found, after accounting for age at clinical examination, a significant association between tau burden and executive function and visuospatial skills. While the strongest models for the objective cognitive measures included A $\beta$  by itself or together with tau (ADNI-MEM), the strongest models for the subjective cognitive measures included either tau only (Planning and Visuospatial) or A $\beta$  only (Memory and Organization). These associations remained significant after adjusting each subjective cognitive model of table 4 with objective cognitive measures such as MMSE ( $p < 0.021$ ) and ADNI-MEM ( $p < 0.027$ ). If further validated, our findings may lay a conceptual foundation for an early prediction of neuropathological designations (e.g., PART and AD trajectories) based on domain-specific SCD characterization. A direct comparison of our finding with previous studies is not straightforward due to the heterogeneity in the selected SCD instruments and age differences across these studies. Currently, there is a lack of a standardized framework to assess SCD across different research groups (Rabin et al., 2015). Consequently, different SCD instruments may not have the same relationships with AD pathologies. Our ADNI CN subjects were  $78 \pm 8$  years of age and presented a more clinically advanced group in



comparison with other studies, such as the clinically normal group of the Harvard Aging Brain Study (Buckley et al., 2017). Future research should include younger subjects (and other cohorts) to help establish the validity of SCD as a prescreening tool for predicting individual pathological profiles (prevalence of A $\beta$ , tau, or both) before conducting PET scans or lumbar punctures to determine evidence of biomarker positivity, thus reducing the costs associated with the selection of appropriate candidates for clinical trials that target a specific pathology (tau or A $\beta$ ).

The post hoc analysis with the weighted two-point correlation curves confirmed the associations of tau with Everyday Visuospatial and Everyday Planning. Moreover, and together with global A $\beta$ , wS<sub>2</sub>-based tau was present in the strongest model for predicting Everyday Organization and Everyday Memory (without interacting with A $\beta$ ). There is no reason to assume that tau should not be associated with subjective memory complaints in normal subjects. However, these associations may have gone undetected with SUVR-based tau measures, particularly when using Everyday Memory instead of more sensitive composite memory scores similar to Buckley et al. (Buckley et al., 2017). The overall associations between wS<sub>2</sub>-based tau and most cognitive measures were higher than the SUVR-based estimates of tau burden, possibly due to several methodological advantages of this metric which may be relevant to early tau characterization in normal subjects when the [<sup>18</sup>F]flortaucipir SUVR signal is low and likely to be affected by the background off-target binding (Baker SL, 2019). However, the results of wS<sub>2</sub>-based texture analysis should be interpreted with caution until future investigations can validate this technique with pathological data.

In summary, this study shows the significance of the subjective cognitive concept in normal and pathological aging using data from clinically normal ADNI subjects. Our analyses did not include longitudinal data. However, as ADNI is an ongoing database, more follow-up scans will be available in the upcoming years to evaluate longitudinal associations between Everyday Cognition domains and AD pathologies and determine how they change across different clinical diagnostic groups.

## Conclusions

In clinically normal subjects, individual domains of subjective cognitive decline have differential associations with A $\beta$  and tau. Unlike subjective cognitive measures, the best prediction models of objective cognitive measures include A $\beta$ . Detailed characterization of subjective cognitive decline may render a valuable tool for pathological prediction (presence of A $\beta$ , tau, or both) and has the potential to improve the identification of individuals with the greatest risk of future dementia.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgment

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California. Funding: This study was supported by NIH grants R00 EB 009106, to S.Sh.

## References

- Amariglio RE, Becker JA, Carmasin J, Wadsworth LP, Lorus N, Sullivan C, Maye JE, Gidicsin C, Pepin LC, Sperling RA, Johnson KA, Rentz DM, 2012 Subjective cognitive complaints and amyloid burden in cognitively normal older individuals. *Neuropsychologia* 50(12), 2880–2886. [PubMed: 22940426]
- Amariglio RE, Buckley RF, Mormino EC, Marshall GA, Johnson KA, Rentz DM, Sperling RA, 2018 Amyloid-associated increases in longitudinal report of subjective cognitive complaints. *Alzheimer's & Dementia: Translational Research & Clinical Interventions* 4, 444–449.
- Amariglio RE, Mormino EC, Pietras AC, Marshall GA, Vannini P, Johnson KA, Sperling RA, Rentz DM, 2015 Subjective cognitive concerns, amyloid- $\beta$ , and neurodegeneration in clinically normal elderly. *Neurology* 85(1), 56–62. [PubMed: 26048028]
- Ashburner J, 2012 SPM: a history. *Neuroimage* 62(2), 791–800. [PubMed: 22023741]
- Baker SL HT, Maass A, LaJoie R, Jagust WJ, 2019 Effect of off-target binding on 18F-Flortaucipir variability in healthy controls across the lifespan. *Journal of Nuclear Medicine* Accepted.
- Baker SL, Maass A, Jagust WJ, 2017 Considerations and code for partial volume correcting [(18)F]-AV-1451 tau PET data. *Data in brief* 15, 648–657. [PubMed: 29124088]
- Barnes LL, Schneider JA, Boyle PA, Bienias JL, Bennett DA, 2006 Memory complaints are related to Alzheimer disease pathology in older persons. *Neurology* 67(9), 1581–1585. [PubMed: 17101887]
- Blair SC, Berge PA, Berryman JG, 1996 Using two-point correlation functions to characterize microgeometry and estimate permeabilities of sandstones and porous glass. *Journal of Geophysical Research: Solid Earth* 101(B9), 20359–20375.
- Buckley RF, Hanseeuw B, Schultz AP, Vannini P, Aghjayan SL, Properzi MJ, Jackson JD, Mormino EC, Rentz DM, Sperling RA, Johnson KA, Amariglio RE, 2017 Region-Specific Association of Subjective Cognitive Decline With Tauopathy Independent of Global  $\beta$ -Amyloid Burden. *JAMA neurology* 74(12), 1455–1463. [PubMed: 28973551]
- Carr DB, Gray S, Baty J, Morris JC, 2000 The value of informant versus individual's complaints of memory impairment in early dementia. *Neurology* 55(11), 1724. [PubMed: 11113230]
- Crane PK, Carle A, Gibbons LE, Insel P, Mackin RS, Gross A, Jones RN, Mukherjee S, Curtis SM, Harvey D, Weiner M, Mungas D, 2012 Development and assessment of a composite score for memory in the Alzheimer's Disease Neuroimaging Initiative (ADNI). *Brain imaging and behavior* 6(4), 502–516. [PubMed: 22782295]
- Dufouil C, Fuhrer R, Alperovitch A, 2005 Subjective Cognitive Complaints and Cognitive Decline: Consequence or Predictor? The Epidemiology of Vascular Aging Study. *Journal of the American Geriatrics Society* 53(4), 616–621. [PubMed: 15817007]

- Farias ST, Mungas D, Reed BR, Cahn-Weiner D, Jagust W, Baynes K, Decarli C, 2008 The measurement of everyday cognition (ECog): scale development and psychometric properties. *Neuropsychology* 22(4), 531–544. [PubMed: 18590364]
- Farrell ME, Chen X, Rundle MM, Chan MY, Wig GS, Park DC, 2018 Regional amyloid accumulation and cognitive decline in initially amyloid-negative adults. *Neurology* 91(19), e1809. [PubMed: 30305451]
- Folstein MF, Robins LN, Helzer JE, 1983 The mini-mental state examination. *Archives of General Psychiatry* 40(7), 812–812. [PubMed: 6860082]
- Gibbons LE, Carle AC, Mackin RS, Harvey D, Mukherjee S, Insel P, Curtis SM, Mungas D, Crane PK, 2012 A composite score for executive functioning, validated in Alzheimer's Disease Neuroimaging Initiative (ADNI) participants with baseline mild cognitive impairment. *Brain imaging and behavior* 6(4), 517–527. [PubMed: 22644789]
- Gilewski MJ, Zelinski EM, Schaie KW, 1990 The Memory Functioning Questionnaire for assessment of memory complaints in adulthood and old age. *Psychology and Aging* 5(4), 482–490. [PubMed: 2278670]
- Glodzik-Sobanska L, Reisberg B, De Santi S, Babb JS, Pirraglia E, Rich KE, Brys M, de Leon MJ, 2007 Subjective Memory Complaints: Presence, Severity and Future Outcome in Normal Older Subjects. *Dementia and Geriatric Cognitive Disorders* 24(3), 177–184. [PubMed: 17641528]
- Harrell FE, 2015 Regression modeling strategies : with applications to linear models, logistic regression, and survival analysis. Springer, Cham.
- Jack CR Jr., Knopman DS, Jagust WJ, Petersen RC, Weiner MW, Aisen PS, Shaw LM, Vemuri P, Wiste HJ, Weigand SD, Lesnick TG, Pankratz VS, Donohue MC, Trojanowski JQ, 2013 Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *The Lancet. Neurology* 12(2), 207–216. [PubMed: 23332364]
- Jagust WJ, Landau SM, Koeppe RA, Reiman EM, Chen K, Mathis CA, Price JC, Foster NL, Wang AY, 2015 The Alzheimer's Disease Neuroimaging Initiative 2 PET Core: 2015. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 11(7), 757–771.
- Jiao Y, Stillinger FH, Torquato S, 2008 Modeling heterogeneous materials via two-point correlation functions. II. Algorithmic details and applications. *Physical Review E* 77(3), 031135.
- Jorm AF, Masaki KH, Davis DG, Hardman J, Nelson J, Markesbery WR, Petrovitch H, Ross GW, White LR, 2004 Memory complaints in nondemented men predict future pathologic diagnosis of Alzheimer disease. *Neurology* 63(10), 1960. [PubMed: 15557525]
- Josephs KA, Murray ME, Tosakulwong N, Whitwell JL, Knopman DS, Machulda MM, Weigand SD, Boeve BF, Kantarci K, Petrucelli L, Lowe VJ, Jack CR Jr., Petersen RC, Parisi JE, Dickson DW, 2017 Tau aggregation influences cognition and hippocampal atrophy in the absence of beta-amyloid: a clinico-imaging-pathological study of primary age-related tauopathy (PART). *Acta neuropathologica* 133(5), 705–715. [PubMed: 28160067]
- Kryscio RJ, Abner EL, Cooper GE, Fardo DW, Jicha GA, Nelson PT, Smith CD, Van Eldik LJ, Wan L, Schmitt FA, 2014 Self-reported memory complaints. *Neurology* 83(15), 1359. [PubMed: 25253756]
- La Joie R, Perrotin A, Egret S, Pasquier F, Tomadesso C, Mezenge F, Desgranges B, de La Sayette V, Chetelat G, 2016 Qualitative and quantitative assessment of self-reported cognitive difficulties in nondemented elders: Association with medical help seeking, cognitive deficits, and  $\beta$ -amyloid imaging. *Alzheimer's & dementia (Amsterdam, Netherlands)* 5, 23–34.
- Landau SM, Mintun MA, Joshi AD, Koeppe RA, Petersen RC, Aisen PS, Weiner MW, Jagust WJ, Alzheimer's Disease Neuroimaging I, 2012 Amyloid deposition, hypometabolism, and longitudinal cognitive decline. *Annals of neurology* 72(4), 578–586. [PubMed: 23109153]
- Landau SM, Thomas BA, Thurfjell L, Schmidt M, Margolin R, Mintun M, Pontecorvo M, Baker SL, Jagust WJ, Alzheimer's Disease Neuroimaging I, 2014 Amyloid PET imaging in Alzheimer's disease: a comparison of three radiotracers. *Eur J Nucl Med Mol Imaging* 41(7), 1398–1407. [PubMed: 24647577]
- Mantel N, 1970 Why Stepdown Procedures in Variable Selection. *Technometrics* 12(3), 621–625.

- Mitchell AJ, Beaumont H, Ferguson D, Yadegarfar M, Stubbs B, 2014 Risk of dementia and mild cognitive impairment in older people with subjective memory complaints: meta-analysis. *Acta Psychiatrica Scandinavica* 130(6), 439–451. [PubMed: 25219393]
- Palmqvist S, Scholl M, Strandberg O, Mattsson N, Stomrud E, Zetterberg H, Blennow K, Landau S, Jagust W, Hansson O, 2017 Earliest accumulation of  $\beta$ -amyloid occurs within the default-mode network and concurrently affects brain connectivity. *Nature Communications* 8(1), 1214.
- Perrotin A, Mormino EC, Madison CM, Hayenga AO, Jagust WJ, 2012 Subjective cognition and amyloid deposition imaging: a Pittsburgh Compound B positron emission tomography study in normal elderly individuals. *Archives of neurology* 69(2), 223–229. [PubMed: 22332189]
- Rabin LA, Smart CM, Crane PK, Amariglio RE, Berman LM, Boada M, Buckley RF, Chetelat G, Dubois B, Ellis KA, Gifford KA, Jefferson AL, Jessen F, Katz MJ, Lipton RB, Luck T, Maruff P, Mielke MM, Molinuevo JL, Naeem F, Perrotin A, Petersen RC, Rami L, Reisberg B, Rentz DM, Riedel-Heller SG, Risacher SL, Rodriguez O, Sachdev PS, Saykin AJ, Slavin MJ, Snitz BE, Sperling RA, Tandetnik C, van der Flier WM, Wagner M, Wolfsgruber S, Sikkes SAM, 2015 Subjective Cognitive Decline in Older Adults: An Overview of Self-Report Measures Used Across 19 International Research Studies. *Journal of Alzheimer's disease : JAD* 48 Suppl 1(0 1), S63–S86. [PubMed: 26402085]
- Reisberg B, Shulman MB, Torossian C, Leng L, Zhu W, 2010 Outcome over seven years of healthy adults with and without subjective cognitive impairment. *Alzheimer's & Dementia* 6(1), 11–24.
- Schöll M, Lockhart SN, Schonhaut DR, O'Neil JP, Janabi M, Ossenkoppele R, Baker SL, Vogel JW, Faria J, Schwimmer HD, Rabinovici GD, Jagust WJ, 2016 PET Imaging of Tau Deposition in the Aging Human Brain. *Neuron* 89(5), 971–982. [PubMed: 26938442]
- Snitz BE, Weissfeld LA, Cohen AD, Lopez OL, Nebes RD, Aizenstein HJ, McDade E, Price JC, Mathis CA, Klunk WE, 2015 Subjective Cognitive Complaints, Personality and Brain Amyloidbeta in Cognitively Normal Older Adults. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry* 23(9), 985–993. [PubMed: 25746485]
- Tewari A, Gokhale AM, Spowart JE, Miracle DB, 2004 Quantitative characterization of spatial clustering in three-dimensional microstructures using two-point correlation functions. *Acta Materialia* 52(2), 307–319.
- Villeneuve S, Rabinovici GD, Cohn-Sheehy BI, Madison C, Ayakta N, Ghosh PM, La Joie R, Arthur-Bentil SK, Vogel JW, Marks SM, Lehmann M, Rosen HJ, Reed B, Olichney J, Boxer AL, Miller BL, Borys E, Jin L-W, Huang EJ, Grinberg LT, DeCarli C, Seeley WW, Jagust W, 2015 Existing Pittsburgh Compound-B positron emission tomography thresholds are too high: statistical and pathological evaluation. *Brain : a journal of neurology* 138(Pt 7), 2020–2033. [PubMed: 25953778]
- Vogel JW, Varga Doležalová M, La Joie R, Marks SM, Schwimmer HD, Landau SM, Jagust WJ, 2017 Subjective cognitive decline and  $\beta$ -amyloid burden predict cognitive change in healthy elderly. *Neurology* 89(19), 2002–2009. [PubMed: 28986416]

**Highlights**

- The best models for predicting SCD domains of memory and organization include A $\beta$ .
- The best models for SCD domains of planning and visuospatial skills include tau.
- Detailed characterization of SCD may help predict the AD pathological designations.

**TABLE 1.**Subject demographics, cognitive performance measures, APOE- $\epsilon$ 4 status, and A $\beta$ -positivity

<b>Parameters</b>	<b>Values</b>
Subjects, n	86
Females, n	39
Age, mean yr $\pm$ SD	78 $\pm$ 8
APOE- $\epsilon$ 4 carriers, n	24
Global [ $^{18}$ F]florbetapir uptake, mean SUVR $\pm$ SD	1.2 $\pm$ 0.2
A $\beta$ -positive (based on 1.11 cutoff for global florbetapir SUVR)	38
<b>Objective cognitive performance tests (mean <math>\pm</math> SD)</b>	
MMSE	28 $\pm$ 2
ADAS-cog	8 $\pm$ 4
ADNI-MEM	0.9 $\pm$ 0.6
ADNI-EF	0.9 $\pm$ 0.8
<b>Self-reported ECog measures (mean <math>\pm</math> SD)</b>	
Everyday Memory	1.7 $\pm$ 0.5
Everyday Language	1.5 $\pm$ 0.5
Everyday Visuospatial Abilities	1.0 $\pm$ 0.3
Everyday Planning	1.2 $\pm$ 0.4
Everyday Organization	1.3 $\pm$ 0.4
Everyday Divided Attention	1.5 $\pm$ 0.6

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 2.**

Associations of regional [<sup>18</sup>F]flortaucipir SUVR values with objective cognitive performance measures and self-reported ECog measures. All associations included age, gender, and APOE-ε4 status as covariates.

	<b>Outcome</b>	<b>Predictor: Regional flortaucipir SUVR</b>	<b>β</b>	<b>SE</b>	<b>P-value</b>	<b>R<sup>2</sup></b>	
	<b>MMSE</b>	Braak I	-3.9	1.45	<b>0.008*</b>	0.20	
		Braak III-IV	-4.7	1.93	<b>0.018*</b>	0.18	
		Braak V-VI	-4.9	2.67	<b>0.07*</b>	0.15	
	<b>ADAS_cog</b>	Braak I	4.70	3.14	0.14	0.22	
		Braak III-IV	2.72	4.21	0.52	0.21	
		Braak V-VI	2.73	5.71	0.63	0.21	
<b>Objective Cognitive Measures</b>	<b>ADNI-EF</b>	Braak I	-1.08	0.63	<b>0.09*</b>	0.18	
		Braak III-IV	-1.48	0.83	<b>0.08*</b>	0.18	
		Braak V-VI	-1.90	1.13	<b>0.10*</b>	0.18	
	<b>ADNI-MEM</b>	Braak I	-0.38	0.52	0.45	0.16	
		Braak III-IV	-0.23	0.68	0.73	0.16	
		Braak V-VI	-0.95	0.92	0.31	0.17	
	<b>Everyday Memory</b>	Braak I	0.63	0.43	0.14	0.16	
		Braak III-IV	0.42	0.57	0.46	0.14	
		Braak V-VI	0.69	0.77	0.37	0.15	
	<b>Everyday Language</b>	Braak I	0.36	0.44	0.42	0.11	
		Braak III-IV	0.33	0.59	0.58	0.10	
		Braak V-VI	0.58	0.79	0.47	0.11	
<b>Subjective Cognitive Measures</b>	<b>Everyday Visuospatial</b>	Braak I	0.43	0.26	<b>0.09*</b>	0.11	
		Braak III-IV	0.25	0.34	0.47	0.09	
		Braak V-VI	0.45	0.46	0.33	0.10	
	<b>Everyday Planning</b>	Braak I	0.78	0.30	<b>0.011*</b>	0.19	
		Braak III-IV	0.76	0.41	<b>0.07*</b>	0.15	
		Braak V-VI	1.30	0.54	<b>0.018*</b>	0.17	
	<b>Everyday Organization</b>	Braak I	0.64	0.34	<b>0.06*</b>	0.08	
		Braak III-IV	0.47	0.46	0.30	0.05	
		Braak V-VI	0.47	0.62	0.45	0.04	
	<b>Everyday Divided Attention</b>	Braak I	-0.07	0.49	0.89	0.11	
		Braak III-IV	0.11	0.66	0.86	0.11	
			Braak V-VI	0.52	0.89	0.56	0.12

Significance codes (Bonferroni-adjusted P-value 0.05/3= 0.016): 0

\*,\*  
0.016

\*,\*  
0.1

\*,\*  
1

**TABLE 3.**

Associations of regional tau accumulation with objective cognitive performance measures and self-reported ECog (age, sex, and APOE- $\epsilon$ 4 status included as covariates).

	Outcome	Predictor: Regional	$\beta$	SE	P-value	R <sup>2</sup>
		<b>florbetapir SUVR</b>				
<b>Objective Cognitive Measures</b>	<b>MMSE</b>	Parietal	-4.45	0.91	<b>&lt;0.0001*</b>	0.33
		Frontal	-3.95	0.99	<b>0.0001*</b>	0.27
		Temporal	-4.25	1.15	<b>0.0004*</b>	0.25
		Cingulate	-3.69	1.00	<b>0.0004*</b>	0.25
	<b>ADAS cog</b>	Parietal	7.57	2.02	<b>0.0003*</b>	0.33
		Frontal	7.58	2.10	<b>0.0005*</b>	0.32
	<b>ADNI-EF</b>	Temporal	8.49	2.42	<b>0.0007*</b>	0.32
		Cingulate	6.95	2.14	<b>0.002*</b>	0.30
		Parietal	-0.96	0.43	<b>0.027*</b>	0.20
		Frontal	-1.08	0.44	<b>0.016*</b>	0.21
	<b>ADNI-MEM</b>	Temporal	-0.93	0.51	<b>0.07*</b>	0.18
		Cingulate	-0.90	0.45	<b>0.05*</b>	0.19
		Parietal	-0.74	0.35	<b>0.03*</b>	0.21
		Frontal	-0.80	0.36	<b>0.03*</b>	0.21
	<b>Everyday Memory</b>	Temporal	-0.85	0.42	<b>0.04*</b>	0.20
		Cingulate	-0.68	0.36	<b>0.06*</b>	0.19
Parietal		0.73	0.29	<b>0.0126*</b>	0.21	
Frontal		0.76	0.29	<b>0.0125*</b>	0.21	
<b>Everyday Language</b>	Temporal	0.78	0.34	<b>0.026*</b>	0.19	
	Cingulate	0.74	0.30	<b>0.016*</b>	0.20	
	Parietal	0.51	0.30	<b>0.098*</b>	0.13	
	Frontal	0.49	0.31	0.12	0.13	
<b>Subjective Cognitive Measures</b>	Temporal	0.59	0.36	0.11	0.13	
	Cingulate	0.44	0.32	0.17	0.12	
	<b>Everyday Visuospatial</b>	Parietal	0.27	0.18	0.13	0.11
	Frontal	0.33	0.18	<b>0.07*</b>	0.12	
<b>Everyday Planning</b>	Temporal	0.25	0.21	0.24	0.10	
	Cingulate	0.29	0.19	0.12	0.11	
	Parietal	0.17	0.21	0.43	0.11	
	Frontal	0.13	0.22	0.57	0.11	
	Temporal	0.08	0.26	0.77	0.10	
	Cingulate	0.11	0.22	0.61	0.11	

Outcome	Predictor: Regional florbetapir SUVR	$\beta$	SE	P-value	R <sup>2</sup>
<b>Everyday</b>	Parietal	0.65	0.22	<b>0.006*</b>	0.13
<b>Organization</b>	Frontal	0.65	0.24	<b>0.008*</b>	0.12
	Temporal	0.74	0.27	<b>0.008*</b>	0.12
	Cingulate	0.67	0.23	<b>0.007*</b>	0.13
<b>Everyday Divided</b>	Parietal	0.05	0.34	0.86	0.11
<b>Attention</b>	Frontal	0.07	0.34	0.85	0.11
	Temporal	0.04	0.41	0.91	0.11
	Cingulate	0.05	0.35	0.89	0.11

Significance codes (Bonferroni-adjusted P-value 0.05/4= 0.0125): 0

\*\*\*  
0.0125

\*\*  
0.1

\*  
1

**TABLE 4:**

The strongest models for predicting each objective cognitive measure (MMSE, ADAS\_Cog, ADNI-EF, ADNI-MEM) and each ECog domain (Memory, Language, Visuospatial, Planning, Organization, Divided Attention).

Outcome	All CN subjects		A $\beta$ -negative CN subjects	
	Best model	Adjusted R <sup>2</sup>	Best model	Adjusted R <sup>2</sup>
Objective Cognitive Measures				
MMSE	A $\beta$ (temp <sup>*</sup> , par <sup>*</sup> ), Age <sup>*</sup>	0.34	Age <sup>*</sup> , A $\beta$ (front <sup>˘</sup> )	0.14
ADAS-Cog	A $\beta$ (par <sup>*</sup> ), Age <sup>*</sup>	0.29	Age <sup>*</sup> , A $\beta$ (temp)	0.18
ADNI-EF	A $\beta$ (front <sup>*</sup> ), Age <sup>*</sup>	0.18	Age <sup>*</sup> , APOE- $\epsilon$ 4, Tau(Braak III-IV <sup>˘</sup> ), A $\beta$ (par <sup>˘</sup> , temp <sup>˘</sup> , cing <sup>*</sup> , front <sup>*</sup> )	0.31
ADNI-MEM	A $\beta$ (front <sup>*</sup> ), Age <sup>*</sup> , Tau(Braak III-IV, Braak V-VI)	0.19	A $\beta$ (front <sup>*</sup> , cing <sup>*</sup> ), Sex <sup>˘</sup> , Age, Tau(Braak III-IV)	0.20
Subjective Cognitive Measures				
Everyday Memory	A $\beta$ (par <sup>*</sup> ), Age, Sex	0.17	A $\beta$ (cing <sup>˘</sup> ), Age <sup>˘</sup> , APOE- $\epsilon$ 4	0.11
Everyday Language	Age, Sex, A $\beta$ (par <sup>˘</sup> )	0.09	None	
Everyday Visuospatial	Age <sup>-</sup> , Tau(Braak I <sup>*</sup> , Braak III-IV <sup>*</sup> , Braak V-VI)	0.11	Age, Tau (Braak I <sup>*</sup> )	0.07
Everyday Planning	Age <sup>*</sup> , Tau(Braak I <sup>*</sup> , Braak III-IV <sup>*</sup> , Braak V-VI <sup>*</sup> )	0.19	Age <sup>-</sup> , Tau (Braak I <sup>*</sup> ), APOE- $\epsilon$ 4	0.18
Everyday Organization	A $\beta$ (par <sup>*</sup> )	0.10	Tau (Braak III-IV <sup>*</sup> , Braak V-VI), A $\beta$ (cing <sup>*</sup> , temp <sup>*</sup> ), APOE- $\epsilon$ 4	0.28
Everyday Divided Attention	Age <sup>-</sup> , APOE- $\epsilon$ 4 <sup>*</sup>	0.08	Tau(Braak III-IV <sup>*</sup> , Braak V-VI <sup>*</sup> ), Sex <sup>˘</sup>	0.16

Blue Cells are models that included only A $\beta$  with p<0.05.

Green cells are models that included only tau (p<0.05).

Red color present models that included both pathologies.

Significance codes: 0

“\*” 0.05

“.” 0.1

“ ” 1