



Published in final edited form as:

Toxicol Appl Pharmacol. 2019 October 01; 380: 114709. doi:10.1016/j.taap.2019.114709.

Differential activity of BPA, BPAF and BPC on zebrafish estrogen receptors *in vitro* and *in vivo*

Caroline Pinto¹, Ruixin Hao¹, Marina Grimaldi², Savini Thrikawala¹, Abdelhay Boulahtouf², Selim Aït-Aïssa³, François Brion³, Jan-Åke Gustafsson⁴, Patrick Balaguer^{5,#,*}, Maria Bondesson^{6,#}

¹Center for Nuclear Receptors and Cell Signaling, Department of Biology and Biochemistry, University of Houston, Houston, TX 77204, USA.

²Institut de Recherche en Cancérologie de Montpellier, INSERM U1194, Institut régional du Cancer de Montpellier, Université de Montpellier, 34298 Montpellier, Cedex 5, France.

³Institut National de l'Environnement Industriel et des risques (INERIS), Unité Ecotoxicologie in vitro et in vivo, UMR-I 02 SEBIO, 60550 Verneuil-en-Halatte, France.

⁴Center for Nuclear Receptors and Cell Signaling, Department of Biology and Biochemistry, University of Houston, Houston, TX 77204, USA; Department of Biosciences and Nutrition, Karolinska Institutet, 14183 Huddinge, Sweden.

⁵Institut de Recherche en Cancérologie de Montpellier, INSERM U1194, Institut régional du Cancer de Montpellier, Université de Montpellier, 34298 Montpellier, Cedex 5, France.

⁶Department of Intelligent Systems Engineering, Indiana University, Bloomington, IN 47408, USA.

Abstract

The high volume production compound bisphenol A (BPA) is of environmental concern largely because of its estrogenic activity. Consequently, BPA analogues have been synthesized to be considered as replacement molecules for BPA. These analogues need to be thoroughly evaluated for their estrogenic activity. Here, we combined mechanism zebrafish-based assays to examine estrogenic and anti-estrogenic activities of BPA and two of its analogues, bisphenol AF (BPAF) and bisphenol C (BPC) *in vitro* and *in vivo*. *In vitro* reporter cell lines were used to investigate agonistic and antagonistic effects of the three bisphenols on the three zebrafish estrogen receptors. The transgenic *Tg(5×ERE:GFP)* and *Cyp19a1b-GFP zebrafish* lines were then used to analyze estrogenic and anti-estrogenic responses of the three bisphenols *in vivo*. BPA, BPAF and BPC were agonists with different potencies for the three zebrafish estrogen receptors *in vitro*. The

*Corresponding author at: IRCM, INSERM U1194, Parc Euromédecine, 208 rue des Apothicaires, 34298 Montpellier Cedex 5, France. patrick.balaguer@inserm.fr.

#Patrick Balaguer and Maria Bondesson share last authorship

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

potent zfER α -mediated activity of BPA and BPAF *in vitro* resulted *in vivo* by activation of GFP expression in zebrafish larvae in the heart (zfER α -dependent) at lower concentrations, and in the liver (zfER β -dependent) at higher concentrations. BPC induced zfER β -mediated luciferase expression *in vitro*, and the zfER β agonism led to activation of GFP expression in the liver and the brain *in vivo*. In addition, BPC acted as a full antagonist on zfER α , and completely inhibited estrogen-induced GFP expression in the heart of the zebrafish larvae. To summarize, applying a combination of zebrafish-based *in vitro* and *in vivo* methods to evaluate bisphenol analogues for estrogenic activity will facilitate the prioritization of these chemicals for further analysis in higher vertebrates as well as the risk assessment in humans.

Keywords

zebrafish; bisphenol; estrogen receptor; transgenic fish; reporter cell lines

Introduction

The environmental pollutant bisphenol A (BPA) has been widely studied during the past decade. BPA has since the 1950s been used to make epoxy resin and polycarbonate plastics, and it is one of the highest volume chemicals produced worldwide. BPA is found in products such as reusable food and beverage containers, lining of cans, dental materials, computers, and thermal paper. Through its estrogenic activity, BPA has been shown to cause deleterious effects in laboratory animals, such as abnormal fetal development (Palanza *et al.*, 2008), alterations in sex differentiation (Rubin *et al.*, 2006), reproduction (Honma *et al.*, 2002), brain function and mood (Leranth *et al.*, 2008; Elsworth *et al.*, 2013) as well as promotion of certain types of cancers (Markey *et al.*, 2001; Munoz-de-Toro *et al.*, 2005; Ho *et al.*, 2006). In humans, epidemiological studies have shown that high BPA levels are linked to heart disease, diabetes and liver toxicity ((Lang *et al.*, 2008) and reviewed in (Vandenberg *et al.*, 2012)). Food, indoor air, indoor dust, wastewater and dermal absorption are all important sources of human exposure to BPA (Vandenberg *et al.*, 2007; Zalko *et al.*, 2011). In addition, BPA is commonly found as a contaminant in wastewater, and may as such have deleterious effects on wildlife (Santos *et al.*, 2016; Xu *et al.*, 2016).

Concurrent with the increasing awareness of potential health problems linked to BPA, a search for safer alternatives has begun. Two of these BPA alternatives, bisphenol AF (BPAF) and bisphenol C (BPC), are the focus of this study. BPAF is a fluorinated derivative of BPA, and it is used both as a crosslinking agent and for polymer synthesis in plastics industry. It has an annual estimated production of 10,000–500,000 pounds in the United States (NTP-NIEHS, 2008). BPAF has been detected in wastewater from water treatment plants and was also shown to be resistant to biodegradation (Sun *et al.*, 2017). BPC is used in the preparation of flame-resistant polycarbonates, and it has also been detected in wastewater (Cesen *et al.*, 2018).

In humans, BPA interferes with estrogen signaling mediated by the two estrogen receptors, ER α and ER β . The canonical estrogen signaling pathway involves estrogen binding to the ligand binding domain (LBD) of the ERs, translocation of the ER-ligand complexes into the

nucleus, followed by recruitment of co-regulator complexes and docking to the estrogen response element (ERE) of target gene promoters (reviewed in (Nilsson *et al.*, 2001)). Similarly to BPA, *in vitro* studies show that several bisphenol analogues are able to bind and activate the human ERs [(Delfosse *et al.*, 2012; Delfosse *et al.*, 2014; Grimaldi *et al.*, 2019) and reviewed in (Chen *et al.*, 2016)]. BPAF has been repeatedly reported to activate ER-dependent transcription at lower concentrations than BPA and other bisphenol derivatives *in vitro* ((Delfosse *et al.*, 2012) and reviewed in (Chen *et al.*, 2016)). While fewer studies have investigated BPC, this chlorinated analogue has been described to have stronger affinity for the ERs than BPA, acting as a full agonist on MCF7 proliferation, and as both an agonist and antagonist in HELN (HeLa ER negative cells stably transfected by the ERE-Luc plasmid) cells stably-expressing the human ERs (HELN hERs cells) (Delfosse *et al.*, 2012).

Zebrafish (*Danio rerio*) is an *in vivo* model to study environmental pollutants. There are three ERs (zfER α , zfER β 2 and zfER β 1) in zebrafish, with an approximate 50% amino-acid sequence identity to the human ER orthologues (Menuet *et al.*, 2002). We have previously generated cell lines stably-transfected with the zfER subtypes to study the potential estrogenic activity of environmental pollutants in zebrafish (Pinto *et al.*, 2014). We demonstrated with these cell models that mammalian estrogenic ligands also modulate the activity of the zfERs, although the selectivity and potency of the ligand often varied for the human and zebrafish orthologues. In zebrafish, the three ERs are all expressed from embryonic stages to adulthood, but with different temporal and organ distributions (Bardet *et al.*, 2002; Chandrasekar *et al.*, 2010; Mouriec *et al.*, 2009). zfER α is detected in the heart valves of the developing zebrafish larvae, while the zfER β 1 subtype is robustly expressed in the liver (Gorelick *et al.*, 2014). In the developing brain, zebrafish embryos express detectable levels of zfER β 2 and zfER β 1, but not zfER α , in different brain regions. The expression of *cyp19a1b* gene was temporally correlated with that of zfER β 2 and zfER β 1 (Mouriec *et al.*, 2009) suggesting that these receptors drive the *cyp19a1b* expression.

During the past few years, several transgenic zebrafish models have been developed to investigate the cell- and/or tissue-specific effects of chemicals on estrogen signaling pathways in developing fish. For instance, in transgenic zebrafish model containing five EREs (*Tg(5 \times ERE:GFP)*) (Gorelick and Halpern, 2011), the GFP expression is clearly detected in tissues expressing the estrogen receptors. Thus, compounds with *in vitro* selectivity for zfER α and zfER β s induce expression of the fluorescent protein in the heart and liver, respectively (Gorelick *et al.*, 2016). For example, PPT specifically activates the zfER β s receptors *in vitro* and induces GFP expression only in the liver of zebrafish larvae, whereas the pan-agonist 17 α -ethinylestradiol induces GFP in both heart and liver (Gorelick and Halpern, 2011; Pinto *et al.*, 2014; Gorelick *et al.*, 2016). Another reporter fish model, the transgenic *cyp19a1b-GFP* zebrafish, expresses GFP under the control of the ER-regulated *cyp19a1b* (brain aromatase) gene in radial glial cells in the brain (Tong *et al.*, 2009). This fish model has been thoroughly evaluated for assessing the estrogenic activity of a panel of environmental contaminants - including natural and synthetic hormones, pharmaceuticals, pesticides, industrial chemicals- in embryos and allowed to demonstrate that numerous contaminants were able to disrupt the brain-specific expression of the *cyp19a1b* gene in an ER-dependent manner (Brion *et al.*, 2012; Cano-Nicolau, *et al.*, 2016; Le Fol *et al.*, 2017).

In this article, we combined *in vitro* and *in vivo* zebrafish mechanism-based reporter gene assays as screening tools to assess the estrogenic properties of BPA, BPAF and BPC. HELN cells expressing the zfERs (HELN zfERs cells) were first used to evaluate the capacity of the selected bisphenols to bind to the three zfERs subtypes and to transactivate zfERs to induce luciferase activity. To further evaluate their estrogenic properties at the organism level, (*Tg(5×ERE:GFP)*) and *tg(cyp19a1b-GFP)* zebrafish embryos were exposed to BPA, BPAF and BPC and the tissue-specific estrogenic responses in the brain, heart and liver of zebrafish embryos were observed and/or quantified and interpreted in light of their *in vitro* estrogenic activities towards the zfER subtypes.

Materials and Methods

Materials

Tissue culture plates were obtained from Greiner Bio-one (Monroe, NC, USA), and media was purchased from Invitrogen (Grand Island, NY, USA). Luciferin (sodium salt) was purchased from Promega (Charbonnières, France). The chemicals 17 β -estradiol (E2), 17 α -ethinylestradiol (EE2), bisphenol A (CAS no. 80–05-7), bisphenol AF (BPAF; CAS no. 1478–61-1), and bisphenol C (BPC; CAS no. 14868–03-2) were purchased from Sigma-Aldrich (Saint-Quentin Fallavier, France). ICI 182,780 (7 α ,17 β -[9[(4,4,5,5,5-pentafluoropentyl)sulfinyl]nonyl]estra-1,3,5(10)-triene-3,17-diol, fulvestrant) was obtained from Zeneca (Macclesfield, UK), and 3,17-Dihydroxy-19-nor-17 α -pregna-1,3,5(10)-triene-21,16 α -lactone (16 α -LE2) was a kind gift of Peter Muhn from Research Laboratories of Schering AG (Berlin, Germany). Compounds were dissolved in dimethyl sulfoxide (DMSO) as 10⁻² M stock solutions.

Cell lines

The establishment and culturing of the HELN zfER α (Esr1), zfER β 1 (Esr2b) and zfER β 2 (Esr2a) reporter cell lines have been previously described (Pinto et al. 2014). Briefly, the HELN-zfER α , β 1 and β 2 cell lines were obtained by transfection of HELN cells (HeLa cells stably transfected with the ERE- β Globin-Luc-SVNeo plasmid) (Delfosse *et al.*, 2012) by the corresponding pSG5-puro plasmids (pSG5-zfER α -puro, pSG5-zfER β 1-puro and pSG5-zfER β 2-puro, respectively).

The HELN zfER cells were maintained in phenol red-free Dulbecco's Modified Eagle's Medium/F12 (DMEM/F12) and supplemented with 5% dextran-coated charcoal-treated fetal bovine serum (FBS-DCC), 1% antibiotic (penicillin/streptomycin), geneticin (1 mg/mL) and puromycin (0.25 μ g/mL) in a 5% CO₂ humidified atmosphere at 37°C.

zfERs binding assays

The assay for measuring zfER binding affinity has been previously described (Pinto *et al.*, 2014). Briefly, the HELN-zfERs cell lines were treated with 3 nM [³H]-E2 in the absence or in the presence of chemicals for 4h at 28°C. At the end of the incubation period, unbound material was aspirated, cells washed with PBS, and [³H]-bound radioactivity counted. Protein concentrations were used to normalize bound radioactivity values expressed in

desintegrations per minute (dpm). Experiments were performed in quadruplicate in at least two independent experiments.

Transactivation assays

Chemicals successive dilutions were performed in cell culture medium, using the same protocol for treatments as described previously (Pinto *et al.*, 2014). The compounds were added 8 h after seeding cells into 96-well plates, and cells were incubated with compounds for 16h at 28°C. Luciferase assays were performed as previously published (Pinto *et al.*, 2014), and experiments were performed in quadruplicates in three independent experiments. Data were expressed as percentage of the maximal activity induced by E2 10 nM. EC50 values were measured using GraphPad Prism (GraphPad Software Inc). Datas were analyzed for significant differences using one-way ANOVA followed by Dunnet's post-comparison test (vs.control) (**p<0.001, *p<0.01, *p<0.05). Differences were considered statistically significant at P <0.05.

Zebrafish maintenance and transgenic zebrafish Tg(5xERE:GFP) treatment

Transgenic fish *Tg(5xERE:GFP)* (Gorelick and Halpern, 2011) were used according to the maintenance and experimental protocols approved by the Institutional Animal Care and Use Committee at the University of Houston. Adult zebrafish were maintained and fed as described previously (Hao *et al.*, 2013).

Embryos from several pairs of adult fish were collected after spawning and cultured in a Petri dish in E3 embryo medium (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl₂, 0.33 mM MgSO₄) at 28.5°C with 14:10 LD. For fluorescent imaging experiments, *Tg(5xERE:GFP)* zebrafish embryos were incubated in E3 medium with 0.003% 1-phenyl-2-thiourea (PTU) (Sigma-Aldrich, St. Louis, MO) starting at 24h post fertilization (hpf) to inhibit pigment formation. Samples of 10 zebrafish larvae at 3 days post fertilization (dpf) were treated for 24 h with EE2, 16 α -LE2 and bisphenols in 24-well plate, and transferred to 96-well plates at a density of one larva per well for imaging. EE2, 16 α -LE2, ICI 182,780 and bisphenols in 100% DMSO were diluted in embryo media, and vehicle concentration did not exceed 0.2% DMSO in treatments. Vehicle only was used as a negative control. The larvae were imaged under 10X magnification lenses using a fluorescent microscope (Olympus IX51) equipped with an Olympus XM10 camera (Olympus, Center Valley, PA).

Treatment of *cyp19a1b*-GFP transgenic zebrafish embryos

Newly fertilized transgenic *cyp19a1b-GFP* zebrafish eggs were collected and treated as previously described (Brion *et al.*, 2012). Briefly, 20 embryos (< 4 h post-fertilization) were exposed from 0 to 4 dpf to DMSO (0.01% v:v), 0.05 nM EE2 (used as positive control) or a range of concentrations of BPA, BPAF or BPC. The exposure medium was renewed totally every 24 h. At the end of the exposure period, the zebrafish embryos were collected for *in vivo* fluorescence imaging of the brain. Each 4 dpf-old zebrafish was photographed in dorsal view under a fluorescence microscope connected to an AxioCam Mrm camera (Zeiss GmbH, Gottingen, Germany). All photographs were taken with the same parameters. The fluorescence of the regions of interest was then quantified by image analysis using an ImageJ macro as previously described (Brion *et al.*, 2012).

RNA extraction and quantitative real-time PCR

Zebrafish larvae were treated with bisphenols and controls from 4 dpf to 6 dpf. Treatments were repeated once at 5 dpf. Total RNA from 40 pooled embryos were collected at 6 dpf using Trizol (Invitrogen Corporation, Carlsbad, CA) and RNeasy spin columns (Qiagen, Chatsworth, CA) according to manufacturer's protocols. DNase I (Qiagen, Chatsworth, CA) was used to remove any DNA in the column. RNA concentration was measured using NanoDrop 1000 spectrophotometer (Agilent Technologies, Palo Alto, CA) and 100 ng of total RNA was used for cDNA synthesis using Superscript III reverse transcriptase (Invitrogen Corporation, Carlsbad, CA) with random hexamer primers.

Expression of Cyp19a1b and Vtg1 were measured by qPCR using an ABI 7500 Fast Real-Time PCR system (Applied Biosystems, Foster City, CA) with iTaq Universal SYBR Green Supermix (Bio-Rad, Hercules, CA) according to the manufacturer's protocol. Primers were synthesized by Integrated DNA Technologies, Inc (Coralville, IA). Primer sequences were for vtg1 Forward (5'-3') ACTACCAACTGGCTGCTTAC and Reverse (5'-3') ACCATCGGCACAGATCTTC; and for cyp19a1b Forward (5'-3') AAAGAGTTACTAATAAAGATCCACCGGTAT and Reverse (5'-3') TCCACAAGCTTTCCATTCA (Hao *et al.*, 2013). Expression levels were normalized to zebrafish β -actin using the Ct method. Student's t-test was performed for statistical analysis using GraphPad Prism version 7 (GraphPad Software, La Jolla, CA).

Results

Transcriptional activity of bisphenols mediated by zfERs *in vitro*

The transcriptional activity of BPA, BPAF and BPC have been reported for human estrogen receptors in HELN cells (Delfosse *et al.*, 2012). To investigate the activity of BPA and its analogues on zebrafish ERs, we treated the HELN cell lines expressing the zfER α , zfER β 1 and zfER β 2 receptors and an ERE-driven luciferase reporter with increasing concentrations of the bisphenols. BPA and BPAF activated the three zfER reporter cells, reaching approximately 53–89% of E2-induced luciferase expression (Figure 1A, B). These bisphenols displayed higher selectivity for zfER α , than for the zfER β subtypes. BPAF was more potent than BPA in inducing luciferase expression, and displayed approximately 3.1–5.6-fold greater potency on the three zfER cells than BPA, with EC50s of 84, 758 and 897 nM for zfER α , zfER β 1 and zfER β 2, respectively (Table 1). BPC activated both zfER α and zfER β 1 at low concentrations, but the luciferase activity reached only 31 and 36% of the maximal luciferase activity, respectively. BPC induced transcription to 59% of maximal activation on zfER β 2 (Figure 1C). The measured EC50s and maximal activation for the BPA, BPAF, BPC and controls are shown in Table 1.

BPA, BPAF and BPC were identified as partial agonists in HELN hERs cells (Delfosse *et al.*, 2012). The partial bisphenols induced-luciferase activity in the human ER reporter cells and the marginal effect of BPC on zfER α and zfER β 1 (figure 1C) suggest that BPC can act as an antagonist for these zfERs. We therefore evaluated the ability of BPC to antagonize the activity of E2. We found that BPC clearly reduced E2-induced transactivation in the zfER α and zfER β cell lines with IC50s of 46, 160 and 136 nM on zfER α , zfER β 1 and zfER β 2,

respectively (Figure 1D; Table 1) thereby revealing a stronger antagonistic activity of BPC towards zfER α and zfER β 1.

The selectivity of BPA, BPAF and BPC for zfER α was confirmed in competitive binding assays with radiolabeled E2 (Table 2). As previously demonstrated for the selective zfER α compound 16 α -LE2 (Pinto *et al.*, 2014), bisphenols displayed stronger binding affinity for zfER α compared to the zfER β subtypes. Additionally, BPAF and BPC displaced radiolabeled E2 at lower concentrations than BPA, confirming the stronger affinity of BPAF and BPC for the zebrafish ERs relative to BPA (Table 2).

Activation of estrogen signaling by bisphenols *in tg ERE-GFP embryos*

To assess the estrogenic effects of BPA, BPAF and BPC *in vivo* in zebrafish embryos, the transgenic fish line *Tg(5xERE:GFP)* containing 5xERE upstream of a *c-fos* promoter and a GFP reporter was used (Gorelick and Halpern, 2011). In order to qualitatively investigate the tissue-specific activity of the bisphenols, we treated 3 dpf zebrafish larvae for 24 h with BPA, BPAF and BPC (Figure 2). The ligand 16 α -LE2, whose zfER α selectivity *in vitro* is associated with *in vivo* GFP expression exclusively in the heart valves at lower doses (10–100 nM) (Figure 2C–D), was used to compare the tissue-specific effects resulting from treatment with the bisphenols. The zfER α -selective agonist activity of BPA and BPAF was confirmed in the zebrafish embryos. At lower concentrations, BPA and BPAF only activated GFP expression in the heart valves, while higher doses of the ligands were required for induction of liver expression (Figure 2G–J and 2K–N). Treatment with BPC induced GFP expression in the liver at 1 μ M, and no heart GFP expression was observed, likely due to its only marginal activity on zfER α (Figure 2O–R).

The stronger potency of BPAF and BPC on the zfERs compared to BPA observed *in our experiments in vitro* was confirmed *in vivo*. Treatment with 100 nM BPAF induced GFP expression in the heart, whereas a dose of 1 μ M BPA was required to achieve a similar effect in the zebrafish embryos. At the hepatic level, GFP expression was induced after exposure to 1 μ M BPAF or 1 μ M BPC, while BPA was only able to activate hepatic GFP expression at 10 μ M (Figure 2G–R). Therefore, the responses observed *in vivo* seem to reflect both the estrogenic activity reported *in vitro* as well as the selectivity of the compounds towards the zfERs. Thus, the stronger potencies of BPAF and BPC compared to BPA seen *in vitro* was also observed at the *in vivo* level.

To investigate the antagonist activity of BPC *in vivo*, we co-treated zebrafish embryos with the estrogenic ligand EE2, which is more stable than E2 *in vivo*, or 16 α -LE2 in combination with BPC (Figure 3 and Figure 4). Treatment with 100 pM EE2 induced GFP expression both in the heart and liver of the embryos, in accordance with its pan-agonist ER activity (Figure 3C) (Pinto *et al.*, 2014; Gorelick *et al.*, 2016). The pure antagonist ICI 182,780 completely blocked both heart and liver GFP expression (Figure 3E). Treatment with 1 μ M BPC induced GFP expression in the liver of the embryos in accordance with its zfER β 2 activity. Co-treatment of EE2 with 1 μ M BPC completely blocked EE2-induced GFP heart expression whereas it only slightly reduced EE2-induced GFP liver expression (Figure 3G). Likewise, BPC was capable of completely inhibiting the GFP expression induced by 10 nM 16 α -LE2 in the heart (Figure 4). These results are in accordance with the *in vitro* antagonist/

agonist activity of BPC, indicating that BPC is an antagonist on zER α and partial agonist on zER β s *in vivo*.

Activation of the brain aromatase promoter in the transgenic cyp19a1b- GFP zebrafish embryos by the three bisphenols

To assess the effect of bisphenols on the expression of the estrogen-sensitive brain aromatase gene, the fluorescence signal in the developing brain of transgenic cyp19a1b-GFP zebrafish embryos was quantified after a 4-day treatment with BPA, BPAF and BPC. All bisphenols induced GFP expression in the brain of the zebrafish embryos (Figure 5A–C). BPA at 10 μ M promoted a 28-fold induction in the GFP signal, whereas no or very weak activity was observed at lower concentrations. Exposure to BPAF or BPC at 1 μ M was sufficient to promote a 18-fold induction in the GFP expression, approximately, and a 2.9-fold and 5.5-fold GFP activation was observed at 0.1 μ M for BPAF and BPC, respectively (Figure 5A).

To assess the potential for the bisphenols to inhibit estrogen-mediated activation of the brain aromatase promoter, the cyp19a1b-GFP zebrafish embryos were treated with 0.05 nM EE2 in the presence of the bisphenols. Under these experimental conditions, BPA, BPAF and BPC were unable to inhibit the EE2-mediated induction of GFP in the zebrafish embryos. These results indicate that BPA, BPAF and BPC act as agonists to induce brain aromatase expression within the radial glial cell context but that the bisphenols lack of ER antagonist activity on the brain aromatase of zebrafish embryos (Figure 5B–C).

Endogenous estrogen target gene activation by bisphenols *in vivo*

To analyze the effect of BPA, BPAF and BPC exposure on the expression of endogenous estrogen target genes, we performed quantitative PCR. We investigated the expression of the estrogen-sensitive markers vtg1 gene, highly expressed in the liver, and the brain cyp19a1b. In concordance with the *in vivo* GFP expression data, BPAF and BPC activated vtg1 and cyp19a1b at lower concentrations than BPA (clearly detected at 1 μ M vs. 10 μ M, respectively) (Figures 6A and 6B). We conclude that the *in vivo* endogenous activation of estrogenic target genes by BPA analogs is in accordance with both the brain and liver *in vivo* GFP activation.

Discussion

Evidence in the literature showing that exposure to BPA causes harmful health effects in humans and wildlife has prompted a search for safer bisphenol alternatives. We assessed the effects of BPA, BPAF and BPC using zER reporter cell lines, and found that BPAF and BPC in comparison to BPA have a stronger binding affinity for the zebrafish ERs, and a lower EC50 to activate them. *In vitro*, BPAF and BPA activated luciferase expression mediated by the three zebrafish estrogen receptors, with stronger affinity for zER α . This zER α selectivity was confirmed by the preferential activation of the GFP reporter in the heart valves *in vivo* in zebrafish larvae. BPC only minimally stimulated the activity of zER α *in vitro*, and partially activated zER β -mediated transcription to levels ranging from 40–60% of the activity of E2. In accordance with its agonist profile *in vitro*, exposure to BPC induced GFP expression in the brain and in the liver of the zebrafish larvae.

Additionally, the stronger affinity of BPAF and BPC for the zebrafish receptors compared to BPA *in vitro* was confirmed *in vivo*, as lower concentrations of these bisphenol analogues were necessary to activate liver and brain GFP expression, as well as promote the induction of vtg1 and cyp19a1b gene expression in zebrafish embryos.

The estrogenic activities of bisphenols have previously been analyzed for human ERs in reporter cell lines (Kitamura *et al.*, 2005; Matsushima *et al.*, 2010; Delfosse *et al.*, 2012; Chen *et al.*, 2016; Grimaldi *et al.*, 2019). The pronounced agonist activity of BPAF has been previously reported for human ERs, with several studies demonstrating that BPAF is more estrogen-active than BPA. In HELN cells expressing the human ERs, both BPAF and BPC are more potent ligands than BPA (Delfosse *et al.*, 2012). Bisphenols show a slightly better affinity for human ER α . This opposite ER selectivity profile between human and zebrafish ERs also observed for several other xenoestrogens, could be due to differences in the constitution and interaction with amino acids in the LBD of human and zebrafish receptors (Pinto *et al.*, 2014).

The *in vivo* estrogenic activity of bisphenols have been studied in fish and rodents. In adult male medaka, BPAF and BPC are more potent than BPA in inducing hepatic upregulation of the expression of vtg and other estrogenic biomarkers (Yamaguchi *et al.*, 2015). Moreover, the bisphenols induce uterine weight gain in ovariectomized rats, and the uterotrophic effect is more pronounced for BPC and BPAF compared to BPA (Conley *et al.*, 2016). These *in vivo* results are in accordance with our results from *in vitro* and transgenic zebrafish models, and provide substantial data to support that BPC and BPAF have a stronger ability than BPA to disrupt estrogen signaling across different vertebrate species.

The present study is, to our knowledge, the first to demonstrate that the chlorinated bisphenol analogue, BPC, has pronounced antagonistic zfER α activity in zebrafish both *in vitro* and *in vivo*. Exposure to BPC resulted in clear inhibition of the EE2- and 16 α -LE2-induced GFP expression in the heart of the zebrafish larvae, in accordance with its zfER α -mediated antagonist activity in the HELN reporter cells. The outcomes of activation or inactivation of estrogen signaling by bisphenols on the development and physiology of the zebrafish heart are unknown. However, it has been reported that estrogenic ligands can alter the basal heart rate in zebrafish embryos. Exposure to estradiol and ICI 182,780 resulted in increased heart rate in zebrafish, an effect that was attributed to modulation of the G protein-coupled estrogen receptor (GPER) pathway (Romano *et al.*, 2016), while treatment with the anti-estrogen tamoxifen caused a decrease in the heart rate of zebrafish embryos (Xia *et al.*, 2016). BPA treatment has been shown to impair the cardiac contractile performance of rodent hearts, an effect associated with altered intracellular calcium handling (Posnack *et al.*, 2015).

Although ER antagonist activity was observed for BPC in the ERE-driven activation of fluorescence in the zebrafish heart, BPC, as well as BPA and BPAF, were unable to inhibit the EE2-induced brain aromatase promoter. zfER β 2 has been reported to be the main ER subtype detected in the brain during zebrafish early developmental stages, while zfER α and zfER β 1 mRNA are in contrast much lower (Mouriec *et al.*, 2009). Studies have also shown that morpholino knockdown of zfER β 2, but not zfER α or zfER β 1, completely blocks

estrogen-mediated upregulation of the aromatase mRNA (Griffin *et al.*, 2013). This data suggests that zER β 2 is the main subtype controlling the activation of brain aromatase during zebrafish early development. The results with BPC support this hypothesis, as BPC would be expected to inhibit the estrogen-mediated activation of aromatase in the cyp19a1b-GFP zebrafish embryos if aromatase expression was indeed controlled by zER α . Inferring from this data, BPC will act as both an agonist and antagonist *in vivo* in a tissue-specific manner.

In addition to perturbing the ER pathway, BPAF and BPC have been reported to interact with other endocrine targets (Delfosse *et al.*, 2014; Grimaldi *et al.*, 2019; Lei *et al.*, 2017; Wang *et al.*, 2014). The increasing and widespread use of these bisphenol analogues and their ability to interact with endocrine molecular targets at lower concentrations than BPA pose a serious concern associated with potential health risks for humans and wildlife. Implementation of green chemistry practices for identification of safer alternatives will improve characterization of chemical hazard and should be useful to guide manufacturers on more sustainable bisphenol alternatives.

In conclusion, we show that a combined strategy using cell-based assays and transgenic zebrafish models can be used to thoroughly assess estrogenic properties of bisphenol derivatives. This approach allows for the quick screening of both estrogenic and anti-estrogenic activities of environmental pollutants, and are sensitive tools to inform dose-response relationships and tissue-specific effects. Characterization of species differences in response to estrogenic compounds can be addressed in the HELN cells expressing human or zebrafish ERs. The transgenic zebrafish models could also be used to address the influence of metabolism and pharmacokinetic parameters on chemical activity. Moreover, these tools are amenable to high-throughput screening, and can be incorporated into screening platforms currently applied to inform perturbations in ER signaling, such as the ones of the U.S. Environmental Protection Agency (EPA)'s ToxCast/Tox21 program.

Acknowledgements

We thank Dr. Daniel A. Gorelick at the Baylor School of Medicine, Houston, TX, for the gift of the *Tg(5 \times ERE:GFP)* fish line.

Funding Information

This work was supported by grants from the U.S. Environmental Protection Agency [Grant # R- 834289], National Institutes of Health [Grant # R21ES020036] (MB/JAG), the French Agency for Food, Environmental and Occupational Health & Safety (ANSES-EVALPE), the French National Research Agency (ANR-NEWPLAST) (PB), the DRC 59 (FB and SA) and the Robert A. Welch Foundation (E-0004) (JAG). The views expressed in the article do not necessarily reflect the views of the funding agencies.

References

- Bardet PL, Horard B, Robinson-Rechavi M, Laudet V, Vanacker JM, 2002 Characterization of oestrogen receptors in zebrafish (*Danio rerio*). *Journal of molecular endocrinology* 28, 153–163. [PubMed: 12063182]
- Brion F, Le Page Y, Piccini B, Cardoso O, Tong SK, Chung BC, Kah O, 2012 Screening estrogenic activities of chemicals or mixtures *in vivo* using transgenic (cyp19a1b-GFP) zebrafish embryos. *Plos One* 7, e36069. [PubMed: 22586461]

- Cano-Nicolau J, Valliant C, Pellegrini E, Charlier TD, Kah O, Coumailleau P, 2016 Estrogenic Effects of Several BPA Analogs in the Developing Zebrafish Brain. *Front Neurosci* 10, 112. [PubMed: 27047331]
- Cesen M, Lenarcic K, Mislej V, Levstek M, Kovacic A, Cimrmancic B, Uranjek N, Kosjek T, Heath D, Dolenc MS, Heath E, 2018 The occurrence and source identification of bisphenol compounds in wastewaters. *Sci Total Environ* 616–617, 744–752.
- Chandrasekar G, Archer A, Gustafsson JA, Andersson Lendahl M, 2010 Levels of 17beta-estradiol receptors expressed in embryonic and adult zebrafish following in vivo treatment of natural or synthetic ligands. *PLoS one* 5, e9678. [PubMed: 20300630]
- Chen D, Kannan K, Tan HL, Zheng ZG, Feng YL, Wu Y, Widelka M, 2016 Bisphenol Analogues Other Than BPA: Environmental Occurrence, Human Exposure, and Toxicity-A Review. *Environ Sci Technol* 50, 5438–5453. [PubMed: 27143250]
- Conley JM, Hannas BR, Furr JR, Wilson VS, Gray LE, 2016 A Demonstration of the Uncertainty in Predicting the Estrogenic Activity of Individual Chemicals and Mixtures From an In Vitro Estrogen Receptor Transcriptional Activation Assay (T47D-KBluc) to the In Vivo Uterotrophic Assay Using Oral Exposure. *Toxicol Sci* 153, 382–395. [PubMed: 27473340]
- Delfosse V, Grimaldi M, Pons JL, Boulahtouf A, le Maire A, Cavailles V, Labesse G, Bourguet W, Balaguer P, 2012 Structural and mechanistic insights into bisphenols action provide guidelines for risk assessment and discovery of bisphenol A substitutes. *Proceedings of the National Academy of Sciences of the United States of America* 109, 14930–14935. [PubMed: 22927406]
- Delfosse V, Grimaldi M, le Maire A, Bourguet W, Balaguer P, 2014 Nuclear Receptor Profiling of Bisphenol-A and Its Halogenated Analogues. *Vitam Horm* 94, 229–251. [PubMed: 24388193]
- Elsworth JD, Jentsch JD, Vandevort CA, Roth RH, Jr DE, Leranath C, 2013 Prenatal exposure to bisphenol A impacts midbrain dopamine neurons and hippocampal spine synapses in non-human primates. *Neurotoxicology* 35, 113–120. [PubMed: 23337607]
- Gorelick DA, Halpern ME, 2011 Visualization of Estrogen Receptor Transcriptional Activation in Zebrafish. *Endocrinology* 152, 2690–2703. [PubMed: 21540282]
- Gorelick DA, Iwanowicz LR, Hung AL, Blazer VS, Halpern ME, 2014 Transgenic Zebrafish Reveal Tissue-Specific Differences in Estrogen Signaling in Response to Environmental Water Samples. *Environ Health Persp* 122, 356–362.
- Gorelick DA, Pinto CL, Hao R, Bondesson M, 2016 Use of Reporter Genes to Analyze Estrogen Response: The Transgenic Zebrafish Model. *Methods in molecular biology* 1366, 315–325. [PubMed: 26585145]
- Griffin LB, January KE, Ho KW, Cotter KA, Callard GV, 2013 Morpholino-mediated knockdown of ERalpha, ERbeta, and ERbetab mRNAs in zebrafish (*Danio rerio*) embryos reveals differential regulation of estrogen-inducible genes. *Endocrinology* 154, 4158–4169. [PubMed: 23928376]
- Grimaldi M, Boulahtouf A, Toporova L, Balaguer P, 2019 Functional profiling of bisphenols for nuclear receptors. *Toxicology*, 420, 39–45. [PubMed: 30951782]
- Hao RX, Bondesson M, Singh AV, Riu A, McCollum CW, Knudsen TB, Gorelick DA, Gustafsson JA, 2013 Identification of Estrogen Target Genes during Zebrafish Embryonic Development through Transcriptomic Analysis. *Plos One* 8, e79020. [PubMed: 24223173]
- Ho SM, Tang WY, Belmonte de Frausto J, Prins GS, 2006 Developmental exposure to estradiol and bisphenol A increases susceptibility to prostate carcinogenesis and epigenetically regulates phosphodiesterase type 4 variant 4. *Cancer research* 66, 5624–5632. [PubMed: 16740699]
- Honma S, Suzuki A, Buchanan DL, Katsu Y, Watanabe H, Iguchi T, 2002 Low dose effect of in utero exposure to bisphenol A and diethylstilbestrol on female mouse reproduction. *Reproductive toxicology* 16, 117–122. [PubMed: 11955942]
- Kitamura S, Suzuki T, Sanoh S, Kohta R, Jinno N, Sugihara K, Yoshihara S, Fujimoto N, Watanabe H, Ohta S, 2005 Comparative study of the endocrine-disrupting activity of bisphenol A and 19 related compounds. *Toxicological sciences : an official journal of the Society of Toxicology* 84, 249–259. [PubMed: 15635150]
- Lang IA, Galloway TS, Scarlett A, Henley WE, Depledge M, Wallace RB, Melzer D, 2008 Association of urinary bisphenol A concentration with medical disorders and laboratory

- abnormalities in adults. *JAMA : the journal of the American Medical Association* 300, 1303–1310. [PubMed: 18799442]
- Le Fol V, Ait-Aissa S, Sonavane M, Porcher JM, Balaguer P, Cravedi JP, Zalko D, Brion F, 2017 *In vitro* and *in vivo* estrogenic activity of BPA, BPF and BPS in zebrafish-specific assays. *Ecotoxicol Environ Saf*,142: 150–156. [PubMed: 28407500]
- Lei B, Xu J, Peng W, Wen Y, Zeng X, Yu Z, Wang Y, Chen T, 2017 In vitro profiling of toxicity and endocrine disrupting effects of bisphenol analogues by employing MCF-7 cells and two-hybrid yeast bioassay. *Environmental toxicology* 32, 278–289. [PubMed: 26916392]
- Leranth C, Hajszan T, Szigeti-Buck K, Bober J, MacLusky NJ, 2008 Bisphenol A prevents the synaptogenic response to estradiol in hippocampus and prefrontal cortex of ovariectomized nonhuman primates. *Proceedings of the National Academy of Sciences of the United States of America* 105, 14187–14191. [PubMed: 18768812]
- Markey CM, Luque EH, Munoz De Toro M, Sonnenschein C, Soto AM, 2001 In utero exposure to bisphenol A alters the development and tissue organization of the mouse mammary gland. *Biology of reproduction* 65, 1215–1223. [PubMed: 11566746]
- Matsushima A, Liu X, Okada H, Shimohigashi M, Shimohigashi Y, 2010 Bisphenol AF is a full agonist for the estrogen receptor ERalpha but a highly specific antagonist for ERbeta. *Environ Health Perspect* 118, 1267–1272. [PubMed: 20427257]
- Menuet A, Pellegrini E, Anglade I, Blaise O, Laudet V, Kah O, Pakdel F, 2002 Molecular characterization of three estrogen receptor forms in zebrafish: binding characteristics, transactivation properties, and tissue distributions. *Biology of reproduction* 66, 1881–1892. [PubMed: 12021076]
- Mouriec K, Lareyre JJ, Tong SK, Le Page Y, Vaillant C, Pellegrini E, Pakdel F, Chung BC, Kah O, Anglade I, 2009 Early regulation of brain aromatase (cyp19a1b) by estrogen receptors during zebrafish development. *Dev Dyn* 238, 2641–2651. [PubMed: 19718764]
- Munoz-de-Toro M, Markey CM, Wadia PR, Luque EH, Rubin BS, Sonnenschein C, Soto AM, 2005 Perinatal exposure to bisphenol-A alters peripubertal mammary gland development in mice. *Endocrinology* 146, 4138–4147. [PubMed: 15919749]
- Nilsson S, Makela S, Treuter E, Tujague M, Thomsen J, Andersson G, Enmark E, Pettersson K, Warner M, Gustafsson JA, 2001 Mechanisms of estrogen action. *Physiological reviews* 81, 1535–1565. [PubMed: 11581496]
- NTP-NIEHS, 2008 Chemical Information Profile for Bisphenol AF.
- Palanza P, Gioiosa L, vom Saal FS, Parmigiani S, 2008 Effects of developmental exposure to bisphenol A on brain and behavior in mice. *Environmental research* 108, 150–157. [PubMed: 18949834]
- Pinto C, Grimaldi M, Boulahtouf A, Pakdel F, Brion F, Ait-Aissa S, Cavailles V, Bourguet W, Gustafsson JA, Bondesson M, Balaguer P, 2014 Selectivity of natural, synthetic and environmental estrogens for zebrafish estrogen receptors. *Toxicol Appl Pharm* 280, 60–69.
- Posnack NG, Brooks D, Chandra A, Jaimes R, Sarvazyan N, Kay M, 2015 Physiological response of cardiac tissue to bisphenol a: alterations in ventricular pressure and contractility. *Am J Physiol-Heart C* 309, H267–H275.
- Romano S, Edwards H, Souder J, Cui X, Gorelick D, 2017 G protein-coupled estrogen receptor regulates heart rate in zebrafish embryos. *PLOS genetics* 13, 11007069.
- Rubin BS, Lenkowski JR, Schaeberle CM, Vandenberg LN, Ronsheim PM, Soto AM, 2006 Evidence of altered brain sexual differentiation in mice exposed perinatally to low, environmentally relevant levels of bisphenol A. *Endocrinology* 147, 3681–3691. [PubMed: 16675520]
- Santos JM, Putt DA, Jurban M, Joiakim A, Friedrich K, Kim H, 2016 Differential BPA levels in sewage wastewater effluents from metro Detroit communities. *Environmental monitoring and assessment* 188, 585. [PubMed: 27663877]
- Sun Q, Wang Y, Li Y, Ashfaq M, Dai L, Xie X, Yu CP, 2017 Fate and mass balance of bisphenol analogues in wastewater treatment plants in Xiamen City, China. *Environ Pollut* 225, 542–549. [PubMed: 28318793]
- Tong SK, Mouriec K, Kuo MW, Pellegrini E, Gueguen MM, Brion F, Kah O, Chung BC, 2009 A cyp19a1b-gfp (aromatase B) transgenic zebrafish line that expresses GFP in radial glial cells. *Genesis* 47, 67–73. [PubMed: 19101983]

- Vandenberg LN, Hauser R, Marcus M, Olea N, Welshons WV, 2007 Human exposure to bisphenol A (BPA). *Reproductive toxicology* 24, 139–177. [PubMed: 17825522]
- Vandenberg LN, Chahoud I, Heindel JJ, Padmanabhan V, Paumgarten FJ, Schoenfelder G, 2012 Urinary, circulating, and tissue biomonitoring studies indicate widespread exposure to bisphenol A. *Ciencia & saude coletiva* 17, 407–434. [PubMed: 22267036]
- Wang S, Rijk JC, Besselink HT, Houtman R, Peijnenburg AA, Brouwer A, Rietjens IM, Bovee TF, 2014 Extending an in vitro panel for estrogenicity testing: the added value of bioassays for measuring antiandrogenic activities and effects on steroidogenesis. *Toxicol Sci* 141, 78–89. [PubMed: 24928889]
- Xia L, Zheng L, Zhou JL, 2016 Transcriptional and morphological effects of tamoxifen on the early development of zebrafish (*Danio rerio*). *J Appl Toxicol* 36, 853–862. [PubMed: 26584595]
- Xu G, Ma S, Tang L, Sun R, Xiang J, Xu B, Bao Y, Wu M, 2016 Occurrence, fate, and risk assessment of selected endocrine disrupting chemicals in wastewater treatment plants and receiving river of Shanghai, China. *Environ Sci Pollut Res Int* 23, 25442–25450. [PubMed: 27699660]
- Yamaguchi A, Ishibashi H, Arizono K, Tominaga N, 2015 In vivo and in silico analyses of estrogenic potential of bisphenol analogs in medaka (*Oryzias latipes*) and common carp (*Cyprinus carpio*). *Ecotox Environ Safe* 120, 198–205.
- Zalko D, Jacques C, Duplan H, Bruel S, Perdu E, 2011 Viable skin efficiently absorbs and metabolizes bisphenol A. *Chemosphere* 82, 424–430. [PubMed: 21030062]

Highlights

- In vitro/in vivo estrogenic activity of BPA, BPAF and BPC in zebrafish
- Selectivity on zfER α of bisphenols
- Antagonism on zfER α of BPC
- Relevance of an integrated strategy to assess EDCs and their substitutes




Fig. 1.

Agonistic and antagonistic activity of bisphenols in HELN zfER reporter cell lines. (A-C) Dose-response curves for BPA, BPAF, and BPC in HELN zfER reporter cell lines. Cells were incubated for 16h with various concentrations of BPA, BPAF, and BPC. (D) Dose responsive curves of BPC in HELN zfER reporter cell lines. Cells were incubated for 16h with various concentrations of BPC in presence of 1 nM E2. Data were expressed as percentage of the maximal activity induced by E2 10 nM. EC50 values were measured using GraphPad Prism (GraphPad Software Inc). Data were analyzed for significant differences using one-way ANOVA followed by Dunnett's post-comparison test (vs. control) (***) $p < 0.001$, ** $p < 0.01$, * $p < 0.05$).




Fig. 2.

In vivo estrogenic activities of bisphenols on ERE reporter fish. *Tg (ERE:GFP)* larvae were treated with 16 α -LE2 or bisphenols at 3 dpf for 24 h. Images were taken at 4 dpf. Lateral view showing GFP expression in the heart valves (red arrows) and liver (white arrow). A. Bright-field image; B. GFP fluorescence image of vehicle-treated zebrafish larvae; C-F. GFP images of zebrafish larvae exposed to increasing concentrations of 16 α -LE2; G-J. GFP images of zebrafish treated with BPA; K-N. Images of zebrafish exposed to BPAF; O-R. GFP images of zebrafish exposed to BPC. *Scale bars*, 200 μ m.




Fig. 3. Anti-estrogenic effects of BPC on ERE reporter fish co-treated with EE2. A. Bright-field image; B. GFP fluorescence image of vehicle-treated zebrafish larvae; C. GFP images of zebrafish larvae exposed to 100 pM EE2; D. Zebrafish exposed to 1 μ M ICI 182,780; E. GFP images of zebrafish co-treated with EE2 and ICI; F. Images of zebrafish exposed to 1 μ M BPC; G. GFP images of zebrafish co-treated with EE2 and BPC. *Scale bars*, 100 μ m.





Fig. 4. Anti-estrogenic effects of BPC on ERE reporter fish co-treated with 16α-LE2. A. Bright-field image; B. GFP fluorescence image of vehicle-treated zebrafish larvae; C. GFP images of zebrafish larvae exposed to 10 nM 16α-LE2; D. GFP images of zebrafish exposed to 3 μM BPC; E. GFP images of zebrafish larvae co-treated with 16α-LE2 and BPC. *Scale bars*, 200 μm.

**Fig. 5.**

In vivo estrogenic activity of bisphenols in the *cyp19a1b-GFP* zebrafish. A. Concentration-dependent induction of GFP by BPA, BPAF and BPC in *cyp19a1b-GFP* transgenic larvae after exposure to the bisphenols from 0 to 4 dpf. Values are expressed as mean \pm SEM (n = 4–13). B. Effect of BPA, BPAF and BPC on EE2-mediated GFP induction in *cyp19a1b-GFP* transgenic larva after exposure from 0 to 4 dpf. Values are expressed as mean \pm SEM (n = 8–14). C. Fluorescence images of *cyp19a1b-GFP* transgenic larvae treated with 0.05 nM EE2, 10 μ M BPA, 1 μ M BPAF and 1 μ M BPC alone and co-treatments of EE2 with the bisphenols. Images were taken in the dorsal view, anterior side to the top.




Fig. 6. Endogenous estrogen target gene activation by bisphenols *in vivo*. Concentration-dependent induction of *vtg1* (A) and *cyp19a1b* (B) mRNA levels after treatment of zebrafish embryos with BPA, BPAF and BPC from 4 to 6 dpf. Transcript levels were normalized to zebrafish β -actin using the Ct method.

Table 1.

Effective concentration (EC₅₀ in nM), IC₅₀ in nM), relative estrogenic potency (REP) and maximal activity (%) of 17β-E₂, 16α-LE₂, BPA and the bisphenol analogues on the zfERs.

	zfERα		zfERβ1		zfERβ2	
Basal activity	18.8 ± 0.3		16.5 ± 0.7		20.2 ± 0.6	
Agonism						
Ligand	EC ₅₀ (REP)	Maximal activity	EC ₅₀	Maximal activity	EC ₅₀	Maximal activity
E ₂	0.08 ± 0.03 (1)	100	0.04 ± 0.01 (1)	100	0.12 ± 0.03 (1)	100
16α-LE ₂	0.65 ± 0.1 (1.2 10 ⁻¹)	100	271 ± 45 (1.5 10 ⁻⁴)	100	69.5 ± 17.5 (1.7 10 ⁻³)	100
BPA	469 ± 42 (1.7 10 ⁻⁴)	79	3326 ± 381 (1.2 10 ⁻⁵)	53.1	2800 ± 562 (4.3 10 ⁻⁵)	56.1
BPAF	84 ± 27 (9.5 10 ⁻⁴)	89.3	758 ± 131 (5.3 10 ⁻⁵)	89.6	897 ± 268 (1.3 10 ⁻⁴)	79.6
BPC	25 ± 7.5 (3.2 10 ⁻³)	31	316 ± 110 (1.3 10 ⁻⁴)	36.1	167 ± 72 (7.2 10 ⁻⁴)	58.5
Antagonism						
	IC ₅₀		IC ₅₀		IC ₅₀	
BPC	46 ± 6.4		160 ± 42		136 ± 75	

Values are the mean ± standard deviations from three separate experiments. Maximal activity is expressed as percentage of luciferase activity induced by 10 nM E₂. REPs were determined as the ratio of E₂ EC₅₀ to that of bisphenol.

Table 2.

Dissociation constant (Kd) values for E2, 16 α -LE2, BPA, BPAF and BPC on the zfERs. Values are the mean \pm standard deviations from three separate experiments.

	zfER α	zfER β 1	zfER β 2
Ligands	Kd (nM)	Kd (nM)	Kd (nM)
E2	2.6 \pm 0.6	1.1 \pm 0.2	2.6 \pm 0.7
16 α -LE2	9.4 \pm 0.9	1109 \pm 126	351 \pm 44
BPA	1245 \pm 265	10040 \pm 2919	6884 \pm 1270
BPAF	428 \pm 85	1023 \pm 184	1993 \pm 628
BPC	78 \pm 16	678 \pm 124	402 \pm 89