



Published in final edited form as:

*Dev Biol.* 2019 November 01; 455(1): 10–18. doi:10.1016/j.ydbio.2019.07.005.

## Meg3-DMR, not the *Meg3* gene, regulates imprinting of the *Dlk1-Dio3* locus

Wende Zhu<sup>1</sup>, Erin M. Botticelli<sup>1</sup>, Rachel E. Kery<sup>1,3</sup>, Yanfei Mao<sup>1,3</sup>, Xin Wang<sup>1,3</sup>, Anli Yang<sup>1,3</sup>, Xianling Wang<sup>1,3</sup>, Jie Zhou<sup>1,3</sup>, Xun Zhang<sup>1</sup>, Roy J. Soberman<sup>2</sup>, Anne Klibanski<sup>1</sup>, Yunli Zhou<sup>1,4</sup>

<sup>1</sup>Neuroendocrine Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114

<sup>2</sup>Nephrology Division, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114

### Abstract

The imprinted *delta like 1 homolog (DLK1) - thyroxine deiodinase type III (DIO3)* locus regulates development and growth. Its imprinting regulation involves two differentially methylated regions (DMRs), intergenic-DMR (IG-DMR) and maternally expressed gene 3-DMR (Meg3-DMR). In mice, a maternal deletion of the IG-DMR leads to LOI in the locus, proving that the IG-DMR is a cis-acting imprinting control region of the locus. However, the Meg3-DMR overlaps with the promoter, exon 1 and intron 1 of the Meg3 gene. Because deletion of the Meg3-DMR inactivates the *Meg3* gene, their roles in imprinting regulation of Meg3-DMR mice is unknown. Therefore, we generated two mouse models: Meg3<sup>(1-4)</sup> and Meg3<sup>(2-4)</sup>, respectively targeting exons 1-4 and exons 2-4 of the *Meg3* gene. A maternal deletion of Meg3<sup>(1-4)</sup> caused embryonic death and LOI in both embryos and placentas, but did not affect methylation status of the IG-DMR. In contrast, mice carrying a maternal deletion of Meg3<sup>(2-4)</sup> were born normally and did not have LOI. These data indicate that it is the Meg3-DMR, not the *Meg3* gene, which regulates imprinting of the *Dlk1-Dio3* locus.

<sup>4</sup>To whom correspondence should be addressed: Yunli Zhou, Neuroendocrine Unit, Massachusetts General Hospital, 55 Fruit Street, Bulfinch 457, Boston, MA 02114, Tel. 617 726-3870; Fax. 617 726-5072; yunli.zhou@mgh.harvard.edu.

<sup>3</sup>Present Addresses:

RE. Kery: Department of Neurobiology and Behavior, The State University of New York at Stony Brook, New York, U.S.A.

Y Mao: Department of Anesthesia and Intensive Care Unit, Xinhua Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

Xin Wang: Department of Histology and Embryology, Zhangshan School of Medicine, Sun Yat-sen University, Guangzhou, China

A Yang: Department of Breast Oncology, Sun Yat-sen University Cancer Center, Guangzhou, Guangdong, China

Xianling Wang: Department of Endocrinology, Chinese PLA General Hospital, Beijing, China J Zhou: Neurosurgery Department, the Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan, China

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declarations of interest: none

## Keywords

DLK1-DIO3 locus; genomic imprinting; IG-DMR; imprinting control region; MEG3; MEG3-DMR

---

## Introduction

Genomic imprinting is an epigenetic phenomenon in which expression of imprinted genes is parental-origin-dependent (1). An imprinted locus is formed by a cluster of imprinted genes in which their expression is tightly regulated by imprinting mechanisms. The *delta like 1 homolog (DLK1) - thyroxine deiodinase type III (DIO3)* locus is located on chromosome 14 in humans and on distal chromosome 12 in mice (2–4). Balanced gene expression in the *DLK1-DIO3* locus plays a pivotal role in development and growth. Loss of imprinting (LOI) in this locus in humans leads to Kagami-Ogata syndrome (5, 6) or Temple syndrome (7). In mice, LOI of the locus causes developmental defects and premature death (8, 9). The locus is also essential in maintaining pluripotentiality of induced pluripotent stem cells (10).

The cis-elements controlling genomic imprinting are differentially methylated regions (DMRs) (11). Two DMRs, intergenic DMR (IG-DMR) and maternally expressed gene 3-DMR (Meg3-DMR), are thought to play important roles regulating imprinting of the *DLK1-DIO3* locus (2, 6). Kagami et al reported that a patient carrying a microdeletion overlapping with the IG-DMR showed LOI in the locus (6). Similarly, Lin et al demonstrated LOI in mice carrying a maternal deletion of the IG-DMR (IG-DMR<sup>-/-</sup>) (Fig. S3) (12, 13). These data proved that the IG-DMR is an imprinting control region (ICR). The MEG3-DMR starts approximately 1.5 kb upstream of the MEG3 gene and extends into the MEG3 intron 1. We previously generated a Meg3-5kb mouse model, in which a 5-kb genomic DNA was targeted including a small portion of the *Meg3* promoter and exons 1 through 5 of the *Meg3* gene (Fig. S3) (14). The maternal deletion of the targeted region resulted in LOI in the locus (14), indicating that the targeted region is necessary for the imprinting regulation of the *Dlk1-Dio3* locus. In agreement, Kagami et al reported that a patient with a microdeletion containing the MEG3-DMR had LOI in *DLK1-DIO3* locus (6). Recently, Sanli et al made deletions in the *Meg3* promoter or intron 1 in embryonic stem cells using CRISPR-Cas9 technique and found that these deletions caused LOI (15). These data support the concept that the MEG3-DMR plays a role in imprinting regulation. However, a contradictory finding was reported by Takahashi et al, who generated a mouse model carrying an approximately 10-kb deletion including *Meg3* exons 1 through 5 (designated as Meg3-10kb) (Fig. S3) (16). Mice carrying a maternal deletion of Meg3-10kb did not demonstrate LOI in the *Dlk1-Dio3* locus, although *Meg3* expression was completely abolished. IG-DMR methylation was not affected by the Meg3-10kb deletion (16). These data imply that the Meg3-DMR does not play a significant role in imprinting regulation. Furthermore, because the Meg3-DMR overlaps with the *Meg3* gene promoter, any deletion in this region inactivates both Meg3-DMR and the *Meg3* gene. Therefore, it is still unknown as to whether the effects on imprinting in the *Dlk1-Dio3* locus were caused by lack of maternal Meg3-DMR, or Meg3 expression.

To address this key unanswered question, we created two mouse models, one with a deletion of exons 1 through 4 of the *Meg3* gene ( $Meg3^{(1-4)}$ ) and the other with a deletion of exons 2 through 4 of the gene ( $Meg3^{(2-4)}$ ). We show that LOI was only observed in embryos with a maternal  $Meg3^{(1-4)}$  deletion ( $Meg3^{(1-4)/+}$ ), not in those with a maternal deletion of  $Meg3^{(2-4)}$ . In addition, the maternal allele of the IG-DMR remains unmethylated in  $Meg3^{(1-4)/+}$  embryos. Our data demonstrate that it is the *Meg3*-DMR, not the *Meg3* gene, that regulates imprinting, likely by regulating expression of maternally expressed genes.

## Materials and Methods

### Mice:

All animals used in this study were approved by the Institutional Animal Care and Use Committee (IACUC) at Massachusetts General Hospital. They were housed at the animal facility of MGH Center for Comparative Medicine. Two strains of mice carrying deletions of the *Meg3* gene were created.  $Meg3^{(1-4)}$  mice contain a deletion of exons 1 to 4 of the *Meg3* gene (Fig. S1)(Supplementary Data). To obtain this strain, a mouse strain carrying floxed exons 1-4 of the *Meg3* gene ( $Meg3^{+/fl(1-4)}$ ) was created using services from inGenious Targeting Laboratories (iTTL, Stony Brook, NY). To generate heterozygous mice carrying a deletion on the paternal allele ( $Meg3^{+/- (1-4)}$ ), male  $Meg3^{+/fl(1-4)}$  mice were mated with female CRE transgenic mice (BALB/c-Tg(CMV-cre)1Cgn/J, Jackson Lab, Bar Harbor, ME).  $Meg3^{(2-4)}$  mice carry a deletion of exons 2 through 4 of the *Meg3* gene (Fig. S2). To obtain this strain, a mouse strain carrying floxed exons 2-4 of the *Meg3* gene ( $Meg3^{+/fl(2-4)}$ ) was created using services from Taconic (TaconicArtemis, Cologne, Germany). To generate heterozygous mice carrying a deletion on the paternal allele ( $Meg3^{+/- (2-4)}$ ), male  $Meg3^{+/fl(2-4)}$  mice were mated with the female aforementioned CRE transgenic mice.

### Gene expression:

Expression of maternally and paternally expressed genes were detected by SYBR quantitative RT-PCR. Total RNAs were isolated from 11.5 dpc embryos and placentas using Qiagen RNeasy Mini Kit with on column DNase treatment (Qiagen, Germantown, MD). For lncRNAs and mRNAs, reverse transcriptions were performed using the ProtoScript® First Strand cDNA Synthesis Kit from New England Biolabs (Ipswich, MA). To detect Rtl1 RNA, the primer used in reverse transcription was 5'GATACTCAAACCTTTTCTGA3'. The primer for Gapdh, 5'TTGAGTGAGTTGTCATATT3', was included as the internal control. The primers used in qPCR are listed in Table S1 and S2. To detect snoRNAs and miRNAs, reverse transcriptions were performed with the qScript mircoRNA DNA synthesis kit from Quantabio (Beverly, MA). The forward primers for small RNAs are listed in Table S2. The reverse primer was the PerfeCTa® Universal PCR Primer obtained from Quantabio. Quantitative PCRs were performed using PowerUp SYBR master mix from ThermoFisher Scientific (Grand Island, NY) per the manufacturer's instruction. Fold changes of gene expression in knockout mice over the WT mice were calculated by comparative Ct method described by Schmittgen et al (17)(33) and detailed in the technical note by Haimes et al at Dharmacon of GE Healthcare (<http://dharmacon.horizondiscovery.com/uploadedFiles/Resources/delta-cq-solaris-technote.pdf>). Gapdh was used as the internal reference gene for lncRNAs and mRNAs. Sno202 was used as the reference for snoRNAs and miRNAs.

### Biallelic expression:

Chromosome-specific gene expression was determined by SNP analysis. In *Dlk1* cDNA (NM\_010052), nucleotide 1207 is a G in C57BL/6 and an A in DBA/2, respectively. In *Rtl1* cDNA (NM\_184109), nt2061 is a T in C57BL/6 and a C in NOD/ShLit. DBA/2J and NOD/ShLitJ mice were obtained from the Jackson Laboratory. Biallelic expression of *Dlk1* in embryos was done as previously described (14). Total RNA was isolated from 11.5 dpc embryos and placentas. Reverse transcriptions were done with the first strand cDNA synthesis kit from New England Biolabs using oligo dT as the RT primer for *Dlk1* and *Rtl1*-RT (5'TCATCTCACTTTCCCTTAAT3') for *Rtl1*. The cDNA fragments containing SNPs for *Dlk1* and *Rtl1* were amplified by PCR. After gel purification, the fragments were sequenced to identify the SNPs. PCR primers for *Dlk1* are 5'TCCTGAAGGTGTCCATGAAAGAGC3' (forward) and 5'AAGCATAGCGTTCCTCGATTCCAC3' (reverse); for *Rtl1* 5'TCTGATACCCACCTTAGGCTGGC3' (forward) and 5'GACGCCGCTTTGATCACTGTCTC3' (reverse).

### DNA methylation analysis:

Methylation status of the IG-DMR was determined by bisulfite sequencing (Bisulfite Sequencing Services, Active Motif, Carlsbad, CA). A region of 500 nt long containing 35 CpG sites, located approximately 12.4 Kb upstream of the *Meg3* gene, was selected for the analysis. Genomic DNAs were isolated from 11.5 dpc embryos, including 3 *Meg3*<sup>(1-4)/+</sup>, 3 *Meg3*<sup>+/ (1-4)</sup> and 3 *Meg3*<sup>+/+</sup>, and their corresponding placentas. After bisulfite treatment, the converted genomic DNAs were amplified by PCR, processed into standard, barcoded Illumina sequencing libraries and sequenced in NextSeq 500 system (Illumina, San Diego, CA). Reads were analyzed using the bismark alignment program (v 0.7.7) (<http://www.bioinformatics.babraham.ac.uk/projects/bismark/>). The mouse chr12 (mm10 assembly) was used as the reference sequence. Between 4.7 and 7.05 million reads were analyzed per sample.

## Results and Discussion

The mouse *Meg3*-DMR is approximately 3.5 kb long overlapping the promoter, exon 1 and intron 1 of the *Meg3* gene. To clarify the role of the *Meg3*-DMR and *Meg3* gene in imprinting regulation of the *Dlk1*-*Dio3* locus, two mouse models were generated. Part of the *Meg3* promoter as well as exons 1 through 4 of the *Meg3* gene were deleted in *Meg3*<sup>(1-4)</sup> mice (Fig. S1). Exons 2 through 4 of the gene were deleted in *Meg3*<sup>(2-4)</sup> mice (Fig. S2). Mice carrying a paternal allele of *Meg3*<sup>(1-4)</sup> (*Meg3*<sup>+/ (1-4)</sup>) or *Meg3*<sup>(2-4)</sup> (*Meg3*<sup>+/ (2-4)</sup>) developed normally. All mice were bred by crossing female *Meg3*<sup>+/ (1-4)</sup> or *Meg3*<sup>+/ (2-4)</sup> with WT male B6 mice. Mice carrying a maternal allele of *Meg3*<sup>(1-4)</sup> (*Meg3*<sup>(1-4)/+</sup>) died embryonically starting at 12.5 day post coitus (dpc). All *Meg3*<sup>(1-4)/+</sup> embryos died by 13.5 dpc (Table S1). In contrast, *Meg3*<sup>(2-4)/+</sup> embryos developed to full term and were born alive. Therefore, we used 11.5 dpc embryos to assess the effects of deletions on imprinting of the *Dlk1*-*Dio3* locus.

There are three *Meg3* transcript variants (Fig. 1A). Variant 3 (NR\_027652.1) was first reported by Schuster-Gossler et al (18). The transcript contains all 10 exons. Variant 1

(NR\_003633.3) and 2 (NR\_027651.2) are derived from multiple sequence entries in NCBI. They contain a long alternative exon 10 (10A). Using quantitative PCR with primers targeting specific variant transcripts, we quantified their expression in WT embryos (Fig. 1B). The Ct values of PCR #1 and 2, detecting all three variants, were similar to those of PCR #2 and 4, detecting variant 3 only, suggesting that the variant 3 is the predominant transcript of all variants. In agreement, the Ct values of PCR #5, 6 and 7, detecting variants 1 and 2, were significantly higher than Ct values from other 4 qPCRs. Their expression levels were less than 6% of total Meg3 expression (Fig. 1C). Compared with the variant 3 levels, the combined expression of variants 1 and 2 were also very low (less than 6%) (Fig. 1C). These data suggest that the variant 3 is the main functional transcript of the Meg3 gene. Therefore, variant 3 was used as the functional product of the Meg3 gene.

To determine Meg3 expression in Meg3 KO embryos, we performed multiple qPCRs to quantify Meg3 transcripts with primers targeting specific exons as indicated in Fig. 1D. In Meg3<sup>(1-4)/+</sup> mice, no Meg3 transcripts were detected with PCR #3 and 18 (Fig. 1E). In Meg3<sup>(2-4)/+</sup> mice, no full-length Meg3 transcript was detected. Instead, multiple short truncated transcripts were detected in embryos, likely due to alternative RNA splicing. Their expression was very low compared to the level at which the full length Meg3 was expressed in the WT littermates (Fig. 1G). The highly possible transcripts were predicted (Fig. 1H). Among them, the most prominent transcript contained exons 1, 8, 9 and 10, which retains approximately 40% of the full-length Meg3 RNA (Fig. 1H). Previously, we have shown that deletion of less than a third of MEG3 RNA in the 5'-, 3'-end or middle of the molecule functionally inactivates MEG3 in p53 activation (19). Therefore, the data suggest that the function of the Meg3 gene in Meg3<sup>(2-4)/+</sup> mice is most likely diminished. Because the function of Meg3 RNA has not been fully understood, however, our data could not rule out the possibility that the low-level expression of truncated Meg3 transcripts is enough to maintain imprinting of the Dlk1-Dio3 locus.

The mouse Dlk1-Dio3 domain contains nearly 60 known maternally expressed genes (MEGs) and 3 known paternally expressed genes (PEGs). We determined expression of 13 MEGs across the maternal locus in embryos and their corresponding placentas (Fig. 2A). In Meg3<sup>(1-4)/+</sup> embryos and placentas, expression of all MEGs was completely abolished compared to their WT littermates (Fig. 2B). Conversely, expression levels of Dlk1, Rtl1 and Dio3 were increased by 1.94, 3.24 and 2.36 fold, respectively, in Meg3<sup>(1-4)/+</sup> embryos (Fig. 2B). Increases in Dlk1 and Rtl1 expression were also observed in Meg3<sup>(1-4)/+</sup> placentas (Fig. 2B). However, the level of Dio3 expression was not changed in Meg3<sup>(1-4)/+</sup> placentas compared to WT placentas (Fig. 2B). Taking advantage of SNPs in the *Dlk1* and *Rtl1* genes among mouse strains, we found that both paternal and maternal alleles of the genes were expressed in Meg3<sup>(1-4)/+</sup> embryos and placentas (Fig. 3A). Therefore, the transcriptional activation of their respective maternal alleles plays a role in increase in expression of Dlk1 and Rtl1. In addition, the maternally expressed miR-127 and miR-136 are anti-sense to Rtl1, which have been suggested to down regulate Rtl1 transcript (20). Therefore, lack of their expression is also likely to play a role in the increase in Rtl1 RNA levels in Meg3<sup>(1-4)/+</sup> embryos and placenta (Fig 2B). These data demonstrated that deletion of the exons 1 through 4 of the Meg3 gene leads to LOI in the *Dlk1-Dio3 locus*.

Contrary to the findings in *Meg3*<sup>(1-4)/+</sup> mice, expression levels of all MEGs downstream of the *Meg3* gene in *Meg3*<sup>(2-4)/+</sup> embryos and placentas were comparable to their respective levels in WT littermates (Fig. 2C). There were no changes in expression of PEGs in *Meg3*<sup>(2-4)/+</sup> embryos and placentas (Fig. 2C). Furthermore, biallelic expression analysis indicated that the maternal alleles of *Dlk1* and *Rtl1* genes were not activated in *Meg3*<sup>(2-4)/+</sup> embryos and placentas (Fig. 3B). These data demonstrate that deletion of exons 2 through 4 does not result in LOI in the *Dlk1-Dio3* locus. Taken together, therefore, the LOI is caused by the deletion of the *Meg3*-DMR not the *Meg3* gene.

We previously generated *Meg3*-5kb mice, in which exons 1-5 of the *Meg3* gene were deleted (14). A maternal deletion of *Meg3*-5kb results in LOI in the *Dlk1-Dio3* locus, which is similar to the finding in *Meg3*<sup>(1-4)/+</sup>. In mat-*Meg3*-5Kb mice, the IG-DMR is completely methylated (14). We examined the methylation status of the IG-DMR core region in *Meg3*<sup>(1-4)</sup> mice (Fig. 4). The methylation status of IG-DMR in *Meg3*<sup>(1-4)/+</sup> embryos was very similar to those found in WT and paternal KO embryos, which were approximately 50% (Fig. 4B). There were also no obvious differences in methylation patterns among their respective placentas (Fig. 4C). These data indicate that deletion of the *Meg3*-DMR does not affect the methylation patterns of the IG-DMR. This raises the possibility that the mechanisms of LOI in these two mouse models are different. Because IG-DMR deletion results in LOI in the *Dlk1-Dio3* locus (12), it is very likely that the reason for LOI in mat-*Meg3*-5kb is methylation of the IG-DMR. This is further supported by the fact that mat-*Meg3*-5kb mice display identical phenotypes to IG-DMR/+ mice, including skeletal muscle defects and death right after birth (12, 14). The *Meg3*-5kb mice were generated using a conventional recombination technology in which the targeted region was replaced by a *neomycin resistant* gene (*Neo<sup>r</sup>*) cassette. The transcription of *Neo<sup>r</sup>* is toward the IG-DMR (14). It has been found that transcription-through is a common mechanism of DNA methylation in imprinted loci, such as the *Gnas* locus (21, 22) and the *Peg3* locus (23). Therefore, it is most likely that IG-DMR methylation in the mat-*Meg3*-5kb mice is caused by transcription from the *Neo<sup>r</sup>* cassette. Taken together, these data indicate that the underlying mechanisms for LOI in the *Dlk1-Dio3* locus between mat-*Meg3*-5kb mice and *Meg3*<sup>(1-4)/+</sup> mice are different: inactivation of IG-DMR in the former and inactivation of *Meg3*-DMR in the latter.

Lin et al reported that methylation of the *Meg3*-DMR was significantly increased in IG-DMR/+ mice (12)(Fig. S3). A similar phenomenon was observed in a Kagami-Ogata patient carrying a microdeletion overlapping the IG-DMR (6). Conversely, the methylation status of the IG-DMR is unaffected by a *Meg3*-DMR deletion as shown in *Meg3*<sup>(1-4)/+</sup> mice (Fig. 4) and in Kagami-Ogata patients with microdeletions containing the 5' part of the MEG3 gene (6, 24). Considering that inactivation of either IG-DMR or *Meg3*-DMR results in LOI of the *Dlk1-Dio3* locus, these data suggest that imprinting regulation by IG-DMR is mediated by the *Meg3*-DMR. Luo et al identified an element 10 kb upstream of the *Meg3* gene involved in regulating expression of MEGs (25). This element, known as the *Meg3*-proximal enhancer, contains binding sites for transcription factor ZFP281 (25, 26). ZFP281 recruits AFF3, a component of RNA polymerase II elongation complex, to the enhancer on the maternal chromosome where the nearby IG-DMR is not methylated (25, 26). This recruitment of AFF3 promotes transcription activation from the *Meg3* promoter (25). On the



paternal chromosome, AFF3 is sequestered to the methylated IG-DMR via ZFP57 (26), which binds specifically to its methylated motifs (27). These data further indicate that the IG-DMR regulates imprinting of the *Dlk1-Dio3* locus through Meg3-DMR.

Data from previous studies suggest that all maternally expressed genes in the *Dlk1-Dio3* domain are transcribed as one large transcript (15, 25, 28). Considering that all MEGs are downstream of the *Meg3* promoter, it is very likely that the *Meg3* promoter controls MEG expression. This is consistent with our finding that all MEG expression is silenced in *Meg3*<sup>(1-4)/+</sup> mice. The Meg3-DMR overlaps with the *Meg3* promoter. Because DNA methylation is a common mechanism in gene silencing, it is logical to hypothesize that the main function of the Meg3-DMR is to modulate *Meg3* promoter activity to control the parental origin-dependent expression of downstream MEGs. It turns on the *Meg3* promoter on the maternal chromosome; while it turns the promoter off on the paternal chromosome. Our results indicate that deletion of the *Meg3-DMR* inactivates the *Meg3* promoter which in turn silences expression of all MEGs (18), implying that the MEG products may play a role in suppression of PEG expression in the *Dlk1-Dio3* locus. Sanli et al observed that Meg3 transcripts colocalized with the *Dlk1* gene on the maternal chromosome and the maternal *Dlk1* promoter was activated in promoter<sup>-/-</sup> and intron1<sup>-/-</sup> cells. They suggested that Meg3 RNAs play a role in regulation of Dlk1 expression (15). However, all ES clones in the study contained deletions in either the *Meg3* promoter or *Meg3* intron 1, which overlap with the *Meg3-DMR*. Therefore, their data did not distinguish whether the effect was caused by lack of Meg3 expression or inactivation of *Meg3-DMR*. In contrast, data from our *Meg3*<sup>(2-4)</sup> mice demonstrated that the reduced Meg3 expression did not affect Dlk1 expression (Fig. 2), which is consistent with data from the Meg3-10kb mice by Takahashi et al (16). The Meg3 expression was completely abolished in mice carrying a maternal Meg3-10kb deletion. However, the Dlk1 expression did not increase at all in those KO mice compared with their WT littermates (16). Furthermore, no changes in Meg3-DMR methylation patterns were observed in *Meg3*<sup>(2-4)/+</sup> embryos compared with those in *Meg3*<sup>+/(2-4)</sup> and WT embryos (Fig. S4). Taken together, these data strongly indicate that Meg3 RNAs are not essential in imprinting regulation.

Anti-Rtl1 encodes seven miRNAs, such as miR-127 and miR-136, which are antisense to Rtl1 (20, 29). These miRNAs were shown to reduce Rtl1 RNA transcripts *in vivo* by RISC-mediated cleavage of RNAs (29). However, all Rtl1 RNAs were transcribed from the paternal allele in mice with the miR-127 deletion (30). These data suggest that these miRNAs are dispensable in Rtl1 imprinting. Approximately 80% of miRNAs in the *Dlk1-Dio3* locus were clustered between miR-379 and miR-410, which including miRNAs encoded by *Mirg* (31). Labialle et al reported that a maternal deletion of the miR-379/miR-410 cluster did not affect expression levels of Dlk1 and Dio3 (32). In agreement, Gao et al found that a smaller deletion between miR-379 and miR-544 did not affect the RNA levels of Dlk1, although the Dlk1 protein levels were increased by 1.6-fold in mice with a maternal deletion (33). These findings indicate that the vast majority of miRNAs in the *Dlk1-Dio3* locus are not involved in inhibition of PEG transcription.

Gene silencing by transcription has been demonstrated in several imprinted loci (34). For example, In the *Igf2r* locus, the paternal allele of *Igf2r* is silenced by the continuous

transcription of long non-coding RNA *Airn* (35, 36). The *Airn* promoter is inside a DMR located in the intron 2 of the *Igf2r* gene. The transcription of *Airn* over the *Igf2r* promoter blocks the transcription initiation from the promoter (35). Similarly, the *Nespa* gene is transcriptionally silenced in the *Gnas* locus by the transcription of its anti-sense *Nespa*s gene (37). In the *Dlk1-Dio3* locus, the paternally expressed genes *Rtl1* and *Dio3* are downstream of the *Meg3*. Therefore, a likewise mechanism may be used by the *Meg3*-DMR to regulate the imprinting, which is to maintain an active transcription from the *Meg3* promoter, whereby to silence *Rtl1* and *Dio3*. This could be tested by inserting a polyadenylation (polyA) cassette before or after the *Rtl1* gene, which stops transcriptions from the *Meg3*-DMR before or after the *Rtl1* gene. This technique has been used to determine transcriptions of *Airn* or *Nespa*s in imprinting regulation (36, 38, 39). Although the resultant RNA transcripts will contain full-length *Meg3* RNAs, we expect that the maternal allele of the *Rtl1* gene will be silenced in models with polyA inserted after the gene, not in models with poly A inserted before the gene. A caveat of this hypothesis does not explain how the transcription from the *Meg3*-DMR silences the *Dlk1* expression. Because not all MEGs within the locus have been thoroughly investigated for their roles in regulating PEG expression, we speculate that *Dlk1* may be silenced by one or more of the MEG products, similar to the inhibition of *Slc22a3* by *Airn* lncRNA in the *Igf2r* locus (40).

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

Funding:

This work was supported in part by the National Institutes of Health (R01 CA193520 to A.K.) and the Jarislowsky Foundation (A.K.). None of the funding sources was involved in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication.

## References

1. Barlow DP, Bartolomei MS. Genomic imprinting in mammals. *Cold Spring Harb Perspect Biol*. 2014 2 1;6(2):a018382. [PubMed: 24492710]
2. da Rocha ST, Edwards CA, Ito M, Ogata T, Ferguson-Smith AC. Genomic imprinting at the mammalian *Dlk1-Dio3* domain. *Trends Genet*. 2008 6;24(6):306–16. [PubMed: 18471925]
3. Takada S, Tevendale M, Baker J, Georgiades P, Campbell E, Freeman T, et al. Delta-like and *gtl2* are reciprocally expressed, differentially methylated linked imprinted genes on mouse chromosome 12. *Curr Biol*. 2000 9 21;10(18):1135–8. [PubMed: 10996796]
4. Miyoshi N, Wagatsuma H, Wakana S, Shiroishi T, Nomura M, Aisaka K, et al. Identification of an imprinted gene, *Meg3/Gtl2* and its human homologue *MEG3*, first mapped on mouse distal chromosome 12 and human chromosome 14q. *Genes Cells*. 2000 3;5(3):211–20. [PubMed: 10759892]
5. Ogata T, Kagami M. Kagami-Ogata syndrome: a clinically recognizable *upd(14)pat* and related disorder affecting the chromosome 14q32.2 imprinted region. *Journal of human genetics*. 2016 2;61(2):87–94. [PubMed: 26377239]
6. Kagami M, O'Sullivan MJ, Green AJ, Watabe Y, Arisaka O, Masawa N, et al. The IG-DMR and the *MEG3*-DMR at human chromosome 14q32.2: hierarchical interaction and distinct functional properties as imprinting control centers. *PLoS Genet*. 2010 6 17;6(6):e1000992. [PubMed: 20585555]




7. Ioannides Y, Lokulo-Sodipe K, Mackay DJ, Davies JH, Temple IK. Temple syndrome: improving the recognition of an underdiagnosed chromosome 14 imprinting disorder: an analysis of 51 published cases. *J Med Genet.* 2014 8;51(8):495–501. [PubMed: 24891339]
8. Georgiades P, Watkins M, Surani MA, Ferguson-Smith AC. Parental origin-specific developmental defects in mice with uniparental disomy for chromosome 12. *Development.* 2000 11;127(21):4719–28. [PubMed: 11023874]
9. Kumamoto S, Takahashi N, Nomura K, Fujiwara M, Kijioka M, Uno Y, et al. Overexpression of microRNAs from the Gtl2-Rian locus contributes to postnatal death in mice. *Hum Mol Genet.* 2017 10 1;26(19):3653–62. [PubMed: 28934383]
10. Stadtfeld M, Apostolou E, Akutsu H, Fukuda A, Follett P, Natesan S, et al. Aberrant silencing of imprinted genes on chromosome 12qF1 in mouse induced pluripotent stem cells. *Nature.* 2010 5 13;465(7295):175–81. Epub 2010/04/27. eng. [PubMed: 20418860]
11. Bartolomei MS, Ferguson-Smith AC. Mammalian genomic imprinting. *Cold Spring Harb Perspect Biol.* 2011 7 1;3(7).
12. Lin SP, Youngson N, Takada S, Seitz H, Reik W, Paulsen M, et al. Asymmetric regulation of imprinting on the maternal and paternal chromosomes at the Dlk1-Gtl2 imprinted cluster on mouse chromosome 12. *Nat Genet.* 2003 9;35(1):97–102. [PubMed: 12937418]
13. Lin SP, Coan P, da Rocha ST, Seitz H, Cavaillie J, Teng PW, et al. Differential regulation of imprinting in the murine embryo and placenta by the Dlk1-Dio3 imprinting control region. *Development.* 2007 1;134(2):417–26. Epub 2006/12/15. eng. [PubMed: 17166925]
14. Zhou Y, Cheunsuchon P, Nakayama Y, Lawlor MW, Zhong Y, Rice KA, et al. Activation of paternally expressed genes and perinatal death caused by deletion of the Gtl2 gene. *Development.* 2010 8;137(16):2643–52. [PubMed: 20610486]
15. Sanli I, Lalevee S, Cammisa M, Perrin A, Rage F, Lleres D, et al. Meg3 Non-coding RNA Expression Controls Imprinting by Preventing Transcriptional Upregulation in cis. *Cell reports.* 2018 4 10;23(2):337–48. [PubMed: 29641995]
16. Takahashi N, Okamoto A, Kobayashi R, Shirai M, Obata Y, Ogawa H, et al. Deletion of Gtl2, imprinted non-coding RNA, with its differentially methylated region induces lethal parent-origin-dependent defects in mice. *Hum Mol Genet.* 2009 5 15;18(10):1879–88. Epub 2009/03/07. eng. [PubMed: 19264764]
17. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc.* 2008;3(6):1101–8. [PubMed: 18546601]
18. Schuster-Gossler K, Bilinski P, Sado T, Ferguson-Smith A, Gossler A. The mouse Gtl2 gene is differentially expressed during embryonic development, encodes multiple alternatively spliced transcripts, and may act as an RNA. *Dev Dyn.* 1998 6;212(2):214–28. [PubMed: 9626496]
19. Zhou Y, Zhong Y, Wang Y, Zhang X, Batista DL, Gejman R, et al. Activation of p53 by MEG3 noncoding RNA. *J Biol Chem.* 2007 8 24;282(34):24731–42. [PubMed: 17569660]
20. Seitz H, Youngson N, Lin SP, Dalbert S, Paulsen M, Bachellerie JP, et al. Imprinted microRNA genes transcribed antisense to a reciprocally imprinted retrotransposon-like gene. *Nat Genet.* 2003 7;34(3):261–2. [PubMed: 12796779]
21. Chotalia M, Smallwood SA, Ruf N, Dawson C, Lucifero D, Frontera M, et al. Transcription is required for establishment of germline methylation marks at imprinted genes. *Genes Dev.* 2009 1 1;23(1):105–17. Epub 2009/01/13. eng. [PubMed: 19136628]
22. Mehta S, Williamson CM, Ball S, Tibbit C, Beechey C, Fray M, et al. Transcription driven somatic DNA methylation within the imprinted Gnas cluster. *PLoS One.* 2015;10(2):e0117378. [PubMed: 25659103]
23. Bretz CL, Kim J. Transcription-driven DNA methylation setting on the mouse Peg3 locus. *Epigenetics.* 2017;12(11):945–52. [PubMed: 28925797]
24. Beygo J, Elbracht M, de Groot K, Begemann M, Kanber D, Platzer K, et al. Novel deletions affecting the MEG3-DMR provide further evidence for a hierarchical regulation of imprinting in 14q32. *Eur J Hum Genet.* 2015 2;23(2):180–8. [PubMed: 24801763]
25. Luo Z, Lin C, Woodfin AR, Bartom ET, Gao X, Smith ER, et al. Regulation of the imprinted Dlk1-Dio3 locus by allele-specific enhancer activity. *Genes Dev.* 2016 1 1;30(1):92–101. [PubMed: 26728555]

26. Wang Y, Shen Y, Dai Q, Yang Q, Zhang Y, Wang X, et al. A permissive chromatin state regulated by ZFP281-AFF3 in controlling the imprinted Meg3 polycistron. *Nucleic Acids Res.* 2017 2 17;45(3):1177–85. [PubMed: 28180295]
27. Quenneville S, Verde G, Corsinotti A, Kapopoulou A, Jakobsson J, Offner S, et al. In embryonic stem cells, ZFP57/KAP1 recognize a methylated hexanucleotide to affect chromatin and DNA methylation of imprinting control regions. *Mol Cell.* 2011 11 4;44(3):361–72. [PubMed: 22055183]
28. Tierling S, Dalbert S, Schoppenhorst S, Tsai CE, Oligier S, Ferguson-Smith AC, et al. High-resolution map and imprinting analysis of the Gtl2-Dnchc1 domain on mouse chromosome 12. *Genomics.* 2006 2;87(2):225–35. [PubMed: 16309881]
29. Davis E, Caiment F, Tordoix X, Cavaille J, Ferguson-Smith A, Cockett N, et al. RNAi-mediated allelic trans-interaction at the imprinted Rtl1/Peg11 locus. *Curr Biol.* 2005 4 26;15(8):743–9. [PubMed: 15854907]
30. Ito M, Sferruzzi-Perri AN, Edwards CA, Adalsteinsson BT, Allen SE, Loo TH, et al. A trans-homologue interaction between reciprocally imprinted miR-127 and Rtl 1 regulates placenta development. *Development.* 2015 7 15;142(14):2425–30. [PubMed: 26138477]
31. Seitz H, Royo H, Bortolin ML, Lin SP, Ferguson-Smith AC, Cavaille J. A large imprinted microRNA gene cluster at the mouse Dlk1-Gtl2 domain. *Genome Res.* 2004 9;14(9):1741–8. [PubMed: 15310658]
32. Labialle S, Marty V, Bortolin-Cavaille ML, Hoareau-Osman M, Pradere JP, Valet P, et al. The miR-379/miR-410 cluster at the imprinted Dlk1-Dio3 domain controls neonatal metabolic adaptation. *EMBO J.* 2014 10 1;33(19):2216–30. [PubMed: 25124681]
33. Gao YQ, Chen X, Wang P, Lu L, Zhao W, Chen C, et al. Regulation of DLK1 by the maternally expressed miR-379/miR-544 cluster may underlie callipyge polar overdominance inheritance. *Proc Natl Acad Sci U S A.* 2015 11 3;112(44):13627–32. [PubMed: 26487685]
34. Kanduri C Long noncoding RNAs: Lessons from genomic imprinting. *Biochim Biophys Acta.* 2016 1;1859(1):102–11. [PubMed: 26004516]
35. Santoro F, Mayer D, Klement RM, Warczok KE, Stukalov A, Barlow DP, et al. Imprinted Igf2r silencing depends on continuous Airn lncRNA expression and is not restricted to a developmental window. *Development.* 2013 3;140(6):1184–95. [PubMed: 23444351]
36. Latos PA, Pauler FM, Koerner MV, Senergin HB, Hudson QJ, Stocsits RR, et al. Airn transcriptional overlap, but not its lncRNA products, induces imprinted Igf2r silencing. *Science.* 2012 12 14;338(6113):1469–72. [PubMed: 23239737]
37. Williamson CM, Ball ST, Dawson C, Mehta S, Beechey CV, Fray M, et al. Uncoupling antisense-mediated silencing and DNA methylation in the imprinted Gnas cluster. *PLoS Genet.* 2011 3;7(3):e1001347. [PubMed: 21455290]
38. Tibbit CJ, Williamson CM, Mehta S, Ball ST, Chotalia M, Nottingham WT, et al. Antisense Activity across the Nesp Promoter is Required for Nespas-Mediated Silencing in the Imprinted Gnas Cluster. *Non-coding RNA.* 2015 11 30;1(3):246–65. [PubMed: 29861426]
39. Santoro F, Pauler FM. Silencing by the imprinted Airn macro lncRNA: transcription is the answer. *Cell Cycle.* 2013 3 1;12(5):711–2. [PubMed: 23422859]
40. Nagano T, Mitchell JA, Sanz LA, Pauler FM, Ferguson-Smith AC, Feil R, et al. The Air noncoding RNA epigenetically silences transcription by targeting G9a to chromatin. *Science.* 2008 12 12;322(5908):1717–20. [PubMed: 18988810]


**Highlights**

- Deletion of Meg3 exons 1-4, overlapping the Meg3-DMR, results in loss of imprinting.
- Deletion of Meg3 exons 2-4, downstream of the Meg3-DMR, does not affect imprinting.
- Meg3-DMR, not the Meg3 gene, regulates imprinting of the Dlk1-Dio3 locus.

**Figure 1.**

Meg3 expression in Meg3<sup>(1-4)</sup> and Meg3<sup>(2-4)</sup> mice. (A) There are three Meg3 transcript variants. Variant 3 contains 10 exons. Variants 1 and 2 contain a large alternative exon 10 (10A). Variant 2 has an alternative exon 5 (5A). Quantitative PCRs used to quantify their expression are indicated as numbers above each variant. (B) The SYBR based qRT-PCR was used to determine expression of the Meg3 variants in 11.5 dpc WT embryos (n=7). Gapdh was used as the internal reference gene. Ct values were presented as mean±SD. The Ct values from qPCR #5, 6 and 7 were compared with those from qPCR #1, 2, 3 and 4 using


ANOVA multiple comparison tests. \*  $p < 0.05$ , considered to be statistically significant. (C) Relative expression levels of Meg3 variants 1 and 2 compared to all three variants (vs qPCR #1 and 2) or to variant 3 (vs qPCR #3 or 4). The values were calculated using  $2^{-Ct}$  method as described in the Material and Methods. Student t-test was used to compare values against the reference qPCR. \*  $p < 0.05$ , considered to be statistically significant. (D) Detection of exons in Meg3 transcript was done by qPCR. For example, exon 1 was detected by qPCR #8; exons 1–2 by qPCR #17. Other exons were similarly detected by their respective qPCRs as indicated. (E) Relative expression levels of Meg3 detected with qPCR #3 and 18 in 11.5 dpc Meg3<sup>(1-4)</sup> embryos compared with their WT littermates. (F) Relative expression levels of Meg3 in 11.5 dpc paternal Meg3<sup>(2-4)</sup> embryos compared with their WT littermates. (G) Relative expression levels of Meg3 in 11.5 dpc maternal Meg3<sup>(2-4)</sup> embryos compared with their WT littermates. The levels of Meg3 transcripts detected by qPCR in KO embryos were normalized against the levels in their respective WT littermates, which were designated as 1. A minimum of 6 embryos for each genotype from at least 2 litters were used for gene expression analysis. Student t-test was used to compare values between KO mice and their WT littermates. \*  $p < 0.05$ , considered to be statistically significant. (H) Expression levels of predicted truncated Meg3 transcripts in Meg3<sup>(2-4)/+</sup> embryos. The percentage of each transcript in KO embryos compared to their WT littermates were deduced from data presented in (G).



**Figure 2.** Maternal deletion of Meg3<sup>(1-4)</sup>, not Meg3<sup>(2-4)</sup>, silences MEGs and increases PEG expression. (A) Schematic representation of locations for selected 14 MEGs and 3 PEGs in the Dlk1-Dio3 locus. (B) Relative expression levels of the selected genes in 11.5 dpc Meg3<sup>(1-4)</sup> embryos (upper panel) or placentas (lower panel) compared with their WT littermates. (C) Relative expression levels of the selected genes in 11.5 dpc Meg3<sup>(2-4)</sup> embryos (upper panel) or placentas (lower panel). Gene expression levels were determined by qRT-PCR. Sno202 was used as the internal reference gene for miRNAs and snoRNAs. Gapdh was the reference gene for lncRNAs and mRNAs. Sno234 is also included as an internal control gene. The expression levels for each gene in KO embryos and placentas




were normalized against the levels in their respective WT littermates, which were designated as 1. For each genotype, a minimum of 7 embryos and their corresponding placentas from at least 2 litters were used for gene expression analysis. Student t-test was used to compare values between KO mice and their WT littermates. \*  $p < 0.05$ , considered to be statistically significant.

**Figure 3.**

Biallelic expression of Dlk1 and Rtl1. Biallelic expression of these two genes were observed in Meg3<sup>(1-4)</sup> mice (A), not in Meg3<sup>(2-4)</sup> mice (B). Chromosome-specific gene expression was determined by SNP. To detect SNP in Dlk1, Meg3<sup>(1-4)</sup> or Meg3<sup>(2-4)</sup> mice in C57BL/6 background (B6<sup>(1-4)</sup> or B6<sup>(2-4)</sup>) were crossed with WT DBA/2J (D2<sup>+/+</sup>) mice. To detect SNP in Rtl1, B6<sup>(1-4)</sup> or B6<sup>(2-4)</sup> mice were crossed with WT NOD/ShLit (NOD<sup>+/+</sup>) mice. Total RNAs were extracted from embryos and placentas of 11.5 dpc. After reverse transcription, cDNA fragments containing SNPs for Dlk1 and Rtl1 were amplified by PCR,

purified and sequenced. For each genotype, a minimum of 4 embryos and their corresponding placentas from at least 2 litters were used for SNP analysis. A purple arrow indicates detected biallelic expression.



**Figure 4.** Deletion of Meg3-DMR does not affect methylation status of IG-DMR. (A) Schematic representation of CpG locations (open circle) in the core region of IG-DMR. (B) Methylation status in embryos of WT, paternal KO (+/ (1-4)) and maternal KO ( (1-4)/+). (C) Methylation status in placentas of WT, paternal KO (+/ (1-4)) and maternal KO ( (1-4)/+). The values were derived from 3 WT, 3 Meg3<sup>(1-4)/+</sup> and 3 Meg3<sup>+/(1-4)</sup> embryos and their corresponding placentas.