

Loss of the Major GABA_A Receptor Subtype in the Brain Is Not Lethal in Mice

Cyrille Sur, Keith A. Wafford, David S. Reynolds, Karen L. Hadingham, Frances Bromidge, Alison Macaulay, Neil Collinson, Gillian O'Meara, Owain Howell, Richard Newman, Janice Myers, John R. Atack, Gerard R. Dawson, Ruth M. McKernan, Paul J. Whiting, and Thomas W. Rosahl

Neuroscience Research Center, Merck Sharp and Dohme Research Laboratories, Harlow, Essex, CM20 2QR, United Kingdom

The $\alpha 1\beta 2\gamma 2$ is the most abundant subtype of the GABA_A receptor and is localized in many regions of the brain. To gain more insight into the role of this receptor subtype in the modulation of inhibitory neurotransmission, we generated mice lacking either the $\alpha 1$ or $\beta 2$ subunit. In agreement with the reported abundance of this subtype, >50% of total GABA_A receptors are lost in both $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice. Surprisingly, homozygotes of both mouse lines are viable, fertile, and show no spontaneous seizures. Initially half of the $\alpha 1^{-/-}$ mice died prenatally or perinatally, but they exhibited a lower mortality rate in subsequent generations, suggesting some phenotypic drift and adaptive changes. Both adult $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice demonstrate normal performances on the rotarod, but $\beta 2^{-/-}$ mice displayed increased locomotor activity. Purkinje cells of the cerebellum primarily express $\alpha 1\beta 2\gamma 2$ receptors, and in

electrophysiological recordings from $\alpha 1^{-/-}$ mice GABA currents in these neurons are dramatically reduced, and residual currents have a benzodiazepine pharmacology characteristic of $\alpha 2$ - or $\alpha 3$ -containing receptors. In contrast, the cerebellar Purkinje neurons from $\beta 2^{-/-}$ mice have only a relatively small reduction of GABA currents. In $\beta 2^{-/-}$ mice expression levels of all six α subunits are reduced by ~50%, suggesting that the $\beta 2$ subunit can coassemble with α subunits other than just $\alpha 1$. Our data confirm that $\alpha 1\beta 2\gamma 2$ is the major GABA_A receptor subtype in the murine brain and demonstrate that, surprisingly, the loss of this receptor subtype is not lethal.

Key words: GABA_A receptor; mouse; cerebellum; radioligand; benzodiazepine; inhibitory current; locomotor activity; rotarod

The GABAergic system is the major contributor of the inhibitory tone throughout the CNS. GABA_A receptors are ligand-gated ion channels that exist as a number of different subtypes. The GABA_A receptor is a pentameric structure, which is formed by the coassembly of subunit polypeptides that exist in a large multigene family (McKernan and Whiting, 1996; Barnard et al., 1998). There are at least 16 different members of the GABA_A receptor gene family, including 6α , 3β , 3γ , δ , ϵ , θ , and π subunits (Whiting et al., 1999). The GABA_A receptor genes are differentially expressed both temporally and spatially throughout the mammalian brain. For example the $\alpha 2$, $\alpha 3$, and $\beta 3$ subunits are the major α and β subunits in the fetal brain, respectively, whereas the $\alpha 1$ and $\beta 2$ subunits are mainly expressed after birth (Zhang et al., 1991; Laurie et al., 1992b). Therefore, in the adult brain the $\alpha 1\beta 2\gamma 2$ subtype is the major subtype accounting for ~43% of all GABA_A receptors, whereas the remaining receptors are made up mostly by $\alpha 2$ - and $\alpha 3$ -containing receptors together with other combinations of quantitatively more minor GABA_A receptor subtypes (McKernan and Whiting, 1996).

Several mouse strains lacking individual GABA_A receptor subunits have been generated to study the physiological role of GABAergic system in the living organism. Mice lacking the $\gamma 2$ subunit die shortly after birth (Gunther et al., 1995), whereas

heterozygotes have a normal life expectancy and demonstrate neophobia in a novel environment (Crestani et al., 1999). The lethality of the $\gamma 2^{-/-}$ mice can be rescued by transgenic overexpression of either the $\gamma 2S$ or $\gamma 2L$ subunit isoforms of the GABA_A receptor indicating that both $\gamma 2$ subunit splice variants can substitute for each other (Baer et al., 2000; Wick et al., 2000). Mice lacking the $\beta 3$ subunit of the GABA_A receptor have cleft palate, epilepsy, and many behavioral characteristics of Angelman syndrome (Culiat et al., 1995; Homanics et al., 1997; DeLorey et al., 1998). Most of the $\beta 3^{-/-}$ mice die as neonates, but the survivors, which are runt until weaning, can achieve normal body size by adulthood. In contrast, mice lacking the $\alpha 6$ subunit of the GABA_A receptor, which is expressed exclusively in cerebellar granule cells, have no major phenotypic abnormalities (Jones et al., 1997). Expression of the δ subunit is inhibited in the $\alpha 6^{-/-}$ mice, suggesting that both subunits form functional GABA_A receptor subtypes in the cerebellar granule cells. Finally, mice deficient for the δ subunit are viable but show attenuated sensitivity to neuroactive steroids and epileptic seizures (Mihalek et al., 1999).

GABA_A receptors are the site of action of a number of clinically important drugs, including benzodiazepines, barbiturates, and anesthetics (Sieghart, 1995; Whiting et al., 1995). The $\alpha 1\beta 2\gamma 2$ subtype is of particular interest in this context because it comprises the major benzodiazepine binding site in the brain. We addressed the question about the physiological role of this receptor subtype by generating mice lacking either the $\alpha 1$ or $\beta 2$ subunit of the GABA_A receptor which are thought to primarily coassemble to form the $\alpha 1\beta 2\gamma 2$ subtype.

Received Dec. 19, 2000; revised Feb. 14, 2001; accepted Feb. 23, 2001.

C.S., K.W., and D.R. contributed equally to different aspects of this work.

Correspondence should be addressed to Dr. Thomas W. Rosahl, Merck Sharp and Dohme Neuroscience Research Center, Terlings Park, Eastwick Road, Harlow, Essex, CM20 2QR, UK. E-mail: thomas_rosahl@merck.com.

Copyright © 2001 Society for Neuroscience 0270-6474/01/213409-10\$15.00/0

MATERIALS AND METHODS

Generation of $\alpha 1^{-/-}$ mice. The GABA_A receptor $\alpha 1$ gene-targeting vector was constructed from the same genomic 129/SvEv λ fixII clone, which has been used for the introduction of the $\alpha 1$ H101R mutation (McKernan et al., 2000). However, for the complete gene knock-out, exon 4 was disrupted at the *MscI* restriction site by cloning the 1.2 kb BstBI+MscI and the 7 kb *EcoRV*+*Bam*HI DNA fragment blunt-ended into the targeting vector. A phosphoglycerate kinase I (PGK) neo and a thymidine kinase (TK) cassette were also engineered blunt-ended into the loxP site containing targeting vector. After linearization with *NotI* the targeting vector was introduced into AB2.2 embryonic stem (ES) cells (Lexicon Genetics) as described (Soriano et al., 1991; Rosahl et al., 1993, 1995). Homologous recombinants were identified by PCR using the primers 5'-ATTAATGGAGAGTGTGGTAATCTTT-3' and 5'-GGATGCGGTGGGCTCTATGGCTTCTGA-3' and were further confirmed by genomic Southern blotting. Correctly targeted ES cell clones were injected into C57BL6 blastocysts, and one of three clones gave rise to highly chimeric males, which transmitted the targeted allele into the germ line. A colony of homozygous and wild-type control animals were established and used for the present study. Some $\alpha 1^{-/-}$ mice were crossed with a *cre*-transgenic mouse (Schwenk et al., 1995) and further interbred to establish $\alpha 1$ homozygotes, which no longer contained the neomycin resistance gene marker.

Mice lacking the $\beta 2$ subunit were generated in a similar way. A 17.5 kb genomic λ /FixII clone containing exons 6, 7, and 8 of the $\beta 2$ subunit was subcloned into pBluescript via the *NotI* sites. An 8.5 kb *SalI* DNA fragment as a long arm and a 1.25 kb *HpaI* + *FspI* DNA fragment as a short arm were cloned into the PGK neo and TK containing modified pBS246 plasmid (T. W. Rosahl and K. L. Hadingham, unpublished observations) resulting in the deletion of exons 6 and 7 in the targeting vector. After linearization of the targeting vector with *NotI* and introduction into AB2.2 ES cells, homologous recombinants were identified using the following PCR primers: 5'-ACCAGTCTGGACCATGAGTCCCA-3' and 5'-GGATGCGGTGGGCTCTATGGCTTCTGA-3'. One of three injected ES cell clones gave rise to a chimera transmitting the gene disruption into the germ line. A colony of $\beta 2^{-/-}$ mice containing the neo gene and some -neo gene by crossing with the deleter mice were generated as for the $\alpha 1^{-/-}$ mice.

Radioligand binding and biochemical analysis. Radioligand binding assays with [³H]Ro15-1788 (87 Ci/mmol; NEN Life Sciences), [³H]Ro15-4513 (21.7 Ci/mmol; NEN Life Sciences) in the presence or absence of 10 μ M diazepam and [³H]muscimol (19.1 Ci/mmol; NEN Life Sciences) were performed on membrane preparations as previously described (Quirk et al., 1994; Sur et al., 1998, 1999a).

Autoradiographic studies of the convulsant binding site of GABA_A receptors were performed using 8 nM [³⁵S]t-butylbicyclophosphorothionate (TBPS) on coronal sections of wild-type and knock-out mouse brains cut at a thickness of 12–16 μ m. Sections were washed in 50 mM Tris—citrate and 200 mM NaBr, pH 7.4, buffer for 10 min and then incubated in the same buffer containing [³⁵S]TBPS or [³⁵S]TBPS plus 10 μ M picrotoxin for nonspecific binding for 90 min at room temperature. Slides were washed twice for 5 min in cold buffer, rinsed in distilled water, and exposed to film for 48 hr.

Autoradiographic analyses of the different benzodiazepine-binding sites were done as previously described (Turner et al., 1991; Sur et al., 1999b) with 2 nM [³H]Ro15-1788 (labels $\alpha 1\beta\gamma 2$, $\alpha 2\beta\gamma 2$, and $\alpha 3\beta\gamma 2$ subtypes), 4 nM [³H]L-655,708 plus 10 μ M zolpidem (labels $\alpha 5\beta\gamma 2$ subtype), and 8 nM [³H]Ro15-4513 plus 20 μ M diazepam (labels $\alpha 4\beta\gamma 2$ and $\alpha 6\beta\gamma 2$ subtypes) on coronal sections (12–16 μ m) from two to four mice per genotype. After 3–8 weeks exposure, autoradiograms were analyzed with a Micro Computer Imaging Device M2 imaging system (Imaging Research, St. Catharines, Ontario, Canada).

Immunoprecipitation studies with selective $\alpha 1$, $\alpha 2$, and $\alpha 3$ antibodies were performed on solubilized receptors as previously described (McKernan et al., 1991) using a 10 nM concentration of [³H]Ro15-1788.

For Western blot analyses of 30 μ g of protein were loaded on a 4–12% Bis-Tris gel (Novex, San Diego, CA). Proteins were then transferred to nitrocellulose membrane (Hybond-C; Amersham Pharmacia Biotech, Little Chalfont, UK) by semidry blotting, and the presence of $\alpha 1$ and $\alpha 3$ subunit was detected by incubation with specific rabbit anti- $\alpha 1$ (10 μ g/ml) and anti- $\alpha 3$ (7 μ g/ml) antibodies and the ECL detection system (Amersham Pharmacia Biotech).

Data analyses and statistics were performed with GraphPad Prism (San Diego, CA).

Electrophysiology. Cerebellum was removed from $\alpha 1^{-/-}$, $\beta 2^{-/-}$, and

wild-type mice at postnatal days 11–17, and the vermal layer was isolated and placed into ice-cold oxygenated dissociation media containing (in mM): 82 Na₂SO₄, 30 K₂SO₄, 5 MgCl₂, 10 HEPES buffer, and 10 glucose at pH 7.4. Tissue was then stirred for 7 min in 10 ml of dissociation media containing 3 mg/ml of protease XXIII (Sigma) at 37°C. The tissue was then washed in warmed oxygenated dissociation media containing 1 mg/ml bovine serum albumin and 1 mg/ml trypsin inhibitor and maintained under oxygenating conditions at room temperature in Tyrode's solution (in mM: 150 NaCl, 4 KCl, 2 CaCl₂, 2 MgCl₂, 10 HEPES, and 10 glucose, pH 7.4). Tissue was withdrawn as needed and triturated with a fire-polished Pasteur pipette to liberate individual cells. Cells were plated onto a glass coverslip and left to settle for at least 30 min before use. Purkinje cell bodies were identified by their characteristic size and morphology. Cells could be used for up to 5 hr after preparation.

Glass coverslips containing the dissociated cells were placed in a perspex recording chamber on the stage of a Nikon Diaphot inverted microscope. Cells were perfused continuously with artificial CSF (aCSF) containing (in mM): 149 NaCl, 3.25 KCl, 2 CaCl₂, 2 MgCl₂, 10 HEPES, 11 D-glucose, D(+)-sucrose, pH 7.4, and observed with phase-contrast optics. Fire-polished patch pipettes were pulled on a WZ, DMZ-Universal puller (Zeitz-Instruments, Munich, Germany) using conventional 120TF-10 electrode glass. Pipette tip diameter was ~1.5–2.5 μ m, with resistances ~4 M Ω . The intracellular solution contained (in mM): 130 CsCl, 10 HEPES, 10 BAPTA-Cs, 5 ATP-Mg, 0.1 leupeptin, and 1 MgCl₂, with 100 μ M NaCO₃, pH-adjusted to 7.3 with CsOH and 320–340 mOsm. Cells were voltage-clamped at -60 mV using an Axopatch 200B amplifier (Axon Instruments, Foster City, CA). Drug solutions were applied to the cells via a multibarrel drug delivery system, which could pivot the barrels into place using a stepping motor. This ensured rapid application and washout of the drug. Drugs were applied to the cell for 5 sec with a 30 sec washout period between applications. Allosteric potentiation of GABA_A receptors was measured relative to a GABA EC₂₀ determined for each cell to account for differences in GABA potency. Modulators were pre-equilibrated for 30 sec before coapplication of GABA.

Behavioral analysis. All mice tested were in a 50% C57BL6, 50% 129SvEv background. They were housed in groups under standard 12 hr light/dark cycle and had *ad libitum* access to standard mouse diet and water. Two- to 3-month-old F4 generation mice were tested for motor coordination on the rotarod, and 6-month-old F5 mice were used to determine spontaneous locomotor activity.

Motor coordination was assessed using the rotarod test. Wild-type ($n = 9$), $\alpha 1^{-/-}$ ($n = 7$), and $\beta 2^{-/-}$ ($n = 6$) mice were first trained until they could remain on a rotarod revolving at 16 rpm for three consecutive 120 sec trials. The next day the mice were placed back on the rotarod for a single trial at 18 rpm (maximum duration 120 sec). The duration a mouse could remain on the rotarod was recorded, and the mouse was then returned to its home cage. The speed of the rotarod was then increased to 21 rpm, and all mice were given another test trial. This process was repeated for rotarod speeds of 24, 27, 30, 33, and 36 rpm. Spontaneous locomotor activity was measured using individual perspex activity chambers (215 × 270 × 210 mm) equipped with two parallel infrared beams running across each end of the base of the chamber. Naive $\alpha 1^{-/-}$ and wild-type mice ($n = 10$ per group) were placed in the activity cages, and cage crosses (i.e., consecutive breaks of each beam) were measured for 1 hr. In a separate experiment naive $\beta 2^{-/-}$ and wild-type mice ($n = 12$ per group) were also assessed for spontaneous locomotor activity for 1 hr. Data were analyzed using a repeated measures ANOVA, and then individual time bins were examined with a two-way ANOVA.

RESULTS

Generation of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice

Exon 4 of the $\alpha 1$ subunit gene of the GABA_A receptor was disrupted, and exons 6 and 7 of the $\beta 2$ subunit gene were deleted in mice by the homologous recombination technique (Rosahl et al., 1993, 1995; McKernan et al., 2000) (Fig. 1*a–h*). A colony of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice were bred and analyzed, whereas the neomycin resistance gene flanked by loxP sites (Fig. 1*c*, *g*, *f*loxed) remained in the targeted locus. These $\alpha 1$ or $\beta 2^{-/-}$ [+neo] mice were kept in a ~50% C57BL6–50% 129SvEv genetic background. Some homozygotes of the F3 generation were crossed with the *cre*




Figure 1. Generation and validation of $\alpha 1$ and $\beta 2$ $-/-$ mice. *a, b, e, f*, Schematic representation of the WT $\alpha 1$ (*a, b*) and $\beta 2$ allele (*e, f*) of the GABA_A receptor and the corresponding targeting vectors. 4, 6, 7, Exons 4, 6, and 7, respectively; 4 Δ , partial deletion of exon 4; neo, neomycin resistance gene; TK, thymidine kinase gene; E, EcoRI restriction site; K, KpnI restriction site; P1, P2, P3, P4, PCR primers for detecting the mutant

containing “deleter” mouse (Schwenk et al., 1995) to remove the neo gene in the offspring (Fig. 1*d,h*). This resulted in a change of the genetic background to ~75% C57BL6–25% 129SvEv. The heterozygous offspring was interbred to produce $\alpha 1$ or $\beta 2$ $-/-$ [$-neo$] and wild-type control mice. Northern blot results using a 45 mer antisense oligonucleotide specific for exon 4 upstream of the gene disruption confirmed complete absence of $\alpha 1$ mRNA in both the $\alpha 1$ $-/-$ [$+neo$] and $\alpha 1$ $-/-$ [$-neo$] mice (Fig. 1*i*). Similarly, a probe for the 5' part of the $\beta 2$ subunit gene demonstrated lack of any $\beta 2$ mRNA in the $\beta 2$ $-/-$ [$+neo$] and $\beta 2$ $-/-$ [$-neo$] mice. No major changes in the expression of the $\alpha 3$, $\alpha 6$, $\beta 3$, and $\gamma 2$ genes were found in either knock-out mouse lines regardless of the presence or absence of the neo gene (Fig. 1*i*) (data not shown). Therefore, all experiments presented in Figures 1*j*–6 were performed on $\alpha 1$ $-/-$ or $\beta 2$ $-/-$ mice containing the neomycin resistance gene. The complete loss of $\alpha 1$ protein in $\alpha 1$ $-/-$ [$+neo$] mice was further substantiated by the absence of a major 50 kDa band on Western blots (Fig. 1*j*). Absence of the $\beta 2$ subunit protein in the $\beta 2$ $-/-$ mice could not be confirmed by Western blotting because no antibody specific for the $\beta 2$ subunit was available.

Pharmacological characterization of GABA_A receptors in $\alpha 1$ $-/-$ and $\beta 2$ $-/-$ mice

Autoradiography with [³⁵S]TBPS, a radioligand that binds to a site putatively located to the channel pore of all GABA_A receptors (Luddens and Korpi, 1995), confirmed a large and widespread loss of GABA_A receptors in cortex (–66 and –46%), septum (–53 and –60%), caudate putamen (–40 and –37%), globus pallidus (–72 and –74%), hippocampus (–53 and –27%), thalamus (–65 and –63%), and cerebellum (–67 and –71%) of both $\alpha 1$ $-/-$ and $\beta 2$ $-/-$ mice, respectively (Fig. 2). These losses were further demonstrated by the 66 and 58% reduction in the total number of GABA_A receptors, as measured by the binding of [³H]muscimol to membranes from the brains of $\alpha 1$ $-/-$ and $\beta 2$ $-/-$ mice (Table 1). Saturation experiments revealed a large reduction in [³H]Ro15–1788 binding sites in the forebrain of $\alpha 1$ $-/-$ (–57 ± 4%; *n* = 3) and $\beta 2$ $-/-$ (–51 ± 4%; *n* = 4) compared with wild-type mice with no change in the affinity of the radioligand (Tables 1, 2). The expression of [³H]Ro15–1788 binding sites was also reduced in the cerebellum of $\alpha 1$ $-/-$ (–67 ± 3%; *n* = 3) and $\beta 2$ $-/-$ (–71 ± 5%; *n* = 4) animals. Surprisingly the reduction of [³H]Ro15–1788 binding in the cerebellum of $\alpha 1$ $-/-$ mice was less than expected because the $\alpha 1\beta\gamma 2$ subtype is reported to be the main benzodiazepine-sensitive GABA_A receptor in this brain region, accounting for 90 ± 3% (*n* = 4) of total [³H]Ro15–1788 sites, as determined by immunoprecipitation with an $\alpha 1$ -specific antisera. Additional immunoprecipitation experiments with an $\alpha 3$ subunit-specific antisera indicated that in wild-type mice the remaining cerebellar [³H]Ro15–

$\alpha 1$ and $\beta 2$ allele, respectively. *c, g*, Targeted allele after homologous recombination for $\alpha 1$ and for $\beta 2$ allele, and after Cre-mediated removal of the neomycin resistance gene (*d, h*), respectively. *i*, Northern blot results using 45 mer oligonucleotides for the $\alpha 1$, $\alpha 3$, $\alpha 6$, $\beta 2$, and $\beta 3$ subunit genes of the GABA_A receptor and neomycin resistance (*neo*) and glyceraldehyde-3-phosphate dehydrogenase gene as probes; 5 μ g of poly(A⁺) selected RNA was loaded per lane; $\alpha 1^{+/+}$ and $\beta 2^{+/+}$: wild-type control samples; $\alpha 1^{-/-}$ [$-neo$] and $\beta 2^{-/-}$ [$-neo$]: samples from $\alpha 1$ and $\beta 2$ homozygotes lacking the neo gene in their genome; $\alpha 1^{-/-}$ [$+neo$] and $\beta 2^{-/-}$ [$+neo$]: samples from $\alpha 1$ and $\beta 2$ homozygotes with the neo gene in their genome. *j*, Western blot demonstrating the absence of $\alpha 1$ polypeptide in the brain of $\alpha 1$ $-/-$ mice.




Figure 2. Loss of GABA_A receptors in the forebrain and cerebellum of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice. Color-coded autoradiograms for [³⁵S]TBPS (8 nM) binding to sections of mouse brain revealed a widespread reduction of GABA_A receptors. Major losses are observed in cortex (−66 and −46%), globus pallidus (−72 and −74%), thalamus (−65 and −63%), and cerebellum (−67 and −71%) for example of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice, respectively. Scale bars, 1 mm.

1788 binding sites ($14 \pm 3\%$; $n = 3$) are contributed by $\alpha 3\beta\gamma 2$ receptors. These observations pointed toward a putative upregulation of $\alpha 3$ subunit containing GABA_A receptors in the cerebellum of $\alpha 1^{-/-}$ mice (see below). Radioligand inhibition binding studies with [³H]Ro15-1788 and a number of benzodiazepine site compounds revealed no significant (Student's *t* test, unpaired, two-tailed) changes in the pharmacology of the remaining GABA_A receptors in both $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mouse lines (Table 2) with the exception of the complete loss of high-affinity binding sites for zolpidem, an $\alpha 1$ -selective compound (Sieghart, 1995), in $\alpha 1^{-/-}$ mice whereas high-affinity zolpidem binding sites still accounted for 49% of total [³H]Ro15-1788 sites in $\beta 2^{-/-}$ brains (Table 2).

Electrophysiological analyses of GABA_A receptors in Purkinje neurons of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice

Cerebellar Purkinje neurons have been shown to express a limited repertoire of GABA_A receptor subunits, predominantly con-

Table 1. Amount (fmol/mg protein) of GABA_A receptors and benzodiazepine binding sites in WT, $\alpha 1^{-/-}$, and $\beta 2^{-/-}$ mice determined by nonlinear regression analysis of saturation data sets

	Wild type	$\alpha 1^{-/-}$ ^a	$\beta 2^{-/-}$ ^b
Muscimol	3014 ± 274	1011 ± 241	1275 ± 210
Ro15-1788:			
Forebrain (F3)	2677 ± 174	1160 ± 219	1326 ± 143
Cerebellum (F3)	1212 ± 123	419 ± 18	339 ± 34

Note that Ro15-1788 values for $\alpha 1^{-/-}$ derived from F3 generation mice. Data are the mean ± SEM of three or four experiments.

^aAll B_{max} are different from wild-type $p < 0.003$ (*t* test; unpaired, two-tailed).

^bAll B_{max} are different from wild-type $p < 0.002$ (*t* test; unpaired, two-tailed).

Table 2. Pharmacological characterization of GABA_A receptors in WT, $\alpha 1^{-/-}$, and $\beta 2^{-/-}$ mice

	K_d^*/K_i (nM)		
	Wild type	$\alpha 1^{-/-}$	$\beta 2^{-/-}$
Muscimol*	30 ± 5	16 ± 6	41 ± 18
Ro15-1788*	2.4 ± 0.6	1.9 ± 0.3	1.8 ± 0.4
Flunitrazepam	9.7 ± 2.2	5.1 ± 1.4	13 ± 3
Ro15-4513	12 ± 3	9.6 ± 2.7	10 ± 3
Zolpidem			
H.A.	35 ± 12	n.d.	23 ± 7
% H.A.	62 ± 15	n.d.	49 ± 14
L.A.	784 ± 312	770 ± 358	727 ± 267

Data are the mean ± SEM of three to five experiments. K_d values were determined from nonlinear regression analysis of full saturation curves. Student's *t* test (unpaired, two-tailed) indicated that compounds have a similar affinity for GABA_A receptors in wild-type than in $\alpha 1^{-/-}$ or $\beta 2^{-/-}$ mice, respectively. n.d., Not detected.

sisting of $\alpha 1\beta 2\gamma 2$ (Laurie et al., 1992a; Persohn et al., 1992; Fritschy and Mohler, 1995). These cells then provided an ideal candidate for the study of the effects of elimination of the $\alpha 1$ or the $\beta 2$ subunits by gene targeting. Whole-cell patch-clamp recordings were made from dissociated cerebellar Purkinje neurons isolated from both the $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice and compared with those of wild-type littermates. Recordings from wild-type neurons revealed the presence of robust GABA-mediated currents in all cells tested with a mean amplitude of 2982 ± 271 nA ($n = 30$) (Fig. 3a) in response to 1 mM GABA application. GABA EC₂₀ currents were significantly enhanced by the benzodiazepine modulators chlordiazepoxide ($148 \pm 8\%$ at 3 μ M; $n = 20$) and the $\alpha 1$ -selective compound zolpidem ($182 \pm 13\%$ at 100 nM; $n = 17$) (Fig. 3a,b). In contrast, 23 of 52 of the cells from the $\alpha 1^{-/-}$ mice did not produce a response to 1 mM GABA. In those cells that did respond to GABA, these currents were significantly reduced in amplitude (mean of 52 cells, 257 ± 54 pA). Comparing measured EC₂₀ values in these cells also demonstrated a significant decrease in the potency of GABA in the $\alpha 1^{-/-}$ cells [3.8 ± 0.3 μ M (20) in wild-type and 19.0 ± 4.6 μ M (5) in $\alpha 1^{-/-}$]. Of those cells with GABA currents large enough to measure an EC₂₀ response, five were studied using the benzodiazepine modulators. The potentiation by chlordiazepoxide was identical to that in wild-type animals ($138 \pm 35\%$), however the effect of 100 nM zolpidem was markedly reduced ($34 \pm 17\%$), suggesting that these receptors contained either $\alpha 2$ or $\alpha 3$ subunits or are formed from $\beta 2\gamma 2$ alone. Similar experiments using the $\beta 2^{-/-}$ animals revealed a different profile. In contrast to the $\alpha 1^{-/-}$ mice, every cell responded to GABA, however the mean current amplitude was reduced compared with wild-type (1234 ± 144 pA; $n = 13$). As with the $\alpha 1^{-/-}$ mice the potentiation of the GABA currents of Purkinje cells from $\beta 2^{-/-}$ mice by chlordiazepoxide was similar to wild-type ($104 \pm 17\%$; $n = 6$), however there was marked potentiation produced by 100 nM zolpidem ($108 \pm 15\%$; $n = 5$). The effect of zolpidem was significantly reduced compared with wild-type mice, suggesting that the receptors present contain a mixed population of $\alpha 1\beta x\gamma 2$ and $\alpha 2/\alpha 3\beta x\gamma 2$. To determine the β -isoform present in the receptors in the $\beta 2^{-/-}$ mice, the effects of the β -subunit-selective agents loreclezole and etomidate were investigated. These compounds selectively potentiate receptors containing a $\beta 2$ or $\beta 3$ subunit, but not $\beta 1$ -containing receptors (Wafford et al., 1994; Hill-Venning et al., 1997). Robust potentiation of the GABA EC₂₀ by both agents was observed in both the




Figure 3. Electrophysiological analysis of GABA currents recorded from cerebellar Purkinje neurons of wild-type and $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice. *a*, Example recordings from wild-type (WT), $\alpha 1^{-/-}$, and $\beta 2^{-/-}$ mice, of the response to 1 mM GABA, and potentiation of submaximal, EC₂₀ currents by 3 μ M chlordiazepoxide or 100 nM zolpidem (this concentration would distinguish $\alpha 1$ from other subtypes). Amplitude is indicated by the scale bar in each case, and drugs were applied as shown by the bar above each response. *b*, Mean current amplitude in response to 1 mM GABA from isolated Purkinje neurons; *n* number is shown in parentheses above each column and includes all cells tested including unresponsive cells. *c*, Mean potentiation of GABA EC₂₀ by 3 μ M chlordiazepoxide; *n* number is shown in parentheses above each column. *d*, Mean potentiation of GABA EC₂₀ by 100 nM zolpidem; *n* number is shown in parentheses above each column.

wild-type and $\beta 2^{-/-}$ mice (Fig. 4), suggesting that the remaining receptors expressed in Purkinje contained predominantly $\beta 3$ subunits.

Behavioral phenotype of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice

After breeding of $\alpha 1$ heterozygotes of the F1 generation, $\alpha 1^{-/-}$ mice were found to be under-represented in the F2 generation, accounting for only 13.4% of all offspring (29 $\alpha 1^{-/-}$ of 217 pups in total). The surviving $\alpha 1$ homozygotes appeared to have lower body weights and were less well groomed before weaning, but improved their appearance with age. Therefore, the behavioral analysis focused on adult animals only. The $\beta 2^{-/-}$ mice were represented in a Mendelian manner in the F2 generation (23 of 95 animals in total) and had no obvious phenotypical abnormalities.

A neurological screen was initially performed on adult F2 mice to determine whether there were any major neurological deficits in these animals. Neither $\alpha 1^{-/-}$ nor $\beta 2^{-/-}$ mice displayed any major deficits on beam balancing and swimming ability tests (data not shown). Two-month-old $\alpha 1^{-/-}$ were smaller and had lower body weights (~30%) than wild-type controls, and this difference persisted until at least 3 months of age. $\alpha 1^{-/-}$ mice were also observed to have a tremor when handled, but this did not impair




Figure 4. Effects of loreclezole, etomidate, and pentobarbital on GABA receptors from WT and $\beta 2^{-/-}$ Purkinje neurons. *a*, Recording from WT and $\beta 2^{-/-}$ neurons showing a GABA response to 1 mM GABA followed by the potentiation of a GABA EC₂₀ response by 3 μ M loreclezole, 30 μ M etomidate, and 100 μ M pentobarbital. Amplitude is indicated by the scale bar, and drugs were applied as shown by the bar above each response. *b–d*, Mean potentiation of the GABA EC₂₀ response by 3 μ M loreclezole (*b*), 30 μ M etomidate (*c*), and 100 μ M pentobarbital (*d*); *n* number is shown in parentheses above each column.

their ability to perform motor tasks. $\beta 2^{-/-}$ mice had normal body weights (data not shown).

A more detailed assessment of motor coordination was performed using the rotarod. A two-factor repeated measures ANOVA indicated that both $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice were as capable of remaining on the rotarod as wild-type controls (genotype: $F_{(2,19)} = 0.05$; $p = 0.95$) (Fig. 5*a*). As expected, performance declined as the revolution speed increased (speed: $F_{(6,114)} = 22.73$; $p < 0.00005$), but there were no deficits seen in either of the knock-out lines. $\alpha 1^{-/-}$ mice displayed a similar level of spontaneous locomotor activity and exploration compared with wild-types (Fig. 5*b*) when placed in the novel environment of activity chambers (genotype: $F_{(1,18)} = 0.02$; $p = 0.66$; repeated measures ANOVA). They also habituated to this environment over a similar time scale as wild-types (time \times genotype interaction: $F_{(29,522)} = 0.90$; $p = 0.51$). In contrast, $\beta 2^{-/-}$ animals (Fig. 5*c*) exhibited a much higher level of activity in this test (genotype: $F_{(1,22)} = 10.63$; $p = 0.0036$), although they did habituate to a similar degree as wild-types (time \times genotype interaction: $F_{(29,638)} = 0.72$; $p = 0.86$).




Figure 5. Behavioral evaluation of $\alpha 1$ and $\beta 2$ $-/-$ mice. *a*, The duration a mouse remained walking on the rotarod revolving at different speeds (18–36 rpm) decreased as the speed was increased. $\alpha 1$ and $\beta 2$ $-/-$ mice did not differ from wild-type mice at any speed examined. Data are the means \pm SEM; $n = 6$ –9. The spontaneous locomotor activity of $\alpha 1$ $-/-$ mice (*b*) did not differ from wild-type mice. In contrast, $\beta 2$ $-/-$ mice (*c*) showed a marked increase in activity ($p < 0.005$) compared with wild-type mice. Data are the mean number of cage crosses in 2 min time bins \pm SEM; $n = 10$ –12.

Table 3. Litter size increases across several generations

Genotype	Generation		
	F3	F4	F5
WT	6.96 \pm 0.65 (23) ^a	6.13 \pm 0.77 (15)	8.35 \pm 0.50 (34)
$\alpha 1$ $-/-$	3.06 \pm 0.40 (16)	4.46 \pm 0.54 (22)	6.46 \pm 0.71 (28) ^b

Litter size (Mean number of pups \pm s.e.m.).

^{a,b} $p < 0.05$ compared with F3 $\alpha 1$ $-/-$; numbers in brackets indicate number of litters analyzed.

Generation-dependent litter size and upregulation of $\alpha 2$ and $\alpha 3$ subunits in $\alpha 1$ $-/-$ mice

To establish an $\alpha 1$ $-/-$ mouse line, homozygotes of the F2 generation were interbred using a variety of breeding pairs and avoiding any brother–sister matings. Similarly, $\beta 2$ $-/-$ and wild-type control lines originating from their F2 littermates were established. However, there were some indications that the $\alpha 1$ $-/-$ surviving pups appeared to be phenotypically less affected in the F4 and F5 generation in comparison with the initial findings of the $\alpha 1$ homozygotes in the F2 generation. Therefore, we analyzed the number of pups in each litter for the $\alpha 1$ $-/-$




Figure 6. Generation-dependent upregulation of $\alpha 2$ and $\alpha 3$ subunits in cerebellum and forebrain of $\alpha 1$ $-/-$ mice. *a*, The loss of [³H]Ro15–1788 binding sites increased in cerebellum (*left panel*) and forebrain (*right panel*) of $\alpha 1$ $-/-$ mice from generation F2 to F5 to finally reach the proportion of [³H]Ro15–1788 sites immunoprecipitated by a selective $\alpha 1$ antibody from wild-type membrane (*filled bars*). *** $p < 0.001$; Student's *t* test. *b*, Evidence for an upregulation of $\alpha 3$ subunit in the cerebellum of $\alpha 1$ $-/-$ mice. Western blot showing an increase in $\alpha 3$ subunit expression in knock-out mice compared with wild-type (*left panel*). Quantitative immunoprecipitation with a $\alpha 3$ antibody demonstrated that $\alpha 3$ subunit-containing GABA_A receptors account for 14 \pm 3% (mean \pm SEM; $n = 3$) and 92 \pm 8% (mean \pm SEM; $n = 4$) of total [³H]Ro15–1788 binding sites in the cerebellum of wild-type and $\alpha 1$ $-/-$ mice, respectively (*right panel*). *c*, Quantitative immunoprecipitation demonstrated a 42% increase in the expression of $\alpha 2$ and $\alpha 3$ subunit-sensitive [³H]Ro15–1788 binding sites in the forebrain of $\alpha 1$ $-/-$ compared with wild-type mice.

mouse line in detail. The litter size from wild-type matings remained constant at approximately seven pups per litter between the F3 and F5 generations (Table 3). In contrast, F2 $\alpha 1^{-/-}$ matings produced significantly smaller litters (approximately three pups per litter), but litter size increased with increasing generation number.

The possibility that this generation-dependent phenotypic change is paralleled by modification in GABA_A receptor expression was investigated with [³H]Ro15-1788 saturation binding, a radioligand that would likely detect upregulation or downregulation of GABA_A receptors (Fig. 6*a,b*). Indeed, in the cerebellum of $\alpha 1^{-/-}$ mice, a generation-dependent reduction in the amount of GABA_A receptors was observed. Specifically, the loss of total [³H]Ro15-1788 binding sites determined by Scatchard analyses was increased from generation F2 (49%; $n = 1$) to generation F3 (67 ± 3%; $n = 3$). The reduction in generation F3 was significantly different ($p < 0.001$; Student's *t* test) from generation F5 (87 ± 1%; $n = 4$) which was equivalent to the amount of wild-type cerebellar $\alpha 1$ receptors sensitive to Ro15-1788 (90 ± 3%; $n = 4$), as determined by immunoprecipitation with an $\alpha 1$ -specific antibody (Fig. 6*a*). The last observation indicated that in F5 generation mice there is no longer an upregulation of benzodiazepine-sensitive GABA_A receptors. A similar trend was demonstrated in forebrains of $\alpha 1^{-/-}$ mice where reduction in [³H]Ro15-1788 sites increased from 45% ($n = 1$) in generation F2 to 57 ± 4% ($n = 3$) in generation F3, and finally reached 66 ± 2% ($n = 4$) in generation F5. In this F5 generation, no upregulation of [³H]Ro15-1788 sites occurred because the recorded reduction was similar to the number of wild-type forebrain $\alpha 1$ receptors sensitive to Ro15-1788 (67 ± 3%; $n = 4$) determined by immunoprecipitation as well as the proportion of high-affinity zolpidem binding sites (62 ± 15%; $n = 4$) found in wild-type mouse (Fig. 6*b*, Table 2). To determine which GABA_A receptor subtype was responsible for this upregulation in the F3 generation, we focused our attention on $\alpha 2$ and $\alpha 3$ subunits that are the most abundantly expressed subunits after the $\alpha 1$ subunit in adult brain and also the predominant embryonic isoforms (Laurie et al., 1992*b*). In cerebellum, Western blot analyses revealed a marked increase in the expression of $\alpha 3$ subunit in $\alpha 1^{-/-}$ compared with wild-type mice (Fig. 6*c*). The faint band observed in wild-type cerebellum is in agreement with $\alpha 3$ receptors accounting for 14 ± 3% ($n = 3$) of total [³H]Ro15-1788 binding sites in this brain region. Furthermore, quantitative immunoprecipitation experiments allowed us to demonstrate that an $\alpha 3$ subunit-specific antiserum precipitated all (92 ± 8%; $n = 4$) available [³H]Ro15-1788 binding sites (Fig. 6*c*). These results indicated that upregulation of $\alpha 3$ subunit in the cerebellum of $\alpha 1^{-/-}$ is solely responsible for the increase in total [³H]Ro15-1788 sites observed in F3 generation. The contribution of $\alpha 2$ and $\alpha 3$ subunits in the upregulation of [³H]Ro15-1788 binding sites in the forebrain of $\alpha 1^{-/-}$ mice was investigated by quantitative immunoprecipitation. Results showed that $\alpha 2$ and $\alpha 3$ subunits account for 17 ± 3% ($n = 4$) and 14 ± 4% ($n = 4$) of total [³H]Ro15-1788 sites in wild-type mice, respectively, and their respective contribution increased to 56 ± 7% ($n = 4$) and 45 ± 7% ($n = 4$) in $\alpha 1^{-/-}$ animals. Given that the total number of [³H]Ro15-1788 sites in the forebrain of F3 generation $\alpha 1^{-/-}$ mice has been shown to represent 43% of that in wild-type mice (Table 1), a net increase of 42% in the expression of both $\alpha 2$ and $\alpha 3$ subunits was estimated (Fig. 6*d*). Putative changes in the expression of minor populations of GABA_A receptors were also investigated by various pharmacological approaches. Quantitative autoradiography

Table 4. Variation in the expression of various GABA_A receptor subtypes in $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice

Subtype	$\alpha 1^{-/-}$	$\beta 2^{-/-}$
$\alpha 1\beta\gamma 2/3$	-100%	-60%
$\alpha 2\beta\gamma 2/3$	0%	-48%
$\alpha 3\beta\gamma 2/3$	0%	-69%
$\alpha 4\beta\gamma 2/3$	0%	-60%
$\alpha 5\beta\gamma 2/3$	0%	-39%
$\alpha 6\beta\gamma 2/3$	-38%	-62%

For $\alpha 1^{-/-}$ mice, data are derived from F5 generation animals. In $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice, data correspond to $\alpha 1$, $\alpha 2$, $\alpha 3$ expression in forebrain, $\alpha 4$ expression in cortex, thalamus, and hippocampus, $\alpha 5$ expression in hippocampus and endopiriform nucleus, and $\alpha 6$ expression in cerebellum. The data were obtained through various methodological approaches such as immunoprecipitation ($\alpha 1-3$ in $\beta 2^{-/-}$), autoradiography ($\alpha 4$, $\alpha 5$), Western blot ($\alpha 1$), and radioligand binding ($\alpha 1-4$ and $\alpha 6$).

with [³H]L-655,708, a $\alpha 5$ subtype-selective ligand, revealed no change in expression of this subunit in CA1-CA3 fields of hippocampus (96% of wild-type), dentate gyrus (99% of wild-type), and endopiriform nucleus (99% of wild-type) (Table 4). Similar autoradiography experiments with [³H]Ro15-4513 plus 20 μ M diazepam showed no changes in the expression of $\alpha 4\beta\gamma 2$ receptors in the thalamus, hippocampus, and cortex of $\alpha 1^{-/-}$ mice. However, binding experiments with 40 nM [³H]Ro15-4513 plus 10 μ M diazepam on cerebellar membranes from F5 generation $\alpha 1^{-/-}$ mice demonstrated a significant reduction (-38 ± 10%; $n = 4$) in the expression of $\alpha 6$ receptors (Table 4). This reduction did not result from the presence of the neomycin cassette in the targeted gene because a similar ($p > 0.42$; Student's *t* test) 26 ± 10% ($n = 4$) reduction in $\alpha 6$ subunit expression was determined in the cerebellum of F6 generation *cre*-excised $\alpha 1^{-/-}$ [-neo] mice.

Effect of deletion of the $\beta 2$ subunit gene on the expression of the various α subunits

To determine whether the compensatory changes in subunit expression observed in $\alpha 1^{-/-}$ mice were because of the loss of $\alpha 1$ subunit or the loss of the $\alpha 1\beta 2\gamma 2$ receptor per se, the expression of the various α subunits was investigated in the $\beta 2^{-/-}$ mice. Immunoprecipitation experiments indicated that $\alpha 1$, $\alpha 2$, and $\alpha 3$ subunits represented 54 ± 12% ($n = 4$), 18 ± 5% ($n = 4$), and 11 ± 1% ($n = 4$) of total forebrain [³H]Ro15-1788 binding sites from $\beta 2^{-/-}$ mice, respectively. These proportions equated to a reduction by 60, 48, and 69% of $\alpha 1$, $\alpha 2$, and $\alpha 3$ subunit-containing receptors (Table 4), given the fact that in $\beta 2^{-/-}$ brains the [³H]Ro15-1788 binding sites represent only 49% (Table 1) of the wild-type complement. Quantitative autoradiography using the $\alpha 5$ subtype-selective ligand [³H]L-655,708 revealed an overall 39% reduction in the expression of $\alpha 5$ subunit with losses of 33% in the hippocampus and 53% in the endopiriform nucleus. Similar experiments with [³H]Ro15-4513 plus 20 μ M diazepam revealed significant losses of $\alpha 4\beta\gamma 2$ receptors in various regions such as the cortex (-55%), the hippocampus (-50%), and the thalamus (-59%). Radioligand binding experiments on mice forebrain with 40 nM [³H]Ro15-4513 plus 10 μ M diazepam confirmed these autoradiographic measures and demonstrated an overall 60 ± 7% ($n = 5$) decrease in $\alpha 4$ -containing receptors (Table 4). Finally, the expression of diazepam-insensitive [³H]Ro15-4513 receptors in the cerebellum of $\beta 2^{-/-}$ mice was reduced by 62 ± 6% ($n = 5$) (Table 4). This downregulation of $\alpha 6$ receptors was again independent of the presence of the neomycin cassette in the targeted gene because diazepam-insensitive [³H]Ro15-4513 sites were

also decreased by $80 \pm 3\%$ ($n = 2$) in the cerebellum of $\beta 2^{-/-}$ [$-neo$] mice.

DISCUSSION

Mouse lines lacking functional GABA_A receptor subunits $\alpha 1$ and $\beta 2$ have been generated to further our understanding of the role of the GABAergic system in the control of inhibitory neurotransmission in the CNS. As expected, a widespread elimination of GABA_A receptors was observed with brain regions having the strongest expression of the $\alpha 1$ and $\beta 2$ subunit experiencing the highest loss of receptors. The consequences of these disruptions in the GABAergic system on behavior and regulation of subunit expression and assembly are discussed.

Lack of major phenotypic deficiency in adult $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice

In both knock-out mouse lines $\sim 60\%$ of the total number of GABA_A receptors were lost, and it was therefore surprising to find that adult $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice do not experience major phenotypic abnormalities or spontaneous seizures. Indeed, targeted disruptions of the $\beta 3$ or $\gamma 2$ subunit of the GABA_A receptor result in a similar substantial loss of GABA_A receptors, but lethality and serious epileptic seizures have been reported for both $\beta 3$ - and $\gamma 2$ -deficient mice (Gunther et al., 1995; Homanics et al., 1997). The balanced reduction in the $\beta 2^{-/-}$ mice of receptors containing each of the six α subunits could explain the seemingly normal appearance of the $\beta 2^{-/-}$ mice. These observations also allude to the putative existence of spare receptors among GABA_A receptor subtype populations. Adult $\alpha 1^{-/-}$ mice, but not $\beta 2^{-/-}$ mice, did exhibit tremor when handled, despite a wide codistribution in the brain of $\alpha 1$ and $\beta 2$ subunit and a similar loss in the total number of GABA_A receptors. It would be of interest to generate an $\alpha 1/\beta 2$ double knock-out mouse, but the close linkage of these two genes makes this an impractical task given current technologies. Although it is likely that there are multiple anatomical and physiological substrates responsible for this phenotypic difference, the different responsiveness to GABA of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ Purkinje cells may be a contributing factor. GABA currents were almost absent in the Purkinje cells of $\alpha 1^{-/-}$, but only halved in those of $\beta 2^{-/-}$ mice compared with wild-type animals. This is in agreement with the predominant expression of $\alpha 1$ subunit in Purkinje neurons, whereas there is equivalent expression of $\beta 2$ and $\beta 3$ subunits (Fritschy et al., 1992; Wisden et al., 1992; Fritschy and Mohler, 1995). However, neither the presumed incomplete inhibitory control of the main cerebellar output, nor the pronounced reduction of $\alpha 1$ and $\alpha 6$ subunits in granule cells of $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ had a detectable impact on the motor capacity of the knock-out mice. Similarly, $\alpha 6^{-/-}/\delta$ deficient mice, in which the cerebellum contains only half the number of GABA_A receptors, have no impairment of motor skills (Jones et al., 1997; Nusser et al., 1999). Altogether, these genetic-based observations indicate that synaptic integration in granule and Purkinje cells is apparently preserved in cerebellum expressing only $\alpha 1$ or $\alpha 6$ or a mix of $\alpha 1$ and $\alpha 6$ (as in $\beta 2^{-/-}$ mice) subunits. Moreover they suggest that $\alpha 1$ and $\alpha 6$ subunits have some functional overlap in addition to common synaptic localization at the surface of granule cells (Nusser et al., 1998). The availability of these different strains of mice should prove very useful to dissect the role of GABAergic inhibition in cerebellar physiology.

Generation-dependent change in phenotype and subunit regulation

Initially, the $\alpha 1^{-/-}$ mice displayed a severely compromised phenotype, with only half of them surviving after birth and a variable phenotype in the survivors. A similar perinatal lethality has also been observed in GABA_A receptor δ subunit knock-out mice (Mihalek et al., 1999). However, the litter size of $\alpha 1^{-/-}$ mice increases in successive generations, which may be the result from selective breeding of the $\alpha 1$ homozygotes, which have a less compromised phenotype and therefore exhibit higher rates of reproduction. Interestingly, the increase in the litter size and a subjectively milder phenotype in the later generations correlates with decreased upregulation of the $\alpha 2/3$ subunit. However, it should be noted that this upregulation of $\alpha 2/3$ subunit was incomplete because $\sim 45\%$ of GABA_A receptors were still missing in the brains of mice from the F2 generation. At present it is not clear how this process originates and is regulated, nor why the $\alpha 2/3$ upregulation results in a subjectively more severe phenotype. A possible explanation would be that several regulatory events take place in $\alpha 1^{-/-}$ mice to compensate for their inability to switch on $\alpha 1$ subunit around the time of birth (Laurie et al., 1992b). Initially, retention of embryonically highly expressed $\alpha 2$ and $\alpha 3$ subunits (Laurie et al., 1992b) could be one event of this adaptive process and that it is then lost in later generations because of putative detrimental effects on animal development. At the molecular level, one could speculate that expression of membrane potential regulating ion channels is changed in a way that relatively normal neuronal inhibition can occur in the absence of $\alpha 1$ subunit-containing GABA_A receptors (Brickley et al., 2001). In such a context, overexpression of $\alpha 2$ and $\alpha 3$ subunit could boost inhibition to a level that impairs normal physiological development. Interestingly, the weight and size of the brain of $\alpha 1^{-/-}$ mice from F3 generation was somewhat smaller ($\sim 15\%$) than in wild-type controls. Although such adaptive phenomena limit our assessment of the role of the targeted protein, their future analysis may provide insights into neuronal regulatory mechanisms as well as on the role of compensatory proteins.

Assembly of GABA_A receptor subunits

Selective subunit knock-out mice represent an ideal system to study the abundance and assembly of subunit that generates native GABA_A receptors (Fritschy et al., 1997; Jones et al., 1997; Nusser et al., 1999). Combining data from analysis of both $\alpha 1^{-/-}$ and $\beta 2^{-/-}$ mice confirms that the combination $\alpha 1\beta 2\gamma 2/3$ is the predominant isoform accounting for 40% of total benzodiazepine-sensitive GABA_A receptors in mouse brain (McKernan et al., 1991; McKernan and Whiting, 1996). A specific decrease of $\alpha 6$ subunit protein by $\sim 30\%$ was found in the cerebellum of both adult $\alpha 1^{-/-}$ [$+neo$] and [$-neo$] mice in the later generations (Table 4). Unlike in the case for the $\alpha 6^{-/-}$ mice (Uusi-Oukari et al., 2000), the presence of the neomycin resistance gene in the targeted locus does not seem to have a major influence on the expression level of its neighboring GABA_A receptor genes in the $\beta 2-\alpha 6-\alpha 1-\gamma 2$ gene cluster (Russek, 1999) in the $\alpha 1$ deficient mice, as documented by normal transcriptional expression of the $\beta 2$, $\alpha 6$, and $\gamma 2$ genes. The coassembly of $\alpha 1$ and $\alpha 6$ subunits is known in rat cerebellum (Pollard et al., 1995; Khan et al., 1996; Jechlinger et al., 1998) where $\alpha 1\alpha 6\beta\gamma 2$ complexes may represent $\sim 40\%$ of diazepam-insensitive Ro15-4513 binding sites (Khan et al., 1996). Thus, the loss of $\alpha 6$ subunit in $\alpha 1^{-/-}$ mice could be explained by the existence of a partnership in which $\alpha 1$ subunit is required for proper expression of $\alpha 1\alpha 6\beta\gamma$ and/or $\alpha 1\alpha 6\beta\delta$. Such a

possibility would be reminiscent of the $\alpha 6/\delta$ partnership (Jones et al., 1997) and support the view that *in vivo* α subunits may play an important role in the assembly of GABA_A receptors (Fritschy et al., 1997; Jones et al., 1997).

The reduction of both total GABA_A receptors and [³H]Ro15-1788 binding sites in both the forebrain and cerebellum of $\beta 2$ -/- mice is consistent with expression levels of this subunit in rat brain (Benke et al., 1994; Li and DeBlas, 1997) and support the conclusion that $\beta 2$ is an abundant neuronal β subunit and there was no significant compensatory upregulation of $\beta 1$ or $\beta 3$ subunit. The $\beta 2$ -/- mice demonstrated a rather ubiquitous association of $\beta 2$ subunit with $\alpha 1$ -6 subunits in agreement with previous biochemical studies (Benke et al., 1994; Jechlinger et al., 1998) and the codistribution of $\beta 2$ and $\alpha 1$ -6 subunits in most brain regions (Persohn et al., 1992; Wisden et al., 1992).

In conclusion, biochemical and pharmacological analyses of $\alpha 1$ -/- and $\beta 2$ -/- mice demonstrated that the subunit combination $\alpha 1\beta 2\gamma 2$ constitutes the most abundant GABA_A receptor subtype in rodent brain, and the $\beta 2$ subunit can coassemble with all α subunits. The loss of half of total GABA_A receptors in both knock-out mouse lines and the subsequent disruption of GABA-mediated inhibition resulted in only mild behavioral deficits in the adult animals. Future studies will focus on the GABA-mediated synaptic transmission and the compensatory mechanisms responsible for the seemingly normal behavior of the $\alpha 1$ -/- and $\beta 2$ -/- mice.

REFERENCES

- Baer K, Essrich C, Balsiger S, Wick MJ, Harris RA, Fritschy J-M, Luscher B (2000) Rescue of $\gamma 2$ subunit-deficient mice by transgenic overexpression of the GABA_A receptor $\gamma 2S$ or $\gamma 2L$ subunit isoforms. *Eur J Neurosci* 12:2639–2643.
- Barnard EA, Skolnick P, Olson RW, Mohler H, Sieghart W, Biggio G, Braestrup C, Bateson AN, Langer SZ (1998) International Union of Pharmacology. XV. Subtypes of γ -aminobutyric acid_A receptors: classification on the basis of subunit structure and receptor function. *Pharmacol Rev* 50:291–313.
- Benke D, Fritschy J-M, Trzeciak A, Bannwarth W, Mohler H (1994) Distribution, prevalence, and drug binding profile of γ -aminobutyric acid type A receptor subtypes differing in the β -subunit variant. *J Biol Chem* 269:27100–27107.
- Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M (2001) Adaptive regulation of neuronal excitability by a voltage-independent K⁺ conductance. *Nature* 409:88–92.
- Crestani F, Lorez M, Baer K, Essrich C, Benke D, Laurent JP, Belzung C, Fritschy J-M, Luscher B, Mohler H (1999) Decreased GABA_A receptor clustering results in enhanced anxiety and a bias for threat cues. *Nat Neurosci* 2:833–839.
- Culiat CT, Stubbs L, Woychik RP, Russell LB, Johnson DK, Rinchik EM (1995) Deficiency of the $\beta 3$ subunit of the type A γ -aminobutyric acid receptor causes cleft palate in mice. *Nat Genet* 11:344–346.
- DeLorey TM, Handforth A, Anagnostaras SG, Homanics GE, Minassian BA, Asatourian A, Fanselow MS, Delgado-Escueta A, Ellison GD, Olson RW (1998) Mice lacking the $\beta 3$ subunit of the GABA_A receptor have the epilepsy phenotype and many of the behavioral characteristics of Angelman syndrome. *J Neurosci* 18:8505–8514.
- Fritschy J-M, Mohler H (1995) GABA_A receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol* 359:154–194.
- Fritschy J-M, Benke D, Mertens S, Oertel WH, Bachi T, Mohler H (1992) Five subtypes of type A γ -aminobutyric acid receptors identified in neurons by double and triple immunofluorescence staining with subunit-specific antibodies. *Proc Natl Acad Sci USA* 89:6726–6730.
- Fritschy JM, Benke D, Johnson DK, Mohler H, Rudolph U (1997) GABA_A-receptor α -subunit is an essential prerequisite for receptor formation *in vivo*. *Neuroscience* 81:1043–1053.
- Gunther U, Benson J, Benke D, Fritschy J-M, Reyes G, Knoflach F, Crestani F, Aguzzi A, Arigoni M, Lang Y, Bluethmann H, Mohler H, Luscher B (1995) Benzodiazepine-insensitive mice generated by targeted disruption of the $\gamma 2$ subunit gene of γ -aminobutyric acid type A receptors. *Proc Natl Acad Sci USA* 92:7749–7753.
- Hill-Venning C, Belelli D, Peters JA, Lambert JJ (1997) Subunit-dependent interaction of the general anaesthetic etomidate with the γ -aminobutyric acid type A receptor. *Br J Pharmacol* 120:749–756.
- Homanics GE, DeLorey TM, Firestone LL, Quinlan JJ, Handforth A, Harrison NL, Krasowski MD, Rick CEM, Korpi ER, Makela R, Brilliant MH, Hagiwara N, Ferguson C, Snyder K, Olsen RW (1997) Mice devoid of γ -aminobutyrate type A receptor $\beta 3$ subunit have epilepsy, cleft palate, and hypersensitive behavior. *Proc Natl Acad Sci USA* 94:4143–4148.
- Jechlinger M, Pelz R, Tretter V, Klausberger T, Sieghart W (1998) Subunit composition and quantitative importance of hetero-oligomeric receptors: GABA_A receptors containing $\alpha 6$ subunits. *J Neurosci* 18:2449–2457.
- Jones A, Korpi ER, McKernan RM, Pelz R, Nusser Z, Makela R, Mellor JR, Pollard S, Bahn S, Stephenson FA, Randall AD, Sieghart W, Somogyi P, Smith AJH, Wisden W (1997) Ligand-gated ion channel subunit partnerships: GABA_A receptor $\alpha 6$ subunit gene inactivation inhibits δ subunit expression. *J Neurosci* 17:1350–1362.
- Khan ZU, Gutierrez A, DeBlas AL (1996) The $\alpha 1$ and $\alpha 6$ subunits can coexist in the same cerebellar GABA_A receptor maintaining their individual benzodiazepine-binding specificities. *J Neurochem* 66:685–691.
- Laurie DJ, Seeburg PH, Wisden W (1992a) The distribution of thirteen GABA-A receptor subunit mRNAs in the rat brain. II. Olfactory bulb and cerebellum. *J Neurosci* 12:1063–1076.
- Laurie DJ, Seeburg PH, Wisden W (1992b) The distribution of thirteen GABA-A receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* 12:4151–4172.
- Li M, DeBlas AL (1997) Coexistence of two β subunit isoforms in the same γ -aminobutyric acid type A receptor. *J Biol Chem* 272:16564–16569.
- Luddens H, Korpi ER (1995) GABA antagonists differentiate between recombinant GABA_A/benzodiazepine receptor subtypes. *J Neurosci* 15:6957–6962.
- McKernan RM, Whiting PJ (1996) Which GABA_A receptor subtypes really occur in the brain? *Trends Neurosci* 19:139–143.
- McKernan RM, Quirk K, Prince R, Cox PA, Gillard NP, Ragan CI, Whiting P (1991) GABA_A receptor subtypes immunopurified from rat brain with α subunit-specific antibodies have unique pharmacological properties. *Neuron* 7:667–676.
- McKernan RM, Rosahl TW, Reynolds DS, Sur C, Wafford KA, Atack JR, Farrar S, Myers J, Cook G, Ferris P, Garrett L, Bristow L, Marshall G, Macaulay A, Brown N, Howell O, Moore KW, Carling RW, Street LJ, Castro JL, Ragan CI, Dawson GR, Whiting PJ (2000) Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA_A receptor $\alpha 1$ subtype. *Nat Neurosci* 3:587–592.
- Mihalek RM, Banerjee PK, Korpi ER, Quinlan JJ, Firestone LL, Mi Z-P, Lagenaar C, Tretter V, Sieghart W, Anagnostaras G, Sage JR, Fanselow MS, Guidotti A, Spigelman I, Li Z, DeLorey TM, Olsen RW, Homanics GE (1999) Attenuated sensitivity to neuroactive steroids in γ -aminobutyrate type A receptor delta subunit knock-out mice. *Proc Natl Acad Sci USA* 96:12905–12910.
- Nusser Z, Sieghart W, Somogyi P (1998) Segregation of different GABA_A receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci* 18:1693–1703.
- Nusser Z, Ahmad Z, Tretter V, Fuchs K, Wisden W, Sieghart W, Somogyi P (1999) Alterations in the expression of GABA_A receptor subunits in cerebellar granule cells after the disruption of the $\alpha 6$ subunit gene. *Eur J Neurosci* 11:1685–1697.
- Persohn E, Malherbe P, Richards JG (1992) Comparative molecular neuroanatomy of cloned GABA_A receptor subunits in the rat CNS. *J Comp Neurol* 326:192–216.
- Pollard S, Thompson CL, Stephenson FA (1995) Quantitative characterization of $\alpha 6$ and $\alpha 1\alpha 6$ subunit-containing native γ -aminobutyric acid_A receptors of adult rat cerebellum demonstrates two α subunits per receptor oligomer. *J Biol Chem* 270:21285–21290.
- Quirk K, Gillard NP, Ragan CI, Whiting PJ, McKernan RM (1994) γ -aminobutyric acid type A receptors in the rat brain can contain both $\gamma 2$ and $\gamma 3$ subunits, but $\gamma 1$ does not exist in combination with another γ subunit. *Mol Pharmacol* 45:1061–1070.
- Rosahl TW, Geppert M, Spillane D, Herz J, Hammer RE, Malenka RC, Sudhof TC (1993) Short term synaptic plasticity is altered in mice lacking synapsin I. *Cell* 75:661–670.
- Rosahl TW, Spillane D, Missler M, Herz J, Selig DK, Wolff JR, Hammer RE, Malenka RC, Sudhof TC (1995) Essential functions of synapsin I and II in synaptic vesicle regulation. *Nature* 375:488–493.
- Russek JJ (1999) Evolution of GABA_A receptor diversity in the human genome. *Gene* 227:213–222.
- Schwenk F, Baron U, Rajewsky KA (1995) *cre*-transgenic mouse strain for the ubiquitous deletion of *loxP*-flanked gene segments including deletion in germ cells. *Nucleic Acid Res* 23:5080–5081.
- Sieghart W (1995) Structure and pharmacology of γ -aminobutyric acid_A receptor subtypes. *Am Soc Pharmacol Exp Ther* 47:181–234.
- Soriano P, Montgomery C, Geske R, Bradley A (1991) Targeted disruption of the *c-src* proto-oncogene leads to osteoporosis in mice. *Cell* 64:693–702.

- Sur C, Quirk K, Dewar D, Atack JR, McKernan RM (1998) Rat and human hippocampal $\alpha 5$ subunit containing γ -aminobutyric acid-A receptors have $\alpha 5\beta 3\gamma 2$ pharmacological characteristics. *Mol Pharmacol* 54:928–933.
- Sur C, Fresu L, Howell O, McKernan RM, Atack JR (1999a) Autoradiographic localization of $\alpha 5$ subunit-containing GABA_A receptors in rat brain. *Brain Res* 822:265–270.
- Sur C, Farrar SJ, Kerby J, Whiting PJ, Atack JR, McKernan RM (1999b) Preferential coassembly of $\alpha 4$ and δ subunits of the γ -aminobutyric acid_A receptor in rat thalamus. *Mol Pharmacol* 56:110–115.
- Turner DM, Sapp DW, Olsen RW (1991) The benzodiazepine/alcohol antagonist Ro15–4513: Binding to a GABA_A receptor subtype that is insensitive to diazepam. *J Pharmacol Exp Ther* 257:1236–1242.
- Uusi-Oukari M, Heikkilä J, Sinkkonen ST, Makela R, Hauer B, Homann GE, Sieghart W, Wisden W, Korpi ER (2000) Long-range interactions in neuronal gene expression: evidence from gene targeting in the GABA_A receptor $\beta 2$ – $\alpha 6$ – $\alpha 1$ – $\gamma 2$ subunit gene cluster. *Mol Cell Neurosci* 16:34–41.
- Wafford KA, Bain CJ, Quirk K, McKernan RM, Wingrove PB, Whiting PJ, Kemp JA (1994) A novel allosteric modulatory site on the GABA_A receptor β subunit. *Neuron* 12:775–782.
- Whiting PJ, McKernan RM, Wafford KA (1995) Structure and pharmacology of vertebrate GABA_A receptor subtypes. *Int Rev Neurobiol* 38:95–138.
- Whiting PJ, Bonnert TP, McKernan RM, Farrar S, LeBourdelle B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghji DJS, Thompson SA, Wafford KA (1999) Molecular and functional diversity of the expanding GABA_A receptor gene family. *Ann NY Acad Sci*: 645–653.
- Wick MJ, Radcliffe RA, Bowers BJ, Mascia MP, Luscher B, Harris RA, Wehner JM (2000) Behavioural changes produced by transgenic overexpression of $\gamma 2L$ and $\gamma 2S$ subunits of the GABA_A receptor. *Eur J Neurosci* 12:2634–2638.
- Wisden W, Laurie DJ, Monyer HM, Seeburg PH (1992) The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon and mesencephalon. *J Neurosci* 12:1040–1062.
- Zhang J-H, Sato M, Tohyama M (1991) Different postnatal ontogenic profiles of neurons containing β ($\beta 1$, $\beta 2$ and $\beta 3$) subunit mRNAs of GABA_A receptor in the rat thalamus. *Dev Brain Res* 58:289–292.