



The Executive Functions in Overweight and Obesity: A Systematic Review of Neuropsychological Cross-Sectional and Longitudinal Studies

Francesca Favieri^{1*}, Giuseppe Forte¹ and Maria Casagrande^{2*}

¹ Department of Psychology, "Sapienza" University of Rome, Rome, Italy, ² Department of Dynamic and Clinical Psychology, "Sapienza" University of Rome, Rome, Italy

OPEN ACCESS

Edited by:

Antonios Dakanalis,
University of Milano Bicocca, Italy

Reviewed by:

Nicholas T. Bello,
Rutgers University, The State
University of New Jersey,
United States
Uku Vainik,
Montreal Neurological Institute and
Hospital, McGill University, Canada
Roger A. Dixon,
University of Alberta, Canada

*Correspondence:

Francesca Favieri
francesca.favieri@uniroma1.it
Maria Casagrande
maria.casagrande@uniroma1.it

Specialty section:

This article was submitted to
Eating Behavior,
a section of the journal
Frontiers in Psychology

Received: 08 May 2019

Accepted: 02 September 2019

Published: 20 September 2019

Citation:

Favieri F, Forte G and Casagrande M
(2019) The Executive Functions in
Overweight and Obesity: A Systematic
Review of Neuropsychological
Cross-Sectional and Longitudinal
Studies. *Front. Psychol.* 10:2126.
doi: 10.3389/fpsyg.2019.02126

Background: The increasing incidence of people affected by overweight or obesity is a significant health problem. The knowledge of the factors which influences the inappropriate eating behaviors causing excessive body fat is an essential goal for the research. Overweight and obesity are significant risk factors for many health diseases, such as cardiovascular problems, diabetes. Recently, many studies have focused on the relationship between body weight and cognitive processes.

Objectives: This systematic review is aimed to investigate the existence and the nature of the relationship between excessive body weight (overweight/obesity) and executive functions, analyzing cross-sectional, and longitudinal studies in order to verify the evidence of a possible causality between these variables.

Methods: The review was carried out according to the PRISMA-Statement, through systematic searches in the scientific databases PubMed, Medline, PsychInfo, and PsycArticles. The studies selected examined performance on executive tasks by participants with overweight or obesity, aged between 5 and 70 years. Studies examining eating disorders or obesity resulting from other medical problems were excluded. Furthermore, the results of studies using a cross-sectional design and those using a longitudinal one were separately investigated.

Results: Sixty-three cross-sectional studies and twenty-eight longitudinal studies that met our inclusion and exclusion criteria were analyzed. The results confirmed the presence of a relation between executive functions and overweight/obesity, although the directionality of this relation was not clear; nor did any single executive function emerge as being more involved than others in this relation. Despite this, there was evidence of a reciprocal influence between executive functions and overweight/obesity.

Conclusions: This systematic review underlines the presence of a relationship between executive functions and overweight/obesity. Moreover, it seems to suggest a bidirectional

trend in this relationship that could be the cause of the failure of interventions for weight reduction. The results of this review highlight the importance of a theoretical model able to consider all the main variables of interest, with the aim to structuring integrated approaches to solve the overweight/obesity problems.

Keywords: executive functions, obesity, overweight, cross-sectional studies, longitudinal studies

INTRODUCTION

Rationale

Obesity and overweight, defined as the accumulation of excessive body fat, are risk factors for many chronic diseases, such as hypertension (Jiang et al., 2016) and diabetes (Hauner, 2017) as well as musculoskeletal (McPhail et al., 2014) and respiratory problems (Littleton, 2012). Prospective studies have shown an association between obesity in adulthood and cognitive impairment in old age (Sanderlin et al., 2017). Moreover, obesity appears to be connected to psychopathologies—such as anxiety disorders and depression (Carpiniello et al., 2009; De Wit et al., 2010; Garipey et al., 2010; Luppino et al., 2010; Carey et al., 2014)—and to social difficulties—such as bullying and social isolation (Kolotkin et al., 2001). The most common risk factors associated with an increase in body weight are poor eating habits and a lack of adequate physical activity (World Health Organization, 2000; Prentice, 2001; Dubbert et al., 2002), which results in a chronic imbalance between individual's needs and energy acquisition (Yumuk et al., 2015).

Conventionally, overweight classifications are made according to the body mass index of an individual (BMI; World Health Organization, 2000). BMI takes into account the weight and height of a person, providing a quantifiable index as the measure of body mass. The WHO considers different severity of overweight: pre-clinical obesity (BMI between 25 and 29.9), obesity class I (BMI between 30 and 34.9), obesity class II (BMI between 35 and 39.9), obesity class III (BMI equal to or higher than 40). An excessive increase in BMI can lead to a higher risk of premature death and a lower quality of life (World Health Organization, 2000). However, some authors focused on other indices that appear to be more sensitive for investigating the relationship between different degrees of overweight and their effects on health, such as waist circumference, waist-to-height ratio, and the body adiposity index (Janssen et al., 2004; Ashwell et al., 2012; Lam et al., 2015). In general, an increase in body fat, in the absence of metabolic and hormonal pathologies (Bray, 1999), is strongly associated with overeating behaviors and excessive ingestion of high-caloric foods that affect the individual's metabolism (McCrory et al., 1999; Ouwens et al., 2003).

The prevalence of obesity or overweight has increased in recent years. In 2014, more than 1.9 billion adults were overweight (World Health Organization, 2015); of these, 600 million were classifiable as obese. Regarding younger people, in 2013 it was estimated that about 42 million children and adolescents between the ages of 5–18 years, and about 12.4% of children below the age of 5 years, were overweight or obese (World Health Organization, 2015; Yumuk et al., 2015). It is

expected that around 60% of the world's population will reach critical BMI values by 2030 (Kelly et al., 2008).

Considering these data, it appears useful to investigate the predisposing and exacerbating factors of increases in BMI and body fat, related to overeating behavior. In line with this need, recent years have seen increasing interest in the cognitive mechanisms involved in overweight or obesity (Liang et al., 2014; Forcano et al., 2018). Furthermore, recent studies (Yang et al., 2018) have focused on the relationship between executive functions and obesity to investigate the existence and nature of this association.

Executive Functions

Executive functions (EFs) is an “umbrella term” (Damasio, 1995; Elliott, 2003; Chan et al., 2008; Diamond, 2013) that includes both complex cognitive processes—such as the resolution of new tasks, the modification of existing behaviors, the planning of new strategies for problem solving, the sequencing of complex actions (Funahashi, 2001; Elliott, 2003), the inhibition of motor or cognitive automatic responses and the control of conflicting information (Diamond, 2013)—and lower-level of cognitive processes, which allow to regulate and control thoughts and actions during goal-directed behavior and involve different cognitive dimensions such as perception and sensation, memory and motivation, attention, reasoning, and problem-solving (Pennington and Ozonoff, 1996).

Although there are various EFs, many studies have centered on three specific processes (Miyake et al., 2000; Diamond, 2013): (i) Cognitive Flexibility (or Shifting), characterized by an attentional shift between tasks or between different mental operations; (ii) Working Memory (or Updating), which includes the updating and monitoring of mental representations in order to respond appropriately to external tasks or stimuli; and (iii) Inhibition, which consists of the voluntary inhibition of dominant or automatic responses for controlling actions, thoughts and emotions, as well as attentional aspects, in order to respond appropriately to the needs of goal-directed behaviors (Miyake et al., 2000; Hofmann et al., 2012; Diamond, 2013). In general, some EFs have been studied more than others due to the presence of cognitive tasks (e.g., Stroop task, Stop Signal Task, Iowa Gambling Task (IGT), Span Task, Maze task) that seem to be more sensitive in the investigation of specific EFs, however there are some limits in their interpretations (Diamond, 2013; Vainik et al., 2013).

Some authors (Grafman and Litvan, 1999; Chan et al., 2008) distinguished between two different groups of EFs: the “cold EFs” and the “hot EFs.” The first—which include verbal reasoning, problem-solving skills, planning, attentional

maintenance, cognitive flexibility, response inhibition, and control of conflicting information—are characterized by the absence of emotional processing of stimuli and do not generate emotional arousal (Chan et al., 2008). The second, the hot EFs—which include expectations of punishment-gratification, social behavior, and decision-making—are characterized by the presence of beliefs and desires, and they include a powerful emotional component (Chan et al., 2008). According to Miyake model, these executive domains would be included in cold EFs. However, in daily life, hot and cold EFs work jointly, and both are necessary to direct our behavior. It is essential to underline that EFs are characterized by individual differences, which during life undergo multiple modifications (Jacques and Marcovitch, 2010; Hall and Marteau, 2014), as the reduction of cognitive flexibility and planning with aging (see Jacques and Marcovitch, 2010), or the alterations in inhibition in psychopathology (Nigg, 2000). These differences and changes can also be traced back to the establishment of healthy behaviors, such as eating habits (Hall and Marteau, 2014).

Executive Functions in Obesity and Overweight

A recent review by Dohle et al. (2018) showed that some studies support the hypothesis that food behaviors affect executive functioning, i.e., healthy eating habits promote the preservation of cognitive functions throughout life (Morris et al., 2005; Smith and Blumenthal, 2016). Other authors are inclined to sustain the opposite point of view, in which cognitive functions are considered as the predictors of food behaviors and, consequently, of body weight changes. According to this view, EFs deficits are considered the cause of inappropriate attitudes to food and represent a trigger for both eating disorders and changes in BMI (Dohle et al., 2018). These different views on the relationship between EFs and eating habits are also observed in the studies that considered the association between obesity/overweight and EFs (Perry, 2004; Pignatti et al., 2006; Davis et al., 2007b; Gonzales et al., 2010).

The theoretical models that consider the relationship between EFs and overweight/obesity are less developed, and usually, they do not focus on specific executive processes. However, it might be interesting to extend this type of studies because they could help in identifying some aspects which are connected to the increase in obesity and related problems. To clarify the nature of the relationship between EFs and overweight/obesity could allow identifying both a causal direction in the relationship between EFs and excessive body weight and the most suitable theoretical model able to explain this relationship.

In general, it could be useful to define whether the studies investigating the relationship between EFs and excessive body weight showed a consensus about the presence of a clear link between the examined variables. In fact, some reviews tried to collect information about this relationship (Fitzpatrick et al., 2013; Vainik et al., 2013; Emery and Levine, 2017; Gettens and Gorin, 2017; Gluck et al., 2017; Yang et al., 2018), but they only confirmed the existence of a correlation between these dimensions, but they were unable to clarify the essence of

this relationship, nor the causality. Identifying whether the EFs represent predictors of weight gain (Smith and Robbins, 2013; Chen et al., 2017), or consequences of the increased body weight (Perry, 2004; Sellbom and Gunstad, 2012) still represents an important goal in research.

A review analyzing the relationship between EFs and overweight aimed to examine studies with different experimental designs (cross-sectional, longitudinal) could help in identifying an eventually causal relationship between variables, as well as it could allow understanding how the interactions that emerged in cross-sectional studies, change over time in longitudinal studies. In our view, this represents an essential goal, because it can be useful to both for structuring interventions aimed at reducing risks related to excessive body weight and/or EFs impairment and contributing to the development of a theoretical model. Moreover, studies analyzing the causality between these variables could be a starting point to identify whether some executive domains are more involved than others during body weight gain. An important aspect to consider is the role of every single executive domain in the relationship with overweight/obesity. To identify whether there is a specific EF or some EFs, which influences or are influenced by the excessive body weight could be useful both for the development of a theoretical model on EFs-overweight relationship and for the definition of risk factors related to excessive body weight or impairment in executive domains. Some studies identified impairment in specific EF domains as decision-making, planning and problem solving (for a review see Fitzpatrick et al., 2013), or in inhibition (Gluck et al., 2017). However, generally, the studies identified alteration of EFs, without well-defined the relationship between the single domain of EFs and excessive body weight. This tendency could be due to the different cognitive tasks used and the high number of methodological designs considered in the studies (Vainik et al., 2013).

Objectives

This systematic review aimed to analyse longitudinal and cross-sectional studies that have investigated the association between EFs and obesity or overweight in the absence of chronic diseases or related eating disorders, trying to add knowledge about the nature of this relationship.

Specifically, the aims of this systematic review are:

- to document the cross-sectional evidence between EFs and overweight/obesity, trying to identify the consensus on the presence of a relationship between EFs and excessive body weight;
- to see if any executive domain has been associated mainly with excessive body weight, considering both positive and negative results;
- to analyse longitudinal studies to assess the causality between EFs and the BMI, considering EFs eventually an outcome or predictor of increase in BMI.

This review represents an attempt to systematize the studies on the relationship between EFs and overweight (Vainik et al., 2013; Emery and Levine, 2017; Yang et al., 2018). The inclusion of longitudinal studies, by also considering different interventions

to reduce weight (as in Thiara et al., 2017) could help to clarify the nature of this relationship better. The final aim of this review is to understand how approaching the problems related to excess body weight.

METHOD

This systematic review was conducted according to the PRISMA-statement (Liberati et al., 2009; Moher et al., 2009). Online registration of the protocol has not been provided.

Research Strategies

The systematic review was conducted using PubMed, PsycINFO, PsycArticles, MedLine databases. The following keywords were used: “Executive Function,” “Inhibition,” “Cognitive Inhibition,” “Selective Attention,” “Updating,” “Working Memory,” “Shifting,” “Cognitive Flexibility,” “BMI,” “Overweight,” “Obesity,” “Overeating,” “Diet.”

The scripts used for the search are presented in **Table 1**.

The starting date of the work was January 8th, 2018. All original, “full-text” papers published in international, peer-reviewed journals up to June 10th, 2018 were considered.

Eligibility Criteria

Selections were made independently by two researchers (FF; MC) and any disagreements resolved by a supervisor (GF). All the studies investigated the relationship between EFs and excessive body weight. Studies including at least one group with overweight or obesity, classified through the international criteria as BMI (World Health Organization, 2000) and BMI percentiles (Flegal et al., 2002), and investigating at least one EF were included. Furthermore, both cross-sectional and longitudinal studies were considered and analyzed separately. For the selection of the articles the following inclusion criteria were used: (a) academic articles published in international, “peer-reviewed” journals; (b) studies written in English; (c) studies on humans with overweight or obesity (BMI higher than 25) at various levels of severity; (d) studies using cognitive tasks to assess EFs; (e) studies including participants aged between 5 and 70 years; (f) cross-sectional and longitudinal studies; (g) studies including different interventions to reduce body weight (bariatric surgery, cognitive remediation therapy, weight-loss programmes that included diets, or physical activity); (h) studies including other psychological variables related to EFs and body weight.

The following exclusion criteria were applied: (a) short report: these type of articles were excluded because after a preliminary analysis of them it was observed that the information reported was too general; (b) studies examining participants with binge eating disorder or other eating disorders; (c) studies focusing on cognitive functions other than executive ones; (d) studies analyzing EFs through self-report questionnaires; (e) studies on obesity of metabolic origin or caused by other medical diseases; (f) studies considering overweight in psychopathological or psychiatric conditions (e.g., depression, schizophrenia, ADHD, etc.). Moreover, for cross-sectional studies, the absence of a normal-weight control group for comparison of executive functioning was an additional exclusion criterion. For the

longitudinal researches, both observational and experimental studies were included.

Additionally, in both cross-sectional and longitudinal studies, differences between groups (with normal-weight and overweight), or between different assessment times (pre, post, follow-up) were mainly commented, although regression analyses (continuous BMI) were also considered. Correlational studies were excluded if the method did not include the presence of different conditions of BMI (including both normal-weight and overweight individuals).

Data Collection Process

According to PICOS (Liberati et al., 2009), the authors extracted from the selected articles information about participants in both the control group and the groups with overweight/obesity (age, BMI, gender), methods (executive tasks used), and main results observed in the EFs tasks.

According to the aims of this review, for the cross-sectional studies, all the results concerning comparisons between overweight/obese groups and normal-weight groups on a cognitive task that assessed one or more EFs were analyzed. For the longitudinal studies, all the results concerning the analysis over time of participants with excessive body weight were examined, including also changes in weight, and executive functioning following either weight-loss programmes or cognitive training. The characteristics of the studies are shown in **Tables 3, 4**.

Quality Assessment

A quality assessment analyzed the eligibility of each article by detecting the quality of the studies. This process was aimed to reduce the risk of bias selection and was conducted using a six-point checklist created explicitly for the screening of the studies of this review. For each point, a maximum score of two (high-quality) could be awarded per article: a score of zero corresponded to a low-quality index, a score of 1 to a medium-quality index and a score of 2 to a high-quality index. To derive an overall quality of score of the study, the mean score of each study was multiplied by 100. Studies with a score <75% were considered with high quality, in line with other qualitative analyses (e.g., Varkevisser et al., 2019). The systematic review excluded studies with very low quality (lower than 50%). **Table 2** shows the six-point quality assessment checklist. **Tables 3, 4** reported the quality assessment for each selected article.

RESULTS

Study Selection

The initial search produced 1,817 articles. After excluding 614 duplicates, 922 articles were rejected according to an analysis of both title and abstract, leaving a final total of 281 studies to be reviewed and subjected to the quality assessment.

At the end of the review process, 88 articles remained. The flow chart (**Figure 1**) shows the study selection process, including the number of studies found, the assessment process and the reasons for the exclusion of the articles.

The 88 selected articles were categorized according to the experimental design. Sixty-three studies used a cross-sectional

TABLE 1 | Script for the systematic research.

	Script
Executive function and Obesity	("executive function"[MeSH Terms] OR ("executive"[All Fields] AND "function"[All Fields]) OR "executive function"[All Fields]) AND (BMI[All Fields] OR ("overweight"[MeSH Terms] OR "overweight"[All Fields]) OR ("obesity"[MeSH Terms] OR "obesity"[All Fields]) OR ("hyperphagia"[MeSH Terms] OR "hyperphagia"[All Fields] OR "overeating"[All Fields]) OR ("diet"[MeSH Terms] OR "diet"[All Fields]))
Inhibition and Obesity	((("inhibition (psychology)"[MeSH Terms] OR ("inhibition"[All Fields] AND ("psychology"[All Fields]) OR "inhibition (psychology)"[All Fields]) OR "inhibition"[All Fields]) OR ("Cogn Int Conf Adv Cogn Technol Appl"[Journal] OR "cognitive"[All Fields]) AND ("inhibition (psychology)"[MeSH Terms] OR "inhibition"[All Fields]) AND ("psychology"[All Fields]) OR "inhibition (psychology)"[All Fields]) OR ("inhibition"[All Fields])) OR (Selective[All Fields] AND ("attention"[MeSH Terms] OR "attention"[All Fields])) AND (BMI[All Fields] OR ("overweight"[MeSH Terms] OR "overweight"[All Fields]) OR ("obesity"[MeSH Terms] OR "obesity"[All Fields]) OR ("hyperphagia"[MeSH Terms] OR "hyperphagia"[All Fields] OR "overeating"[All Fields]) OR ("diet"[MeSH Terms] OR "diet"[All Fields]))
Working Memory and Obesity	(Updating[All Fields] OR ("memory, short-term"[MeSH Terms] OR ("memory"[All Fields] AND "short-term"[All Fields]) OR "short-term memory"[All Fields] OR ("working"[All Fields] AND "memory"[All Fields]) OR "working memory"[All Fields]) AND (BMI[All Fields] OR ("overweight"[MeSH Terms] OR "overweight"[All Fields]) OR ("obesity"[MeSH Terms] OR "obesity"[All Fields]) OR ("hyperphagia"[MeSH Terms] OR "hyperphagia"[All Fields] OR "overeating"[All Fields]) OR ("diet"[MeSH Terms] OR "diet"[All Fields]))
Cognitive Flexibility and Obesity	(Shifting[All Fields] OR ("Cogn Int Conf Adv Cogn Technol Appl"[Journal] OR "cognitive"[All Fields]) AND ("pliability"[MeSH Terms] OR "pliability"[All Fields] OR "flexibility"[All Fields])) AND (BMI[All Fields] OR ("overweight"[MeSH Terms] OR "overweight"[All Fields]) OR ("obesity"[MeSH Terms] OR "obesity"[All Fields]) OR ("hyperphagia"[MeSH Terms] OR "hyperphagia"[All Fields] OR "overeating"[All Fields]) OR ("diet"[MeSH Terms] OR "diet"[All Fields]))

TABLE 2 | Checklist for quality assessment.

1) The use of standardized executive tasks*.	0 = No standardized tasks; 1 = Use of some non-standardized tasks; 2 = Use of all standardized tasks.
2) Controlling of psychological (e.g. depression, anxiety, emotional dysregulation) and/or physiological variables (e.g. blood values, hormonal and inflammatory aspects).	0 = No control of variables; 1 = Control of psychological or physiological variables; 2 = Control of both psychological and physiological variables.
3) The use of international guidelines for BMI classification.	0 = No international guidelines; 1 = Shared guidelines (i.e. CDC); 2 = International guidelines.
4) Quality of the method description (about executive variables).	0 = Procedures and assessment tools are not well indicated; 1 = Procedures and assessment tools are partially described; 2 = Procedure and assessment tools are well described.
5) Quality of results description (about executive variables).	0 = Executive functioning is not included in the results; 1 = Executive functioning is partially included in the results; 2 = Executive functioning is included in the results.
6) Quality of discussion and conclusion (about executive variables).	0 = Executive functioning is not included in either discussion or conclusion; 1 = Executive functioning is not well included in discussion and conclusion; 2 = Executive functioning is included in both discussion and conclusion.

*Behavioral tasks widely used in literature for the analysis of a specific executive function.

design, and twenty-eight studies used a longitudinal design (see **Tables 3, 4**). Three studies (Deckers et al., 2017; Demos et al., 2017; Vantieghem et al., 2018) used both cross-sectional and longitudinal design. These studies considered the differences between participants with normal-weight and participants with overweight or obesity and analyzed the differences in executive performances during the time. For this reason, they were considered in both sections of the review.

Quality Assessment for Risk Bias

Seventy-nine per cent of the studies ($N = 70$) were of high quality, while 20% ($N = 18$) were of low quality. **Figure 2** shows the percentage of studies per quality level for each point on the assessment tool. Overall, studies showed higher quality in their results and discussion sections. Conversely, lower scores were found for the control of psychological and physiological variables (**Figure 2**). The selection of the articles for the systematic review

was justified by the good quality of each study, explicitly considering the results on EFs. In general, despite a large number of the selected studies, the high quality of the studies may have reduced the risks of misinterpretation of the results.

Cross-Sectional Studies

Systematic searching gave 63 cross-sectional studies that met the inclusion criteria (see **Table 3**). Of these studies, twenty-nine involved adult participants (aged over 30 years), twenty examined adolescents (aged 12–22 years), two studies looked at young adults (aged 23–30 years) and finally twelve studies investigated the relationship between EFs and excessive body weight in children (aged >12 years) (see **Table 3**).

Only nine studies had a higher proportion of males than females (Pignatti et al., 2006; Verdejo-García et al., 2010; Gentier et al., 2013; Mole et al., 2015; Qavam et al., 2015; Reyes et al., 2015; Alarcón et al., 2016; Tsai et al., 2016; Wu et al.,

TABLE 3 | Cross-sectional studies.

Study	Participants					Cognitive task	Method			EF Domain	
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)		Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	Quality of the study
STUDY ON ADULT POPULATION											
Ariza et al. (2012)	OB ¹	42	31.81 (6.51)	67	38.3 (7.59)	TMT ³	–	OB equal to NW	OB equal to NW	OB equal to NW	83.3%
	NW ²	42	29.67 (6.97)	69	22.07 (1.97)	SCWT ⁴ WCST ⁵ Letter–Number Sequence					
Bongers et al. (2015)	OB	185	35.19 (7.59)	71	38.18 (6.17)	Stop–Signal Task	OB equal to NW	OB equal to NW	–	–	75.0%
	NW	134	33.04 (8.15)	74	22.35 (1.63)	Delay Discounting Task (food cue)					
Brogan et al. (2011)	OB	42	52.24 (10.89)	71	41.45 (9.17)	IGT ⁶	OB poor than NW	–	–	–	83.3%
	NW	50	47.34 (16.34)	66	24.36 (3.78)						
Catoira et al. (2016)	OB	81	30	100	35.81	WCST	–	OB poor than NW	–	OB equal to NW	91.7%
	NW	32	26.5	100	22.56	TMT SCWT Verbal Fluency					
Cohen et al. (2011)	OW	42	58.9 (8.3)	48	31.8 (6.8)	SCWT	–	OB poor than NW	OB poor than NW	OB poor than NW	66.7%
	NW	107	61.2 (8.0)	52	24.1 (1.4)	WCST TMT Digit Span					
Danner et al. (2012)	OB	18	44.56 (13.36)	100	30.84 (3)	IGT	OB poor than NW	–	–	–	83.3%
	OB–BED ⁷	19	38.05 (10.97)	100	28.74 (6.25)						
	NW	30	36.13 (14.09)	100	22.32 (1.96)						
Dassen et al. (2018a)	OB	82	41.12 (12.62)	64.4	38.94 (5.24)	2–Back Task	–	OB poor than NW	OB poor than NW	OB equal to NW	66.7%
	NW	71	43.40 (13.44)	77.5	22.63 (1.53)	Stop–Signal Task TMT					
Deckers et al. (2017)	OB	545	58 (15)	58	31.2 (3.9)	Concept Shifting Test	–	–	–	OB poor than NW	83.3%
	NW	1262	48.9 (16.2)	46	24.9 (2.5)						
Demos et al. (2017)	OB	37	46.95 (7.9)	100	33.5 (3.9)	Food Choice Decision Making Task	OB poor than NW	–	–	–	83.3%
	NW	30	43.97 (8.9)	100	22.7 (1.8)						
Fagundo et al. (2012)	OB	52	40.5 (11.1)	100	39.8 (7.4)	WCST	OB poor than NW	OB poor than NW	–	OB poor than NW	75.0%
	AN ⁸	35	28.1 (8.2)	100	17.2 (1.4)	SCWT					
	NW	137	24.8 (7)	100	21.5 (2.7)	IGT					
Frank et al. (2014)	OB	11	42.6 (4)	100	40.2 (0.8)	Working Memory Task (food cue)	–	–	OB equal to NW	–	50.0%
	ExOB ⁹	9	42 (2.8)	100	27.1 (0.9)						
	NW	11	36.6 (3.8)	100	21.4 (0.5)						

(Continued)

TABLE 3 | Continued

Study	Participants					Cognitive task	Method			EF Domain		Quality of the study
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)		Global EF/other EFs	Inhibition	Updating	Cognitive flexibility		
Galioto et al. (2013)	OB	81	51.78 (16.96)	55.9	34.67 (5.59)	Digit Span Maze Test Switching of Attention Task	OB poor than NW	–	OB poor than NW	OB poor than NW	83.3%	
	OW ¹⁰	210	50 (17.24)	37.5	27.12 (1.45)							
	NW	288	44.72 (18.37)	58	22.35 (1.73)							
Gameiro et al. (2017)	OB	76	43.24 (9.05)	68	>30	WCST Go/No-Go Task Color Trait Test Verbal Fluency Motor Series	–	OB poor than NW	–	OB poor than NW	75.0%	
	NW	38	40.53 (10.75)	71	<25							
Gonzales et al. (2010)	OB	12	48.5 (8.6)	50	34.4 (3.5)	Digit Span COWAT ¹¹ TMT n-Back Task	–	OB equal to OW equal to NW	OB equal to OW equal to NW	OB equal to OW equal to NW	100.0%	
	OW	11	52 (5.1)	45	27.4 (1.4)							
	NW	9	51.8 (4.3)	77	22.4 (2.2)							
Gunstad et al. (2007) [1]	OW	140	32.40 (9.10)	46.4	28.4 (4.42)	Verbal Interference Task Switching of Attention Task Maze Test	OB poor than NW	OB poor than NW	–	OB poor than NW	75.0%	
	NW	178	31.56 (8.71)	55.1	22.09 (1.71)							
Gunstad et al. (2007) [2]	OW	58	60.4 (7.62)	55.1	29.17 (3.54)	Verbal Interference Task Switching of Attention Task Maze Task	OB poor than NW	OB poor than NW	–	OB poor than NW	75.0%	
	NW	32	58.34 (6.62)	53.4	23.09 (1.59)							
Hendrick et al. (2012)	OB	13	34.8 (9.6)	100	33.2 (2.6)	Stop-Signal Task	–	OB equal to OW equal to NW	–	–	83.3%	
	OW	12	33.2 (16.7)	100	25.6 (2)							
	NW	18	26.2 (6.7)	100	20.2 (1)							
Lasselin et al. (2016)	OB–	29	39.4 (10.5)	62	40.7 (3.7)	IED ¹⁴	–	–	–	OB–HighCR poor than OB–LowCR; NW	83.3%	
	LowCR ¹²	37	37.9 (9)	89	42 (3.8)							
	OB–	20	38.9 (10.1)	90	22 (3)							
	HighCR ¹³											
Loeber et al. (2012)	OB	20	47.9 (12.5)	65	38.8 (6.3)	Go/No-Go Task [food cue] Dot Probe Task (food cue)	–	OB equal to NW	–	–	75.0%	
	NW	20	44.9 (11.7)	60	22.6 (1.1)							

(Continued)

TABLE 3 | Continued

Study	Participants					Cognitive task	Method			EF Domain	
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)		Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	Quality of the study
Mole et al. (2015)	OB	30	44.06 (9.7)	37	32.72 (3.41)	Delay Discounting Task Stop-Signal Task Information Sampling Task	OB poor than NW	OB equal to NW	–	–	83.3%
	NW	30	43.59 (10.01)	37	24.11 (2.89)						
Navas et al. (2016)	OB	20	32.15 (5.96)	55	35.5 (2.6)	The Ehel of Fortune Task IGT	OB poor than OW; NW	–	–	–	83.3%
	OW	21	35 (6.31)	52	27.34 (1.59)						
	NW	38	33.18 (6.59)	58	22.21 (1.70)						
Perpiñá et al. (2017)	OB	27	47.78 (11.46)	85.2	43.92 (10.04)	WCST IGT	OB poor than NW	–	–	OB poor than NW	75.0%
	NW	39	31.9 (13.54)	76.9	23.21 (3.48)						
Pignatti et al. (2006)	OB	34	43.40 (8.13)	42	42.17 (6)	IGT	OB poor than NW	–	–	–	66.7%
	NW	20	46.65 (16.33)	50	22.16 (1.83)						
Restivo et al. (2017)	OB- Bar ¹⁵	25	43.9 (10.7)	92	44.7 (2.9)	COWAT SCWT WCST Color Trail Test PASAT ¹⁷	–	OB- Bar; OB- BarDDM poor than NW	OB- Bar; OB- BarDDM poor than NW	OB- Bar; OB- BarDDM poor than NW	100.0%
	OB- Bar ¹⁵	21	43.2 (10.9)	90	43.7 (4.8)						
	OB- BarDDM ¹⁶	20	43.8 (11)	90	22.4 (2)						
	NW										
Schiff et al. (2016)	OB	23	36.2 (9.5)	78	36.2 (5.7)	Temporal Discounting Task TMT FAB ¹⁸ Simple RT Task Choice RT Task Sterburg Task Simon Task	OB equal to NW	OB equal to NW	OB equal to NW	OB equal to NW	83.3%
	NW	23	33.8 (8.9)	78	22.4 (2.2)						
Spitoni et al. (2017)	OB	24	49.8 (13.66)	79	41.1 (8.03)	BADS ¹⁹ -Rule shift Cards Hayling Sentence Completion Task	–	OB poor than NW	–	–	91.7%
	NW	37	35.7 (11.2)	65	22.5 (3.01)						
Stanek et al. (2013)	OB	152	43.45 (11.28)	84	45.23 (6.91)	Digit Span Switching of Attention Task Verbal Interferences Maze Test	OB poor than NW	OB poor than NW	OB poor than NW	OB poor than NW	83.3%
	NW	580	47.66 (18)	55	25.84 (4.97)						
Stingl et al. (2012)	OB	34	36.5 (9.5)	70	30.4 (3.2)	N-Back Visual Task (food cue)	–	–	OB poor than NW	–	75.0%
	NW	34	38.4 (11)	70	22 (2.1)						

(Continued)

TABLE 3 | Continued

Study	Participants					Method			EF Domain		
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)	Cognitive task	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	Quality of the study
Van der Oord et al. (2018)	OB	39	42.82 (13.23)	82.1	39.7 (5.31)	Stop-Signal Task	OB equal to	OW equal to	OW equal to	–	75.0%
	NW	25	44.9 (15.32)	72	22.94 (1.43)	IGT Chessboard Working Memory Task	NW	NW	NW	–	
Voon et al. (2014)	OB	30	42.97 (8.59)	–	32.72 (3.41)	Premature	–	OB equal to	–	–	58.3%
	NW	30	43.59 (10.01)	–	24.11 (2.89)	Responding Task	–	NW	–	–	
STUDY ON ADOLESCENTS											
Alarcón et al. (2016)	OB	18	14.4 (0.4)	33	%Score	WS-WM ²⁰	–	–	OB poor than	–	75.0%
	OW	46	13.8 (0.2)	46	96.9 (0.3)				OW; NW	–	
	NW	88	14.2 (0.1)	45	90 (0.4) 58.9 (1.8)					–	
Bauer and Manning (2016)	OW	74	15.59 (1.30)	100	%Score	Visual Working	–	–	OW poor than	–	75.0%
	NW	84	15.57 (1.24)	100	>85° <85°	Memory Task	–	–	NW	–	
Calvo et al. (2014)	OB	30	21.21 (2.45)	60	36.36 (6.17)	Go/No-Go Task	–	OB poor than	OB poor than	–	75.0%
	NW	32	21.06 (2.32)	53.1	21.66 (1.78)	Running Memory Continuous Performance Task Standard Continuous Performance Task	–	NW	NW	–	
Delgado-Rico et al. (2012)	OW	42	14.19 (1.38)	67	29.15 (4.51)	SCWT	–	OW equal to	–	OW equal to	66.7%
	NW	21	14.14 (1.46)	48	19.84 (2.64)	(Stroop- Switching Performance)	–	NW	–	NW	
Fields et al. (2013)	OB	21	14.86 (0.85)	52	>95°	Delay Discounting	OB; OW poor	OB equal to	–	–	83.3%
	OW	20	15.2 (0.67)	55	85° - 95°	Task	than NW	OW equal to	–	–	
	NW	20	15 (0.86)	60	5° - 85°	Go/No-Go Task Conner's Continuous Performance Test	–	NW	–	–	
Galioto Wiedemann et al. (2014)	OB	36	21.2 (2.9)	61.1	36.4 (5.7)	Go/No-Go Task	–	OB poor than	OB poor than	–	75.0%
	NW	36	20.7 (2)	50	22 (1.7)	Running Memory Continuous Performance Task	–	NW	NW	–	
Kittel et al. (2017)	OB	22	14.82 (2.63)	82	%score	IGT	OB equal to	OB; OB-Bed	–	OB equal to	75.0%
	OB-Bed	22	14.91 (2.22)	82	98.91 (2.3)	SCWT	OB-Bed	poor than NW	–	OB-Bed	
	NW	22	15.23 (2.39)	82	99.16 (0.57) 58.91 (24.03)		equal to NW	–	–	equal to NW	

(Continued)

TABLE 3 | Continued

Study	Participants					Cognitive task	Method			EF Domain	
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)		Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	Quality of the study
Maayan et al. (2011)	OB	54	17.5 (1.59)	63.6	39.86 (9.46)	SCWT	–	OB poor than	OB poor than	OB poor than	91.7%
	NW	37	17.32 (1.59)	56.8	21.67 (2.49)	TMT COWAT WRAML-WM ²¹		NW	NW	NW	
Moreno-López et al. (2012)	OW	36	14.22 (1.4)	72	28.53 (4.97)	SCWT	–	OB equal to	–	–	75.0%
	NW	16	14.13 (.136)	56	20.26 (2.8)			NW			
Nederkoorn et al. (2006)	OB–Bed	15	13.7	67	33 (4.3)	Stop–Signal Task	OB poor than	OB poor than	–	–	66.7%
	OB–	15	13.9	60	33.5 (4.4)	Door Opening Task	NW	NW			
	NBed NW	31	13.7	61	19.3 (2.0)						
Qavam et al. (2015)	OB	40	[15–18]	0	%Score	TOL ²²	OB poor than	–	–	–	66.7%
	OW	40		0	>95°		OW; NW. OW				
	NW	40		0	85°–95° 5°–85°		poor than NW				
Sellaro and Colzato (2017) [1]	OW	17	23.4 (0.8)	75	27.7 (0.6)	Stop–Signal Task	–	OW poor than	–	–	91.7%
	NW	22	21.2 (0.6)	77	21.9 (0.4)			NW			
Sellaro and Colzato (2017) [2]	OW	19	22.9 (1)	58	28.7 (0.6)	Simon Task	–	OW poor than	–	–	91.7%
	NW	24	20.5 (0.5)	79	21.7 (0.4)			NW			
Steenbergen and Colzato (2017)	OW	26	20.27 (0.44)	73	27.58 (0.41)	Switching of Attention Task	–	–	–	OW poor than	83.3%
	NW	26	20.36 (0.41)	81	21.67 (0.25)					NW	
Sweat et al. (2017)	OB	108	19.6 (1.54)	63	35.57 (4.97)	SWCT	–	OB equal to	–	OB equal to	83.3%
	NW	54	19.39 (1.52)	53.7	21.45 (1.87)	TMT TOL		NW		NW	
Vantieghem et al. (2018)	OB	62	15.8 (1.8)	71	39.9 (8.19)	SCWT	–	OB poor than	–	–	83.3%
	NW	30	16 (1.1)	47	20.95 (2.11)			NW			
Verbeken et al. (2014)	OW	64	13.59 (1.62)	54.2	Adjusted BMI (%)	HDT ²³	OW poor than	–	–	–	58,3%
	NW	66	12.42 (1.16)		145.37 (16.27) 102.56 (8.99)		NW				
Verdejo-García et al. (2010)	OW	27	14.3 (1.2)	41	31.58 (7.08)	SCWT	OW poor than	OW poor than	–	OW poor than	83.3%
	NW	34	15.29 (0.91)	38	21.01 (1.97)	Five–Digit Test TMT IGT	NW	NW		NW	
Weller et al. (2008) [1]	OB	29	19.6 (2.9)	100	38.4 (6.6)	Delay Discounting Task	OB poor than	–	–	–	75.0%
	NW	26	20 (2.6)	100	21.9 (2.3)		NW				
Weller et al. (2008) [2]	OB	19	19.2 (1.3)	0	35.4 (4.8)	Delay Discounting Task	OB equal to	–	–	–	75.0%
	NW	21	19.4 (1.5)	0	22.3 (1.2)		NW				

(Continued)

TABLE 3 | Continued

Study	Participants					Method			EF Domain		
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)	Cognitive task	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	Quality of the study
Wu et al. (2016)	OB NW	19 20	21.3 (2.6)	74 70	33 (2.9) 22.2 (2.2)	SCWT TMT Verbal Fluency Digit Span	–	OB equal to NW	OB poor than NW	OB poor than NW	58.3%
Yau et al. (2014)	OB NW	30 30	17.64 (1.62) 17.22 (1.55)	57 63	35.47 (5.88) 21.12 (2.18)	TMT WCST SCWT COWAT	–	OB equal to NW	OB poor than NW	OB poor than NW	91.7%
STUDY ON YOUNG ADULTS											
Coppin et al. (2014)	OB OW NW	17 16 16	25.17 (4.39) 24.94 (4.55) 24.25 (4.25)	53 44 56	36.02 (6.54) 27.63 (1.49) 22.43 (1.45)	CCPT ²⁴	–	–	OB, OW poor than NW	–	91.7%
Yadava and Sharma (2014)	UW NW NW2 OW OB	39 50 58 58 25	26.9 [20–42]	100	<18.5 18.5–22.9 23–24.9 25–29.9 30	Digit Symbol Test SCWT Ascending Digit Task	–	OB poor than NW	OB poor than NW	OB poor than NW	75.0%
STUDY ON CHILDS											
Blanco-Gómez et al. (2015)	OB OW NW	39 149 316	[6–10]	49 53 50	%Score >97 95–97 <95	Children's Color Traits Test (1,2) Five Digit Test	–	OB poor than OW; NW	–	OB poor than OW; NW	83.3%
Bozkurt et al. (2017)	OB NW	92 55	11.85 (2.43) 11.9 (2.96)	56 54	29.73 (2.33) 21.07 (1.81)	FTT ²⁵ SDC ²⁶ SCWT SAT ²⁷ CPT ²⁸	OB poor than NW	OB poor than NW	OB poor than NW	OB poor than NW	83.3%
Gentier et al. (2013)	OB NW	19 19	9.8 (1.5) 9.9 (1.5)	47 47	Cut-off (Cole et al. 2000) 21.62 (3.51) 16.48 (1.76)	Four Choice Reaction Time Task	OB poor than NW	–	–	–	58.3%
Goldschmidt et al. (2018)	OW– LC ²⁹ OW–C ³⁰ NW–C ³¹	26 34 15	10.2 (0.9) 10.8 (1.1) 10.4 (1.1)	61 56 60	z-score 2.08 (0.47) 2.02 (0.47)	Flanker Task DCCST ³² IGT TOL List Sorting	OB–LC; OB–C poor than NW–C	OB–LC equal to OB–C equal to NW–C	OB–LC; OB–C poor than NW–C	–	75.0%

(Continued)

TABLE 3 | Continued

Study	Participants					Method			EF Domain		
	Group	N	Age M (SD)	Sex (% female)	BMI M (SD)	Cognitive task	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	Quality of the study
Kamijo et al. (2012a)	OB	30	9 (0.5)	100	%score	Go/No-Go Task	-	OB poor than NW	-	-	75.0%
	OW	26	8.7 (0.6)	100	>95°						
	NW	70	8.9 (0.6)	100	>85° >5°						
Kamijo et al. (2012b)	OB	37	9 (0.5)	51	%score	Go/No-Go Task	-	OB poor than NW	-	-	75.0%
	NW	37	8.9 (0.5)	51	>95° 5°-85°						
Kamijo et al. (2012c)	OB	37	8.9 (0.6)	54	%score	Flanker Task	-	OB poor than NW	-	-	75.0%
	NW	37	8.8 (0.6)	54	98 (1.4) 56.8 (19.9)						
Pearce et al. (2018) [1]	OB	41	13.3 (3.4)	54	%Score	BART ³³	OB equal to NW	-	-	-	75.0%
	NW	37	13.1 (2.7)	30	98.8 (1.2) 58.3 (26.1)						
Pearce et al. (2018) [2]	OB	29	11.4 (2.6)	48	%Score	Stop-Signal Task N-Back Task	-	OB equal to NW	OB equal to NW	-	75.0%
	NW	30	11.9 (2.6)	40	98.5 (1.3) 60.7 (25.4)						
Reyes et al. (2015)	OW	93	10.2 (1)	44	z-score	SCWT Go/No-Go Task	-	WB poor than NW	-	-	58.3%
	NW	92	10.3 (0.2)	46	1.9 (0.6) 0.1 (0.5)						
Skoranski et al. (2013)	OB	28	12.8 (2.4)	79	%Score	Arrow Task	-	OB poor than NW	-	-	58.3%
	NW	32	12.8 (2.5)	47	>85° 5°-85°						
Tsai et al. (2016)	OB	26	(month)	31	%Score	Posner Paradigm Task	-	OB poor than NW	-	-	58.3%
	NW	26	114.58 (3.69) 113.73 (3.85)	31	>95° 5°-85°						
Wu et al. (2017)	OB	44	12.38 (1.22)	32	>30	Digit Span Memory Task (digits; digit-food cue; digit-cartoon)	-	-	OW poor than NW	-	66.7%
	NW	23	11.78 (1)	26	25-30						
	OW	92	11.93 (0.92)	56	<25						

1OB, Obese; 2NW, Normal-Weight; 3TMT, Trail Making Test; 4SCWT, Stroop Color-Word Task; 5WCST, Wisconsin Card Sorting Test; 6IGT, Iowa Gambling Task; 7OB-BED, Obese with Binge Eating Disorder; 8AN, Anorexia Nervosa; 9ExOB, Normal-weight people who were previously obese; 10OW, Overweight; 11COWAT, Controlled Oral Word Association Task; 12OB-LowCR, Obese with low sensitivity to C-reactive protein; 13OB-HighCR, Obese with high sensitivity to C-reactive protein; 14IED, Intra/Extra-dimensional set shift test; 15OB-Bar, Obese and on the waiting list for bariatric intervention; 16OB-BarDDM, Obese and on the waiting list for bariatric intervention with Major Depressive Disorder; 17PASAT, Paced Auditory Serial Attention Test; 18FAB, Frontal Assessment Battery; 19BADS, Behavioral Assessment of the Dysexecutive Syndrome; 20WS-WM, Working Memory Task of Wechsler Scale; 21WRAML-WM, Wide-Range Assessment of Memory and Learning-Working Memory; 22TOL, Tower of London; 23HDT, Hungry Donkey Task; 24CCPT, Conditioned Cue Preference Test; 25FTT, Finger-Tapping Test; 26SDC, Symbol Digit Coding; 27SAT, Shifting Attention Test; 28CPT, Continuous Performance Test; 29OW-LC, Overweight with high loss of control; 30OW-C, Overweight with low loss of control; 31NW-C, Normal-weight with low loss of control; 32DCCST, Dimensional Change Card Sort task; 33BART, Balloon Analog Risk Task.

TABLE 4 | Longitudinal Studies.

Study	Participants					Method			EF Domain				Quality of the study
	Group	N	Age M (SD)	Sex (% female) ^a	BMI ^b M(SD)	Treatment	Cognitive task	Program /follow-up	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	
*Allom et al. (2018)	OB-CRT ¹ OB-C ²	42 38	41.39 (7.85)	86	39.76 (7.53)	Cognitive Remediation Therapy	WCST ³ TMT ⁴	3 months	OB-CRT ↑ OB-C =	-	.	OB-CRT ↑ OB-C =	83.3%
Alosco et al. (2014a)	OB-AD ⁵ OB-NAD ⁶	14 80	40 (11.42) 45.1 (10.99)	21.4 15	T1.45.17 (5.02) T2.37.85 (5.43) T1.46.07 (5.33) T2.38.06 (4.86)	Bariatric Surgery [Alzheimer History]	TMT Maze Task	12 weeks	OB- AD=OB- NAD=	-	.	OB- AD=OB- NAD=	75.0%
Alosco et al. (2014b)	OB-Bar ⁷ OB-C	63 23	42.29 (11.42) 41.13 (12.55)	90.5 95.7	T1.46.5 (5.26) T2.31.34 (6.42) T1.40.9 (5.24) T2.40.9 (5.64)	Bariatric Surgery	Digit Span Switching Attention Task Maze Task	24 months	OB-Bar ↑ OB-C =	-	OB-Bar ↑ OB-C =	OB-Bar ↑ OB-C =	91.7%
Alosco et al. (2014c)	OB-Bar	78	43.5 (10.59)	82.1	T1.46.63 (5.28) T2.30.51 (5.39)	Bariatric Surgery	Switching of Attention Task Maze Task	12 months	OB-Bar ↑	-	-	OB-Bar ↑	66.7%
Alosco et al. (2014d)	OB-Bar	50	44.08 (10.76)	92	T1.46.61 (5.27) T2.32.35 (6.57) T3.33.02 (6.27)	Bariatric Surgery	Digit Span Switching of Attention Task Verbal Interference Maze Task	I. 36 months II. 48 months (LD) ⁸	I.OB-Bar ↑ II. OB-Bar ↑ OB-Bar ↑	I.OB-Bar ↑ II. OB-Bar ↑	I.OB-Bar ↑ II. OB-Bar ↑	I.OB-Bar ↑ II. OB-Bar ↑	83.3%
Alosco et al. (2015)	OB-Bar	84	43.86 (10.39)	83.3	T1.46.88 (6.08) T2.30.05 (5.39)	Bariatric Surgery	Digit Span Switching of Attention Task Verbal Interference Task	12 months	-	OB-Bar ↑	OB-Bar ↑	OB-Bar ↑	83.3%
*Augustijn et al. (2018)	OB	T1.32 T2.30	9.6 (1.1)	T1.56 T2.60	Z scores T1. 2.7 (0.3) T2.2.0 (0.4)	Weight Loss Program	CANTAB ⁹	6–10 months	OW↑	OW↑	OW↑	OW=	75.0%
Bryan and Tiggemann (2001)	OB-WL ¹⁰ OB-C	42 21	48.9 (8.2) 50.9 (7.3)	100 100	T1.34.1 (4.3) T1.35.2 (4.8)	Weight Loss Program	TMT WCST Self-Ordered Piniting Task Initial Letter Fluency Excluded Letter Fluency Digit Span	12 weeks	-	OB-WL ↑ OB-C=	OB-WL =OB-C=	OB-WL =OB-C=	91.7%
*Dassen et al. (2018b)	OW-WMT ¹¹ OW-C	T1.51 T2.34 T1.40 T2.36	47.97 (10.69)	74.7	T1.30.96 (3.64) T2.29.95 (3.46) T1.30.49 (3.97) T2.30.17 (4.14)	Working Memory Training	2-Back Task	25 session	-	-	OW-WMT↑	-	66.7%

(Continued)

TABLE 4 | Continued

Study	Participants					Method			EF Domain				Quality of the study
	Group	N	Age M (SD)	Sex (% female) ^a	BMI ^b M(SD)	Treatment	Cognitive task	Program /follow-up	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	
Davis et al. (2007a)	OW-HE ¹² OW-LE ¹³ OW-NE ¹⁴	32 33 29	9.2 (0.84)	60	z-score 2.1 (0.4)	Weight Loss Program: Aerobic Exercise	CAS ¹⁵ : Planning Subscales for EF	15 weeks	OW-HE↑	-	-	-	66.7%
Davis et al. (2011)	OW-HE OW-LE OW-NE	56 55 60	9.3 (1.0)	56	z-score 2.1 (0.4)	Weight Loss Program: Aerobic exercise	CAS: Planning Subscales for EF	13 weeks	OW-HE↑	-	-	-	75.0%
Deckers et al. (2017)	OB NW	T1.545 T2.190 T1.1262 T2.834	T1.58 (15) T2.48.9 (16.2) T1.48.9 (16.2) T2.46.7 (14.9)	I.58 II.59 I.46 II.43	T1.31.2 (3.9) T2.28.7 (2.4) T1.24.9 (2.5) T2.24.8 (2.4)	-	Concept Shifting Test	6 years 12 years	OB= NW=	-	-	OB=NW=	83.3%
Demos et al. (2017)	OB-WL NW	37 30	46.95 (7.9) 43.97 (8.9)	100 100	T1.33.5 (3.9) T1.22.7 (1.8)	Weight Loss Program	Food Choice Decision Making Task	12-16 weeks	OB-WL↑	-	-	-	83.3%
Galioto et al. (2015)	OB-Bar	72	43.55 (10.21)	81.7	T1.46.32 (5.51) T2.30.18 (5.25)	Bariatric Surgery	Digit Span Switching of Attention Task Verbal Interference Verbal Fluency	12 months	OB-Bar↑	OB-Bar↑	OB-Bar↑	OB-Bar↑	91.7%
*Galioto et al. (2016)	OB	23	50.35 (15.11)	68	44.21 (8.82)	Weight Loss Program	Dot Counting Task N-Back Task Set Shifting Task Unstructured Task Flanker Task	8 weeks	-	OB↑	OB=	OB↑	100.0%
*Kulendran et al. (2014)	OB-WL	53	14.28 (1.15)	60	T1.33.75 (7.9)	Weight Loss Program	Stop-Signal Task Delay Discounting Task	2-8 weeks	-	OB-WL↑	-	-	83.3%
*Kulendran et al. (2017)	OB-Bar	45	43.42 (13.06)	31	T1.44.25 (6.34) T2.35.51 (7.08)	Bariatric Surgery	Stop-Signal Task (food-cue) Temporal Discounting Task	6 months	-	OB-Bar↑	-	-	75.0%

(Continued)

TABLE 4 | Continued

Study	Participants					Method			EF Domain				Quality of the study
	Group	N	Age M (SD)	Sex (% female) ^a	BMI ^b M(SD)	Treatment	Cognitive task	Program /follow-up	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	
*Pauli-Pott et al. (2010)	OW	111	11.1 (2.0)	57	95 ^o percentile	Weight Loss Program	Go/No-Go Task Interference Task	1 year	-	OW↑	-	-	91.7%
Pearce et al. (2017)	OB-Bar	10	17 (1.37)	60	T1.47.18 (6.98)	Bariatric Surgery	Verbal N-Back Test Ballon analog risk task	4 months	OB-Bar=	-	OB-Bar=	(DM area shows a reduction of activation in OB-Bar after the surgery)	83.3%
	OB-C	14	16.42 (1.35)	71	T1.45.32 (8.19)				OB-C=		OB-C=		
	NW	12	16.51 (1.27)	50	T1.21.57 (2.59)				NW=		NW=		
Raman et al. (2018)	OB-CRT	42	40.6 (2.4)	86	39.2 (7.4)	Computerized Cognitive Remediation Therapy	WCST TMT	8 weeks	-	-	-	OB-CRT ↑	83.3%
	OB-C	38	42.2 (8.8)		40.3 (7.8)			3 months				OB-C =	
*Spitznagel et al. (2013)	OB-Bar	84	44.75 (9.99)	79.8	T1.46.13 (5.80) T2.37.46 (4.99) T3.31.07 (6.44)	Bariatric Surgery	Switching of Attention Task Digit Span Maze Task	I.12 weeks II. 12 months	I.OB-Bar= II.OB-Bar↑	-	I.OB-Bar= II.OB-Bar↑	I.OB-Bar= II.OB-Bar↑	75.0%
*Spitznagel et al. (2014)	OB-Bar	55	45 (10.28)	87.3	T1.45.11 (5.11) T2.37.23 (4.76) T3.31.69 (5.84)	Bariatric Surgery	Digit Span Switching of Attention Verbal Interference Verbal Fluency Maze Task	12 weeks 36 months	OB-Bar↑	OB-Bar =	OB-Bar↑	OB-Bar =	83.3%
*Stinson et al. (2018)	OW	46	37.2 (10.2)	24	28.3 (6.7)	-	IGT WCST SCWT	32 ± 25 months	OW=	OW↓	-	OW=	100.0%
Vantieghem et al. (2018)	OB-WL	62	15.8 (1.8)	71	T1.39.9 (8.19)	Weight Loss Program	SCWT	30 weeks	-	OB-WL↑	-	-	83.3%
	NW	30	16 (1.1)	47	T2.32.21 (7.14) 20.95 (2.11)								
Verbeken et al. (2013)	OB-EFT ¹⁸	22	11.50 (1.60)	50	Adjusted BMI	Executive Function Training	Corsi Block-Tapping Task Stop-Signal Task	Post-Test	OB-EFT ↑	-	-	-	58.3%
	OB-C	22	11.41 (1.93)	41	T1. 131.58 (21.70)			8 weeks	OB-C=				
					T1. 132.91 (15.98)			12 weeks					
Witbracht et al. (2012)	OB	29	32.7 (9.2)	100	32 (2.6)	Weight Loss Program	IGT	12 weeks	OB↑	-	-	-	83.3%

(Continued)

TABLE 4 | Continued

Study	Group	N	Participants		BMI ^b M(SD)	Treatment	Method		EF Domain				Quality of the study
			Age M (SD)	Sex (% female) ^a			Cognitive task	Program /follow-up	Global EF/other EFs	Inhibition	Updating	Cognitive flexibility	
Xie et al. (2017)	OB-WL	30	15.07 (0.83)	27	T1:32.83 (3.84)	Weight Loss Program	Flanker Task	4 weeks	-	OB-WL↑	-	-	75.0%
	OB-C	28	15.18 (0.39)	36	T2:29.19 (3.52) T1:30.90 (1.95) T2:30.47 (2.13)					OB-C=	-	-	
*Xu et al. (2017)	OB-WL	31	18.2 (3.2)	39	34.4 (4.8)	Weight Loss Program	SCWT	4 weeks program	-	OB-WL ↑	-	-	83.3%

^aEF-predicted weight loss. ^bPercentage of females. ^cBody Mass Index. ↑, Better performance after treatment; ↓, Worse performance after treatment. No differences. ¹OB-CRT, Obese and in Cognitive Remediation Therapy Treatment; ²OB-C, Obese-Control (No treatment group); ³WCST, Wisconsin Card Sorting Test; ⁴TMT, Trail Making Test; ⁵OB-AD, Obese with history of Alzheimer's; ⁶OB-NAD, Obese with no history of Alzheimer's; ⁷OB-Bar, Obese and subjected to bariatric surgery; ⁸LD, Loss Data; ⁹CANTAB, Cambridge Neuropsychological Test Automated Battery; ¹⁰OB-WL, Obese and subjected to a weight-loss programme; ¹¹OB-WMT, Obese and subjected to Working Memory Training; ¹²OW-HE, Overweight and subjected to high-exercise training; ¹³OW-LW, Overweight and subjected to low-exercise training; ¹⁴OW-NE, Overweight with no exercise training; ¹⁵CAS, Cognitive Assessment System; ¹⁶SCWT, Stroop Color-Word Task; ¹⁷IGT, Iowa Gambling Task (for decision-making); ¹⁸OB-EFT, Obese and subjected to Executive Function Training.

2017). Furthermore, Weller et al. (2008) performed two different analyses to examine samples of males and females independently.

All the studies used BMI and the related WHO classification to assign participants to different overweight groups. For children and adolescents, the guidelines for using percentiles recommended by WHO or Center for Disease Control and Prevention (CDC) were employed (Flegal et al., 2002; de Onis et al., 2007), except in two studies (Reyes et al., 2015; Goldschmidt et al., 2018) where z scores for CDC classification were used (Harrington et al., 2013).

Studies focused mainly on differences in executive functioning between individuals with obesity and normal-weight; thirteen analyzed differences between participants with normal-weight and overweight; thirteen studies investigated differences in performance between participants with obesity, overweight, and normal-weight (see Table 3).

Most of the studies reported a significant difference between the groups in executive functioning, confirming the relationship between excessive body weight and executive dysfunctions. Only thirteen studies reported no differences (Gonzales et al., 2010; Ariza et al., 2012; Delgado-Rico et al., 2012; Hendrick et al., 2012; Loeber et al., 2012; Moreno-López et al., 2012; Frank et al., 2014; Voon et al., 2014; Bongers et al., 2015; Reyes et al., 2015; Schiff et al., 2016; Sweat et al., 2017; Van der Oord et al., 2018).

EXECUTIVE FUNCTIONS IN CROSS-SECTIONAL STUDIES

Cognitive Flexibility


The tasks most commonly used to assess cognitive flexibility were the Wisconsin Card Sorting Test (WCST) (Milner, 1963), the Trail Making Test (TMT, AB) (Reitan, 1958) and the Switching of Attention Task (Rogers and Monsell, 1995) (see Table 3).

Twenty-seven studies assessed the differences between groups on cognitive flexibility (see Table 3); only eight of them found no differences in cognitive flexibility between normal-weight and overweight/obese groups (Gonzales et al., 2010; Ariza et al., 2012; Delgado-Rico et al., 2012; Catoira et al., 2016; Schiff et al., 2016; Kittel et al., 2017; Sweat et al., 2017; Dassen et al., 2018a). In general, the results showed greater difficulty in performing tasks involving this function in participants with obesity compared to those with normal-weight. Furthermore, the study by Blanco-Gómez et al. (2015) highlighted a further difference: compared with participants with overweight, participants with obesity showed higher flexibility deficits.

Inhibition

The most common cognitive tasks used to measure inhibitory control were the Stroop Color-Word Task (Stroop, 1935) and the Stop-Signal Task (Lappin and Eriksen, 1966) (see Table 3).

Forty-five studies investigated the relationship between inhibitory control and excessive body weight (see Table 3). Of these, seventeen studies reported no differences between the groups (Gonzales et al., 2010; Ariza et al., 2012; Delgado-Rico et al., 2012; Hendrick et al., 2012; Loeber et al., 2012; Moreno-López et al., 2012; Stingl et al., 2012; Fields et al., 2013; Voon et al., 2014; Yau et al., 2014; Bongers et al., 2015; Schiff et al.,



2016; Wu et al., 2016; Goldschmidt et al., 2018; Pearce et al., 2018; Van der Oord et al., 2018). The remaining studies reported lower inhibitory control in obese than in normal-weight participants.

Working Memory

The Digit Span Test (in particular the Backwards version) (Reynolds, 1997), and the N-Back Test (Kane et al., 2007) were

used in various versions (see **Table 3**) to investigate differences in working memory performance.

Of the twenty-four studies that analyzed the relation between overweight/obesity and working memory (see **Table 2**), six observed no differences between groups (Gonzales et al., 2010; Ariza et al., 2012; Frank et al., 2014; Restivo et al., 2017; Pearce et al., 2018; Van der Oord et al., 2018). The remaining studies found that participants with overweight/obesity performed worse




FIGURE 2 | Percentage of the studies and quality levels for each point of tool assessment.

than normal-weight participants on working memory tasks; moreover, obese participants performed worse than participants with overweight (Coppin et al., 2014; Alarcón et al., 2016).

Decision-Making, Planning, and Problem-Solving

The tasks used to assess decision-making, planning and problem-solving were the IGT (Bechara et al., 2005) and the Delay Discounting Task (Richards et al., 1999) (see **Table 3**).

Twenty-six studies (see **Table 3**) investigated differences in performances between groups on tasks involving complex EFs such as decision-making, planning and problem-solving. Among these studies, only six (Bongers et al., 2015; Mole et al., 2015; Schiff et al., 2016; Kittel et al., 2017; Pearce et al., 2018; Van der Oord et al., 2018), individuals with obesity performed worse than those with normal-weight on decision making, planning and risk-taking. Furthermore, Schiff et al. (2016), despite of they observed no clear between differences in decision-making, found that the group with obesity responded differently in terms of gratification mechanisms connected with food (as measured by the Temporal Discounting Task), in fact, they showed more sensitivity to reward stimuli than normal-weight group. Weller et al. (2008), using the Delay Discounting task, found that women affected by obesity, compared to women with normal-weight, preferred an immediate reward than a major one after some time. This difference was not observed in men (Weller et al., 2008).

DISCUSSION

The analysis of the cross-sectional studies confirmed the existence of a relationship between overweight/obesity and EFs,

even if it did not indicate the direction of this relationship. Many types of cognitive tasks were used to investigate executive functioning, but, despite this heterogeneity, the results were consistent. However, the very different demands of the tasks used did not allow determining whether one single EF is more closely involved than the others in the relationship with overweight/obesity, though the most analyzed EF related to excessive body weight is Inhibition (see **Table 3**). The studies that failed to confirm a relationship between EFs and overweight/obesity used a small sample size (Hendrick et al., 2012; Schiff et al., 2016), or a high number of cognitive tasks (Gonzales et al., 2010).

The present systematic review included studies that take into account people with different ages considering from children to the elderly. This choice was aimed to investigate whether the relationship between EFs and overweight/obesity presents similar characteristics, regardless of the age of the participants. The results of the review confirmed the relationship between EFs and overweight both in studies examining adults and young adults (Gunstad et al., 2007; Fagundo et al., 2012; Coppin et al., 2014) and in those that looked at children (Yadava and Sharma, 2014; Bozkurt et al., 2017) and adolescents (Nederkoorn et al., 2006; Galioto Wiedemann et al., 2014). These results prevent us from making inferences about the causality of this relationship over a lifespan but highlight the existence of a negative relationship between executive performances and overweight, regardless of the age considered.

Many studies tried to control for certain variables (gender, age, and education) that might influence executive performance, by matching samples or controlling the effects of these variables through statistical analysis (Gunstad et al., 2007;

Deckers et al., 2017; Kittel et al., 2017; Perpiñá et al., 2017). This methodological aspect highlighted the existence of some dimensions (e.g., demographical variables as gender or educational level) that might influence the relation between body weight and EFs; therefore, considering these variables can contribute to further strengthen the results (Kittel et al., 2017).

Generally, the analyzed studies used suitable inclusion criteria that allow excluding individuals with chronic medical conditions, psychological diseases or eating disorders, in order to avoid an effect of these dimensions on the observed results (Fagundo et al., 2012; Galioto et al., 2013; Galioto Wiedemann et al., 2014). Moreover, in some studies, physiological differences between participants with normal-weight and overweight/obesity were reported. In particular, participants with severe obesity showed worse values, in blood pressure, cholesterol levels, insulin resistance (Maayan et al., 2011; Perpiñá et al., 2017) and levels of glycolic metabolism activation although, in the absence of pathological medical conditions in line with well-known results (Heymsfield and Wadden, 2017).

Both psychopathological and physiological aspects related to obesity, and specifically with severe obesity, have an impact on the executive functioning and consequently with the performances in executive tasks; therefore these variables should be controlled in further studies.

Although the cross-sectional studies showed no clear direction in the overweight–executive functioning relationship, many of the authors have advanced various hypotheses (Gonzales et al., 2010; Galioto Wiedemann et al., 2014). For example, Kamijo et al. (2012a) hypothesized that ineffective inhibitory control of the prefrontal cortex would cause excessive consumption of calories that is directly associated with an increase in body fat. Moreover, other authors considered also the dopaminergic mechanism involved in executive processing as related to weight variations (Arnsten and Li, 2005). Neuroimaging studies of individuals with obesity have shown an association between the hypoactivation of dopaminergic D2-receptors and a decrease in neural metabolism in the areas most involved in executive functioning (Volkow et al., 2011). Furthermore, dopamine is also implicated in the reward system (Volkow et al., 2011; Smith and Robbins, 2013). This neural system resulted impaired in individuals with excessive body weight, and alterations of this system could influence the approach to food in terms of favoring the consumption of high-calorie foods to achieve higher gratification (Schiff et al., 2016). All these findings could support theoretical models on the genesis of obesity (Davis et al., 2007b; Smith and Robbins, 2013) that view changes in executive functioning as one of the leading causes of weight gain. The hypothesis of executive dysfunctions as a cause of inappropriate eating behavior could partially support the theoretical model of Food Addiction, in which the excessive consumption of food is characterized by behavioral aspects similar to those defining other substance addiction diseases (Wang et al., 2004; Smith and Robbins, 2013).

Nevertheless, other authors viewed executive deficits as a consequence of obesity, recognizing it as a cause of neurophysiological and metabolic diseases, such as changes in

insulin sensitivity (Gonzales et al., 2010), inflammatory processes as a result of body fat accumulation (Lasselín et al., 2016), and changes in cerebrovascular blood flow (Verdejo-García et al., 2010; Qavam et al., 2015). These alterations could be the cause of structural changes (e.g., a reduction of the orbitofrontal cortex) (Cohen et al., 2011) or functional changes (e.g., reduced functional connectivity of executive networks) (Tsai et al., 2016) in the cerebral areas involved in executive functioning. This vision seems to be in line with the Neuroinflammation Model (Perry, 2004) in which high BMI appears to result in systemic inflammation, which negatively affects cognitive functions including executive ones (C-reactive protein and interleukin would play an essential role in this process; Bourassa and Sbarra, 2017), and with the model proposed by Sellbom and Gunstad (2012) in which the changing in blood flow and metabolism of the frontal lobes as well as the atrophy of the frontal and temporal lobes would cause an impairment in inhibitory control resulting in an increase in overeating behaviors (Sellbom and Gunstad, 2012).

The consistent results confirming the relationship between EFs and obesity suggests that even a moderate increase in body weight may be associated with a decrease in executive performances (Verdejo-García et al., 2010; Cohen et al., 2011; Sellaro and Colzato, 2017). These views are supported by results obtained comparing groups of participants with normal-weight, overweight and obesity, in which differences in performances also emerged between overweight and obesity conditions (Galioto Wiedemann et al., 2014; Wu et al., 2017).

Another aspect highlighted by the cross-sectional studies is the role of certain psychological variables related to BMI (Catoira et al., 2016; Restivo et al., 2017) that appear to modulate the relationship between EFs and excessive body weight. Indeed, the presence of high levels of anxiety and depression in individuals with obesity, even in the absence of established psychopathologies, appears to result in worse executive performances (Restivo et al., 2017). These findings could be linked to the theoretical model of Emotionally-Driven Eating (Dallman, 2010), which postulated that overeating, related to overweight, is a dysfunctional attempt to regulate emotions in people characterized by a deficit in emotion regulation.

Longitudinal Studies

Our systematic search allows selecting twenty-eight longitudinal studies investigating executive functioning in individuals with overweight or obesity (see **Table 4**). Of these, eighteen examined adult participants (aged more than 30 years), five looked at children (aged <12 years) and five involved adolescents (aged 12–22 years) (see **Table 4**).

All studies used BMI to classify overweight and obesity, although z-scores (Davis et al., 2007a, 2011; Augustijn et al., 2018), percentiles (Pauli-Pott et al., 2010), or adapted BMI scores (Verbeken et al., 2014) were used in studies involving children.

Five studies (Alosco et al., 2014a; Kulendran et al., 2017; Xie et al., 2017; Xu et al., 2017; Stinson et al., 2018) reported having a significantly higher percentage of males than females in their sample.

Twelve studies (Bryan and Tiggemann, 2001; Davis et al., 2007a, 2011; Pauli-Pott et al., 2010; Witbracht et al., 2012; Kulendran et al., 2014; Galioto et al., 2016; Demos et al., 2017; Xie et al., 2017; Xu et al., 2017; Augustijn et al., 2018; Vantieghem et al., 2018) analyzed the effects of non-invasive programmes aimed at weight-loss on the relationship between BMI and EFs: some interventions integrated various modalities of treatment, specifically diet and physical activity (Pauli-Pott et al., 2010; Kulendran et al., 2014; Galioto et al., 2016; Demos et al., 2017; Xie et al., 2017; Xu et al., 2017; Vantieghem et al., 2018); while others focused only on diet programmes (Bryan and Tiggemann, 2001; Witbracht et al., 2012) or physical activity (Davis et al., 2007a, 2011). Furthermore, two studies (Kulendran et al., 2014; Augustijn et al., 2018) provided residential interventions, with treatment lasting from four (Davis et al., 2011) to fifty-2 weeks (Pauli-Pott et al., 2010). In all the studies, at least two measurements were taken: one before and one after the procedure.

Ten studies examined the effects of bariatric surgery on the executive functioning in participants with severe obesity (Spitznagel et al., 2013, 2014; Alosco et al., 2014a,c,d, 2015; Galioto et al., 2015; Kulendran et al., 2017; Pearce et al., 2017). The analysis of EFs was performed before surgery and at follow-up, with time intervals ranging from 12 weeks (Spitznagel et al., 2013, 2014) to 48 months (Alosco et al., 2014d). In some cases, more than one follow-up was carried out (Spitznagel et al., 2013, 2014; Alosco et al., 2014d).

All the studies investigating weight reduction in participants with obesity reported a general improvement in EF performances. Only Pearce et al. (2017) failed to detect any significant changes in performances.

Four studies assessed the effects of cognitive interventions on EFs in obese participants, showed a general improvement in executive performances associated with a reduction in body weight. Specifically, two studies evaluated the benefits of Cognitive Remediation Therapy (Alosco et al., 2014b; Allom et al., 2018), one assessed the impact of an intervention focused on Working Memory (Galioto et al., 2015) and one focused on the effects of a treatment aimed at strengthening cognitive functions in general (Verbeke et al., 2014).

Two further studies analyzed the trend over time of body weight and executive functioning in adults with obesity (Deckers et al., 2017; Stinson et al., 2018) without introducing weight reduction programs and reported inconsistent results. Deckers et al. (2017) found no relationship between weight changes and executive performance, while Stinson et al. (2018) found evidence of the role of EFs, specifically of reduced inhibitory control, in maintaining high body weight.

Eleven studies (Pauli-Pott et al., 2010; Spitznagel et al., 2013, 2014; Kulendran et al., 2014; Galioto et al., 2015, 2016; Xu et al., 2017; Augustijn et al., 2018; Dassen et al., 2018a; Stinson et al., 2018) investigated the predictive role of performance on executive tasks on body weight changes, and observed that appropriate executive functioning predicted a reduction in body weight in participants with obesity or overweight.

EXECUTIVE FUNCTIONS IN LONGITUDINAL STUDIES

Cognitive Flexibility

The tasks most commonly used to assess cognitive flexibility were the WCST, TMT, and Switching of Attention Task (see **Table 4**).

Of the eleven studies that investigated the relationship between cognitive flexibility and obesity (see **Table 4**), six (Bryan and Tiggemann, 2001; Alosco et al., 2014c; Spitznagel et al., 2014; Deckers et al., 2017; Augustijn et al., 2018; Stinson et al., 2018) failed to confirm this relationship. Those that found an association between obesity and executive functioning reported an improvement in performance as a result of weight reduction. Furthermore, negative performance appeared to be associated with less weight reduction over time (Spitznagel et al., 2013; Augustijn et al., 2018).

Inhibition

The Stroop Color-Word Task and Stop-Signal Task were the tests most commonly used to investigate cognitive and motor inhibition (see **Table 4**).

Thirteen studies reported a relationship between BMI and cognitive inhibition (Bryan and Tiggemann, 2001; Pauli-Pott et al., 2010; Alosco et al., 2014d, 2015; Kulendran et al., 2014, 2017; Galioto et al., 2015, 2016; Xie et al., 2017; Xu et al., 2017; Augustijn et al., 2018; Stinson et al., 2018; Vantieghem et al., 2018). Some of these studies showed that inhibition control predicted a reduction of body weight considering both bariatric surgery (Kulendran et al., 2017) and weight loss programs (Pauli-Pott et al., 2010; Kulendran et al., 2014; Galioto et al., 2016; Xu et al., 2017; Augustijn et al., 2018; Stinson et al., 2018). Other studies showed an improvement in the inhibition after bariatric surgery (Alosco et al., 2014d, 2015; Galioto et al., 2015) or weight-loss programs (Bryan and Tiggemann, 2001; Xie et al., 2017; Vantieghem et al., 2018).

Working Memory

The task most often used to investigate working memory was the Digit Span Test (Reynolds, 1997) (see **Table 4**).

Eight studies reported a negative relationship between working memory and body weight (Spitznagel et al., 2013, 2014; Alosco et al., 2014b,d, 2015; Galioto et al., 2015; Augustijn et al., 2018; Dassen et al., 2018b). Indeed some authors found an improvement of the performance in working memory tasks after bariatric surgery (Alosco et al., 2014b,d, 2015; Galioto et al., 2015), while other authors found a predictive role of working memory performance in the outcome of weight reduction programs (Augustijn et al., 2018; Dassen et al., 2018b) or bariatric surgery (Spitznagel et al., 2013, 2014); better performance predicted success of interventions. Conversely, three studies found no relationship between obesity and working memory (Bryan and Tiggemann, 2001; Galioto et al., 2016; Pearce et al., 2017).

Decision-Making

Decision-making, as measured using the Iowa Gambling Task, did not appear to be directly associated with weight reduction

in patients with obesity and overweight (Witbracht et al., 2012; Stinson et al., 2018). Only Demos et al. (2017) observed an improvement in decision-making following a reduction in body weight, but these authors used a task that employed food-related stimuli.

DISCUSSION

The results of the longitudinal studies confirmed the findings reported in cross-sectional studies, highlighting a relationship between executive functioning and overweight/obesity even if the direction of this relationship remains unclear.

Studies that analyzed the effects of treatments aimed at reducing body weight showed a general improvement in executive tasks as a result. This improvement appeared to occur both in adult populations (Bryan and Tiggemann, 2001; Witbracht et al., 2012) and in children and adolescents (Davis et al., 2011; Kulendran et al., 2014; Vantieghem et al., 2018). Moreover, studies focused on the ability of executive functioning to predict the success of weight-loss interventions found that higher executive functioning could be the cause of BMI reduction (see **Table 4**). Not all EFs appear to be related to obesity. Such inconsistency in the results could be due to different versions of the tasks used to evaluate EFs, as shown in the studies analyzed decision-making.

As for the cross-sectional studies, the authors interpreted the results based on two different types of theoretical models. One hypothesizes that excessive body weight is the cause of changes in executive functioning, according to results showing an improvement in executive tasks following treatment for weight loss (Davis et al., 2007a, 2011; Alosco et al., 2014a,b,c,d, 2015; Verbeken et al., 2014; Galioto et al., 2015; Demos et al., 2017; Xie et al., 2017; Vantieghem et al., 2018). In the other theoretical view, EFs are considered as predictors of eating behaviors related to excessive body weight, like overeating. Studies assessing the effects of strengthening EFs in participants with overweight or obesity (Verbeken et al., 2014; Allom et al., 2018; Dassen et al., 2018b; Raman et al., 2018) have observed both an increase in executive functioning and a reduction in BMI. This reduction may be due to improved eating behavior as a result of adequate working memory, cognitive flexibility, and inhibitory control. These enforcement functions would promote healthier behaviors, reducing the risk associated with obesity, and further improve weight reduction (Allom et al., 2018). Studies that have shown the predictive role of the EFs on the success of weight-loss treatments (Pauli-Pott et al., 2010; Spitznagel et al., 2013, 2014; Kulendran et al., 2014, 2017; Galioto et al., 2016; Xu et al., 2017; Augustijn et al., 2018) confirmed the critical role of executive functioning in the occurrence of obesity.

Concerning bariatric surgery, the effects of weight-loss on executive performances resulted only at the follow-up (Spitznagel et al., 2013; Alosco et al., 2014a; Pearce et al., 2017). This result could be interpreted in two ways. On the one hand, it may suggest that a reduction in body fat favors improvement in executive functioning (Alosco et al., 2014b) as a consequence of the resolution of metabolic alterations related to excessive

BMI; on the other hand, better performance at baseline could lead to an improvement in healthy eating habits (Spitznagel et al., 2013; Pearce et al., 2017), linked to a reduction of BMI over time. This last interpretation is supported by the results observed at the follow-up that showed a higher reduction in BMI in participants presenting better EFs performance at baseline (Spitznagel et al., 2013; Pearce et al., 2017). Lastly, it is interesting to note that control groups with obesity that did not benefit from the treatments (Bryan and Tiggemann, 2001; Alosco et al., 2014c; Pearce et al., 2017; Xie et al., 2017) did not show improvement in performance on cognitive tasks in the follow up assessment. These results confirm that a reduction in body fat leads to gains in executive functioning, although the groups that did not benefit from the treatment did not show a further executive decline.

Despite these findings, short-term follow-ups showed no evidence of a causal relationship of EFs on obesity. These studies did not observe significant differences between participants with obesity who have reduced their body weight and those who maintained their condition unchanged (Deckers et al., 2017). In line with these results, we can conclude that the relationship between EFs and excessive body weight appears robust even when longitudinal studies are considered. However, even considering the results of longitudinal studies appear challenging to determine the direction of this relationship, and further studies are needed.

GENERAL DISCUSSION

Only in recent years, the studies focused their attention on the relationship between excessive body weight and EFs (Fitzpatrick et al., 2013). This relationship appears to be confirmed by most of the studies, both cross-sectional (e.g., Verdejo-García et al., 2010; Cohen et al., 2011; Maayan et al., 2011; Dassen et al., 2018a) and longitudinal (e.g., Spitznagel et al., 2013; Alosco et al., 2014d, 2015; Augustijn et al., 2018), analyzed in this systematic review, despite the heterogeneity of the tasks used and the methodological framework adopted. Functional and neuroimaging studies confirmed changes in the cortical areas involved in executive functioning in participants with obesity (Stingl et al., 2012; Alarcón et al., 2016; Tsai et al., 2016) even when cognitive tasks failed to highlight any significant differences in performance between obesity and normal-weight conditions (Hendrick et al., 2012; Frank et al., 2014; Pearce et al., 2017).

The choice to selected studies which considered different aged made us possible to highlight a similar pattern in the relationship between EFs and overweight/obesity in children (Blanco-Gómez et al., 2015; Tsai et al., 2016) and adults (Cohen et al., 2011; Deckers et al., 2017), despite the individual differences linked to age.

This systematic review allowed us to observe poor performance on executive function tasks also in people with overweight, not only in those with obesity (Verdejo-García et al., 2010; Sellaro and Colzato, 2017), although only a few studies have investigated the condition of overweight (BMI between 25 and 30) compared to normal-weight (BMI lower than 25) and obesity (BMI higher than 30). These results should be explored in

further studies to verify how executive functioning is expressed at the different stages of overweight and to understand if the early intervention could prevent the worsening of the increase in adiposity.

As previously reported, the results of these studies have been interpreted according to two different theoretical models. At the conclusion of this systematic review, no single theoretical model appears to prevail. The empirical data seem to support both theoretical models: the one postulating the influence of executive system dysfunctions on obesity (Drewnowski, 1997; Goldstone et al., 2009; Smith and Robbins, 2013), the other viewing impairment of executive functioning as a consequence of the obesity (Ricca et al., 2009; Pieper and Laugero, 2013).

Other longitudinal studies are needed to disentangle the relationship between obesity and executive dysfunctions. These studies could either examine the eating behavior and BMI of people with low executive functioning over time or monitor the executive functionality of people with overweight who become obese over time. Finally, the possibility that the relationship between executive dysfunctions and overweight/obesity could be bidirectional cannot be excluded; in fact, many studies seem to suggest that the bidirectionality is the real nature of this relation (Spitznagel et al., 2013; Augustijn et al., 2018; Raman et al., 2018).

This systematic review has some strengths, such as the decision to exclude studies of children of preschool age (younger than 5 years) and those over 70 years. This decision was taken for various reasons. First, the EFs and underlying neural areas of preschool-age children are immature and still developing (Diamond, 2013); moreover, during this period children are still introjecting eating habits learned from the external environment (Guxens et al., 2009; Gregory et al., 2010). However, previous studies have shown a specific predictive value of EFs performance in preschool children concerning weight and eating behaviors (Park et al., 2014), and for this reason, it would be interesting to study this specific age group separately. The decision to not include studies of people over 70 years of age was influenced by the “obesity paradox” hypothesis (Artham et al., 2008; Park et al., 2014), which recognizes the health benefits to older people in having a higher BMI. Furthermore, impairment of EFs in older people can be associated with the aging process (Fjell et al., 2016). Although analysis of the relationship between EFs and overweight/obesity in these two age groups could be interesting, their inclusion in the present study would have led to extreme heterogeneity.

LIMITATIONS

This systematic review was not able to identify if one specific EF had a more significant role than another on the analyzed relationship. This result could represent a limitation of this study because it has not allowed us to establish whether differences in performance were due to changes in some functions rather than others. This limitation is due mainly to the heterogeneity of cognitive tasks (Yang et al., 2018). Another limit could be represented by the selection of participants from 5–70 years; in fact, also if the results are coherent, it is known that the

brain continues to develop from childhood to young adulthood, and the differences related to aging could influence the relation between cognitive aspects and weight changes. These age-related differences may have covered possible results that could indicate a causal direction between the variables.

Considering the longitudinal studies, the most extended follow-up period—of 4 years—was performed by Alosco et al. (2014d), though with considerable data loss. No other study investigated the relationship between EFs and body weight following body loss treatment over such a long time. This aspect represents a further limitation of the results, i.e., it is not clear whether the improvements were sustained over time or whether a subsequent reversal of the trend occurs, which might have been the reason behind the drop-out from treatment among bariatric patients. Besides, a possible change in the trend over time could indicate that it is the executive damage that influences the success of weight-loss interventions.

Another limitation is represented by not having included in the systematic review the analysis of psychological variables that could modulate the relationship between EFs and excessive body weight. In fact, a few of the selected studies controlled the psychological dimensions, like anxiety, depression, or emotional regulation. It would be interesting to carry out an analysis of these dimensions. One of the studies that considered some emotional components showed that emotions could modulate the relationship between executive functioning and obesity. However, others found no effect of psychological variables on this relationship (Yau et al., 2014). Even so, examining these psychological variables might also lead to a better understanding of the Emotionally-Driven Eating Model (Chen et al., 2017) in individuals with no eating disorders.

Further limitations are due to the limited samples considered by both cross-sectional and longitudinal studies and by the higher prevalence of females among the participants that do not allow to generalize the results. Weller et al. (2008) found different results between males and females on EF performances, and further studies would be useful to analyse gender differences.

A significant limitation of this work concerns the lack of studies comparing participants with overweight and obesity separately. This comparison would have allowed us to examine the relationship between different severities of excessive body weight and the impairment of executive functioning, and to identify the nature of this relationship. Furthermore, a specific focus on participants with overweight would also have led to determine the cognitive characteristics that might serve as warning signs of the development of obesity.

Finally, a meta-analysis measuring the statistical power of the results obtained from the studies analyzed in this study could help to interpret better the results obtained in this systematic review.

CONCLUSIONS

The analysis of studies on the relationship between executive functioning and excessive body weight did not give us decisive responses to all the questions advanced by this systematic review but clarified a large part of the issues on this topic.

A consistent relationship between executive functioning and overweight/obesity has been confirmed, but it remains unclear whether a general executive dysfunction is involved or whether one EF is more implicated than others.

Although it was not possible to confirm a specific theoretical model on the relationship between EFs and overweight/obesity, the association between these dimensions appears to be the result of a complex interaction between different factors that influence both people's attitude to food and eating and their executive functioning. Prolonged inappropriate food intake related to the maintenance of excessive body weight leads to poorer performance on executive tasks. Furthermore, executive impairment exacerbates inappropriate behaviors, leading to increased body fat (Stinson et al., 2018). Both these aspects are associated with a real risk of cognitive impairment in old age (Sanderlin et al., 2017) and difficulty in responding appropriately to external stimuli that is typical of executive dysfunctions and which would negatively affect the life of obese individuals. It is essential to intervene in both these dimensions to reduce the impact of obesity on quality of life.

It would be interesting in future to evaluate the effectiveness of long-term interventions involving weight-loss programmes. The success of weight-loss interventions may be strictly linked to an improvement in executive functioning because effective executive skills would allow healthier lifestyles. In this context, it might be useful to examine the integrated model of Sellbom and Gunstad (2012) and the Emotionally-Driven Eating model (Gianini et al., 2013; Wagner et al., 2013) in terms of the relationship between BMI and cognitive functioning variables such as mood and emotional regulation that were not often analyzed in the studies reviewed here.

The leading role of this systematic review was to underline the powerful connection between cognitive aspects, specifically

EFs, and excessive body weight, highlighting the importance of considering the nature of the link between these variables in studies on overeating and obesity. A relevant suggestion that emerges by this study is the need for longitudinal studies which, starting from the analysis of EFs, monitor the BMI over time.

It could be essential for structuring intervention aimed at enhancing EFs to prevent the drop-out rate among patients with severe obesity who fail to benefit for a long time from the effects of treatments (Galioto et al., 2016; Xu et al., 2017), by favoring the long-term maintenance of the lower weight achieved. It needs to reduce the risk of further weight gain in people with overweight, thereby preventing the occurrence of severe obesity. Moreover, an integrated approach that also takes emotion regulation and mood into account could be the best strategy for countering dysfunctional eating behaviors and executive functioning; therefore it would be necessary to develop an integrated theoretical model that should jointly consider EFs, eating behavior, emotion regulation, and mood in the field of overweight and obesity.

AUTHOR CONTRIBUTIONS

MC and FF conception of the review, the literature search, and the writing of the manuscript. GF resolved disagreements in the choice of the articles and also contributed to the discussions of the analyzed searches and the revisions of the manuscript. The authors revised, read, and approved the submitted version.

FUNDING

This work was supported by Ph.D Founding Program, Departement of Psychology, Sapienza University of Rome.

REFERENCES

- Alarcón, G., Ray, S., and Nagel, B. J. (2016). Lower working memory performance in overweight and obese adolescents is mediated by white matter microstructure. *J. Int. Neuropsychol. Soc.* 22, 281–292. doi: 10.1017/S135561771500q5
- Allom, V., Mullan, B., Smith, E., Hay, P., and Raman, J. (2018). Breaking bad habits by improving executive function in individuals with obesity. *BMC Public Health* 18:505. doi: 10.1186/s12889-018-5392-y
- Alosco, M. L., Galioto, R., Spitznagel, M. B., Strain, G., Devlin, M., Cohen, R., et al. (2014a). Cognitive function after bariatric surgery: evidence for improvement 3 years after surgery. *Am. J. Surg.* 207, 870–876. doi: 10.1016/j.amjsurg.2013.05.018
- Alosco, M. L., Spitznagel, M. B., Strain, G., Devlin, M., Cohen, R., Crosby, R. D., et al. (2014c). The effects of cystatin C and alkaline phosphatase changes on cognitive function 12-months after bariatric surgery. *J. Neurol. Sci.* 345, 176–180. doi: 10.1016/j.jns.2014.07.037
- Alosco, M. L., Spitznagel, M. B., Strain, G., Devlin, M., Cohen, R., Crosby, R. D., et al. (2015). Improved serum leptin and ghrelin following bariatric surgery predict better postoperative cognitive function. *J. Clin. Neurol.* 11, 48–56. doi: 10.3988/jcn.2015.11.1.48
- Alosco, M. L., Spitznagel, M. B., Strain, G., Devlin, M., Cohen, R., Paul, R., et al. (2014b). Improved memory function two years after bariatric surgery. *Obesity* 22, 32–38. doi: 10.1002/oby.20494
- Alosco, M. L., Spitznagel, M. B., Strain, G., Devlin, M., Crosby, R. D., Mitchell, J. E., et al. (2014d). Family history of Alzheimer's disease limits improvement in cognitive function after bariatric surgery. *SAGE Open Med.* 2:2050312114539477. doi: 10.1177/2050312114539477
- Ariza, M., Garolera, M., Jurado, M. A., Garcia-Garcia, I., Hernan, I., Sánchez-Garre, C., et al. (2012). Dopamine genes (DRD2/ANKK1-TaqA1 and DRD4-7R) and executive function: their interaction with obesity. *PLoS ONE* 7:e41482. doi: 10.1371/journal.pone.0041482
- Arnsten, A. F., and Li, B. M. (2005). Neurobiology of executive functions: catecholamine influences on prefrontal cortical functions. *Biol. Psychiatry* 57, 1377–1384. doi: 10.1016/j.biopsych.2004.08.019
- Artham, S. M., Lavie, C. J., Milani, R. V., and Ventura, H. O. (2008). The obesity paradox: impact of obesity on the prevalence and prognosis of cardiovascular diseases. *Postgrad. Med.* 120, 34–41. doi: 10.3810/pgm.2008.07.1788
- Ashwell, M., Gunn, P., and Gibson, S. (2012). Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obesity Rev.* 13, 275–286. doi: 10.1111/j.1467-789X.2011.00952.x
- Augustijn, M. J. C. M., D'Hondt, E., Van Acker, L., De Guchteneere, A., Lenoir, M., Caeyenberghs, K., et al. (2018). Role of motor competence and executive functioning in weight loss: a study in children with obesity. *J. Dev. Behav. Pediatr.* 39, 642–651. doi: 10.1097/DBP.0000000000000589
- Bauer, L. O., and Manning, K. J. (2016). Challenges in the detection of working memory and attention decrements among overweight

- adolescent girls. *Neuropsychobiology* 73, 43–51. doi: 10.1159/000442670
- Bechara, A., Damasio, H., Tranel, D., and Damasio, A. R. (2005). The Iowa gambling task and the somatic marker hypothesis: some questions and answers. *Trends Cogn. Sci.* 9, 159–162. doi: 10.1016/j.tics.2005.02.002
- Blanco-Gómez, A., Ferré, N., Luque, V., Cardona, M., Gispert-Llauradó, M., Escribano, J., et al. (2015). Being overweight or obese is associated with inhibition control in children from six to ten years of age. *Acta Paediatr.* 104, 619–625. doi: 10.1111/apa.12976
- Bongers, P., van de Giessen, E., Roefs, A., Nederkoorn, C., Booi, J., van den Brink, W., et al. (2015). Being impulsive and obese increases susceptibility to speeded detection of high-calorie foods. *Health Psychol.* 34, 677–685. doi: 10.1037/hea0000167
- Bourassa, K., and Sbarra, D. A. (2017). Body mass and cognitive decline are indirectly associated via inflammation among aging adults. *Brain Behav. Immun.* 60, 63–70. doi: 10.1016/j.bbi.2016.09.023
- Bozkurt, H., Özer, S., Yilmaz, R., Sönmezgöz, E., Kazancı, Ö., Erbaş, O., et al. (2017). Assessment of neurocognitive functions in children and adolescents with obesity. *Appl. Neuropsychol.* 6, 262–268. doi: 10.1080/21622965.2016.1150184
- Bray, G. A. (1999). Etiology and pathogenesis of obesity. *Clin. Cornerstone* 2, 1–15. doi: 10.1016/S1098-3597(99)90001-7
- Brogan, A., Hevey, D., O'Callaghan, G., Yoder, R., and O'Shea, D. (2011). Impaired decision making among morbidly obese adults. *J. Psychosom. Res.* 70, 189–196. doi: 10.1016/j.jpsychores.2010.07.012
- Bryan, J., and Tiggeemann, M. (2001). The effect of weight-loss dieting on cognitive performance and psychological well-being in overweight women. *Appetite* 36, 147–156. doi: 10.1006/appe.2000.0389
- Calvo, D., Galioto, R., Gunstad, J., and Spitznagel, M. B. (2014). Uncontrolled eating is associated with reduced executive functioning. *Clin. Obes.* 4, 172–179. doi: 10.1111/cob.12058
- Carey, M., Small, H., Yoong, S. L., Boyes, A., Bisquera, A., and Sanson-Fisher, R. (2014). Prevalence of comorbid depression and obesity in general practice: a cross-sectional survey. *Br. J. Generic Pract.* 64, e122–e127. doi: 10.3399/bjgp14X677482
- Carpiniello, B., Pinna, F., Pillai, G., Nonnoi, V., Pisano, E., Corrias, S., et al. (2009). Obesity and psychopathology. A study of psychiatric comorbidity among patients attending a specialist obesity unit. *Epidemiol. Psychiatr. Sci.* 18, 119–127. doi: 10.1017/S1121189X00001007
- Catoira, N. P., Tapajóz, F., Allegri, R. F., Lafjer, J., Rodríguez Cámara, M., Iturry, M. L., et al. (2016). Obesity, metabolic profile, and inhibition failure: young women under scrutiny. *Physiol. Behav.* 157, 87–93. doi: 10.1016/j.physbeh.2016.01.040
- Chan, R. C., Shum, D., Touloupoulou, T., and Chen, E. Y. (2008). Assessment of executive functions: review of instruments and identification of critical issues. *Arch. Clin. Neuropsychol.* 23, 201–216. doi: 10.1016/j.acn.2007.08.010
- Chen, P. A., Chavez, R. S., and Heatherton, T. F. (2017). Structural integrity between executive control and reward regions of the brain predicts body fat percentage in chronic dieters. *Cogn. Neurosci.* 8, 162–166. doi: 10.1080/17588928.2016.1235556
- Cohen, J. I., Yates, K. F., Duong, M., and Convit, A. (2011). Obesity, orbitofrontal structure and function are associated with food choice: a cross-sectional study. *BMJ Open* 1:e000175. doi: 10.1136/bmjopen-2011-000175
- Coppin, G., Nolan-Poupart, S., Jones-Gotman, M., and Small, D. M. (2014). Working memory and reward association learning impairments in obesity. *Neuropsychologia* 65, 146–155. doi: 10.1016/j.neuropsychologia.2014.10.004
- Dallman, M. F. (2010). Stress-induced obesity and the emotional nervous system. *Trends Endocrinol. Metabol.* 21, 159–165. doi: 10.1016/j.tem.2009.10.004
- Damasio, A. R. (1995). On some functions of the human prefrontal cortex. *Ann. N. Y. Acad. Sci.* 769, 241–252. doi: 10.1111/j.1749-6632.1995.tb38142.x
- Danner, U. N., Ouweland, C., van Haastert, N. L., Hornsveld, H., and de Ridder, D. T. (2012). Decision-making impairments in women with binge eating disorder in comparison with obese and normal weight women. *Eur. Eating Disord. Rev.* 20, e56–e62. doi: 10.1002/erv.1098
- Dassen, F. C., Houben, K., Van Breukelen, G. J., and Jansen, A. (2018b). Gamified working memory training in overweight individuals reduces food intake but not body weight. *Appetite* 124, 89–98. doi: 10.1016/j.appet.2017.05.009
- Dassen, F. C. M., Houben, K., Allom, V., and Jansen, A. (2018a). Self-regulation and obesity: the role of executive function and delay discounting in the prediction of weight loss. *J. Behav. Med.* 41, 806–818. doi: 10.1007/s10865-018-9940-9
- Davis, C., Patte, K., Levitan, R., Reid, C., Tweed, S., and Curtis, C. (2007b). From motivation to behaviour: a model of reward sensitivity, overeating, and food preferences in the risk profile for obesity. *Appetite* 48, 12–19. doi: 10.1016/j.appet.2006.05.016
- Davis, C. L., Tomporowski, P. D., Boyle, C. A., Waller, J. L., Miller, P. H., Naglieri, J. A., et al. (2007a). Effects of aerobic exercise on overweight children's cognitive functioning: a randomized controlled trial. *Res. Q. Exerc. Sport* 78, 510–519. doi: 10.1080/02701367.2007.10599450
- Davis, C. L., Tomporowski, P. D., McDowell, J. E., Austin, B. P., Miller, P. H., Yanasak, N. E., et al. (2011). Exercise improves executive function and achievement and alters brain activation in overweight children: a randomized, controlled trial. *Health Psychol.* 30, 91–98. doi: 10.1037/a0021766
- de Onis, M. D., Onyango, A. W., Borghi, E., Siyam, A., Nishida, C., and Siekmann, J. (2007). Development of a WHO growth reference for school-aged children and adolescents. *Bull. World Health Organ.* 85, 660–667. doi: 10.2471/BLT.07.043497
- De Wit, L., Luppino, F., van Straten, A., Penninx, B., Zitman, F., and Cuijpers, P. (2010). Depression and obesity: a meta-analysis of community-based studies. *Psychiatr Res.* 178, 230–235. doi: 10.1016/j.psychres.2009.04.015
- Deckers, K., van Bortel, M. P. J., Verhey, F. R. J., and Köhler, S. (2017). Obesity and cognitive decline in adults: effect of methodological choices and confounding by age in a longitudinal study. *J. Nutri. Health Aging* 21, 546–553. doi: 10.1007/s12603-016-0757-3
- Delgado-Rico, E., Río-Valle, J. S., González-Jiménez, E., Campoy, C., and Verdejo-García, A. (2012). BMI predicts emotion-driven impulsivity and cognitive inflexibility in adolescents with excess weight. *Obesity* 20, 1604–1610. doi: 10.1038/oby.2012.47
- Demos, K. E., McCaffery, J. M., Thomas, J. G., Mailloux, K. A., Hare, T. A., and Wing, R. R. (2017). Identifying the mechanisms through which behavioral weight-loss treatment improves food decision-making in obesity. *Appetite* 114, 93–100. doi: 10.1016/j.appet.2017.03.013
- Diamond, A. (2013). Executive functions. *Annu. Rev. Psychol.* 64, 135–168. doi: 10.1146/annurev-psych-113011-143750
- Dohle, S., Diel, K., and Hofmann, W. (2018). Executive functions and the self-regulation of eating behavior: a review. *Appetite* 124, 4–9. doi: 10.1016/j.appet.2017.05.041
- Drewnowski, A. (1997). Taste preferences and food intake. *Annu. Rev. Nutr.* 17, 237–253. doi: 10.1146/annurev.nutr.17.1.237
- Dubbert, P. M., Carithers, T., Hall, J. E., Barbour, K. A., Clark, B. L., Sumner, A. E., et al. (2002). Obesity, physical inactivity, and risk for cardiovascular disease. *Am. J. Med. Sci.* 324, 116–126. doi: 10.1097/0000441-200209000-00002
- Elliott, R. (2003). Executive functions and their disorders: Imaging in clinical neuroscience. *Br. Med. Bull.* 65, 49–59. doi: 10.1093/bmb/65.1.49
- Emery, R. L., and Levine, M. D. (2017). Questionnaire and behavioral task measures of impulsivity are differentially associated with body mass index: a comprehensive meta-analysis. *Psychol. Bull.* 143, 868–902. doi: 10.1037/bul0000105
- Fagundo, A. B., De la Torre, R., Jiménez-Murcia, S., Agüera, Z., Granero, R., Tárrega, S., et al. (2012). Executive functions profile in extreme eating/weight conditions: from anorexia nervosa to obesity. *PLoS ONE* 7:e43382. doi: 10.1371/journal.pone.0043382
- Fields, S. A., Sabet, M., and Reynolds, B. (2013). Dimensions of impulsive behavior in obese, overweight, and healthy-weight adolescents. *Appetite* 70, 60–66. doi: 10.1016/j.appet.2013.06.089
- Fitzpatrick, S., Gilbert, S., and Serpell, L. (2013). Systematic review: are overweight and obese individuals impaired on behavioural tasks of executive functioning? *Neuropsychol. Rev.* 23, 138–156. doi: 10.1007/s11065-013-9224-7
- Fjell, A. M., Sneve, M. H., Grydeland, H., Storsve, A. B., and Walhovd, K. B. (2016). The disconnected brain and executive function decline in aging. *Cereb. Cortex* 27, 2303–2317. doi: 10.1093/cercor/bhw082
- Flegal, K. M., Wei, R., and Ogden, C. (2002). Weight-for-stature compared with body mass index-for-age growth charts for the United States from the Centers for Disease control and prevention. *Am. J. Clin. Nutr.* 75, 761–766. doi: 10.1093/ajcn/75.4.761

- Forcano, L., Mata, F., de la Torre, R., and Verdejo-García, A. (2018). Cognitive and neuromodulation strategies for unhealthy eating and obesity: systematic review and discussion of neurocognitive mechanisms. *Neurosci. Biobehav. Rev.* 87, 161–191. doi: 10.1016/j.neubiorev.2018.02.003
- Frank, S., Wilms, B., Veit, R., Ernst, B., Thurnheer, M., Kullmann, S., et al. (2014). Altered brain activity in severely obese women may recover after Roux-en Y gastric bypass surgery. *Int. J. Obes.* 38, 341–348. doi: 10.1038/ijo.2013.60
- Funahashi, S. (2001). Neuronal mechanisms of executive control by the prefrontal cortex. *Neurosci. Res.* 39, 147–165. doi: 10.1016/S0168-0102(00)00224-8
- Galioto, R., Allosco, M. L., Meister, J., and Spitznagel, M. B. (2014). Self-reported physical activity is associated with cognitive function in lean, but not obese individuals. *Clin. Obes.* 4, 309–315. doi: 10.1111/cob.12071
- Galioto, R., Allosco, M. L., Spitznagel, M. B., Strain, G., Devlin, M., Cohen, R., et al. (2015). Glucose regulation and cognitive function after bariatric surgery. *J. Clin. Exp. Neuropsychol.* 37, 402–413. doi: 10.1080/13803395.2015.1023264
- Galioto, R., Bond, D., Gunstad, J., Pera, V., Rathier, L., and Tremont, G. (2016). Executive functions predict weight loss in a medically supervised weight loss programme. *Obesity Sci. Pract.* 2, 334–340. doi: 10.1002/osp4.70
- Galioto, R. M., Allosco, M. L., Spitznagel, M. B., Stanek, K. M., and Gunstad, J. (2013). Cognitive reserve preserves cognitive function in obese individuals. *Aging Neuropsychol. Cogn.* 20, 684–699. doi: 10.1080/13825585.2012.762972
- Gameiro, F., Perea, M. V., Ladera, V., Rosa, B., and Garcia, R. (2017). Executive functioning in obese individuals waiting for clinical treatment. *Psicothema* 29, 61–66. doi: 10.7334/psicothema2016.202
- Garipey, G., Nitka, D., and Schmitz, N. (2010). The association between obesity and anxiety disorders in the population: a systematic review and meta-analysis. *Int. J. Obes.* 34, 407–419. doi: 10.1038/ijo.2009.252
- Gentier, I., Augustijn, M., Deforche, B., Tanghe, A., De Bourdeaudhuij, I., Lenoir, M., et al. (2013). A comparative study of performance in simple and choice reaction time tasks between obese and healthy-weight children. *Res. Dev. Disabil.* 34, 2635–2641. doi: 10.1016/j.ridd.2013.04.016
- Gettens, K. M., and Gorin, A. A. (2017). Executive function in weight loss and weight loss maintenance: a conceptual review and novel neuropsychological model of weight control. *J. Behav. Med.* 40, 687–701. doi: 10.1007/s10865-017-9831-5
- Gianini, L. M., White, M. A., and Masheb, R. M. (2013). Eating pathology, emotion regulation, and emotional overeating in obese adults with binge eating disorder. *Eat. Behav.* 14, 309–313. doi: 10.1016/j.eatbeh.2013.05.008
- Gluck, M. E., Viswanath, P., and Stinson, E. J. (2017). Obesity, appetite, and the prefrontal cortex. *Curr. Obes. Rep.* 6, 380–388. doi: 10.1007/s13679-017-0289-0
- Goldschmidt, A. B., O'Brien, S., Lavender, J. M., Pearson, C. M., Le Grange, D., and Hunter, S. J. (2018). Executive functioning in a racially diverse sample of children who are overweight and at risk for eating disorders. *Appetite* 124, 43–49. doi: 10.1016/j.appet.2017.03.010
- Goldstone, A. P., Precht de Hernandez, C. G., Beaver, J. D., Muhammed, K., Croese, C., Bell, G., et al. (2009). Fasting biases brain reward systems towards high-calorie foods. *Eur. J. Neurosci.* 30, 1625–1635. doi: 10.1111/j.1460-9568.2009.06949.x
- Gonzales, M. M., Tarumi, T., Miles, S. C., Tanaka, H., Shah, F., and Haley, A. P. (2010). Insulin sensitivity as a mediator of the relationship between BMI and working memory-related brain activation. *Obesity* 18, 2131–2137. doi: 10.1038/oby.2010.183
- Grafman, J., and Litvan, I. (1999). Importance of deficits in executive functions. *Lancet* 354, 1921–1923. doi: 10.1016/S0140-6736(99)90438-5
- Gregory, J. E., Paxton, S. J., and Brozovic, A. M. (2010). Maternal feeding practices, child eating behaviour and body mass index in preschool-aged children: a prospective analysis. *Int. J. Behav. Nutri. Phys. Activity* 7:55. doi: 10.1186/1479-5868-7-55
- Gunstad, J., Paul, R. H., Cohen, R. A., Tate, D. F., Spitznagel, M. B., and Gordon, E. (2007). Elevated body mass index is associated with executive dysfunction in otherwise healthy adults. *Compr. Psychiatr.* 48, 57–61. doi: 10.1016/j.comppsy.2006.05.001
- Guxens, M., Mendez, M. A., Julvez, J., Plana, E., Forns, J., Basagaña, X., et al. (2009). Cognitive function and overweight in preschool children. *Am. J. Epidemiol.* 170, 438–446. doi: 10.1093/aje/kwp140
- Hall, P. A., and Marteau, T. M. (2014). Executive function in the context of chronic disease prevention: theory, research and practice. *Prevent. Med.* 68, 44–50. doi: 10.1016/j.ypmed.2014.07.008
- Harrington, D. M., Staiano, A. E., Broyles, S. T., Gupta, A. K., and Katzmarzyk, P. T. (2013). BMI percentiles for the identification of abdominal obesity and metabolic risk in children and adolescents: evidence in support of the CDC 95th percentile. *Eur. J. Clin. Nutr.* 67, 218–222. doi: 10.1038/ejcn.2012.203
- Hauener, H. (2017). “Obesity and diabetes,” in *Textbook of Diabetes*, eds R. I. G. Holt, C. Cockram, A. Flyvbjerg, and B. J. Goldstein, 215–228.
- Hendrick, O. M., Luo, X., Zhang, S., and Li, C. S. (2012). Saliency processing and obesity: a preliminary imaging study of the stop signal task. *Obesity* 20, 1796–1802. doi: 10.1038/oby.2011.180
- Heymsfield, S. B., and Wadden, T. A. (2017). Mechanisms, pathophysiology, and management of obesity. *N. England J. Med.* 376, 254–266. doi: 10.1056/NEJMra1514009
- Hofmann, W., Schmeichel, B. J., and Baddeley, A. D. (2012). Executive functions and self-regulation. *Trends Cogn. Sci.* 16, 174–180. doi: 10.1016/j.tics.2012.01.006
- Jacques, S., and Marcovitch, S. (2010). “Development of executive function across the life span,” in *The Handbook of Life-Span Development, Vol. 1: Cognition, Biology, and Methods*, eds W. F. Overton and R. M. Lerner (Hoboken, NJ: John Wiley & Sons Inc.), 431–466. doi: 10.1002/9780470880166.hlsd001013
- Janssen, I., Katzmarzyk, P. T., and Ross, R. (2004). Waist circumference and not body mass index explains obesity-related health risk. *Am. J. Clin. Nutr.* 79, 379–384. doi: 10.1093/ajcn/79.3.379
- Jiang, S. Z., Lu, W., Zong, X. F., Ruan, H. Y., and Liu, Y. (2016). Obesity and hypertension. *Exp. Ther. Med.* 12, 2395–2399. doi: 10.3892/etm.2016.3667
- Kamijo, K., Khan, N. A., Pontifex, M. B., Scudder, M. R., Drollette, E. S., Raine, L. B., et al. (2012a). The relation of adiposity to cognitive control and scholastic achievement in preadolescent children. *Obesity* 20, 2406–2411. doi: 10.1038/oby.2012.112
- Kamijo, K., Pontifex, M. B., Khan, N. A., Raine, L. B., Scudder, M. R., Drollette, E. S., et al. (2012b). The association of childhood obesity to neuroelectric indices of inhibition. *Psychophysiology* 49, 1361–1371. doi: 10.1111/j.1469-8986.2012.01459.x
- Kamijo, K., Pontifex, M. B., Khan, N. A., Raine, L. B., Scudder, M. R., Drollette, E. S., et al. (2012c). The negative association of childhood obesity to cognitive control of action monitoring. *Cereb. Cortex* 24, 654–662. doi: 10.1093/cercor/bhs349
- Kane, M. J., Conway, A. R. A., Miura, T. K., and Colflesh, G. J. (2007). Working memory, attention control, and the N-back task: a question of construct validity. *J. Exp. Psychol.* 33:615. doi: 10.1037/0278-7393.33.3.615
- Kelly, T., Yang, W., Chen, C. S., Reynolds, K., and He, J. (2008). Global burden of obesity in 2005 and projections to 2030. *Int. J. Obes.* 32, 1431–1437. doi: 10.1038/ijo.2008.102
- Kittel, R., Schmidt, R., and Hilbert, A. (2017). Executive functions in adolescents with binge-eating disorder and obesity. *Int. J. Eating Disord.* 50, 933–941. doi: 10.1002/eat.22714
- Kolotkin, R. L., Meter, K., and Williams, G. R. (2001). Quality of life and obesity. *Obesity Rev.* 2, 219–229. doi: 10.1046/j.1467-789X.2001.00040.x
- Kulendran, M., Borovoi, L., Purkayastha, S., Darzi, A., and Vlaev, I. (2017). Impulsivity predicts weight loss after obesity surgery. *Surg. Obesity Related Dis.* 13, 1033–1040. doi: 10.1016/j.soard.2016.12.031
- Kulendran, M., Vlaev, I., Sugden, C., King, D., Ashrafian, H., Gately, P., et al. (2014). Neuropsychological assessment as a predictor of weight loss in obese adolescents. *Int. J. Obes.* 38, 507–512. doi: 10.1038/ijo.2013.198
- Lam, B. C. C., Koh, G. C. H., Chen, C., Wong, M. T. K., and Fallows, S. J. (2015). Comparison of body mass index (BMI), body adiposity index (BAI), waist circumference (WC), waist-to-hip ratio (WHR) and waist-to-height ratio (WHtR) as predictors of cardiovascular disease risk factors in an adult population in Singapore. *PLoS ONE* 10:e0122985. doi: 10.1371/journal.pone.0122985
- Lappin, J. S., and Eriksen, C. W. (1966). Use of a delayed signal to stop a visual reaction-time response. *J. Exp. Psychol.* 72, 805–811. doi: 10.1037/h0021266
- Lasselain, J., Magne, E., Beau, C., Aubert, A., Dexpert, S., Carrez, J., et al. (2016). Low-grade inflammation is a major contributor of impaired attentional set shifting in obese subjects. *Brain Behav. Immun.* 58, 63–68. doi: 10.1016/j.bbi.2016.05.013
- Liang, J., Matheson, B. E., Kaye, W. H., and Boutelle, K. N. (2014). Neurocognitive correlates of obesity and obesity-related behaviors in children and adolescents. *Int. J. Obes.* 38, 494–506. doi: 10.1038/ijo.2013.142

- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Götzsche, P. C., Ioannidis, J. P., et al. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 6:e1000100. doi: 10.1371/journal.pmed.1000100
- Littleton, S. W. (2012). Impact of obesity on respiratory function. *Respirology* 17, 43–49. doi: 10.1111/j.1440-1843.2011.02096.x
- Loeber, S., Grosshans, M., Korucuoglu, O., Vollmert, C., Vollstädt-Klein, S., Schneider, S., et al. (2012). Impairment of inhibitory control in response to food-associated cues and attentional bias of obese participants and normal-weight controls. *Int. J. Obes.* 36, 1334–1339. doi: 10.1038/ijo.2011.184
- Luppino, F. S., de Wit, L. M., Bouvy, P. F., Stijnen, T., Cuijpers, P., Penninx, B. W., et al. (2010). Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch. Gen. Psychiatry* 67, 220–229. doi: 10.1001/archgenpsychiatry.2010.2
- Maayan, L., Hoogendoorn, C., Sweat, V., and Convit, A. (2011). Disinhibited eating in obese adolescents is associated with orbitofrontal volume reductions and executive dysfunction. *Obesity* 19, 1382–1387. doi: 10.1038/oby.2011.15
- McCrary, M. A., Fuss, P. J., Hays, N. P., Vinken, A. G., Greenberg, A. S., and Roberts, S. B. (1999). Overeating in America: association between restaurant food consumption and body fatness in healthy adult men and women ages 19 to 80. *Obes. Res.* 7, 564–571. doi: 10.1002/j.1550-8528.1999.tb00715.x
- McPhail, S. M., Schippers, M., Marshall, A. L., Waite, M., and Kuipers, P. (2014). Perceived barriers and facilitators to increasing physical activity among people with musculoskeletal disorders: a qualitative investigation to inform intervention development. *Clin. Interv. Aging* 9, 2113–2122. doi: 10.2147/CLIA.S72731
- Milner, B. (1963). Effects of different brain lesions on card sorting: the role of the frontal lobes. *Arch. Neurol.* 9, 90–100. doi: 10.1001/archneur.1963.00460070100010
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., and Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: a latent variable analysis. *Cogn. Psychol.* 41, 49–100. doi: 10.1006/cogp.1999.0734
- Moher, D., Liberati, A., Tetzlaff, J., and Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann. Intern. Med.* 151, 264–269. doi: 10.7326/0003-4819-151-4-200908180-00135
- Mole, T. B., Irvine, M. A., Worbe, Y., Collins, P., Mitchell, S. P., Bolton, S., et al. (2015). Impulsivity in disorders of food and drug misuse. *Psychol. Med.* 45, 771–782. doi: 10.1017/S0033291714001834
- Moreno-López, L., Soriano-Mas, C., Delgado-Rico, E., Rio-Valle, J. S., and Verdejo-García, A. (2012). Brain structural correlates of reward sensitivity and impulsivity in adolescents with normal and excess weight. *PLoS ONE* 7:e49185. doi: 10.1371/journal.pone.0049185
- Morris, M. C., Evans, D. A., Tangney, C. C., Bienias, J. L., and Wilson, R. S. (2005). Fish consumption and cognitive decline with age in a large community study. *Arch. Neurol.* 62:1849–1853. doi: 10.1001/archneur.62.12.noc50161
- Navas, J. F., Vilar-López, R., Perales, J. C., Steward, T., Fernández-Aranda, F., and Verdejo-García, A. (2016). Altered decision-making under risk in obesity. *PLoS ONE* 11:e0155600. doi: 10.1371/journal.pone.0155600
- Nederkorn, C., Braet, C., Van Eijs, Y., Tanghe, A., and Jansen, A. (2006). Why obese children cannot resist food: the role of impulsivity. *Eat. Behav.* 7, 315–322. doi: 10.1016/j.eatbeh.2005.11.005
- Nigg, J. T. (2000). On inhibition/disinhibition in developmental psychopathology: views from cognitive and personality psychology and a working inhibition taxonomy. *Psychol. Bull.* 126:220. doi: 10.1037/0033-2909.126.2.220
- Ouwens, M. A., van Strien, T., and van der Staak, C. P. (2003). Tendency toward overeating and restraint as predictors of food consumption. *Appetite* 40, 291–298. doi: 10.1016/S0195-6663(03)00006-0
- Park, J., Ahmadi, S. F., Streja, E., Molnar, M. Z., Flegal, K. M., Gillen, D., et al. (2014). Obesity paradox in end-stage kidney disease patients. *Prog. Cardiovasc. Dis.* 56, 415–425. doi: 10.1016/j.pcad.2013.10.005
- Pauli-Pott, U., Albayrak, O., Hebebrand, J., and Pott, W. (2010). Does inhibitory control capacity in overweight and obese children and adolescents predict success in a weight-reduction program? *Eur. Child Adolescent Psychiatry* 19, 135–141. doi: 10.1007/s00787-009-0049-0
- Pearce, A. L., Mackey, E., Cherry, J. B. C., Olson, A., You, X., Magge, S. N., et al. (2017). Effect of adolescent bariatric surgery on the brain and cognition: a pilot study. *Obesity* 25, 1852–1860. doi: 10.1002/oby.22013
- Pearce, A. L., Mackey, E., Nadler, E. P., and Vaidya, C. J. (2018). Sleep health and psychopathology mediate executive deficits in pediatric obesity. *Childhood Obesity* 14, 189–196. doi: 10.1089/chi.2017.0281
- Pennington, B. F., and Ozonoff, S. (1996). Executive functions and developmental psychopathology. *J. Child Psychol. Psychiatry* 37, 51–87. doi: 10.1111/j.1469-7610.1996.tb01380.x
- Perpiñá, C., Segura, M., and Sánchez-Reales, S. (2017). Cognitive flexibility and decision-making in eating disorders and obesity. *Eating and Weight Disorders-Studies on Anorexia. Bulimia Obesity* 22, 435–444. doi: 10.1007/s40519-016-0331-3
- Perry, V. H. (2004). The influence of systemic inflammation on inflammation in the brain: implications for chronic neurodegenerative disease. *Brain Behav. Immun.* 18, 407–413. doi: 10.1016/j.bbi.2004.01.004
- Pieper, J. R., and Laugero, K. D. (2013). Preschool children with lower executive function may be more vulnerable to emotional-based eating in the absence of hunger. *Appetite* 62, 103–109. doi: 10.1016/j.appet.2012.11.020
- Pignatti, R., Bertella, L., Albani, G., Mauro, A., Molinari, E., and Semenza, C. (2006). Decision-making in obesity: a study using the Gambling Task. *Eat. Weight Disord. Stud. Anorexia Bulimia Obesity* 11, 126–132. doi: 10.1007/BF03327557
- Prentice, A. M. (2001). Overeating: the health risks. *Obesity Res.* 9, 234S–238S. doi: 10.1038/oby.2001.124
- Qavam, S. E., Anisan, A., Fathi, M., and Pourabbasi, A. (2015). Study of relationship between obesity and executive functions among high school students in Bushehr, Iran. *J. Diabetes Metab. Disord.* 14:79. doi: 10.1186/s40200-015-0211-9
- Raman, J., Hay, P., Tchanturia, K., and Smith, E. (2018). A randomised controlled trial of manualized cognitive remediation therapy in adult obesity. *Appetite* 123, 269–279. doi: 10.1016/j.appet.2017.12.023
- Reitan, R. M. (1958). Validity of the Trail Making Test as an indicator of organic brain damage. *Percept. Mot. Skills* 8, 271–276. doi: 10.2466/pms.1958.8.3.271
- Restivo, M. R., McKinnon, M. C., Frey, B. N., Hall, G. B., Syed, W., and Taylor, V. H. (2017). The impact of obesity on neuropsychological functioning in adults with and without major depressive disorder. *PLoS ONE* 12:e0176898. doi: 10.1371/journal.pone.0176898
- Reyes, S., Peirano, P., Peigneux, P., Lozoff, B., and Algarin, C. (2015). Inhibitory control in otherwise healthy overweight 10-year-old children. *Int. J. Obes.* 39, 1230–1235. doi: 10.1038/ijo.2015.49
- Reynolds, C. R. (1997). Forward and backward memory span should not be combined for clinical analysis. *Arch. Clin. Neuropsychol.* 12, 29–40. doi: 10.1093/arclin/12.1.29
- Ricca, V., Castellini, G., Lo Sauro, C., Ravaldi, C., Lapi, F., Mannucci, E., et al. (2009). Correlations between binge eating and emotional eating in a sample of overweight subjects. *Appetite* 53, 418–421. doi: 10.1016/j.appet.2009.07.008
- Richards, J. B., Zhang, L., Mitchell, S. H., and De Wit, H. (1999). Delay or probability discounting in a model of impulsive behavior: effect of alcohol. *J. Exp. Anal. Behav.* 71, 121–143. doi: 10.1901/jeab.1999.71-121
- Rogers, R. D., and Monsell, S. (1995). Costs of a predictable switch between simple cognitive tasks. *J. Exp. Psychol.* 124:207. doi: 10.1037/0096-3445.124.2.207
- Sanderlin, A. H., Todem, D., and Bozoki, A. C. (2017). Obesity and co-morbid conditions are associated with specific neuropsychiatric symptoms in mild cognitive impairment. *Front. Aging Neurosci.* 9:164. doi: 10.3389/fnagi.2017.00325
- Schiff, S., Amodio, P., Testa, G., Nardi, M., Montagnese, S., Caregaro, L., et al. (2016). Impulsivity toward food reward is related to BMI: evidence from intertemporal choice in obese and normal-weight individuals. *Brain Cogn.* 110, 112–119. doi: 10.1016/j.bandc.2015.10.001
- Sellaro, R., and Colzato, L. S. (2017). High body mass index is associated with impaired cognitive control. *Appetite* 113, 301–309. doi: 10.1016/j.appet.2017.03.008
- Sellbom, K. S., and Gunstad, J. (2012). Cognitive function and decline in obesity. *J. Alzheimer Dis.* 30, S89–S95. doi: 10.3233/JAD-2011-111073
- Skoranski, A. M., Most, S. B., Lutz-Stehl, M., Hoffman, J. E., Hassink, S. G., and Simons, R. F. (2013). Response monitoring and cognitive control in childhood obesity. *Biol. Psychol.* 92, 199–204. doi: 10.1016/j.biopsycho.2012.09.001
- Smith, D. G., and Robbins, T. W. (2013). The neurobiological underpinnings of obesity and binge eating: a rationale for adopting the food addiction model. *Biol. Psychiatry* 73, 804–810. doi: 10.1016/j.biopsycho.2012.08.026

- Smith, P. J., and Blumenthal, J. A. (2016). Dietary factors and cognitive decline. *J. Prevent. Alzheimer Dis.* 3, 53–64. doi: 10.14283/jpad.2015.71
- Spitoni, G. F., Ottaviani, C., Petta, A. M., Zingaretti, P., Aragona, M., Sarnicola, A., et al. (2017). Obesity is associated with lack of inhibitory control and impaired heart rate variability reactivity and recovery in response to food stimuli. *Int. J. Psychophysiol.* 116, 77–84. doi: 10.1016/j.ijpsycho.2017.04.001
- Spitznagel, M. B., Alosco, M., Galioto, R., Strain, G., Devlin, M., Sysko, R., et al. (2014). The role of cognitive function in postoperative weight loss outcomes: 36-month follow-up. *Obesity Surg.* 24, 1078–1084. doi: 10.1007/s11695-014-1205-2
- Spitznagel, M. B., Garcia, S., Miller, L. A., Strain, G., Devlin, M., Wing, R., et al. (2013). Cognitive function predicts weight loss after bariatric surgery. *Surg. Obesity Related Dis.* 9, 453–459. doi: 10.1016/j.soard.2011.10.008
- Stanek, K. M., Strain, G., Devlin, M., Cohen, R., Paul, R., Crosby, R. D., et al. (2013). Body mass index and neurocognitive functioning across the adult lifespan. *Neuropsychology* 27, 141–151. doi: 10.1037/a0031988
- Steenbergen, L., and Colzato, L. S. (2017). Overweight and cognitive performance: high body mass index is associated with impairment in reactive control during task switching. *Front. Nutri.* 4:51. doi: 10.3389/fnut.2017.00051
- Stingl, K. T., Kullmann, S., Ketterer, C., Heni, M., Häring, H. U., Fritsche, A., et al. (2012). Neuronal correlates of reduced memory performance in overweight subjects. *Neuroimage* 60, 362–369. doi: 10.1016/j.neuroimage.2011.12.012
- Stinson, E. J., Krakoff, J., and Gluck, M. E. (2018). Depressive symptoms and poorer performance on the Stroop Task are associated with weight gain. *Physiol. Behav.* 186, 25–30. doi: 10.1016/j.physbeh.2018.01.005
- Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *J. Exp. Psychol.* 18, 643–662. doi: 10.1037/h0054651
- Sweat, V., Yates, K. F., Migliaccio, R., and Convit, A. (2017). Obese adolescents show reduced cognitive processing speed compared with healthy weight peers. *Childhood Obesity* 13, 190–196. doi: 10.1089/chi.2016.0255
- Thiara, G., Cigliobianco, M., Muravsky, A., Paoli, R. A., Mansur, R., Hawa, R., et al. (2017). Evidence for neurocognitive improvement after bariatric surgery: a systematic review. *Psychosomatics* 58, 217–227. doi: 10.1016/j.psym.2017.02.004
- Tsai, C. L., Chen, F. C., Pan, C. Y., and Tseng, Y. T. (2016). The neurocognitive performance of visuospatial attention in children with obesity. *Front. Psychol.* 7:1033. doi: 10.3389/fpsyg.2016.01033
- Vainik, U., Dagher, A., Dubé, L., and Fellows, L. K. (2013). Neurobehavioural correlates of body mass index and eating behaviours in adults: a systematic review. *Neurosci. Biobehav. Rev.* 37, 279–299. doi: 10.1016/j.neubiorev.2012.11.008
- Van der Oord, S., Braet, C., Cortese, S., and Claes, L. (2018). Testing the dual pathway model of ADHD in obesity: a pilot study. *Eating Weight Disord. Stud. Anorexia Bulimia Obesity* 23, 507–512. doi: 10.1007/s40519-017-0375-z
- Vantieghem, S., Bautmans, I., De Guchteneere, A., Tanghe, A., and Probyn, S. (2018). Improved cognitive functioning in obese adolescents after a 30-week inpatient weight loss program. *Pediatr. Res.* 1, 267–271. doi: 10.1038/s41390-018-0047-3
- Varkevisser, R. D. M., van Stralen, M. M., Kroeze, W., Ket, J. C. F., and Steenhuis, I. H. M. (2019). Determinants of weight loss maintenance: a systematic review. *Obesity Rev.* 20, 171–211. doi: 10.1111/obr.12772
- Verbeken, S., Braet, C., Bosmans, G., and Goossens, L. (2014). Comparing decision making in average and overweight children and adolescents. *Int. J. Obes.* 38, 547–551. doi: 10.1038/ijo.2013.235
- Verbeken, S., Braet, C., Goossens, L., and Van der Oord, S. (2013). Executive function training with game elements for obese children: a novel treatment to enhance self-regulatory abilities for weight-control. *Behav. Res. Ther.* 51, 290–299. doi: 10.1016/j.brat.2013.02.006
- Verdejo-García, A., Pérez-Expósito, M., Schmidt-Río-Valle, J., Fernández-Serrano, M. J., Cruz, F., Pérez-García, M., et al. (2010). Selective alterations within executive functions in adolescents with excess weight. *Obesity* 18, 1572–1578. doi: 10.1038/oby.2009.475
- Volkow, N. D., Wang, G. J., and Baler, R. D. (2011). Reward, dopamine and the control of food intake: implications for obesity. *Trends Cogn. Sci.* 15, 37–46. doi: 10.1016/j.tics.2010.11.001
- Voon, V., Irvine, M. A., Derbyshire, K., Worbe, Y., Lange, I., Abbott, S., et al. (2014). Measuring “waiting” impulsivity in substance addictions and binge eating disorder in a novel analogue of rodent serial reaction time task. *Biol. Psychiatr.* 75, 148–155. doi: 10.1016/j.biopsych.2013.05.013
- Wagner, D. D., Altman, M., Boswell, R. G., Kelley, W. M., and Heatherton, T. F. (2013). Self-regulatory depletion enhances neural responses to rewards and impairs top-down control. *Psychol. Sci.* 24, 2262–2271. doi: 10.1177/0956797613492985
- Wang, G. J., Volkow, N. D., Telang, F., Jayne, M., Ma, J., Rao, M., et al. (2004). Exposure to appetitive food stimuli markedly activates the human brain. *Neuroimage* 21, 1790–1797. doi: 10.1016/j.neuroimage.2003.11.026
- Weller, R. E., Cook, E. W. III, Avsar, K. B., and Cox, J. E. (2008). Obese women show greater delay discounting than healthy-weight women. *Appetite* 51, 563–569. doi: 10.1016/j.appet.2008.04.010
- Witbracht, M. G., Laugero, K. D., Van Loan, M. D., Adams, S. H., and Keim, N. L. (2012). Performance on the Iowa Gambling Task is related to magnitude of weight loss and salivary cortisol in a diet-induced weight loss intervention in overweight women. *Physiol. Behav.* 106, 291–297. doi: 10.1016/j.physbeh.2011.04.035
- World Health Organization (2000). *Obesity: Preventing and Managing the Global Epidemic* (No. 894). World Health Organization; 894: i–xii, 1–253.
- World Health Organization (2015). *World Health Statistics 2015*. Geneva, WHO, 2015, pp 101–111. Available online at: www.who.int/gho/publications/world_health_statistics/2015/en/
- Wu, N., Chen, Y., Yang, J., and Li, F. (2017). Childhood obesity and academic performance: the role of working memory. *Front. Psychol.* 8:611. doi: 10.3389/fpsyg.2017.00611
- Wu, X., Nussbaum, M. A., and Madigan, M. L. (2016). Executive function and measures of fall risk among people with obesity. *Percept. Mot. Skills* 122, 825–839. doi: 10.1177/0031512516646158
- Xie, C., Wang, X., Zhou, C., Xu, C., and Chang, Y. K. (2017). Exercise and dietary program-induced weight reduction is associated with cognitive function among obese adolescents: a longitudinal study. *Peer J* 5:e3286. doi: 10.7717/peerj.3286
- Xu, X., Deng, Z. Y., Huang, Q., Zhang, W. X., Qi, C. Z., and Huang, J. A. (2017). Prefrontal cortex-mediated executive function as assessed by Stroop task performance associates with weight loss among overweight and obese adolescents and young adults. *Behav. Brain Res.* 321, 240–248. doi: 10.1016/j.bbr.2016.12.040
- Yadava, A., and Sharma, N. R. (2014). Cognitive functioning in relation to body mass index. *J. Indian Acad. Appl. Psychol.* 40, 203–210. doi: 10.1093%2Fjgerona%2F40x152
- Yang, Y., Shields, G. S., Guo, C., and Liu, Y. (2018). Executive function performance in obesity and overweight individuals: a meta-analysis and review. *Neurosci. Biobehav. Rev.* 84, 225–244. doi: 10.1016/j.neubiorev.2017.11.020
- Yau, P. L., Kang, E. H., Javier, D. C., and Convit, A. (2014). Preliminary evidence of cognitive and brain abnormalities in uncomplicated adolescent obesity. *Obesity* 22, 1865–1871. doi: 10.1002/oby.20801
- Yumuk, V., Tsigos, C., Fried, M., Schindler, K., Busetto, L., and Micic, D. (2015). European guidelines for obesity management in adults. *Obesity Facts* 8, 402–424. doi: 10.1159/000442721

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Favieri, Forte and Casagrande. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.