






Review

Purine-Metabolising Enzymes and Apoptosis in Cancer

Marcella Camici ^{1,*}, Mercedes Garcia-Gil ^{2,†}, Rossana Pesi ¹, Simone Allegrini ¹ and Maria Grazia Tozzi ¹

¹ Dipartimento di Biologia, Unità di Biochimica, Via S. Zeno 51, 56127 Pisa, Italy

² Dipartimento di Biologia, Unità di Fisiologia Generale, Via S. Zeno 31, 56127 Pisa, Italy

* Correspondence: marcella.camici@unipi.it; Tel.: +39-050 2211458

† These authors equally contributed to the work.

Received: 24 July 2019; Accepted: 7 September 2019; Published: 12 September 2019



Abstract: The enzymes of both de novo and salvage pathways for purine nucleotide synthesis are regulated to meet the demand of nucleic acid precursors during proliferation. Among them, the salvage pathway enzymes seem to play the key role in replenishing the purine pool in dividing and tumour cells that require a greater amount of nucleotides. An imbalance in the purine pools is fundamental not only for preventing cell proliferation, but also, in many cases, to promote apoptosis. It is known that tumour cells harbour several mutations that might lead to defective apoptosis-inducing pathways, and this is probably at the basis of the initial expansion of the population of neoplastic cells. Therefore, knowledge of the molecular mechanisms that lead to apoptosis of tumoural cells is key to predicting the possible success of a drug treatment and planning more effective and focused therapies. In this review, we describe how the modulation of enzymes involved in purine metabolism in tumour cells may affect the apoptotic programme. The enzymes discussed are: ectosolic and cytosolic 5'-nucleotidases, purine nucleoside phosphorylase, adenosine deaminase, hypoxanthine-guanine phosphoribosyltransferase, and inosine-5'-monophosphate dehydrogenase, as well as recently described enzymes particularly expressed in tumour cells, such as deoxynucleoside triphosphate triphosphohydrolase and 7,8-dihydro-8-oxoguanine triphosphatase.

Keywords: purine salvage; apoptosis; CD73; cN-II; ADA; PNP; HPRT; IMPDH; SAMHD1; MTH1

1. Introduction

Intracellular purine nucleotide concentration is determined and maintained through two distinct pathways, both depending on a common metabolite: 5-phosphoribosyl-1-pyrophosphate (PRPP) (Figure 1). The de novo pathway consists of 10 reactions catalysed by six enzymes, some of which are allosterically regulated mainly by PRPP and purine nucleotides [1]. Furthermore, the six enzymes can cluster near mitochondria and microtubules to form dynamic multienzyme complexes referred to as “purinosomes” [2]. The purinosome formation causes a strong increase in the rate of purine synthesis. This regulatory mechanism ensures that, during proliferation and in the absence of preformed purine ring to be salvaged, the supply of purine compounds is secured. When preformed purine rings are available, they can be converted in one step into the corresponding nucleoside monophosphates through the action of the salvage pathway enzymes adenine phosphoribosyltransferase (APRT) and hypoxanthine-guanine phosphoribosyltransferase (HPRT) which utilize PRPP as a co-substrate (Figure 1) [1]. The availability of preformed purine ring prevents purinosome formation [2]. Regulation both at genetic and protein levels ensures the correct amount of nucleotides to sustain replicative and metabolic needs [3]. Therefore, it is not surprising that alterations of replicative rate and eventually activation of apoptosis are a consequence of dysfunctions of purine metabolism. Furthermore, an

imbalance in purine supply can affect also mitochondrial proliferation and function, leading to metabolic changes and apoptosis [4]. Nevertheless, in some cases, the relationship between enzyme dysfunction and proliferation rate, metabolic alterations, and apoptosis is not as simple and direct as expected. The knowledge of the consequences of the alteration of enzymes involved in purine metabolism is essential not only for the understanding of “where and how” these enzymes impact metabolic pathways, but also to uncover whether they can be targets of antineoplastic drugs or be responsible for drug resistance. This review presents the most recent reports on the impact of purine catabolism and salvage enzymes (Figure 2) in the activation of apoptosis and the implied molecular mechanisms. Furthermore, the possible applications of this knowledge to anti-tumour therapy are discussed.




Figure 1. De novo and salvage pathways for purine nucleotide biosynthesis. Cyan background: de novo synthesis; yellow background: salvage synthesis. The figure outlines the central role played by PRPP, needed for both de novo and salvage pathways. Hyp: hypoxanthine; Gua: guanine; Ade: adenine; Rib-5-P: ribose-5-phosphate; PRPP: 5-phosphoribosyl-1-pyrophosphate; Gln: glutamine; Gly: glycine; THF: tetrahydrofolate; Asp: aspartate; S-AMP: succinyl-AMP; XMP: xanthosine-5'-monophosphate; IMP: inosine-5'-monophosphate; GMP: guanosine-5'-monophosphate; AMP: adenosine-5'-monophosphate.




Figure 2. Pathways of purine metabolism. 1: Ectosomal 5'-nucleotidase; 2: Adenosine deaminase; 3: Cytosolic 5'-nucleotidase II; 4: Purine nucleoside phosphorylase; 5: Hypoxanthine-guanine phosphoribosyltransferase; 6: IMP dehydrogenase. Inset: deoxyribonucleoside triphosphates (dNTP) are converted into the respective deoxynucleosides (dNs) by a one-step reaction catalysed by a sterile alpha motif and histidine-aspartate (HD) domain-containing protein 1 (enzyme 7). An increase in reactive oxygen species (ROS) brings about an increase in 8-oxo-dGTP, converted into the monophosphate by human MutT homolog 1 (enzyme 8). Ado: adenosine; Guo: guanosine; Ino: inosine.

2. Ectosomal 5'-Nucleotidase

Ectosomal 5'-nucleotidase (CD73) (EC 3.1.3.5) is a 70-kDa glycosylated protein bound to the outer surface of the plasma membrane by a glycosylphosphatidylinositol anchor. The enzyme is overexpressed in a variety of tumours [5] and in some of them, is associated with a highly invasive cancer phenotype, drug resistance, and tumour-promoting functions [6]. CD73 catalyses the dephosphorylation of extracellular AMP to adenosine (Ado), which plays important roles in many physiological and pathophysiological conditions through G-protein coupled Ado receptors (A1, A2A, A2B, and A3) [7,8]. The roles played by Ado are complex since its interaction with different receptors achieves different results; in particular, A1 and A3 receptors are coupled with G_i -proteins determining a decrease in cyclic AMP (cAMP), while A2A and A2B receptors are coupled with G_s -proteins causing an increase in intracellular cAMP [9] (Figure 3). Therefore, the role played by CD73 depends on the nature and distribution of the Ado receptors in a particular kind of cell. In addition, CD73 plays also

many different roles in cell physiology not related to its catalytic activity [10]. Analysing enzymatic and non-enzymatic functions of CD73, it was concluded that both faces were involved in the aggressive behaviour of cancer cells. The enzymatic function seems to be primarily involved in invasion, whereas the non-enzymatic action of the protein contributes to cell adhesion and migration through activation of focal adhesion kinase [10]. Among the tumours in which CD73 is upregulated, breast cancer is the most studied [11]. In some cases of this type of cancer, an involvement of CD73 enzymatic activity was demonstrated, since supply of Ado could be a substitute for enzyme upregulation in promoting proliferation and motility [12]. In other cases, the involvement of different not yet described mechanisms independent of enzyme activity was demonstrated [13]. In all cases, an activation of the Akt/GSK-3 β pathway was found to be involved in the tumour growth and motility promoted by CD73 [13]. In human cervical cancer cells, an increase in cell proliferation and motility was associated to CD73 overexpression, but the mechanism was independent of enzyme activity. In fact, CD73 inhibitors were unable to prevent the increase in proliferation in cells that overexpressed the enzyme. In addition, an increase in Ado, that could be expected when the enzyme is overexpressed, induced a decreased cell proliferation [14]. In this type of tumour, an activation of the Akt pathway was demonstrated as well [14]. In colorectal cancer, a downregulation of miR-30a was shown to determine an increase of CD73 expression in tumour cells, which promoted proliferation and inhibited apoptosis. MiR-30a is one of the most important tumour-suppressor factors in various human cancers and its level is significantly decreased in several tumours [15]. Since, as stated before, many effects exerted by the expression of CD73 on tumours are mediated by the participation of the enzyme activity in the conversion of extracellular ATP into Ado, such effects are dependent on the amount and nature of the Ado receptors expressed by tumour cells and other cells present in the tumour microenvironment. In this regard, in contrast to reports for other cancer types (see above), in an *in vivo* study on medulloblastoma, overexpression of CD73 reduced tumour growth and vascularization, and also promoted differentiation and initiated apoptosis, supposedly by the accumulation of Ado which interacted with A1 receptor [16]. The level of CD73 expression is very important not only for tumour growth and motility, but also for the success of the therapeutic approach. In fact, during chemotherapy, various immunogenic mediators accumulate in the tumour microenvironment, included ATP. Extracellular ATP, in this case, is released by the cells undergoing intrinsically or extrinsically activated apoptosis, through pannexin-1 channels and functions as a “find me” signal for P2Y family receptors expressed by macrophages and dendritic cells [17]. In this mechanism, the rapid ATP degradation catalysed by several extracellular enzymes is a determinant for the immunogenic activation, since the accumulated AMP, which cannot interact with P2Y receptors, can generate Ado through CD73, which mainly mediates the immune-escape of tumour cells interacting with A2A receptors [18]. Moreover, also in the case of regulatory T-cells (Tregs) in a tumour environment, the presence of several signals triggers apoptosis, and apoptotic Treg cells achieved superior immuno-suppression via an oxidative stress-associated mechanism [19]. Therefore, the induction of apoptosis in a tumour environment in which cells highly expressing CD73 and Ado receptors A2A are present is not always a successful therapeutic approach. Recently, CD73 has been targeted for the synthesis of new inhibitory compounds which prevent extracellular Ado formation from AMP and the consequent immune-escape [20].

3. Cytosolic 5'-Nucleotidase II

Cytosolic 5'-nucleotidase II (cN-II) (EC 3.1.3.5) is an ubiquitously expressed and highly conserved enzyme that hydrolyses purine nucleoside monophosphates (preferentially IMP and GMP) into their corresponding nucleosides and inorganic phosphate. Its enzymatic activity and biochemical features have been well characterised, and the reader is referred to a recent review for further investigation of these issues [21]. The first observation on its prognostic value in nucleoside analogue-treated patients with acute myeloid leukaemia indicated that patients with high cN-II expression in leukaemic blasts showed a poorer outcome with respect to those with a lower expression [22]. Since then, the implication of cN-II in cancer cells and in the response to anticancer treatment has been extensively

demonstrated [23,24]. Indeed, shRNA-based cell models with downregulated cN-II are more sensitive to purine nucleoside and nucleobase analogues, as compared to control cells [25]. The enzyme is highly expressed in tumour cells, and cN-II expression in human neuroblastoma cells and in lung cancer cells correlated with cell proliferation [26,27], whereas its inhibition in human breast cancer cells was associated with a better defence towards reactive oxygen species (ROS) and a better adaptability to glucose deprivation in culture media [28]. In a cell model of lung cancer (A549), an activation of p53 and inactivation of Akt following cN-II partial silencing was demonstrated [27]. Indeed, in an astrocytoma cell line (ADF), transitory cN-II silencing was followed by caspase-3 activation and apoptosis [29]. Furthermore, the siRNA-mediated inhibition of cN-II expression in murine skeletal muscles induced an increase in the AMP/ATP ratio and a subsequent activation of AMP-activated protein kinase (AMPK) [30], even though this was not confirmed in cN-II deficient mice [31]. In parallel to these biological studies, several genetic as well as genome wide association studies have identified the cN-II encoding gene *NT5C2* or some genetic variants as being associated to various pathological conditions such as hereditary spastic paraplegia 45, psychiatric disorders, a disturbance of blood pressure, and a decrease in body mass index [32]. Fluctuation of cN-II expression has been demonstrated to influence the concentration of intracellular nucleotides, depending on the type of cell and the technique utilised for enzyme silencing. In some cases, the alteration of nucleotide concentration is not statistically significant, but the effect on metabolic and proliferative features is still present [27,31,33,34]. These observations indicate that cN-II regulates several cellular pathways through a mechanism at least partially independent of its impact on intracellular nucleotide concentration.

4. Adenosine Deaminase

Adenosine deaminase (ADA) (E.C.3.5.4.4) catalyses the deamination of Ado and deoxyadenosine (dAdo) to inosine and deoxyinosine, respectively. There are two isoenzymes of ADA in human tissues, ADA1 and ADA2 [35]. ADA1 is ubiquitous, has a similar affinity for both substrates, and can interact with membrane proteins such as the dipeptidyl dipeptidase-4 (CD26), A1 [36] and A2A receptors [37]. ADA1 also acts as an ectoenzyme which catalyses the deamination of Ado and dAdo in the extracellular space. ADA2 is the main ADA isoenzyme found in human serum [35]. Low ADA activity was found in prostate [38] and gastric tumour tissues [39] and in lymphocytes of patients suffering from different pathologies, such as gynaecological [40], renal [41], head and neck [42], and gastric tumours [43,44], as well as Hodgkin's lymphoma [45]. Low ADA activity in lymphocytes, as outlined later on in this section, could account for the decreased cellular immune function in cancer patients. Conversely, ADA activity was increased in cancerous tissue from breast [46–48], kidney [49], and colorectal tumours [50], in serum of patients with bladder [51] ovarian [52], laryngeal [53], and head and neck squamous cell carcinomas [54,55] and in lymphocytes of patients with chronic lymphocytic leukaemia (CLL) [56]. High ADA activity might be advantageous to the cancer cells by causing, in association with purine nucleoside phosphorylase (PNP), an increase in hypoxanthine, a substrate for the salvage pathway (Figure 2). In addition, increased ADA activity might be a compensatory mechanism against toxic accumulation of its substrates. Indeed, Ado and dAdo are known to induce apoptosis, and ADA inhibition is an antitumoural strategy. Insights from the apoptosis-induced effect of ADA deficiency were obtained from studies regarding severe combined immunodeficiency (SCID) caused by mutations in the *ADA1* gene. The primary cause of lymphotoxicity in ADA-SCID is considered to be the accumulation of dAdo and dATP. In the absence of ADA activity, Ado and dAdo accumulate both in the extracellular compartments and inside the cells. dAdo is then phosphorylated by deoxycytidine kinase (dCK) and/or ADK to dAdo monophosphate, which in turn is converted to dAdo triphosphate (dATP). Intracellular dATP might generate DNA strand breaks and inhibit ribonucleotide reductase, leading to DNA synthesis impairment and apoptosis [57]. Deoxycoformycin (dCF), a powerful inhibitor of ADA [58], has been used alone or in combination with other drugs for the treatment of several types of lymphatic leukaemia [59–62]. dATP accumulation induced by dCF in

hairy cell leukaemia led to activation of p53, release of cytochrome c from mitochondria and activation of apoptotic protease-activating factor 1 (Apaf-1), and therefore caspase-9 and caspase-3 activation [63].

The combination of dAdo and dCF was found to be toxic for several tumoural cell lines such as rat hepatoma cells [64], and human colon carcinoma cell lines LoVo and HT29 [65–67]. The treatment with dAdo and dCF in combination resulted in the activation of the apoptotic mitochondrial pathway in LoVo, human astrocytoma, and neuroblastoma cell lines [67–70] with cytochrome c release and caspase-3 activation. Activation of caspase-8, and of both caspase-9 and -8 has also been found in astrocytoma and in neuroblastoma cells, respectively [69,70]. In astrocytoma cells, but not in neuroblastoma cells, a reduction in the production of lactate preceded the effect of dAdo and dCF on cell viability, suggesting a decreased glycolytic capacity. In both cell lines, dAdo must be phosphorylated in order to exert its cytotoxic effect; however, a decrease in the energy charge was observed in astrocytoma, but not in neuroblastoma cells [69,70].




Figure 3. Effectors of adenosine receptor-mediated apoptosis. The figure shows the different types of G-proteins associated with the four adenosine receptors and illustrates the apoptosis effectors found in several models by using agonists and antagonists of the receptors, or receptor silencing. The numbers in brackets refer to the respective reference. Note that adenosine receptors can be involved in survival in other cell types [71].

The involvement of the four Ado receptors in apoptosis of cancer cells has been reported and reviewed recently [71,72]. Apoptosis can occur through A1 [73–75], A2A [76–78], A2B [79,80], and A3 [81–89] receptors (Figure 3). Extracellular Ado can enter the cells through Ado transporters, and as intracellular Ado is converted by ADK to AMP which can activate AMPK, an energy sensor of the cells involved both in survival and cancer suppression (for a recent review see [90]). AMPK is responsible for apoptosis in some human gastric [91,92] cancer cells and astrocytoma cells among others [74]. In human mesothelioma cells this pathway led to upregulation of p53 [93]. p53 exerts its pro-apoptotic effect by transcription-dependent and transcription-independent actions. The targets of p53 transactivation are pro-apoptotic members of the Bcl-2 family (Bax, Bid, Puma, Noxa), as well as other apoptotic effector proteins (Apaf-1, caspase-8, caspase-6), cell death receptors, and cell death ligands. The p53 protein also acts directly in the mitochondria facilitating the oligomerization of Bax and Bak and interacting with anti-apoptotic Bcl-2, Bcl-xL, and Mcl-1 proteins [94]. The ADA inhibitor erythro-9-(2-hydroxy-3-nonyl)adenine (EHNA) but not dCF,

induced apoptosis in malignant pleural mesothelioma [95] by increasing intracellular Ado that needed to be converted to AMP, since ADK inhibition neutralised the effect. EHNA also inhibits cyclic nucleotide phosphodiesterase [96], therefore, it was postulated that this enzyme might have a role in the cytotoxicity observed in these cells. In addition, Ado-induced apoptosis involves the intrinsic and extrinsic pathways resulting in caspase-3 activation, and caspase-independent pathways leading to accumulation of Apoptosis Inducing Factor (AIF) (or its homologous AIF-homologous mitochondrion-associated inducer of death, AMID) in the nucleus [71,97]. AIF can be released from mitochondria, migrates to the nucleus where it can recruit nucleases, or organize a DNA-degrading complex [98]. In human hepatoma cells, Ado upregulated AMID, and promoted its translocation to the nucleus where it may induce DNA degradation [99]. Ado accumulation also reverses the action of S-adenosylhomocysteine hydrolase, increasing S-adenosylhomocysteine, which inhibits the transmethylation reactions. S-adenosylhomocysteine has been suggested to be involved in Ado-induced apoptosis in hepatoma HepG2 by altering gene expression [100]. It is interesting to note that dCF, through Ado receptor-dependent mechanisms, was also able to decrease the aggressiveness of cancer cells by modulating migration and invasion and by regulating endothelial cell permeability [101].

5. Purine Nucleoside Phosphorylase

Human purine nucleoside phosphorylase (hPNP) (EC 2.4.2.1), an ubiquitously expressed homotrimer, catalyses the phosphorolysis of 6-oxo-ribo- and 6-oxo-2'-deoxyribo-nucleosides to the corresponding bases and pentose-1-phosphate, but does not accept Ado and dAdo as substrates [102]. Conversely, the homoexameric *Escherichia coli* PNP (ePNP) efficiently acts on adenine-based nucleosides [103].

The importance of hPNP for T-cell functions was recognised more than 40 years ago [104] with the discovery that an inherited deficiency of this enzyme caused a severe T-cell lymphopaenia. In fact, the inhibition of PNP allows for the phosphorylation of 6-oxo-2'-deoxyribo-nucleosides which are channelled towards nucleotide synthesis instead of degradation. In particular, T-cell specific toxicity results from the inherently high phosphorylation of dGuo and the slow catabolism of dGMP in this cell type [105,106]. The major kinase activities responsible for the phosphorylation of dGuo appear to be cytosolic dCK and mitochondrial deoxyguanosine kinase [106]. dGMP is then converted to dGTP in two non-rate limiting kinase reactions. The overproduction of dGTP perturbs the deoxyribonucleotide pool, inhibits DNA synthesis, and induces cell death through mechanisms that will be discussed later on in this section. These observations provided a rationale for the development of hPNP inhibitors for the treatment of leukaemia. Among these inhibitors, forodesine (also known as immucillin H), a transition state analogue inhibitor of hPNP [107], has been thoroughly investigated as antineoplastic agent, in particular against T-cell mediated disorders. In PNP-deficient mice the mitochondrial deoxyguanosine kinase activity appears to be the responsible for the accumulation of dGTP in mitochondria, which leads to cell death with mitochondrial damage [108]. This abnormal accumulation interferes with mitochondrial DNA synthesis and repair [109], leading to p53 activation and apoptosis [110,111], possibly related to forodesine-induced ROS production and loss of mitochondrial membrane potential [112]. In vitro experiments performed with cultured human leukaemia cells demonstrated the toxicity of the combination of forodesine and dGuo [113,114]. In CLL, results indicate that forodesine induces apoptosis both through a p53-dependent [113] and a p53-independent [112] pathway. Balakrishnan et al. [113] reported studies carried out in leukaemic lymphocytes obtained from patients with CLL. The treatment with dGuo and forodesine in combination led to an increase in dGTP content, which correlated with a stabilization through Ser15 phosphorylation of p53, with an upregulation of downstream protein p21. A correlation between dGTP accumulation and caspase-8, -9, and -3 activation and subsequent polyADP-ribose polymerase (PARP) cleavage was also observed. Furthermore, all these effects appeared to be tumour-specific, as normal T and B lymphocytes isolated from healthy subjects did not undergo apoptosis. Alonso et al. [112] showed that forodesine in combination with dGuo induced a dose-dependent cell death in CLL cells, but the

response was independent of deletions in 17p13 (TP53) and 11q22-q23 (ATM), genetic aberrations acquired in advanced disease and associated with drug resistance and short survival [115]. Therefore, forodesine activated the mitochondrial apoptotic pathway, acting by a p53-independent mechanism in tumours with no functional p53. Interestingly, the authors found an upregulation of the p53 homolog p73 at transcriptional and translational level in CLL cells with p53 deletion, thus suggesting that this antileukaemic agent might be beneficial in CLL patients with impaired p53-dependent apoptotic pathway [115]. For the clinical practice of forodesine against lymphoblastic leukaemia, the reader is referred to previously published reviews and articles [116–118].

Gene-directed enzyme prodrug therapy (GDEPT) is a strategy of prodrug delivery which comprises a three component-system: an inactive drug (prodrug), a gene coding for the enzyme that converts the inactive prodrug into the active drug, and a vector [119]. For the choice of the vector and the specific gene delivery to the target tumour cell, the reader is referred to previously published reports [119,120]. In the early 1990s Parker et al. developed a gene therapy strategy based on the selective expression of the ePNP gene in tumour cells [121,122]. The bacterial enzyme, unlike mammalian PNP, can accept Ado and its analogues as substrates [103] and can be used to cleave nontoxic purine nucleoside analogues to very cytotoxic adenine analogues [122]. The toxic purine base analogues generated by ePNP readily diffuse across cell membrane, therefore killing not only the tumour cell in which they are generated, but also many surrounding tumour cells that do not express ePNP (bystander effect). The first prodrug used with ePNP was 9- β -D-[2-deoxyribofuranosyl]-6-methylpurine (MeP-dR), which, being a poor substrate for human salvage enzymes, such as dCK and hPNP, is not cytotoxic to human cells [123]. Therefore, its toxicity is exerted in tumour cells, where ePNP GDEPT is directed: in fact, 6-methylpurine (MeP), the product of MeP-dR phosphorolysis, is activated to the cytotoxic compound by cellular HPRT and/or APRT activities (see Section 6) and then incorporated both in RNA and DNA. Two other dAdo analogues, 2-fluoro-2'-deoxyadenosine (F-dAdo) and 9- β -arabinofuranosyl-2-fluoroadenine (F-araAde), have also received attention as substrates of ePNP. They are transformed by ePNP to 2-fluoroadenine (F-Ade), an adenine analogue approximately 100-fold more potent than MeP [122], which is converted to the cytotoxic mononucleotide by APRT. With F-araAde being an insoluble compound, it is administered as F-araAde-5'-monophosphate (F-araAMP), which must be dephosphorylated by plasma phosphatases, before entering the cell (Figure 4). An important feature of the ePNP GDEPT is its inhibition of protein and RNA/DNA synthesis, making it a good therapeutic approach for solid tumours, which exhibit a high fraction of non-replicating cells [123–125]. The ePNP/F-araAMP GDEPT in combination with docetaxel was used against multidrug-resistant ovarian cancer cells [126] and castration-resistant prostate cancer cells [127]. In both cases the involvement of apoptosis in tumour cell death was established. In particular, in ovarian cancer cells, a down-regulation of anti-apoptotic Bcl-2 protein and upregulation of pro-apoptotic Bax, Bik, and Bok proteins were achieved with the treatment. Furthermore, a caspase-mediated proteolysis of PARP was observed [126]. Krohne et al. [128] demonstrated that ePNP/F-araAMP induced hepatocellular carcinoma cell death predominantly by apoptosis, which occurred earlier in p53-positive HepG2 cells, as compared to p53-negative Hep3B cells. However, the efficiency of tumour cell death was similar in both cell lines. Moreover, they also demonstrated that the induction of apoptosis was independent of the Fas/FasL signalling pathway. They concluded that the ePNP GDEPT could be advantageous against tumours carrying p53 mutations or resistant to Fas-mediated apoptosis [128].




Figure 4. Metabolism of F-araAMP in normal and tumoural cells subjected to *Escherichia coli* PNP (ePNP)/F-araAMP GDEPT. 9- β -arabinofuranosyl-2-fluoroadenosine 5'-monophosphate (F-araAMP) is cleaved by plasma phosphatases into 9- β -arabinofuranosyl-2-fluoroadenine (F-araAde), which enters both normal and tumoural cells. Inside the cell, it is activated to the monophosphate by cellular deoxycytidine kinase (dCK). In tumour cells, the presence of ePNP allows for an additional activation pathway, which proceeds through a phosphorolytic cleavage to give 2-fluoroadenine (F-Ade), which is activated by cellular adenine phosphoribosyltransferase (APRT). Through the action of ribonucleotide reductase (RR), F-ADP is converted to the respective deoxynucleotide, thus interfering also on DNA synthesis. GDEPT: gene-directed enzyme prodrug therapy.

6. Hypoxanthine Guanine Phosphoribosyltransferase

HPRT (EC 2.4.2.8) plays a central role in the purine salvage pathway. The enzyme catalyses the conversion of guanine and hypoxanthine to the respective nucleoside monophosphates, by using PRPP as donor of the phosphoribosyl moiety [129]. Adenine is not a substrate of the enzyme; indeed, for the salvage of this purine base, an additional APRT activity (EC 2.4.2.7) is required [130]. The relevance of HPRT in the process of activation of thiopurines for the treatment of malignancies, rheumatic diseases, inflammatory bowel disease, and other pathologies is well documented [131,132]. In fact, HPRT catalyses the conversion of 6-thioguanine to 6-thioguanosine monophosphate and represents the first step towards the production of deoxy-6-thioguanosine 5'-triphosphate. This modified deoxynucleoside triphosphate is incorporated into DNA and is further methylated by S-adenosylmethionine to form S6-methylthioguanine, which, during DNA replication, pairs with thymine and its normal partner, cytosine [133]. The mispairs induced by 6-thioguanine are processed by the DNA mismatch repair leading to cell-cycle arrest followed by apoptosis [133,134]. In colorectal cancer cell lines proficient in mismatch repair, an interconnection between autophagy and apoptosis has been demonstrated. In fact, thiopurines induced autophagy as a survival mechanism which antagonises apoptosis. On the other hand, apoptosis was the major cell death mechanism, since caspase inhibition protected the cells from death induced by thiopurines. Interestingly, the inhibition of apoptosis promoted autophagic

response [135]. Thus, a tight interplay between apoptosis and autophagy controls cell fate in response to thiopurines. In fact, thiopurines induced mitochondrial depolarization and increased ROS production, which activated mitophagy for the degradation of damaged mitochondria. However, when mitochondrial depolarization exceeded the threshold for the activation of the cell death machinery, cells were directed to apoptotic death [135]. Therefore, ROS play a relevant role in thiopurine-induced cell death and represent a critical factor that links autophagy and apoptosis through mitochondria.

7. Inosine 5'-Monophosphate Dehydrogenase

Inosine 5'-monophosphate dehydrogenase (IMPDH) (EC 1.1.1.205) is a cytoplasmic enzyme that catalyses the NAD-dependent oxidation of inosine monophosphate (IMP) to xanthosine monophosphate (XMP), the first and rate-limiting step toward the de novo synthesis of GTP [136]. This enzyme plays a central role in purine metabolism, since guanylates are needed not only for nucleic acid synthesis, but also for cellular regulation, such as signal transduction, energy transfer, and microtubule dynamic instability [137]. Two isoforms of IMPDH have been identified: type 1 and type 2 [138]. Human IMPDH type 1 (hIMPDH1) is ubiquitously expressed and its activity is maintained at near constant level in normal and neoplastic cells, whereas the human IMPDH type 2 (hIMPDH2) isoform appears upregulated in proliferating cells [139]. The disproportionate increase of hIMPDH2 activity in neoplastic cells has made this isoform a key target for the development of anticancer drugs. The underlying rationale for the increased activity of hIMPDH2 may be that, although guanylates can be salvaged from guanine by HPRT, the level of circulating guanine is low in dividing cells and this route is probably insufficient to satisfy the needs of guanylates. Indeed, both natural and synthetic inhibitors of IMPDH are used as anticancer [139], antiviral [140], immunosuppressive, and antimicrobial agents [141–146]. In Tuberous Sclerosis Complex 2 (TSC-2) deficient cell and tumour models with aberrantly elevated mammalian Target of Rapamycin Complex 1 (mTORC1) signalling and actively producing ribosomes, IMPDH inhibition, by limiting the nucleotide pool, leads to selective replication stress, DNA damage, and apoptosis [147]. Moreover, a selective vulnerability to IMPDH inhibition was reported in a subset of small cell lung cancers, expressing high level of MYC, a transcription factor which appears to regulate purine biosynthesis by activating transcription of *IMPDH* [148]. Recently, it was found that an upregulation of IMPDH2 drives GTP biosynthesis, enhances rRNA and tRNA synthesis, and stimulates nucleolar hypertrophy in glioblastoma cells, thus connecting these features of malignant transformation [149]. Several research groups reported the non-enzymatic functions of IMPDH; IMPDH binds to DNA and RNA in vivo, in a manner independent of its catalytic activity [150]. Indeed, IMPDH was found in the nuclei of human cells, and several experiments suggest that IMPDH has a role in replication, transcription and translation. For example, IMPDH interacts with polyribosomes through a subdomain and is associated with translating rhodopsin mRNA [151]. Moreover, the evidence that recruitment of IMPDH to actively transcribed genes mediates C-terminal domain phosphorylation of RNA polymerase II in yeast supported a new cellular function of IMPDH [152]. IMPDH also acts as a cell-cycle-regulated transcription repressor. At the end of S-phase, the enzyme accumulates in the nucleus and represses proliferation genes such as those coding histones and E2f, a master driver of the G1/S transition [153]. In human embryonic kidney HEK293 and human colon cancer HCT116 cells, p53 regulated purine metabolism through a p53-miR-34a-IMPDH pathway (miR-34a is a master regulator of tumour suppression) and consequently, miR-34a-mediated inhibition of IMPDH perturbed the GTP-dependent Ras signalling pathway [137]. Inhibition of IMPDH activity has also been shown to induce differentiation in some cancer cell lines [154–157], while apoptosis was induced in other cell lines [158–161]. Over the past 30 years, several IMPDH inhibitors have been tested on many cancer models. Apoptosis triggered by these inhibitors occurs through multiple pathways depending on the type of compound and the used tumoural model. Figure 5 summarizes the apoptotic pathways triggered by the best known IMPDH inhibitors [136,137,162–170]. In all the experiments, apoptotic mechanisms were reversed by the simultaneous addition of guanosine, that, through the salvage pathway, circumvents the IMPDH inhibition by restoration of the guanylate pool.

(caspase-independent apoptosis) [136,162,163]; (3): upregulation of p53 with (a) downregulation of Bcl-2 and Bcl-xL, with consequent inhibition of p27 and survivin, cytochrome c (Cyt c) release, and activation of caspase-9, caspase-3 and polyADP-ribose polymerase (PARP; intrinsic apoptotic pathway) [159–161,164,165], (b) upregulation of PUMA and BIM with consequent SMAC/DIABLO release from mitochondria, inhibition of Inhibitor of Apoptosis (IAPs) (a caspase-3 inhibitor) with activation of caspase-3 [164], and (c) activation of caspase-2 with cleavage of Bid into truncated Bid (t-Bid) and AIF/Endo G release from mitochondria [164]; (4): synergistic effect of IMPDH inhibitors with TRAIL through binding with death receptors (DR4 and DR5) which recruit initiator caspase-8 via the adaptor protein FADD. Activated caspase-8 stimulates apoptosis via two parallel cascades: direct cleavage and activation of caspase-3, or cleavage of Bid into t-Bid which translocates to mitochondria, inducing cytochrome c release, with sequential activation of caspase-9 and -3 (extrinsic apoptotic pathway) [164]. TRAIL: tumour necrosis factor-related apoptosis-inducing ligand; FADD: Fas-associated protein with death domain.

8. Sterile Alpha Motif and HD Domain-Containing Protein 1

Sterile alpha motif and HD domain-containing protein 1 (SAMHD1) was identified as a mammalian deoxynucleoside triphosphate triphosphohydrolase (dNTPase, EC 3.1.5.-), which plays a key role in the regulation of dNTP homeostasis through hydrolysis of intracellular dNTPs to give a free deoxynucleoside and a triphosphate [171,172]. SAMHD1 has been also identified as an important host restriction factor that inhibits the infection of several retroviruses and DNA viruses by diminishing the intracellular dNTP pool needed for their replication [173]. SAMHD1 has also been implicated in the regulation of immune responses [174]. Homozygous mutations in *SAMHD1* gene have been identified in 17% of patients with Aicardi-Goutières syndrome (AGS), an autoimmune disorder that is attributed to excessive accumulation of small DNA fragments in brain. Additionally, homozygous deletion in the *SAMHD1* gene was also identified in atypical AGS patients [175]. *SAMHD1* somatic mutations have been identified in several cancers, including solid cancers such as glioblastoma [176], colorectal [177], lung [178], and pancreatic cancers [179] and blood-related malignancies such as CLL [180] and myeloma [181]. Additionally, SAMHD1 mRNA and/or protein expression is also significantly downregulated in CLL [180] and breast [182] and lung cancers [183]. Crystallographic studies have demonstrated that SAMHD1 is a homotetrameric enzyme allosterically activated by dNTPs [184]. Each monomer contains one catalytic site specific for dNTPs and two allosteric sites: site one has a very high affinity for GTP and dGTP and site two is specific for deoxyribose and binds all four dNTPs with lower affinity than site one [185]. The active site of SAMHD1 can accommodate various base modifications, indeed, SAMHD1 hydrolyses 2-amino-2'-dATP, O6-methyl-dGTP, 5-methyl-2'-dCTP and 2-thio-dTTP as well as the canonical substrates, suggesting a role of the enzyme in the metabolism of nucleoside analogues of therapeutic interest [186]. Furthermore, monomeric and dimeric SAMHD1 bind to single-stranded nucleic acids [187,188], while the tetrameric form of the enzyme is active as a dNTPase [188]. On this basis, Seamon et al. [188] proposed a model in which the level of cellular dNTPs operates as a switch between the two types of SAMHD1 activity. Under conditions of low dNTP concentrations, the monomeric and dimeric forms would prevail, with high affinity for single-stranded RNA, whereas when dNTP concentrations are high, the tetrameric dNTPase would be predominant. These observations suggest that the single-stranded nucleic acid binding activity of SAMHD1, in addition to its dNTPase action, are likely determinants involved in retrovirus restriction and immune activation. Recent progress indicates that gene mutations and epigenetic mechanisms lead to downregulation of SAMHD1 activity or expression in multiple cancers. Impaired SAMHD1 function can cause increased dNTP pool resulting in genomic instability and cell-cycle progression, thereby facilitating cancer cell proliferation [189]. In line with this conclusion, exogenous expression of SAMHD1 inhibited proliferation and induced apoptosis in cutaneous T-cell lymphoma-derived HuT78 cells, through increased activation of extrinsic apoptotic signalling mediators, and sensitised the cells to FasL-stimulated apoptosis [190]. In addition, in an acute myeloid leukaemia-derived human

leukaemic cell line (THP 1), *SAMHD1* knockout induced cell proliferation and reduced apoptosis and this effect was in part due to activation of PI3K-Akt-p27 signalling axis. Furthermore, the *SAMHD1* knockout attenuated the ability of THP-1 cells to form subcutaneous tumour cells in xenografted immunodeficient mice. This effect correlated with significantly increased expression of tumour necrosis factor α (TNF- α) in cancer, which may suggest that TNF- α -mediated inflammation could account for the decreased tumorigenicity in vivo [191].

9. Human MutT Homolog 1

Human MutT homolog 1 (MTH1) is a widely expressed monomeric enzyme that hydrolyses oxidised purine nucleoside triphosphates, such as 8-oxo-dGTP, 8-oxo-dATP and 2-hydroxy-dATP, to monophosphates and pyrophosphate (7,8-dihydro-8-oxoguanine triphosphatase, EC 3.6.1.55) [192]. It is mostly present in cytosol and, in lesser amounts, in mitochondria and in nuclei. Higher expression levels of the enzyme were found in thymus, testis, embryo, proliferating blood lymphocytes [193], and in cancer cells when compared with normal ones [194]. Its enzymatic activity, together with its cellular distribution and the overexpression in cancers, lead to hypothesise for MTH1 a possible role in the sanitization of nucleotide pools both for nuclear and mitochondrial DNA/RNA replication and transcription processes, preventing the misincorporation of oxidised nucleotides into nucleic acids (Figure 6, Panel 1). In fact, cancer cells are characterised by a marked alteration of the redox status with a consequent increase in ROS [195]. A small increase in ROS levels promotes proliferation and cancer cell survival [196,197], whereas large amounts of ROS inhibit proliferation [198] and promote senescence [199,200] and apoptosis in several cancer models [200–203]. While it is widely accepted that MTH1 is dispensable in healthy cells [194], the role of the enzyme in removing oxidised dNTP in cancer cells, thus promoting their survival, and the consequent possibility to use this enzyme as a chemotherapeutic target, is controversial. These discordances, arising probably from the use in the different reports of different experimental conditions (inhibitors, siRNA, shRNA, or CRISPR technique) and models (cell cultures, xenograft, etc.), have been reviewed in the past few years and the reader is referred to these extensive reviews for a close examination [194,204–206]. We will focus on the mechanisms by which targeting MTH1 could lead, according to the supporters of this thesis, to apoptosis and/or senescence of cancer cells. Downregulation or inhibition of MTH1 leads to the incorporation in nuclear and mitochondrial DNA of oxidised nucleotides (Figure 6, Panel 2), which may be removed by a base excision repair (BER) process [194,207]. It has been proposed that wild type p53 is able to stimulate the BER response at different levels [208–210]. However, the excessive incorporation in DNA of oxidised nucleotides, may cause an overload of the finely coordinated BER system, leading eventually to the production of either abasic sites or single strand brakes or double strand brakes [194]. A connection between this DNA damage and different detrimental effects (stop of cell cycle in G1, senescence, apoptosis p53, caspase-3 mediated or parthanatos) has been proposed [111,194,200,206,211] (Figure 6, panel 2a). Conversely, if the BER process does not work properly, or in models containing null or mutant p53 [212,213], tumorigenesis can be promoted via the accumulation of mutations caused by the mispairing of 8-oxo-dGTP and 2-hydroxy-dATP (Figure 6, Panel 2b). In addition, 8-oxo-dGTP promotes chain termination following insertion by telomerase into the repeated telomeric sequence TTAGGG [214], and in oxidant conditions, cancer cells with very short telomers are quite sensitive to MTH1 downregulation [214] (Figure 6, Panel 2c).




Figure 6. Scheme of possible effects of therapeutic use of human MutT homolog 1 (MTH1). Panel 1: Cancer cell proliferation is stimulated by the accumulation of ROS, which favours the formation of 8-oxo-dGTP and 2-hydroxy-dATP. In order to avoid the potentially toxic effects of the incorporation of oxo-derivatives of nucleotides, the cells upregulate MTH1 sanitizing enzyme. Panel 2: downregulation or inhibition of MTH1 leads to an abnormal incorporation of oxidised nucleotides in DNA. Panel 2a: in order to deal with this excessive accumulation of oxo-nucleotides in DNA, the base excision repair (BER) system is, in general, upregulated. If the misincorporation of oxo-nucleotides exceeds the capability of BER, abasic sites (AS) or single strand breaks (SSB) or double strand breaks (DSB) accumulate, leading to several serious consequences. Panel 2b: if BER is downregulated, incorporated oxo-nucleotides are not repaired. In this case no cytotoxic effect is reported, but a promotion of tumourigenesis can occur. Panel 2c: the misincorporation of 8-oxo-dGTP by telomerase into the repeated telomeric sequence TTAGGG leads to the inhibition of the polymerase activity of the enzyme.

10. Concluding Remarks

A correct balance of NTPs and dNTPs is necessary for the prevention of multiple pathologies. A healthy cell must maintain two asymmetric and spatial-temporal dNTP pools; one for nuclear DNA (nDNA) synthesis and repair and another for mitochondrial DNA (mDNA) replication and repair. Disruptions in dNTP balance are associated with enhanced mutagenesis, leading to genomic instability which promotes cancer [215] and may have a role in metabolic disease [216]. Cytosolic dNTP pool concentrations positively correlate with the cell cycle. In fact, the quantity of dNTPs at the beginning of S-phase is not sufficient for a complete DNA duplication [217]. The increase of dNTPs during the S-phase is necessary for faithful nDNA replication. mDNA is replicated continuously in post-mitotic cells, and faithful maintenance of mDNA also depends on correctly balanced dNTPs [218]. Thus, both proliferating and non-proliferating cells need to fine-tune nucleotide and dNTP synthesis to allow for both nDNA and mDNA replication and repair to maintain the health of the cell. Indeed, starvation or mTORC1 inhibition leads to selective autophagy of ribosomes, providing a source of nucleosides to be used by the salvage pathway enzymes for the synthesis of nucleotides, thus contributing to starved cell survival [219]. An imbalance of NTPs may have metabolic consequences and also generate an imbalance of dNTPs that can be synthesised by salvage of deoxynucleosides and by NDP reduction catalysed by ribonucleotide reductase. A plethora of signals concur to activate DNA repair mechanisms or to promote apoptosis in cells in which NTP pools are altered. The knowledge of the pathways linking enzyme expression or inhibition and apoptosis is fundamental to find new chemotherapeutic approaches able to activate apoptosis in cancer cells. Enzyme inhibition or silencing techniques were utilised to demonstrate the involvement of several interconverting or salvage enzymes in the determination of nucleotide imbalance. Some enzymes described in this review have been demonstrated or suspected to also have a role distinct from catalytic activity. In fact, ectosolic 5'-nucleotidase promotes cancer not only through the production of ectosolic Ado: cytosolic 5'-nucleotidase regulates proliferation and apoptosis in a way not so easily attributable to its regulation of intracellular nucleotide concentration, IMPDH also acts as transcription factor, and, finally, SAMHD1 binds to single-stranded RNA. All these observations make it difficult to directly link enzyme dysfunction with NTP imbalance and activation of apoptosis. Unfortunately, in many papers the nucleotide imbalance was inferred by the alteration of enzyme expression or inhibition and the consequent activation of the apoptotic programme, while the effective nucleotide concentration was not measured. Furthermore, in the case of enzymes involved in the Ado metabolism it is not so easy to distinguish between the effects exerted by Ado itself and those mediated by the interaction with its receptors. In addition, dAdo that does not interact with Ado receptors is able to trigger apoptosis through the formation of dATP and possibly through other still unknown mechanisms. In any case, the knowledge of the catalytic capacity of the enzymes involved in the pathways of Figure 2 and of their substrate specificity and regulation opened the way to novel pharmacological approaches to cancer therapy. Furthermore, the sometimes-surprising results obtained by their inhibition or silencing indicate that much more needs to be unravelled about purine salvage and interconverting pathways.

Author Contributions: Conceptualization, M.C., M.G.-G., and M.G.T.; Writing—Original Draft Preparation, M.G.-G., R.P., S.A., and M.G.T.; Writing—Review and Editing, M.C.

Funding: This work was supported by local funding from the University of Pisa (ex 60%) to M.C., M.G.-G. and M.G.T.

Acknowledgments: We wish to thank A. Wallwork for his helpful reading of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Ipata, P.L.; Balestri, F.; Camici, M.; Tozzi, M.G. Molecular mechanisms of nucleoside recycling in the brain. *Int. J. Biochem. Cell Biol.* **2011**, *43*, 140–145. [[CrossRef](#)] [[PubMed](#)]

2. An, S.; Kumar, R.; Sheets, E.D.; Benkovic, S.J. Reversible compartmentalization of de novo purine biosynthetic complexes in living cells. *Science* **2008**, *320*, 103–106. [[CrossRef](#)] [[PubMed](#)]
3. Pedley, A.M.; Benkovic, S.J. A new view into the regulation of purine metabolism: The purinosome. *Trends Biochem. Sci.* **2017**, *42*, 141–154. [[CrossRef](#)] [[PubMed](#)]
4. Rampazzo, C.; Miazzi, C.; Franzolin, E.; Pontarin, G.; Ferraro, P.; Frangini, M.; Reichard, P.; Bianchi, V. Regulation by degradation, a cellular defense against deoxyribonucleotide pool imbalances. *Mutat. Res.* **2010**, *703*, 2–10. [[CrossRef](#)] [[PubMed](#)]
5. Zhang, B. CD73 promotes tumor growth and metastasis. *Oncoimmunology* **2012**, *1*, 67–70. [[CrossRef](#)] [[PubMed](#)]
6. Gao, Z.W.; Dong, K.; Zhang, H.Z. The roles of CD73 in cancer. *Biomed. Res. Int.* **2014**, *2014*, 460654. [[CrossRef](#)] [[PubMed](#)]
7. Regateiro, F.S.; Cobbold, S.P.; Waldmann, H. CD73 and adenosine generation in the creation of regulatory microenvironments. *Clin. Exp. Immunol.* **2013**, *171*, 1–7. [[CrossRef](#)] [[PubMed](#)]
8. Kazemi, M.H.; Raoofi Mohseni, S.; Hojjat-Farsangi, M.; Anvari, E.; Ghalamfarsa, G.; Mohammadi, H.; Jadidi-Niaragh, F. Adenosine and adenosine receptors in the immunopathogenesis and treatment of cancer. *J. Cell Physiol.* **2018**, *233*, 2032–2057. [[CrossRef](#)] [[PubMed](#)]
9. Sousa, J.B.; Fresco, P.; Diniz, C.; Goncalves, J. Adenosine receptor ligands on cancer therapy: A review of patent literature. *Recent Pat. Anticancer Drug Discov.* **2018**, *13*, 40–69. [[CrossRef](#)] [[PubMed](#)]
10. Sadej, R.; Skladanowski, A.C. Dual, enzymatic and non-enzymatic, function of ecto-5'-nucleotidase (eN, CD73) in migration and invasion of A375 melanoma cells. *Acta Biochim. Pol.* **2012**, *59*, 647–652. [[CrossRef](#)]
11. Stagg, J.; Divisekera, U.; McLaughlin, N.; Sharkey, J.; Pommey, S.; Denoyer, D.; Dwyer, K.M.; Smyth, M.J. Anti-CD73 antibody therapy inhibits breast tumor growth and metastasis. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 1547–1552. [[CrossRef](#)] [[PubMed](#)]
12. Yang, X.; Pei, S.; Wang, H.; Jin, Y.; Yu, F.; Zhou, B.; Zhang, H.; Zhang, D.; Lin, D. Tiamulin inhibits breast cancer growth and pulmonary metastasis by decreasing the activity of CD73. *BMC Cancer* **2017**, *17*, 255. [[CrossRef](#)] [[PubMed](#)]
13. Yu, J.; Wang, X.; Lu, Q.; Wang, J.; Li, L.; Liao, X.; Zhu, W.; Lv, L.; Zhi, X.; Yu, J.; et al. Extracellular 5'-nucleotidase (CD73) promotes human breast cancer cells growth through AKT/GSK-3beta/beta-catenin/cyclinD1 signaling pathway. *Int. J. Cancer* **2018**, *142*, 959–967. [[CrossRef](#)] [[PubMed](#)]
14. Gao, Z.W.; Wang, H.P.; Lin, F.; Wang, X.; Long, M.; Zhang, H.Z.; Dong, K. CD73 promotes proliferation and migration of human cervical cancer cells independent of its enzyme activity. *BMC Cancer* **2017**, *17*, 135. [[CrossRef](#)] [[PubMed](#)]
15. Lu, N.; Lin, T.; Wang, L.; Qi, M.; Liu, Z.; Dong, H.; Zhang, X.; Zhai, C.; Wang, Y.; Liu, L.; et al. Association of SOX4 regulated by tumor suppressor miR-30a with poor prognosis in low-grade chondrosarcoma. *Tumour Biol.* **2015**, *36*, 3843–3852. [[CrossRef](#)]
16. Cappellari, A.R.; Pillat, M.M.; Souza, H.D.; Dietrich, F.; Oliveira, F.H.; Figueiro, F.; Abujamra, A.L.; Roesler, R.; Lecka, J.; Sevigny, J.; et al. Ecto-5'-Nucleotidase overexpression reduces tumor growth in a xenograph medulloblastoma model. *PLoS ONE* **2015**, *10*, e0140996. [[CrossRef](#)]
17. Boyd-Tressler, A.M.; Lane, G.S.; Dubyak, G.R. Up-Regulated ectonucleotidases in fas-associated death domain protein- and receptor-interacting protein kinase 1-deficient jurkat leukemia cells counteract extracellular ATP/AMP accumulation via Pannexin-1 channels during chemotherapeutic drug-induced apoptosis. *Mol. Pharm.* **2017**, *92*, 30–47. [[CrossRef](#)]
18. Antonioli, L.; Blandizzi, C.; Pacher, P.; Hasko, G. Immunity, inflammation and cancer: A leading role for adenosine. *Nat. Rev. Cancer* **2013**, *13*, 842–857. [[CrossRef](#)]
19. Maj, T.; Wang, W.; Crespo, J.; Zhang, H.; Wang, W.; Wei, S.; Zhao, L.; Vatan, L.; Shao, I.; Szeliga, W.; et al. Oxidative stress controls regulatory T cell apoptosis and suppressor activity and PD-L1-blockade resistance in tumor. *Nat. Immunol.* **2017**, *18*, 1332–1341. [[CrossRef](#)]
20. Dumontet, C.; Peyrottes, S.; Rabeson, C.; Cros-Perrial, E.; Geant, P.Y.; Chaloin, L.; Jordheim, L.P. CD73 inhibition by purine cytotoxic nucleoside analogue-based diphosphonates. *Eur. J. Med. Chem.* **2018**, *157*, 1051–1055. [[CrossRef](#)]
21. Tozzi, M.G.; Pesi, R.; Allegrini, S. On the physiological role of cytosolic 5'-nucleotidase II (cN-II): Pathological and therapeutical implications. *Curr. Med. Chem.* **2013**, *20*, 4285–4291. [[CrossRef](#)] [[PubMed](#)]

22. Galmarini, C.M.; Graham, K.; Thomas, X.; Calvo, F.; Rousselot, P.; El Jafaari, A.; Cros, E.; Mackey, J.R.; Dumontet, C. Expression of high Km 5'-nucleotidase in leukemic blasts is an independent prognostic factor in adults with acute myeloid leukemia. *Blood* **2001**, *98*, 1922–1926. [[CrossRef](#)] [[PubMed](#)]
23. Cividini, F.; Pesi, R.; Chaloin, L.; Allegrini, S.; Camici, M.; Cros-Perrial, E.; Dumontet, C.; Jordheim, L.P.; Tozzi, M.G. The purine analog fludarabine acts as a cytosolic 5'-nucleotidase II inhibitor. *Biochem. Pharm.* **2015**, *94*, 63–68. [[CrossRef](#)] [[PubMed](#)]
24. Jordheim, L.P.; Chaloin, L. Therapeutic perspectives for cN-II in cancer. *Curr. Med. Chem.* **2013**, *20*, 4292–4303. [[CrossRef](#)] [[PubMed](#)]
25. Jordheim, L.P.; Puy, J.Y.; Cros-Perrial, E.; Peyrottes, S.; Lefebvre, I.; Perigaud, C.; Dumontet, C. Determination of the enzymatic activity of cytosolic 5'-nucleotidase cN-II in cancer cells: Development of a simple analytical method and related cell line models. *Anal. Bioanal. Chem.* **2015**, *407*, 5747–5758. [[CrossRef](#)] [[PubMed](#)]
26. Cividini, F.; Cros-Perrial, E.; Pesi, R.; Machon, C.; Allegrini, S.; Camici, M.; Dumontet, C.; Jordheim, L.P.; Tozzi, M.G. Cell proliferation and drug sensitivity of human glioblastoma cells are altered by the stable modulation of cytosolic 5'-nucleotidase II. *Int. J. Biochem. Cell Biol.* **2015**, *65*, 222–229. [[CrossRef](#)] [[PubMed](#)]
27. Pesi, R.; Petrotto, E.; Colombaioni, L.; Allegrini, S.; Garcia-Gil, M.; Camici, M.; Jordheim, L.P.; Tozzi, M.G. Cytosolic 5'-Nucleotidase II silencing in a human lung carcinoma cell line opposes cancer phenotype with a concomitant increase in p53 phosphorylation. *Int. J. Mol. Sci.* **2018**, *19*, 2115. [[CrossRef](#)] [[PubMed](#)]
28. Bricard, G.; Cadassou, O.; Cassagnes, L.E.; Cros-Perrial, E.; Payen-Gay, L.; Puy, J.Y.; Lefebvre-Tournier, I.; Tozzi, M.G.; Dumontet, C.; Jordheim, L.P. The cytosolic 5'-nucleotidase cN-II lowers the adaptability to glucose deprivation in human breast cancer cells. *Oncotarget* **2017**, *8*, 67380–67393. [[CrossRef](#)]
29. Careddu, M.G.; Allegrini, S.; Pesi, R.; Camici, M.; Garcia-Gil, M.; Tozzi, M.G. Knockdown of cytosolic 5'-nucleotidase II (cN-II) reveals that its activity is essential for survival in astrocytoma cells. *Biochim. Biophys. Acta* **2008**, *1783*, 1529–1535. [[CrossRef](#)]
30. Kulkarni, S.S.; Karlsson, H.K.; Szekeres, F.; Chibalin, A.V.; Krook, A.; Zierath, J.R. Suppression of 5'-nucleotidase enzymes promotes AMP-activated protein kinase (AMPK) phosphorylation and metabolism in human and mouse skeletal muscle. *J. Biol. Chem.* **2011**, *286*, 34567–34574. [[CrossRef](#)]
31. Kviklyte, S.; Vertommen, D.; Yerna, X.; Andersen, H.; Xu, X.; Gailly, P.; Bohlooly, Y.M.; Oscarsson, J.; Rider, M.H. Effects of genetic deletion of soluble 5'-nucleotidases NT5C1A and NT5C2 on AMPK activation and nucleotide levels in contracting mouse skeletal muscles. *Am. J. Physiol. Endocrinol. Metab.* **2017**, *313*, e48–e62. [[CrossRef](#)] [[PubMed](#)]
32. Jordheim, L.P. Expanding the clinical relevance of the 5'-nucleotidase cN-II/NT5C2. *Purinergic Signal.* **2018**, *14*, 321–329. [[CrossRef](#)] [[PubMed](#)]
33. Cividini, F.; Filoni, D.N.; Pesi, R.; Allegrini, S.; Camici, M.; Tozzi, M.G. IMP-GMP specific cytosolic 5'-nucleotidase regulates nucleotide pool and prodrug metabolism. *Biochim. Biophys. Acta* **2015**, *1850*, 1354–1361. [[CrossRef](#)] [[PubMed](#)]
34. Allegrini, S.; Filoni, D.N.; Galli, A.; Collavoli, A.; Pesi, R.; Camici, M.; Tozzi, M.G. Expression of bovine cytosolic 5'-Nucleotidase (cN-II) in yeast: Nucleotide pools disturbance and its consequences on growth and homologous recombination. *PLoS ONE* **2013**, *8*, e63914. [[CrossRef](#)] [[PubMed](#)]
35. Gakis, C. Adenosine deaminase (ADA) isoenzymes ADA1 and ADA2: Diagnostic and biological role. *Eur. Respir. J.* **1996**, *9*, 632–633. [[CrossRef](#)] [[PubMed](#)]
36. Franco, R.; Casado, V.; Ciruela, F.; Saura, C.; Mallol, J.; Canela, E.I.; Lluís, C. Cell surface adenosine deaminase: Much more than an ectoenzyme. *Prog. Neurobiol.* **1997**, *52*, 283–294. [[CrossRef](#)]
37. Moreno, E.; Canet, J.; Gracia, E.; Lluís, C.; Mallol, J.; Canela, E.I.; Cortes, A.; Casado, V. Molecular evidence of adenosine deaminase linking adenosine A2A receptor and CD26 proteins. *Front. Pharm.* **2018**, *9*, 106. [[CrossRef](#)] [[PubMed](#)]
38. Biri, H.; Ozturk, S.; Kacmaz, M.; Karaca, K.; Tokucoglu, H.; Durak, I. Activities of DNA turnover and free radical metabolizing enzymes in cancerous human prostate tissue. *Cancer Investig.* **1999**, *17*, 314–319. [[CrossRef](#)]
39. Namiot, Z.; Stasiewicz, J.; Namiot, A.; Kemon, A.; Kralisz, M.; Gorski, J. Adenosine deaminase activity in patients with the intestinal type of gastric carcinoma. *Cancer Lett.* **1996**, *109*, 199–202. [[CrossRef](#)]
40. Specchia, G.; Pavone, V.; Maggio, F.; Lojudice, L.; Iacobazzi, A.; Detullio, L.; Cagnazzo, G.; Liso, V. Adenosine-Deaminase activity in peripheral lymphocytes of patients with gynecologic malignancies. *Boll. Inst. Sieroter. Milan.* **1985**, *64*, 404–407.

41. Sufrin, G.; Tritsch, G.L.; Mittelman, A.; Moore, R.H.; Murphy, G.P. Adenosine-Deaminase activity in patients with renal adenocarcinoma. *Cancer* **1977**, *40*, 796–802. [[CrossRef](#)]
42. Dasmahapatra, K.S.; Hill, H.Z.; Dasmahapatra, A.; Suarez, S. Evaluation of adenosine-deaminase activity in patients with head and neck-cancer. *J. Surg. Res.* **1986**, *40*, 368–373. [[CrossRef](#)]
43. Kojima, O.; Majima, T.; Uehara, Y.; Yamane, T.; Fujita, Y.; Takahashi, T.; Majima, S. Alteration of adenosine-deaminase levels in peripheral-blood lymphocytes of patients with gastric-cancer. *Jpn. J. Surg.* **1985**, *15*, 130–133. [[CrossRef](#)]
44. Russo, M.; Giancane, R.; Apice, G.; Galanti, B. Adenosine-Deaminase and purine nucleoside phosphorylase activities in peripheral lymphocytes from patients with solid tumors. *Br. J. Cancer* **1981**, *43*, 196–200. [[CrossRef](#)]
45. Murray, J.L.; Perezsoler, R.; Bywaters, D.; Hersh, E.M. Decreased adenosine-deaminase (Ada) and 5′nucleotidase (5nt) activity in peripheral-blood T-Cells in Hodgkin disease. *Am. J. Hematol.* **1986**, *21*, 57–66. [[CrossRef](#)]
46. Camici, M.; Tozzi, M.G.; Allegrini, S.; Delcorso, A.; Sanfilippo, O.; Daidone, M.G.; Demarco, C.; Ipata, P.L. Purine salvage enzyme-activities in normal and neoplastic human tissues. *Cancer Biochem. Bioph.* **1990**, *11*, 201–209.
47. Aghaei, M.; Karami-Tehrani, F.; Salami, S.; Atri, M. Adenosine deaminase activity in the serum and malignant tumors of breast cancer: The assessment of isoenzyme ADA1 and ADA2 activities. *Clin. Biochem.* **2005**, *38*, 887–891. [[CrossRef](#)]
48. Mahajan, M.; Tiwari, N.; Sharma, R.; Kaur, S.; Singh, N. Oxidative stress and its relationship with adenosine deaminase activity in various stages of breast cancer. *Indian J. Clin. Biochem.* **2013**, *28*, 51–54. [[CrossRef](#)]
49. Durak, I.; Beduk, Y.; Kavutcu, M.; Suzer, O.; Yaman, O.; Ozturk, H.S.; Canbolat, O.; Ulutepe, S. Activity of the enzymes participating in purine metabolism of cancerous and noncancerous human kidney tissues. *Cancer Invest.* **1997**, *15*, 212–216. [[CrossRef](#)]
50. Eroglu, A.; Canbolat, O.; Demirci, S.; Kocaoglu, H.; Eryavuz, Y.; Akgul, H. Activities of adenosine deaminase and 5′-nucleotidase in cancerous and noncancerous human colorectal tissues. *Med. Oncol.* **2000**, *17*, 319–324. [[CrossRef](#)]
51. Pirincci, N.; Gecit, I.; Gunes, M.; Yuksel, M.B.; Kaba, M.; Tanik, S.; Demir, H.; Aslan, M. Serum adenosine deaminase, catalase and carbonic anhydrase activities in patients with bladder cancer. *Clinics* **2012**, *67*, 1443–1446. [[CrossRef](#)]
52. Urunsak, I.F.; Gulec, U.K.; Paydas, S.; Seydaoglu, G.; Guzel, A.B.; Vardar, M.A. Adenosine deaminase activity in patients with ovarian neoplasms. *Arch. Gynecol. Obs.* **2012**, *286*, 155–159. [[CrossRef](#)]
53. Sharma, S.D.; Desai, P.B.; Metgudmath, R.B. Evaluation of serum adenosine deaminase and retinol in patients with laryngeal cancer. *Indian J. Pharm. Biol. Res.* **2013**, *1*, 5. [[CrossRef](#)]
54. Lal, H.; Munjal, S.K.; Wig, U.; Saini, A.S. Serum enzymes in head and neck cancer III. *J. Laryngol. Otol.* **1987**, *101*, 1062–1065. [[CrossRef](#)]
55. Mishra, R.; Agarwal, M.K.; Chansuria, J.P. Serum adenosine deaminase levels as an index of tumor growth in head and neck malignancy. *Indian J. Otolaryngol. Head Neck Surg.* **2000**, *52*, 360–363. [[CrossRef](#)]
56. Ghaderi, B.; Amini, S.; Maroofi, F.; Jalali, C.; Javanmardi, M.; Roshani, D.; Abdi, M. Adenosine deaminase activity in chronic lymphocytic leukemia and healthy subjects. *Iran. J. Cancer Prev.* **2016**, *9*, e5069. [[CrossRef](#)]
57. Whitmore, K.V.; Gaspar, H.B. Adenosine deaminase deficiency—More than just an immunodeficiency. *Front. Immunol.* **2016**, *7*, 314. [[CrossRef](#)]
58. Agarwal, R.P. Recovery of 2′-deoxycoformycin-inhibited adenosine deaminase of mouse erythrocytes and leukemia L1210 in vivo. *Cancer Res.* **1979**, *39*, 1425–1427.
59. Dohner, H.; Ho, A.D.; Thaler, J.; Stryckmans, P.; Sonneveld, P.; de Witte, T.; Lechner, K.; Lauria, F.; Bodewadt-Radzun, S.; Suci, S.; et al. Pentostatin in prolymphocytic leukemia: Phase II trial of the European organization for research and treatment of cancer leukemia cooperative study group. *J. Natl. Cancer Inst.* **1993**, *85*, 658–662. [[CrossRef](#)]
60. Willis, C.R.; Goodrich, A.; Park, K.; Waselenko, J.K.; Lucas, M.; Reese, A.; Diehl, L.F.; Grever, M.R.; Byrd, J.C.; Flinn, I.W. A phase I/II study examining pentostatin, chlorambucil, and theophylline in patients with relapsed chronic lymphocytic leukemia and non-Hodgkin’s lymphoma. *Ann. Hematol.* **2006**, *85*, 301–307. [[CrossRef](#)]
61. Kay, N.E.; LaPlant, B.R.; Pettinger, A.M.; Call, T.G.; Leis, J.F.; Ding, W.; Parikh, S.A.; Conte, M.J.; Bowen, D.A.; Shanafelt, T.D. Cumulative experience and long term follow-up of pentostatin-based chemoimmunotherapy trials for patients with chronic lymphocytic leukemia. *Expert Rev. Hematol.* **2018**, *11*, 337–349. [[CrossRef](#)]

62. Tedeschi, A.; Rossi, D.; Motta, M.; Quaresmini, G.; Rossi, M.; Coscia, M.; Anastasia, A.; Rossini, F.; Cortelezzi, A.; Nador, G.; et al. A phase II multi-center trial of pentostatin plus cyclophosphamide with ofatumumab in older previously untreated chronic lymphocytic leukemia patients. *Haematologica* **2015**, *100*, e501–e504. [[CrossRef](#)]
63. Johnston, J.B. Mechanism of action of pentostatin and cladribine in hairy cell leukemia. *Leuk. Lymphoma* **2011**, *52*, 43–45. [[CrossRef](#)]
64. Hunt, S.W., 3rd; Hoffee, P.A. Adenosine deaminase from deoxycoformycin-sensitive and -resistant rat hepatoma cells. Purification and characterization. *J. Biol. Chem.* **1982**, *257*, 14239–14244.
65. Camici, M.; Turriani, M.; Tozzi, M.G.; Turchi, G.; Cos, J.; Alemany, C.; Miralles, A.; Noe, V.; Ciudad, C.J. Purine enzyme profile in human colon-carcinoma cell-lines and differential sensitivity to deoxycoformycin and 2'-Deoxyadenosine in combination. *Int. J. Cancer* **1995**, *62*, 176–183. [[CrossRef](#)]
66. Bemi, V.; Tazzini, N.; Banditelli, S.; Giorgelli, F.; Pesi, R.; Turchi, C.; Mattana, A.; Sgarrella, F.; Tozzi, M.G.; Camici, M. Deoxyadenosine metabolism in a human colon-carcinoma cell line (LoVo) in relation to its cytotoxic effect in combination with deoxycoformycin. *Int. J. Cancer* **1998**, *75*, 713–720. [[CrossRef](#)]
67. Giannecchini, M.; D'Innocenzo, B.; Pesi, R.; Sgarrella, F.; Iorio, M.; Collecchi, P.; Tozzi, M.G.; Camici, M. 2'-deoxyadenosine causes apoptotic cell death in a human colon carcinoma cell line. *J. Biochem. Mol. Toxic* **2003**, *17*, 329–337. [[CrossRef](#)]
68. Garcia-Gil, M.; Tozzi, M.G.; Allegrini, S.; Folcarelli, S.; Della Sala, G.; Voccoli, V.; Colombaioni, L.; Camici, M. Novel metabolic aspects related to adenosine deaminase inhibition in a human astrocytoma cell line. *Neurochem. Int.* **2012**, *60*, 523–532. [[CrossRef](#)]
69. Garcia-Gil, M.; Tozzi, M.G.; Balestri, F.; Colombaioni, L.; Camici, M. Mitochondrial damage and apoptosis induced by adenosine deaminase inhibition and deoxyadenosine in human neuroblastoma cell lines. *J. Cell Biochem.* **2016**, *117*, 1671–1679. [[CrossRef](#)]
70. Garcia-Gil, M.; Tozzi, M.G.; Varani, S.; Della Verde, L.; Petrotto, E.; Balestri, F.; Colombaioni, L.; Camici, M. The combination of adenosine deaminase inhibition and deoxyadenosine induces apoptosis in a human astrocytoma cell line. *Neurochem. Int.* **2015**, *80*, 14–22. [[CrossRef](#)]
71. Soleimani, A.; Bahreyni, A.; Roshan, M.K.; Soltani, A.; Ryzhikov, M.; Shafiee, M.; Soukhtanloo, M.; Jaafari, M.R.; Mashkani, B.; Hassanian, S.M. Therapeutic potency of pharmacological adenosine receptors agonist/antagonist on cancer cell apoptosis in tumor microenvironment, current status, and perspectives. *J. Cell. Physiol.* **2019**, *234*, 2329–2336. [[CrossRef](#)]
72. Thirupathi, A.; Chang, Y.Z. Role of AMPK and its molecular intermediates in subjugating cancer survival mechanism. *Life Sci.* **2019**, *227*, 30–38. [[CrossRef](#)]
73. Saito, M.; Yaguchi, T.; Yasuda, Y.; Nakano, T.; Nishizaki, T. Adenosine suppresses CW2 human colonic cancer growth by inducing apoptosis via A(1) adenosine receptors. *Cancer Lett.* **2010**, *290*, 211–215. [[CrossRef](#)]
74. Sai, K.; Yang, D.; Yamamoto, H.; Fujikawa, H.; Yamamoto, S.; Nagata, T.; Saito, M.; Yamamura, T.; Nishizaki, T. A(1) adenosine receptor signal and AMPK involving caspase-9/-3 activation are responsible for adenosine-induced RCR-1 astrocytoma cell death. *Neurotoxicology* **2006**, *27*, 458–467. [[CrossRef](#)]
75. Chen, Y.; Yang, S.H.; Hueng, D.Y.; Syu, J.P.; Liao, C.C.; Wu, Y.C. Cordycepin induces apoptosis of C6 glioma cells through the adenosine 2A receptor-p53-caspase-7-PARP pathway. *Chem. Biol. Interact.* **2014**, *216*, 17–25. [[CrossRef](#)]
76. Merighi, S.; Mirandola, P.; Milani, D.; Varani, K.; Gessi, S.; Klotz, K.N.; Leung, E.; Baraldi, P.G.; Borea, P.A. Adenosine receptors as mediators of both cell proliferation and cell death of cultured human melanoma cells. *J. Invest. Derm.* **2002**, *119*, 923–933. [[CrossRef](#)]
77. Tamura, K.; Kanno, T.; Fujita, Y.; Gotoh, A.; Nakano, T.; Nishizaki, T. A(2a) adenosine receptor mediates HepG2 cell apoptosis by downregulating Bcl-X(L) expression and upregulating Bid expression. *J. Cell Biochem.* **2012**, *113*, 1766–1775. [[CrossRef](#)]
78. Yasuda, Y.; Saito, M.; Yamamura, T.; Yaguchi, T.; Nishizaki, T. Extracellular adenosine induces apoptosis in Caco-2 human colonic cancer cells by activating caspase-9/-3 via A(2a) adenosine receptors. *J. Gastroenterol.* **2009**, *44*, 56–65. [[CrossRef](#)]
79. Hajiahmadi, S.; Panjehpour, M.; Aghaei, M.; Shabani, M. Activation of A2b adenosine receptor regulates ovarian cancer cell growth: Involvement of Bax/Bcl-2 and caspase-3. *Biochem. Cell Biol.* **2015**, *93*, 321–329. [[CrossRef](#)]

80. Jafari, S.M.; Joshaghani, H.R.; Panjehpour, M.; Aghaei, M. A2B adenosine receptor agonist induces cell cycle arrest and apoptosis in breast cancer stem cells via ERK1/2 phosphorylation. *Cell Oncol.* **2018**, *41*, 61–72. [[CrossRef](#)]
81. Abedi, H.; Aghaei, M.; Panjehpour, M.; Hajiahmadi, S. Mitochondrial and caspase pathways are involved in the induction of apoptosis by IB-MECA in ovarian cancer cell lines. *Tumour Biol.* **2014**, *35*, 11027–11039. [[CrossRef](#)]
82. Jafari, S.M.; Panjehpour, M.; Aghaei, M.; Joshaghani, H.R.; Enderami, S.E. A3 adenosine receptor agonist inhibited survival of breast cancer stem cells via GLI-1 and ERK1/2 pathway. *J. Cell Biochem.* **2017**, *118*, 2909–2920. [[CrossRef](#)]
83. Cohen, S.; Stemmer, S.M.; Zozulya, G.; Ochaion, A.; Patoka, R.; Barer, F.; Bar-Yehuda, S.; Rath-Wolfson, L.; Jacobson, K.A.; Fishman, P. CF102 an A3 adenosine receptor agonist mediates anti-tumor and anti-inflammatory effects in the liver. *J. Cell. Physiol.* **2011**, *226*, 2438–2447. [[CrossRef](#)]
84. Kanno, T.; Gotoh, A.; Fujita, Y.; Nakano, T.; Nishizaki, T. A(3) adenosine receptor mediates apoptosis in 5637 human bladder cancer cells by G(q) protein/PKC-dependent AIF upregulation. *Cell. Physiol. Biochem.* **2012**, *30*, 1159–1168. [[CrossRef](#)]
85. Kanno, T.; Nakano, T.; Fujita, Y.; Gotoh, A.; Nishizaki, T. Adenosine induces apoptosis in SBC-3 human lung cancer cells through A(3) adenosine receptor-dependent AMID upregulation. *Cell. Physiol. Biochem.* **2012**, *30*, 666–677. [[CrossRef](#)]
86. Nagaya, H.; Gotoh, A.; Kanno, T.; Nishizaki, T. A3 adenosine receptor mediates apoptosis in in vitro RCC4-VHL human renal cancer cells by up-regulating AMID expression. *J. Urol.* **2013**, *189*, 321–328. [[CrossRef](#)]
87. Aghaei, M.; Panjehpour, M.; Karami-Tehrani, F.; Salami, S. Molecular mechanisms of A3 adenosine receptor-induced G1 cell cycle arrest and apoptosis in androgen-dependent and independent prostate cancer cell lines: Involvement of intrinsic pathway. *J. Cancer Res. Clin. Oncol.* **2011**, *137*, 1511–1523. [[CrossRef](#)]
88. Otsuki, T.; Kanno, T.; Fujita, Y.; Tabata, C.; Fukuoka, K.; Nakano, T.; Gotoh, A.; Nishizaki, T. A3 adenosine receptor-mediated p53-dependent apoptosis in Lu-65 human lung cancer cells. *Cell. Physiol. Biochem.* **2012**, *30*, 210–220. [[CrossRef](#)]
89. Kim, S.G.; Ravi, G.; Hoffmann, C.; Jung, Y.J.; Kim, M.; Chen, A.; Jacobson, K.A. p53-Independent induction of Fas and apoptosis in leukemic cells by an adenosine derivative, CI-IB-MECA. *Biochem. Pharm.* **2002**, *63*, 871–880. [[CrossRef](#)]
90. Jiang, X.; Tan, H.Y.; Teng, S.; Chan, Y.T.; Wang, D.; Wang, N. The role of AMP-Activated protein kinase as a potential target of treatment of hepatocellular carcinoma. *Cancers* **2019**, *11*, 647. [[CrossRef](#)]
91. Tsuchiya, A.; Nishizaki, T. Anticancer effect of adenosine on gastric cancer via diverse signaling pathways. *World J. Gastroenterol.* **2015**, *21*, 10931–10935. [[CrossRef](#)]
92. Saitoh, M.; Nagai, K.; Nakagawa, K.; Yamamura, T.; Yamamoto, S.; Nishizaki, T. Adenosine induces apoptosis in the human gastric cancer cells via an intrinsic pathway relevant to activation of AMP-Activated protein kinase. *Biochem. Pharmacol.* **2004**, *67*, 2005–2011. [[CrossRef](#)]
93. Nogi, Y.; Kanno, T.; Nakano, T.; Fujita, Y.; Tabata, C.; Fukuoka, K.; Gotoh, A.; Nishizaki, T. AMP converted from intracellularly transported adenosine upregulates p53 expression to induce malignant pleural mesothelioma cell apoptosis. *Cell. Physiol. Biochem.* **2012**, *30*, 61–74. [[CrossRef](#)]
94. Zuckerman, V.; Wolyniec, K.; Sionov, R.V.; Haupt, S.; Haupt, Y. Tumour suppression by p53: The importance of apoptosis and cellular senescence. *J. Pathol.* **2009**, *219*, 3–15. [[CrossRef](#)]
95. Nakajima, Y.; Kanno, T.; Nagaya, T.; Kuribayashi, K.; Nakano, T.; Gotoh, A.; Nishizaki, T. Adenosine deaminase inhibitor EHNA exhibits a potent anticancer effect against malignant pleural mesothelioma. *Cell. Physiol. Biochem.* **2015**, *35*, 51–60. [[CrossRef](#)]
96. Haynes, J.; Killilea, D.W.; Peterson, P.D.; Thompson, W.J. Erythro-9-(2-hydroxy-3-nonyl)adenine inhibits cyclic-3',5'-guanosine monophosphate-stimulated phosphodiesterase to reverse hypoxic pulmonary vasoconstriction in the perfused rat lung. *J. Pharm. Exp.* **1996**, *276*, 752–757.
97. Tsuchiya, A.; Kanno, T.; Saito, M.; Miyoshi, Y.; Gotoh, A.; Nakano, T.; Nishizaki, T. Intracellularly transported adenosine induces apoptosis in [corrected] MCF-7 human breast cancer cells by accumulating AMID in the nucleus. *Cancer Lett.* **2012**, *321*, 65–72. [[CrossRef](#)]
98. Bano, D.; Prehn, J.H.M. Apoptosis-Inducing factor (AIF) in physiology and disease: The tale of a repented natural born killer. *EBioMedicine* **2018**, *30*, 29–37. [[CrossRef](#)]

99. Yang, D.; Yaguchi, T.; Nagata, T.; Gotoh, A.; Dovat, S.; Song, C.; Nishizaki, T. AMID mediates adenosine-induced caspase-independent HuH-7 cell apoptosis. *Cell Physiol. Biochem.* **2011**, *27*, 37–44. [[CrossRef](#)]
100. Hermes, M.; Osswald, H.; Kloor, D. Role of S-adenosylhomocysteine hydrolase in adenosine-induced apoptosis in HepG2 cells. *Exp. Cell Res.* **2007**, *313*, 264–283. [[CrossRef](#)]
101. Kutryb-Zajac, B.; Koszalka, P.; Mierzejewska, P.; Bulinska, A.; Zabielska, M.A.; Brodzik, K.; Skrzypkowska, A.; Zelazek, L.; Pelikant-Malecka, I.; Slominska, E.M.; et al. Adenosine deaminase inhibition suppresses progression of 4T1 murine breast cancer by adenosine receptor-dependent mechanisms. *J. Cell Mol. Med.* **2018**, *22*, 5939–5954. [[CrossRef](#)]
102. Erion, M.D.; Takabayashi, K.; Smith, H.B.; Kessi, J.; Wagner, S.; Honger, S.; Shames, S.L.; Ealick, S.E. Purine nucleoside phosphorylase. 1. Structure-function studies. *Biochemistry* **1997**, *36*, 11725–11734. [[CrossRef](#)]
103. Bennett, E.M.; Li, C.; Allan, P.W.; Parker, W.B.; Ealick, S.E. Structural basis for substrate specificity of *Escherichia coli* purine nucleoside phosphorylase. *J. Biol. Chem.* **2003**, *278*, 47110–47118. [[CrossRef](#)]
104. Giblett, E.R.; Ammann, A.J.; Wara, D.W.; Sandman, R.; Diamond, L.K. Nucleoside-phosphorylase deficiency in a child with severely defective T-cell immunity and normal B-cell immunity. *Lancet* **1975**, *1*, 1010–1013. [[CrossRef](#)]
105. Madrid-Marina, V.; Lestan, B.; Nowak, P.J.; Fox, I.H.; Spychala, J. Altered properties of human T-lymphoblast soluble low Km 5'-nucleotidase: Comparison with B-lymphoblast enzyme. *Leuk. Res.* **1993**, *17*, 231–240. [[CrossRef](#)]
106. Huang, M.; Wang, Y.; Gu, J.; Yang, J.; Noel, K.; Mitchell, B.S.; Schramm, V.L.; Graves, L.M. Determinants of sensitivity of human T-cell leukemia CCRF-CEM cells to immucillin-H. *Leuk. Res.* **2008**, *32*, 1268–1278. [[CrossRef](#)]
107. Evans, G.B.; Furneaux, R.H.; Lewandowicz, A.; Schramm, V.L.; Tyler, P.C. Exploring structure-activity relationships of transition state analogues of human purine nucleoside phosphorylase. *J. Med. Chem.* **2003**, *46*, 3412–3423. [[CrossRef](#)]
108. Arpaia, E.; Benveniste, P.; Di Cristofano, A.; Gu, Y.; Dalal, I.; Kelly, S.; Hershfield, M.; Pandolfi, P.P.; Roifman, C.M.; Cohen, A. Mitochondrial basis for immune deficiency. Evidence from purine nucleoside phosphorylase-deficient mice. *J. Exp. Med.* **2000**, *191*, 2197–2208. [[CrossRef](#)]
109. Zhu, C.; Johansson, M.; Permert, J.; Karlsson, A. Enhanced cytotoxicity of nucleoside analogs by overexpression of mitochondrial deoxyguanosine kinase in cancer cell lines. *J. Biol. Chem.* **1998**, *273*, 14707–14711. [[CrossRef](#)]
110. Moll, U.M.; Zaika, A. Nuclear and mitochondrial apoptotic pathways of p53. *FEBS Lett.* **2001**, *493*, 65–69. [[CrossRef](#)]
111. Oka, S.; Ohno, M.; Tsuchimoto, D.; Sakumi, K.; Furuichi, M.; Nakabeppu, Y. Two distinct pathways of cell death triggered by oxidative damage to nuclear and mitochondrial DNAs. *EMBO J.* **2008**, *27*, 421–432. [[CrossRef](#)]
112. Alonso, R.; Lopez-Guerra, M.; Upshaw, R.; Bantia, S.; Smal, C.; Bontemps, F.; Manz, C.; Mehrling, T.; Villamor, N.; Campo, E.; et al. Forodesine has high antitumor activity in chronic lymphocytic leukemia and activates p53-independent mitochondrial apoptosis by induction of p73 and BIM. *Blood* **2009**, *114*, 1563–1575. [[CrossRef](#)]
113. Balakrishnan, K.; Nimmanapalli, R.; Ravandi, F.; Keating, M.J.; Gandhi, V. Forodesine, an inhibitor of purine nucleoside phosphorylase, induces apoptosis in chronic lymphocytic leukemia cells. *Blood* **2006**, *108*, 2392–2398. [[CrossRef](#)]
114. Gandhi, V.; Balakrishnan, K. Pharmacology and mechanism of action of forodesine, a T-cell targeted agent. *Semin. Oncol.* **2007**, *34*, S8–S12. [[CrossRef](#)]
115. Dohner, H.; Stilgenbauer, S.; Benner, A.; Leupolt, E.; Krober, A.; Bullinger, L.; Dohner, K.; Bentz, M.; Lichter, P. Genomic aberrations and survival in chronic lymphocytic leukemia. *N. Engl. J. Med.* **2000**, *343*, 1910–1916. [[CrossRef](#)]
116. Korycka, A.; Blonski, J.Z.; Robak, T. Forodesine (BCX-1777, Immucillin H)—A new purine nucleoside analogue: Mechanism of action and potential clinical application. *Mini Rev. Med. Chem.* **2007**, *7*, 976–983. [[CrossRef](#)]
117. Balakrishnan, K.; Ravandi, F.; Bantia, S.; Franklin, A.; Gandhi, V. Preclinical and clinical evaluation of forodesine in pediatric and adult B-cell acute lymphoblastic leukemia. *Clin. Lymphoma Myeloma Leuk.* **2013**, *13*, 458–466. [[CrossRef](#)]

118. Ito, Y.; Makita, S.; Tobinai, K. Development of new agents for peripheral T-cell lymphoma. *Expert Opin. Biol. Ther.* **2019**, *19*, 197–209. [[CrossRef](#)]
119. Zhang, J.; Kale, V.; Chen, M. Gene-directed enzyme prodrug therapy. *AAPS J.* **2015**, *17*, 102–110. [[CrossRef](#)]
120. Afshar, S.; Asai, T.; Morrison, S.L. Humanized ADEPT comprised of an engineered human purine nucleoside phosphorylase and a tumor targeting peptide for treatment of cancer. *Mol. Cancer* **2009**, *8*, 185–193. [[CrossRef](#)]
121. Parker, W.B.; King, S.A.; Allan, P.W.; Bennett, L.L., Jr.; Secrist, J.A., 3rd; Montgomery, J.A.; Gilbert, K.S.; Waud, W.R.; Wells, A.H.; Gillespie, G.Y.; et al. In vivo gene therapy of cancer with E. coli purine nucleoside phosphorylase. *Hum. Gene Ther.* **1997**, *8*, 1637–1644. [[CrossRef](#)]
122. Parker, W.B.; Allan, P.W.; Shaddix, S.C.; Rose, L.M.; Speegle, H.F.; Gillespie, G.Y.; Bennett, L.L., Jr. Metabolism and metabolic actions of 6-methylpurine and 2-fluoroadenine in human cells. *Biochem. Pharm.* **1998**, *55*, 1673–1681. [[CrossRef](#)]
123. Parker, W.B.; Sorscher, E.J. Use of E. coli purine nucleoside phosphorylase in the treatment of solid tumors. *Curr. Pharm. Des.* **2017**, *23*, 22. [[CrossRef](#)]
124. Martiniello-Wilks, R.; Wang, X.Y.; Voeks, D.J.; Dane, A.; Shaw, J.M.; Mortensen, E.; Both, G.W.; Russell, P.J. Purine nucleoside phosphorylase and fludarabine phosphate gene-directed enzyme prodrug therapy suppresses primary tumour growth and pseudo-metastases in a mouse model of prostate cancer. *J. Gene Med.* **2004**, *6*, 1343–1357. [[CrossRef](#)]
125. Arvidsson, Y.; Sumantran, V.; Watt, F.; Uramoto, H.; Funa, K. Neuroblastoma-specific cytotoxicity mediated by the Mash1-promoter and E. coli purine nucleoside phosphorylase. *Pediatr. Blood Cancer* **2005**, *44*, 77–84. [[CrossRef](#)]
126. Singh, P.P.; Joshi, S.; Russell, P.J.; Nair, S.; Khatri, A. Purine nucleoside phosphorylase mediated molecular chemotherapy and conventional chemotherapy: A tangible union against chemoresistant cancer. *BMC Cancer* **2011**, *11*, 368. [[CrossRef](#)]
127. Singh, P.P.; Joshi, S.; Russell, P.J.; Verma, N.D.; Wang, X.; Khatri, A. Molecular chemotherapy and chemotherapy: A new front against late-stage hormone-refractory prostate cancer. *Clin. Cancer Res.* **2011**, *17*, 4006–4018. [[CrossRef](#)]
128. Krohne, T.U.; Shankara, S.; Geissler, M.; Roberts, B.L.; Wands, J.R.; Blum, H.E.; Mohr, L. Mechanisms of cell death induced by suicide genes encoding purine nucleoside phosphorylase and thymidine kinase in human hepatocellular carcinoma cells in vitro. *Hepatology* **2001**, *34*, 511–518. [[CrossRef](#)]
129. Arnold, W.J.; Kelley, W.N. Human hypoxanthine-guanine phosphoribosyltransferase. Purification and subunit structure. *J. Biol. Chem.* **1971**, *246*, 7398–7404.
130. Wilson, J.M.; O'Toole, T.E.; Argos, P.; Shewach, D.S.; Daddona, P.E.; Kelley, W.N. Human adenine phosphoribosyltransferase. Complete amino acid sequence of the erythrocyte enzyme. *J. Biol. Chem.* **1986**, *261*, 13677–13683.
131. Sahasranaman, S.; Howard, D.; Roy, S. Clinical pharmacology and pharmacogenetics of thiopurines. *Eur. J. Clin. Pharm.* **2008**, *64*, 753–767. [[CrossRef](#)]
132. Moon, W.; Loftus, E.V., Jr. Review article: Recent advances in pharmacogenetics and pharmacokinetics for safe and effective thiopurine therapy in inflammatory bowel disease. *Aliment. Pharm.* **2016**, *43*, 863–883. [[CrossRef](#)]
133. Swann, P.F.; Waters, T.R.; Moulton, D.C.; Xu, Y.Z.; Zheng, Q.; Edwards, M.; Mace, R. Role of postreplicative DNA mismatch repair in the cytotoxic action of thioguanine. *Science* **1996**, *273*, 1109–1111. [[CrossRef](#)]
134. Yan, T.; Berry, S.E.; Desai, A.B.; Kinsella, T.J. DNA mismatch repair (MMR) mediates 6-thioguanine genotoxicity by introducing single-strand breaks to signal a G2-M arrest in MMR-proficient RKO cells. *Clin. Cancer Res.* **2003**, *9*, 2327–2334.
135. Chaabane, W.; Appell, M.L. Interconnections between apoptotic and autophagic pathways during thiopurine-induced toxicity in cancer cells: The role of reactive oxygen species. *Oncotarget* **2016**, *7*, 75616–75634. [[CrossRef](#)]
136. Isakovic, A.M.; Dulovic, M.; Markovic, I.; Kravic-Stevovic, T.; Bumbasirevic, V.; Trajkovic, V.; Isakovic, A. Autophagy suppression sensitizes glioma cells to IMP dehydrogenase inhibition-induced apoptotic death. *Exp. Cell Res.* **2017**, *350*, 32–40. [[CrossRef](#)]
137. Kim, H.R.; Roe, J.S.; Lee, J.E.; Hwang, I.Y.; Cho, E.J.; Youn, H.D. A p53-inducible microRNA-34a downregulates Ras signaling by targeting IMPDH. *Biochem. Biophys. Res. Commun.* **2012**, *418*, 682–688. [[CrossRef](#)]

138. Natsumeda, Y.; Ohno, S.; Kawasaki, H.; Konno, Y.; Weber, G.; Suzuki, K. Two distinct cDNAs for human IMP dehydrogenase. *J. Biol. Chem.* **1990**, *265*, 5292–5295.
139. Shah, C.P.; Kharkar, P.S. Newer human inosine 5'-monophosphate dehydrogenase 2 (hIMPDH2) inhibitors as potential anticancer agents. *J. Enzym. Inhib. Med. Chem.* **2018**, *33*, 972–977. [[CrossRef](#)]
140. Nair, V.; Shu, Q. Inosine monophosphate dehydrogenase as a probe in antiviral drug discovery. *Antivir. Chem. Chemother.* **2007**, *18*, 245–258. [[CrossRef](#)]
141. Bentley, R. Mycophenolic acid: A one hundred year odyssey from antibiotic to immunosuppressant. *Chem. Rev.* **2000**, *100*, 3801–3826. [[CrossRef](#)]
142. Knight, S.R.; Morris, P.J. Does the evidence support the use of mycophenolate mofetil therapeutic drug monitoring in clinical practice? A systematic review. *Transplantation* **2008**, *85*, 1675–1685. [[CrossRef](#)]
143. Shah, C.P.; Kharkar, P.S. Inosine 5'-monophosphate dehydrogenase inhibitors as antimicrobial agents: Recent progress and future perspectives. *Future Med. Chem.* **2015**, *7*, 1415–1429. [[CrossRef](#)]
144. Shu, Q.; Nair, V. Inosine monophosphate dehydrogenase (IMPDH) as a target in drug discovery. *Med. Res. Rev.* **2008**, *28*, 219–232. [[CrossRef](#)]
145. Pankiewicz, K.W.; Petrelli, R.; Singh, R.; Felczak, K. Nicotinamide adenine dinucleotide based therapeutics, update. *Curr. Med. Chem.* **2015**, *22*, 3991–4028. [[CrossRef](#)]
146. Cuny, G.D.; Suebsuwong, C.; Ray, S.S. Inosine-5'-monophosphate dehydrogenase (IMPDH) inhibitors: A patent and scientific literature review (2002–2016). *Expert Opin. Pat.* **2017**, *27*, 677–690. [[CrossRef](#)]
147. Valvezan, A.J.; Turner, M.; Belaid, A.; Lam, H.C.; Miller, S.K.; McNamara, M.C.; Baglini, C.; Housden, B.E.; Perrimon, N.; Kwiatkowski, D.J.; et al. mTORC1 couples nucleotide synthesis to nucleotide demand resulting in a targetable metabolic vulnerability. *Cancer Cell* **2017**, *32*, 624–638. [[CrossRef](#)]
148. Huang, F.; Ni, M.; Chalishazar, M.D.; Huffman, K.E.; Kim, J.; Cai, L.; Shi, X.; Cai, F.; Zacharias, L.G.; Ireland, A.S.; et al. Inosine monophosphate dehydrogenase dependence in a subset of small cell lung cancers. *Cell Metab.* **2018**, *28*, 369–382. [[CrossRef](#)]
149. Kofuji, S.; Hirayama, A.; Eberhardt, A.O.; Kawaguchi, R.; Sugiura, Y.; Sampetean, O.; Ikeda, Y.; Warren, M.; Sakamoto, N.; Kitahara, S.; et al. IMP dehydrogenase-2 drives aberrant nucleolar activity and promotes tumorigenesis in glioblastoma. *Nat. Cell Biol.* **2019**, *21*, 1003–1014. [[CrossRef](#)]
150. McLean, J.E.; Hamaguchi, N.; Belenky, P.; Mortimer, S.E.; Stanton, M.; Hedstrom, L. Inosine 5'-monophosphate dehydrogenase binds nucleic acids in vitro and in vivo. *Biochem. J.* **2004**, *379*, 243–251. [[CrossRef](#)]
151. Mortimer, S.E.; Xu, D.; McGrew, D.; Hamaguchi, N.; Lim, H.C.; Bowne, S.J.; Daiger, S.P.; Hedstrom, L. IMP dehydrogenase type 1 associates with polyribosomes translating rhodopsin mRNA. *J. Biol. Chem.* **2008**, *283*, 36354–36360. [[CrossRef](#)]
152. Park, J.H.; Ahn, S.H. IMP dehydrogenase is recruited to the transcription complex through serine 2 phosphorylation of RNA polymerase II. *Biochem. Biophys. Res. Commun.* **2010**, *392*, 588–592. [[CrossRef](#)]
153. Kozhevnikova, E.N.; van der Knaap, J.A.; Pindyurin, A.V.; Ozgur, Z.; van Ijcken, W.F.; Moshkin, Y.M.; Verrijzer, C.P. Metabolic enzyme IMPDH is also a transcription factor regulated by cellular state. *Mol. Cell* **2012**, *47*, 133–139. [[CrossRef](#)]
154. Tricot, G.J.; Jayaram, H.N.; Lapis, E.; Natsumeda, Y.; Nichols, C.R.; Kneebone, P.; Heerema, N.; Weber, G.; Hoffman, R. Biochemically directed therapy of leukemia with tiazofurin, a selective blocker of inosine 5'-phosphate dehydrogenase activity. *Cancer Res.* **1989**, *49*, 3696–3701.
155. Laliberte, J.; Yee, A.; Xiong, Y.; Mitchell, B.S. Effects of guanine nucleotide depletion on cell cycle progression in human T lymphocytes. *Blood* **1998**, *91*, 2896–2904.
156. Kiguchi, K.; Collart, F.R.; Henning-Chubb, C.; Huberman, E. Cell differentiation and altered IMP dehydrogenase expression induced in human T-lymphoblastoid leukemia cells by mycophenolic acid and tiazofurin. *Exp. Cell Res.* **1990**, *187*, 47–53. [[CrossRef](#)]
157. Kiguchi, K.; Collart, F.R.; Henning-Chubb, C.; Huberman, E. Induction of cell differentiation in melanoma cells by inhibitors of IMP dehydrogenase: Altered patterns of IMP dehydrogenase expression and activity. *Cell Growth Differ.* **1990**, *1*, 259–270.
158. Hunakova, L.; Bies, J.; Sedlak, J.; Duraj, J.; Jakubikova, J.; Takacsova, X.; Novotny, L.; Chorvath, B. Differential sensitivity of ovarian carcinoma cell lines to apoptosis induced by the IMPDH inhibitor benzamide riboside. *Neoplasma* **2000**, *47*, 274–279.

159. Moosavi, M.A.; Yazdanparast, R.; Sanati, M.H.; Nejad, A.S. 3-Hydrogenkwadaphnin targets inosine 5'-monophosphate dehydrogenase and triggers post-G1 arrest apoptosis in human leukemia cell lines. *Int. J. Biochem. Cell Biol.* **2005**, *37*, 2366–2379. [[CrossRef](#)]
160. Pan, H.; Hu, Q.; Wang, J.; Liu, Z.; Wu, D.; Lu, W.; Huang, J. Myricetin is a novel inhibitor of human inosine 5'-monophosphate dehydrogenase with anti-leukemia activity. *Biochem. Biophys. Res. Commun.* **2016**, *477*, 915–922. [[CrossRef](#)]
161. Yang, H.; Fang, Z.; Wei, Y.; Bohannon, Z.S.; Ganan-Gomez, I.; Pierola, A.A.; Paradiso, L.J.; Iwamura, H.; Garcia-Manero, G. Preclinical activity of FF-10501-01, a novel inosine-5'-monophosphate dehydrogenase inhibitor, in acute myeloid leukemia. *Leuk. Res.* **2017**, *59*, 85–92. [[CrossRef](#)]
162. Khanna, N.; Jayaram, H.N.; Singh, N. Benzamide riboside induced mitochondrial mediated apoptosis in human lung cancer H520 cells. *Life Sci.* **2004**, *75*, 179–190. [[CrossRef](#)]
163. Takebe, N.; Cheng, X.; Fandy, T.E.; Srivastava, R.K.; Wu, S.; Shankar, S.; Bauer, K.; Shaughnessy, J.; Tricot, G. IMP dehydrogenase inhibitor mycophenolate mofetil induces caspase-dependent apoptosis and cell cycle inhibition in multiple myeloma cells. *Mol. Cancer* **2006**, *5*, 457–466. [[CrossRef](#)]
164. Meli, M.; Tolomeo, M.; Grifantini, M.; Franchetti, P.; Cappellacci, L.; Simoni, D.; Invidiata, F.P.; Aiello, S.; Dusonchet, L. The synergistic apoptotic effects of thiophenfurin, an inosine monophosphate dehydrogenase inhibitor, in combination with retinoids in HL60 cells. *Oncol. Rep.* **2007**, *17*, 185–192. [[CrossRef](#)]
165. Ishitsuka, K.; Hideshima, T.; Hamasaki, M.; Raje, N.; Kumar, S.; Podar, K.; Le Gouill, S.; Shiraiishi, N.; Yasui, H.; Roccaro, A.M.; et al. Novel inosine monophosphate dehydrogenase inhibitor VX-944 induces apoptosis in multiple myeloma cells primarily via caspase-independent AIF/Endo G pathway. *Oncogene* **2005**, *24*, 5888–5896. [[CrossRef](#)]
166. Kim, M.E.; Ha, T.K.; Yoon, J.H.; Lee, J.S. Myricetin induces cell death of human colon cancer cells via BAX/BCL2-dependent pathway. *Anticancer Res.* **2014**, *34*, 701–706.
167. Floryk, D.; Thompson, T.C. Antiproliferative effects of AVN944, a novel inosine 5-monophosphate dehydrogenase inhibitor, in prostate cancer cells. *Int. J. Cancer* **2008**, *123*, 2294–2302. [[CrossRef](#)]
168. Messina, E.; Gazzaniga, P.; Micheli, V.; Guaglianone, M.R.; Barbato, S.; Morrone, S.; Frati, L.; Agliano, A.M.; Giacomello, A. Guanine nucleotide depletion triggers cell cycle arrest and apoptosis in human neuroblastoma cell lines. *Int. J. Cancer* **2004**, *108*, 812–817. [[CrossRef](#)]
169. Li, M.; Su, B.S.; Chang, L.H.; Gao, Q.; Chen, K.L.; An, P.; Huang, C.; Yang, J.; Li, Z.F. Oxymatrine induces apoptosis in human cervical cancer cells through guanine nucleotide depletion. *Anticancer Drugs* **2014**, *25*, 161–173. [[CrossRef](#)]
170. Gu, J.J.; Gathy, K.; Santiago, L.; Chen, E.; Huang, M.; Graves, L.M.; Mitchell, B.S. Induction of apoptosis in IL-3-dependent hematopoietic cell lines by guanine nucleotide depletion. *Blood* **2003**, *101*, 4958–4965. [[CrossRef](#)]
171. Goldstone, D.C.; Ennis-Adeniran, V.; Hedden, J.J.; Groom, H.C.; Rice, G.I.; Christodoulou, E.; Walker, P.A.; Kelly, G.; Haire, L.F.; Yap, M.W.; et al. HIV-1 restriction factor SAMHD1 is a deoxynucleoside triphosphate triphosphohydrolase. *Nature* **2011**, *480*, 379–382. [[CrossRef](#)]
172. Powell, R.D.; Holland, P.J.; Hollis, T.; Perrino, F.W. Aicardi-Goutieres syndrome gene and HIV-1 restriction factor SAMHD1 is a dGTP-regulated deoxynucleotide triphosphohydrolase. *J. Biol. Chem.* **2011**, *286*, 43596–43600. [[CrossRef](#)]
173. Baldauf, H.M.; Pan, X.; Erikson, E.; Schmidt, S.; Daddacha, W.; Burggraf, M.; Schenkova, K.; Ambiel, I.; Wabnitz, G.; Gramberg, T.; et al. SAMHD1 restricts HIV-1 infection in resting CD4(+) T cells. *Nat. Med.* **2012**, *18*, 1682–1687. [[CrossRef](#)]
174. Rice, G.I.; Bond, J.; Asipu, A.; Brunette, R.L.; Manfield, I.W.; Carr, I.M.; Fuller, J.C.; Jackson, R.M.; Lamb, T.; Briggs, T.A.; et al. Mutations involved in Aicardi-Goutieres syndrome implicate SAMHD1 as regulator of the innate immune response. *Nat. Genet.* **2009**, *41*, 829–832. [[CrossRef](#)]
175. Leshinsky-Silver, E.; Malinger, G.; Ben-Sira, L.; Kidron, D.; Cohen, S.; Inbar, S.; Bezaleli, T.; Levine, A.; Vinkler, C.; Lev, D.; et al. A large homozygous deletion in the SAMHD1 gene causes atypical Aicardi-Goutieres syndrome associated with mtDNA deletions. *Eur. J. Hum. Genet.* **2011**, *19*, 287–292. [[CrossRef](#)]
176. Parsons, D.W.; Jones, S.; Zhang, X.; Lin, J.C.; Leary, R.J.; Angenendt, P.; Mankoo, P.; Carter, H.; Siu, I.M.; Gallia, G.L.; et al. An integrated genomic analysis of human glioblastoma multiforme. *Science* **2008**, *321*, 1807–1812. [[CrossRef](#)]

177. Cancer Genome Atlas Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **2012**, *487*, 330–337. [[CrossRef](#)]
178. Liu, J.; Lee, W.; Jiang, Z.; Chen, Z.; Jhunjhunwala, S.; Haverty, P.M.; Gnad, F.; Guan, Y.; Gilbert, H.N.; Stinson, J.; et al. Genome and transcriptome sequencing of lung cancers reveal diverse mutational and splicing events. *Genome Res.* **2012**, *22*, 2315–2327. [[CrossRef](#)]
179. Jones, S.; Zhang, X.; Parsons, D.W.; Lin, J.C.; Leary, R.J.; Angenendt, P.; Mankoo, P.; Carter, H.; Kamiyama, H.; Jimeno, A.; et al. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science* **2008**, *321*, 1801–1806. [[CrossRef](#)]
180. Clifford, R.; Louis, T.; Robbe, P.; Ackroyd, S.; Burns, A.; Timbs, A.T.; Wright Colopy, G.; Dreau, H.; Sigaux, F.; Judde, J.G.; et al. SAMHD1 is mutated recurrently in chronic lymphocytic leukemia and is involved in response to DNA damage. *Blood* **2014**, *123*, 1021–1031. [[CrossRef](#)]
181. Walker, B.A.; Wardell, C.P.; Melchor, L.; Hulkki, S.; Potter, N.E.; Johnson, D.C.; Fenwick, K.; Kozarewa, I.; Gonzalez, D.; Lord, C.J.; et al. Intracлонаl heterogeneity and distinct molecular mechanisms characterize the development of t(4; 14) and t(11; 14) myeloma. *Blood* **2012**, *120*, 1077–1086. [[CrossRef](#)]
182. Sjoblom, T.; Jones, S.; Wood, L.D.; Parsons, D.W.; Lin, J.; Barber, T.D.; Mandelker, D.; Leary, R.J.; Ptak, J.; Silliman, N.; et al. The consensus coding sequences of human breast and colorectal cancers. *Science* **2006**, *314*, 268–274. [[CrossRef](#)]
183. Wang, J.L.; Lu, F.Z.; Shen, X.Y.; Wu, Y.; Zhao, L.T. SAMHD1 is down regulated in lung cancer by methylation and inhibits tumor cell proliferation. *Biochem. Biophys. Res. Commun.* **2014**, *455*, 229–233. [[CrossRef](#)]
184. Zhu, C.; Gao, W.; Zhao, K.; Qin, X.; Zhang, Y.; Peng, X.; Zhang, L.; Dong, Y.; Zhang, W.; Li, P.; et al. Structural insight into dGTP-dependent activation of tetrameric SAMHD1 deoxynucleoside triphosphate triphosphohydrolase. *Nat. Commun.* **2013**, *4*, 2722. [[CrossRef](#)]
185. Miazzi, C.; Ferraro, P.; Pontarin, G.; Rampazzo, C.; Reichard, P.; Bianchi, V. Allosteric regulation of the human and mouse deoxyribonucleotide triphosphohydrolase sterile alpha-motif/histidine-aspartate domain-containing protein 1 (SAMHD1). *J. Biol. Chem.* **2014**, *289*, 18339–18346. [[CrossRef](#)]
186. Amie, S.M.; Bambara, R.A.; Kim, B. GTP is the primary activator of the anti-HIV restriction factor SAMHD1. *J. Biol. Chem.* **2013**, *288*, 25001–25006. [[CrossRef](#)]
187. Beloglazova, N.; Flick, R.; Tchigvintsev, A.; Brown, G.; Popovic, A.; Nocek, B.; Yakunin, A.F. Nuclease activity of the human SAMHD1 protein implicated in the Aicardi-Goutieres syndrome and HIV-1 restriction. *J. Biol. Chem.* **2013**, *288*, 8101–8110. [[CrossRef](#)]
188. Seamon, K.J.; Sun, Z.; Shlyakhtenko, L.S.; Lyubchenko, Y.L.; Stivers, J.T. SAMHD1 is a single-stranded nucleic acid binding protein with no active site-associated nuclease activity. *Nucleic Acids Res.* **2015**, *43*, 6486–6499. [[CrossRef](#)]
189. Kohnken, R.; Kodigepalli, K.M.; Wu, L. Regulation of deoxynucleotide metabolism in cancer: Novel mechanisms and therapeutic implications. *Mol. Cancer* **2015**, *14*, 176. [[CrossRef](#)]
190. Kodigepalli, K.M.; Li, M.; Liu, S.L.; Wu, L. Exogenous expression of SAMHD1 inhibits proliferation and induces apoptosis in cutaneous T-cell lymphoma-derived HuT78 cells. *Cell Cycle* **2017**, *16*, 179–188. [[CrossRef](#)]
191. Kodigepalli, K.M.; Bonifati, S.; Tirumuru, N.; Wu, L. SAMHD1 modulates in vitro proliferation of acute myeloid leukemia-derived THP-1 cells through the PI3K-Akt-p27 axis. *Cell Cycle* **2018**, *17*, 1124–1137. [[CrossRef](#)]
192. Fujikawa, K.; Kamiya, H.; Yakushiji, H.; Nakabeppu, Y.; Kasai, H. Human MTH1 protein hydrolyzes the oxidized ribonucleotide, 2-hydroxy-ATP. *Nucleic Acids Res.* **2001**, *29*, 449–454. [[CrossRef](#)]
193. Oda, H.; Nakabeppu, Y.; Furuichi, M.; Sekiguchi, M. Regulation of expression of the human MTH1 gene encoding 8-oxo-dGTPase. Alternative splicing of transcription products. *J. Biol. Chem.* **1997**, *272*, 17843–17850. [[CrossRef](#)]
194. Rai, P.; Sobol, R.W. Mechanisms of MTH1 inhibition-induced DNA strand breaks: The slippery slope from the oxidized nucleotide pool to genotoxic damage. *DNA Repair* **2019**, *77*, 18–26. [[CrossRef](#)]
195. Trachootham, D.; Zhou, Y.; Zhang, H.; Demizu, Y.; Chen, Z.; Pelicano, H.; Chiao, P.J.; Achanta, G.; Arlinghaus, R.B.; Liu, J.; et al. Selective killing of oncogenically transformed cells through a ROS-mediated mechanism by beta-phenylethyl isothiocyanate. *Cancer Cell* **2006**, *10*, 241–252. [[CrossRef](#)]
196. Storz, P. Reactive oxygen species in tumor progression. *Front. Biosci. J. Virtual Libr.* **2005**, *10*, 1881–1896. [[CrossRef](#)]

197. Ishikawa, K.; Takenaga, K.; Akimoto, M.; Koshikawa, N.; Yamaguchi, A.; Imanishi, H.; Nakada, K.; Honma, Y.; Hayashi, J. ROS-generating mitochondrial DNA mutations can regulate tumor cell metastasis. *Science* **2008**, *320*, 661–664. [[CrossRef](#)]
198. Ames, B.N. Dietary carcinogens and anticarcinogens. Oxygen radicals and degenerative diseases. *Science* **1983**, *221*, 1256–1264. [[CrossRef](#)]
199. Rai, P.; Onder, T.T.; Young, J.J.; McFaline, J.L.; Pang, B.; Dedon, P.C.; Weinberg, R.A. Continuous elimination of oxidized nucleotides is necessary to prevent rapid onset of cellular senescence. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 169–174. [[CrossRef](#)]
200. Gad, H.; Koolmeister, T.; Jemth, A.S.; Eshtad, S.; Jacques, S.A.; Strom, C.E.; Svensson, L.M.; Schultz, N.; Lundback, T.; Einarsdottir, B.O.; et al. MTH1 inhibition eradicates cancer by preventing sanitation of the dNTP pool. *Nature* **2014**, *508*, 215–221. [[CrossRef](#)]
201. Youn, C.K.; Jun, J.Y.; Hyun, J.W.; Hwang, G.; Lee, B.R.; Chung, M.H.; Chang, I.Y.; You, H.J. hMTH1 depletion promotes oxidative-stress-induced apoptosis through a Noxa- and caspase-3/7-mediated signaling pathway. *DNA Repair* **2008**, *7*, 1809–1823. [[CrossRef](#)]
202. Russo, M.T.; Blasi, M.F.; Chiera, F.; Fortini, P.; Degan, P.; Macpherson, P.; Furuichi, M.; Nakabeppu, Y.; Karran, P.; Aquilina, G.; et al. The oxidized deoxynucleoside triphosphate pool is a significant contributor to genetic instability in mismatch repair-deficient cells. *Mol. Cell Biol.* **2004**, *24*, 465–474. [[CrossRef](#)]
203. Huber, K.V.; Salah, E.; Radic, B.; Gridling, M.; Elkins, J.M.; Stukalov, A.; Jemth, A.S.; Gokturk, C.; Sanjiv, K.; Stromberg, K.; et al. Stereospecific targeting of MTH1 by (S)-crizotinib as an anticancer strategy. *Nature* **2014**, *508*, 222–227. [[CrossRef](#)]
204. Nakabeppu, Y.; Oka, S.; Sheng, Z.; Tsuchimoto, D.; Sakumi, K. Programmed cell death triggered by nucleotide pool damage and its prevention by MutT homolog-1 (MTH1) with oxidized purine nucleoside triphosphatase. *Mutat. Res.* **2010**, *703*, 51–58. [[CrossRef](#)]
205. Rampazzo, C.; Tozzi, M.G.; Dumontet, C.; Jordheim, L.P. The druggability of intracellular nucleotide-degrading enzymes. *Cancer Chemother. Pharm.* **2016**, *77*, 883–893. [[CrossRef](#)]
206. Samaranyake, G.J.; Huynh, M.; Rai, P. MTH1 as a chemotherapeutic target: The elephant in the room. *Cancers* **2017**, *9*, 47. [[CrossRef](#)]
207. Svilar, D.; Goellner, E.M.; Almeida, K.H.; Sobol, R.W. Base excision repair and lesion-dependent subpathways for repair of oxidative DNA damage. *Antioxid. Redox Signal.* **2011**, *14*, 2491–2507. [[CrossRef](#)]
208. Zhou, J.; Ahn, J.; Wilson, S.H.; Prives, C. A role for p53 in base excision repair. *EMBO J.* **2001**, *20*, 914–923. [[CrossRef](#)]
209. Achanta, G.; Huang, P. Role of p53 in sensing oxidative DNA damage in response to reactive oxygen species-generating agents. *Cancer Res.* **2004**, *64*, 6233–6239. [[CrossRef](#)]
210. Chatterjee, A.; Mambo, E.; Osada, M.; Upadhyay, S.; Sidransky, D. The effect of p53-RNAi and p53 knockout on human 8-oxoguanine DNA glycosylase (hOgg1) activity. *FASEB J.* **2006**, *20*, 112–114. [[CrossRef](#)]
211. Wang, R.; Li, C.; Qiao, P.; Xue, Y.; Zheng, X.; Chen, H.; Zeng, X.; Liu, W.; Boldogh, I.; Ba, X. OGG1-initiated base excision repair exacerbates oxidative stress-induced parthanatos. *Cell Death Dis.* **2018**, *9*, 628. [[CrossRef](#)]
212. Patel, A.; Burton, D.G.; Halvorsen, K.; Balkan, W.; Reiner, T.; Perez-Stable, C.; Cohen, A.; Munoz, A.; Giribaldi, M.G.; Singh, S.; et al. MutT Homolog 1 (MTH1) maintains multiple KRAS-driven pro-malignant pathways. *Oncogene* **2015**, *34*, 2586–2596. [[CrossRef](#)]
213. Abbas, H.H.K.; Alhamoudi, K.M.H.; Evans, M.D.; Jones, G.D.D.; Foster, S.S. MTH1 deficiency selectively increases non-cytotoxic oxidative DNA damage in lung cancer cells: More bad news than good? *BMC Cancer* **2018**, *18*, 423. [[CrossRef](#)]
214. Fouquerel, E.; Lormand, J.; Bose, A.; Lee, H.T.; Kim, G.S.; Li, J.; Sobol, R.W.; Freudenthal, B.D.; Myong, S.; Opresko, P.L. Oxidative guanine base damage regulates human telomerase activity. *Nat. Struct. Mol. Biol.* **2016**, *23*, 1092–1100. [[CrossRef](#)]
215. Mathews, C.K. DNA precursor metabolism and genomic stability. *FASEB J.* **2006**, *20*, 1300–1314. [[CrossRef](#)]
216. Shimizu, I.; Yoshida, Y.; Suda, M.; Minamino, T. DNA damage response and metabolic disease. *Cell Metab.* **2014**, *20*, 967–977. [[CrossRef](#)]
217. Kohalmi, S.E.; Glatke, M.; McIntosh, E.M.; Kunz, B.A. Mutational specificity of DNA precursor pool imbalances in yeast arising from deoxycytidylate deaminase deficiency or treatment with thymidylate. *J. Mol. Biol.* **1991**, *220*, 933–946. [[CrossRef](#)]

218. Pai, C.C.; Kearsey, S.E. A critical balance: dNTPs and the maintenance of genome stability. *Genes* **2017**, *8*, 57. [[CrossRef](#)]
219. Wyant, G.A.; Abu-Remaileh, M.; Frenkel, E.M.; Laqtom, N.N.; Dharamdasani, V.; Lewis, C.A.; Chan, S.H.; Heinze, I.; Ori, A.; Sabatini, D.M. NUFIP1 is a ribosome receptor for starvation-induced ribophagy. *Science* **2018**, *360*, 751–758. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).