

Review

Transforming Growth Factor-Beta (TGF β) Signaling Pathway in Cholangiocarcinoma

Panagiotis Papoutsoglou [†], Corentin Louis [†] and Cédric Coulouarn ^{* ID}

Inserm, Univ Rennes, Inra, Institut NuMeCan (Nutrition Metabolisms and Cancer), UMR_S 1241, 35033 Rennes, France

* Correspondence: cedric.coulouarn@inserm.fr; Tel.: +33-223-233-881

[†] These authors contributed equally to this work.

Received: 5 July 2019; Accepted: 19 August 2019; Published: 23 August 2019



Abstract: Cholangiocarcinoma is a deadly cancer worldwide, associated with a poor prognosis and limited therapeutic options. Although cholangiocarcinoma accounts for less than 15% of liver primary cancer, its silent nature restricts early diagnosis and prevents efficient treatment. Therefore, it is of clinical relevance to better understand the molecular basis of cholangiocarcinoma, including the signaling pathways that contribute to tumor onset and progression. In this review, we discuss the genetic, molecular, and environmental factors that promote cholangiocarcinoma, emphasizing the role of the transforming growth factor β (TGF β) signaling pathway in the progression of this cancer. We provide an overview of the physiological functions of TGF β signaling in preserving liver homeostasis and describe how advanced cholangiocarcinoma benefits from the tumor-promoting effects of TGF β . Moreover, we report the importance of noncoding RNAs as effector molecules downstream of TGF β during cholangiocarcinoma progression, and conclude by highlighting the need for identifying novel and clinically relevant biomarkers for a better management of patients with cholangiocarcinoma.

Keywords: TGF β ; cholangiocarcinoma; liver cancer; signaling; noncoding RNA

1. Cholangiocarcinoma

1.1. Epidemiology and Risk Factors

Cholangiocarcinomas (CCAs) comprise heterogeneous hepatobiliary tumors with cholangiocyte differentiation features. CCA is the second most common hepatic malignant tumor after hepatocellular carcinoma (HCC). CCAs are classified as intrahepatic (iCCA), perihilar (pCCA), and distal (dCCA), according to their anatomic location [1]. CCA subtypes differ by their epidemiology, etiology, pathogenesis, and thus, clinical management and targeted therapeutic options. Although challenges in the classification of CCA made it difficult to quantify, a gradual increase in CCA incidence was reported worldwide during the last few decades [2]. The discrepancy in incidence of sporadic CCA worldwide is associated with well-established risk factors. Higher incidence of CCAs in Eastern countries reflects a geographical disparity in the prevalence of risk factors [1]. For instance, hepatobiliary flukes such as *Clonorchis sinensis* and *Opisthorchis viverrini* are both common risk factors in Southeast Asia, where CCA is recognized as a non-rare cancer [3,4]. Hepatitis B and C have also been identified as risk factors for CCAs, especially for iCCA. While hepatitis C is prevalent in Western countries, hepatitis B is strongly associated with CCAs in Asia [5]. Furthermore, the association between primary sclerosing cholangitis (PSC) and CCA is well-established. Studies from Western countries reported that PSC patients developed CCAs with a prevalence ranging from 5% to 15% and a yearly developmental rate of CCAs ranging from 0.5% to 1.5% [6]. Other risk factors including hepatolithiasis, metabolic

syndrome, alcohol, smoking, and diabetes are also suspected to be involved in CCA onset, all of these factors contributing to generate a pro-inflammatory environment in biliary tracts [5,7]. As a result of its silent nature, CCA is generally diagnosed at the advanced stage, when therapeutic options are limited. Surgery is currently the best option for CCA treatment, even though tumor size, metastasis, and lymph node invasion make it unfeasible in more than 65% cases [8]. For the resectable early stage CCAs, survival at five years ranges from 15% to 40%, but is associated with a high risk of recurrence [9]. However, for unresectable advanced CCAs, overall survival (OS) is below 15 months [9]. The lack of a clear and global picture of cellular and molecular alterations, which occur in aggressive CCAs, might account for this unfavorable clinical outcome. Improvement in CCA outcome relies on efforts toward a better understanding of cholangiocarcinogenesis mechanisms to develop efficient targeted therapies, as well as the identification of reliable biomarkers for early detection.

1.2. Molecular Pathogenesis

CCA is frequently associated with drastic changes in the tumor microenvironment, including intense extracellular matrix remodeling and inflammation, which modulate the activity of signaling pathways involved in tumor onset and progression [10,11]. These alterations lead to an aberrant expression and/or activation of key cytokines, tyrosine kinases, and ultimately transcription factors which control cell fate [12–14]. As an example, an increase of interleukin 6 (IL6) secretion by CCAs and desmoplastic stromal cells lead to the activation of STAT3, a latent cytoplasmic transcription factor. IL6 binds to the dimerized GP130 receptors associated with Janus family kinases (JAK) including JAK1, JAK2, and TYK2, leading to STAT3 phosphorylation and activation [15]. STAT3 acts not only as an activator of transcription but also as a signal transducer. Its activation modulates a variety of genes involved in cell survival, proliferation, and migration. An elevated expression of STAT3 in CCA tumor tissues has been identified as an independent prognostic factor for OS and disease-free survival (DFS) [16]. There is also evidence demonstrating that epidermal growth factor receptor (EGFR) contributes to CCA progression by disturbing cell–cell adhesion and cell motility, triggering epithelial to mesenchymal transition (EMT) and thus promoting a pro-metastatic process [17]. EGFR activation by its ligands (e.g., EGF, TGFA, AREG) initiates several signal transduction cascades, including extracellular signal-regulated kinases (ERK) 1/2 and serine/threonine kinase 1 (AKT1), which are both implicated in cell proliferation and migration [18].

Developmental pathways are well-conserved axes required for biliary tract cell differentiation and proliferation. Unsurprisingly, dysregulations of these pathways have been described in CCAs. Recent evidence showed that a persistent activation of Notch signaling is associated with iCCA [19]. A study using a mouse model of iCCA revealed that the Notch axis was critical in hepatocyte conversion into biliary lineage, and therefore, an enhanced activity of this pathway may contribute to malignant conversion of hepatocytes into CCAs [20]. Hedgehog (HH) is another developmental pathway involved in critical cell fate decision, including apoptosis, stem cell maintenance, and wound healing [21]. HH pathway was identified as a key player in tumor initiation in several cancers, including CCAs [22,23]. El Khatib et al. investigated the effects of blocking the HH pathway using cyclopamine in vitro on human CCA cells, and in vivo using xenograft of CCA cells in mice. Cyclopamine is a steroidal alkaloid isolated from *Veratrum californicum*. It plays a critical role in embryonic development by hindering the HH pathway. Such a muting of the HH pathway resulted in a blockage of CCA cell migration and invasion. In CCA xenografts, cyclopamine treatment led to a significant inhibition of tumor growth, highlighting the importance of the HH pathway in CCAs and the clinical relevance of its inhibition [24].

High throughput strategies identified numerous genetic, epigenetic, and genomic alterations in CCA, and highlighted specific targetable signaling pathways. Among them, mutations in isocitrate dehydrogenases genes (*IDH1* and *IDH2*) and chromatin-remodeling genes, including AT-rich interaction domain 1A (*ARID1A*), as well as gene fusions involving fibroblast growth factor receptor 2 (*FGFR2*) were frequently detected in CCAs [25–28]. Mutations of the tumor suppressor *TP53* are also commonly

found in CCAs (44% of cases), so are mutations of *KRAS* (17% of cases) and *SMAD4* (17% of cases) [29]. According to the same study, seven genes (*TP53*, *SMAD4*, *KRAS*, *RNF43*, *NDC80*, *ROBO2*, and *GNAS*) scored as the top mutated genes in *Opisthorchis viverrini*-associated CCAs, with false discovery rates ranging from 2.1×10^{-5} to 0.29. More recently, protein tyrosine phosphatase non-receptor 3 (*PTPN3*) was reported to be frequently mutated in iCCA and associated with tumor recurrence [30].

1.3. Targeted Therapies

Although significant progress in understanding the molecular basis of CCA pathogenesis has been achieved, there is no approved molecular targeted therapy that significantly improves patient survival. Thus, there is a critical need of designing innovative therapeutic strategies and biomarkers for a better management of patients with CCA, which remains a deadly cancer with a very poor prognosis [9]. Previous genomic characterizations of the stroma in iCCA highlighted clinically relevant biomarkers predictive of patient survival, some of them being related to the transforming growth factor β (TGF β) signaling [11,31,32]. Several clinical trials are still in progress based on genetic alterations observed in CCAs. For example, the therapeutic effect of the multi-targeted tyrosine kinase inhibitor, ponatinib, has been tested in two patients with advanced CCA tumors associated with activating *FGFR2* gene fusions. This trial resulted in an efficient anti-tumor response characterized notably by shrinkage of metastatic lymph nodes [33]. Promising results using IDH1 and IDH2 inhibitors have been also reported. Two mutant IDH inhibitors, enasidenib (AG-221) and ivosidenib (AG-120), have been approved in ongoing trials on Acute Myeloid Leukemia (AML), and their therapeutic benefits are now studied in other malignancies including CCA [34].

2. Transforming Growth Factor Beta (TGF β) Pathway

2.1. TGF β Signaling: From Receptor Activation to Transcription Of Target Genes

Members of the TGF β family are pleiotropic cytokines that exhibit important roles in tissue homeostasis, cell differentiation, and embryonic development. Extracellular TGF β ligands bind to transmembrane type I and type II TGF β receptors (T β RI and T β RII, respectively), thereby initiating a signaling cascade that ultimately leads to altered expression of protein-coding and noncoding target genes [35,36]. Initiation of the TGF β pathway involves binding of ligands to T β RII, formation of a heterotetrameric complex between T β RII and T β RI, and phosphorylation of T β RI by T β RII, the latter possessing serine/threonine kinase activity in its intracellular domain. Phosphorylation of T β RI turns on its serine/threonine kinase activity, resulting in the phosphorylation of the members of the SMAD family, SMAD2 and SMAD3, which interact, at their carboxy-terminal domain, with the common mediator SMAD4 and transduce the signal into the nucleus. The transport of the activated SMAD complex into the nucleus is mediated by proteins of the nuclear pore, such as importin- β 1 and CAN/Nup214, and requires the GTPase RAN [37]. In the nucleus, the SMAD complexes associate with various coactivator or corepressor factors to regulate gene transcription in a positive or negative manner, respectively (Figure 1).




Figure 1. Basics of transforming growth factor β (TGF β) signaling. Binding of TGF β ligands to their receptors initiates the signal, through the phosphorylation of T β RI by T β RII. Then, T β RI transmits the signal to SMAD2 and SMAD3 by phosphorylating their MH2 domains. This phosphorylation enables SMAD2 and SMAD3 activation and the formation of a complex with SMAD4, which, in turn, enters into the nucleus through nuclear pores with the assistance of importins. In the nucleus, the activated SMAD complex regulates the expression of target genes, in a positive or negative manner, depending on its association with transcriptional coactivators or corepressors. The signaling is subjected to negative regulation by SMAD7, which prevents SMAD2/3 activation and induces degradation of T β RI, and by Ski/SnoN, which interferes with the formation of active SMAD2/3/4 complex. MH1: MAD homology 1 domain, L: Linker domain, MH2: MAD homology 2 domain, T β RI: type I TGF β receptor, T β RII: type II TGF β receptor.

Structurally, the transcription factors SMAD2, SMAD3, and SMAD4 consist of three main domains, an N-terminal (or MH1), a central (or linker), and a C-terminal (or MH2) domain [38]. The MH1 domain enables SMAD3 and SMAD4 to directly recognize and bind to specific SMAD-binding elements (SBEs) at promoter or enhancer DNA regions [39]. The linker domain is subjected to extensive post-translational modifications that influence stability and interactions with proteins from other signaling pathways. The MH2 domain is important for the interaction of SMAD2 and SMAD3 with T β RI, SMAD4, and other protein partners.

In parallel to TGF β -induced SMAD activation, TGF β receptors can also induce non-SMAD pathways, such as MAP kinase (MAPK), phosphatidylinositol-3' kinase (PI3K), and the cell polarity regulator PAR6, which can mediate its biological functions [40]. The various physiological actions of TGF β signaling require control mechanisms to ensure that the magnitude of the signal meets the temporal, spatial, or developmental needs of the cellular systems. Thus, several mechanisms prevent an overactivation of the TGF β signaling. Notably, SMAD7 and Sloan Kettering Institute proto-oncogene (SKI)/SKI-related novel gene (SnoN) proteins are the best described negative regulators of TGF β

signaling. SMAD7 is a member of the SMAD family and its inhibitory role on TGF β signaling is exerted in two ways. First, SMAD7 antagonizes with SMAD2/3 for binding to T β RI receptors, thereby limiting the phosphorylation of SMAD2/3. Second, SMAD7 promotes the ubiquitin-dependent proteosomal degradation of T β RI receptor by recruiting E3-ubiquitin ligases (SMURFs) to the receptor [41]. SKI and SnoN proteins interact with SMAD2/3 or SMAD4 and interfere with the formation of an active SMAD2/3-SMAD4 complex, thus also interfering with the propagation of the TGF β signal [41] (Figure 1).

2.2. Crosstalk of TGF β Signaling with Other Signaling Pathways

TGF β -mediated cellular responses are frequently achieved by the cooperation of TGF β with other signaling pathways. Thus, the crosstalk between TGF β and a plethora of pathways, such as WNT, HIPPO, NF- κ B, Notch, hedgehog, JAK/STAT, MAPK, and PI3K-AKT, is supported by the current literature [42]. Interestingly, most of these pathways are deregulated in CCAs [43]. As an example, TGF β and HIPPO pathways are functionally associated to regulate CCA progression. Hyperactivation of yes-associated protein-1 (Yap1) and transcriptional coactivator with PDZ-binding motif (Taz), due to a genetic depletion of Mps One Binder kinase activator (Mob)1a/1b in mouse liver, results in an increased incidence of combined HCC-CCA and iCCA. These effects are accompanied by increased levels of TGF β 2 and TGF β 3 ligands. Interestingly, a positive correlation between YAP1 and SMAD2 activation has been shown in patients with HCC-CCA and iCCA [44]. Interleukin-6 (IL-6) and TGF β pathways also converge to potentiate malignancy and resistance to chemotherapy in biliary tract cancer [45].

2.3. Physiological Responses in the Liver

TGF β signaling dictates transcriptional programs, which influence diverse physiological processes, such as cell cycle arrest, apoptosis, EMT, and immune surveillance. When the tight regulatory mechanisms of TGF β signaling activity are circumvented, pathological conditions, such as fibrotic diseases and tumorigenesis, may arise due to excessive and uncontrolled activity of the pathway. For example, the pathogenesis of liver fibrosis involves extravagant production and deposition of extracellular matrix (ECM) components i.e., collagen, produced at the liver tissue, in response to chronic liver damage [46]. At the onset of liver fibrosis, a combinatorial action of inflammatory responses, infiltrating immune cells, and cytokine signaling (e.g., TGF β) triggers the activation of hepatic stellate cells (HSC) and their transition towards myofibroblasts [47]. Failure of recovery, as a consequence of constant liver injury, favors a persistent pro-fibrotic microenvironment, ultimately leading to severe fibrotic disease. Both SMAD-dependent and non-SMAD signaling can mediate the effects of TGF β in HSC, such as the induction of connective tissue growth factor (CTGF) [48].

In epithelial cells, TGF β blocks cell proliferation, mainly through the induction of cyclin-dependent kinase (CDK) inhibitors, such as P21, P15, and P27, and the reduction of the oncogene MYC [49]. The pro-apoptotic effects of TGF β are well-described in normal liver or HCC cell lines. For example, TGF β induces apoptosis of Hep3B HCC cell line, a process that involves the activation of caspases. In the case of HUH7 cells, TGF β -mediated apoptosis engages the activation of caspases and the downregulation of anti-apoptotic proteins, such as BCL-XL and XIAP [50]. In other cellular models, the involvement of the c-Jun N-terminal kinase (JNK) pathway [51], the induction of the tumor suppressor TGF β inducible early growth response protein 1 (TIEG1) [52], or the death-associated protein kinase (DAPK) [53] have been reported as mediators of apoptosis downstream of TGF β .

TGF β also plays a central role in promoting EMT, a process whereby epithelial cells lose their well-defined morphology and adhesion and acquire mesenchymal traits, allowing them to migrate from their original site [54]. EMT is crucial for normal embryogenesis, however, during tumor progression, epithelial carcinomas can profit from hyperactivation of EMT-inducing signaling pathways, such as WNT and TGF β , to gain migratory properties, enabling them to metastasize. Many reports pinpoint the positive contribution of EMT to metastasis [55], although a few of them support that EMT is

not a determinant factor for pancreatic [56] or lung metastasis [57], but is associated with increased resistance of these cancers to chemotherapeutic agents. In fact, EMT has been also linked to the generation of cancer stem cells (CSCs), defined as undifferentiated cancer cells, which possess stem cell characteristics, contribute to tumor heterogeneity, and show low sensitivity to chemotherapy [58]. Therefore, by positively regulating EMT, TGF β is also capable of conferring stemness features to cancer cells, as described in HCC [59]. At the molecular level, TGF β restricts epithelial phenotypes either by downregulating components of tight junctions, such as E-cadherin, or by redistributing them away from the cell membrane, thereby leading to their disassembly. In addition, TGF β induces the expression of EMT transcription factors (SNAIL, SLUG, ZEB1, ZEB2, TWIST) and mesenchymal markers (fibronectin, vimentin, collagens) to achieve EMT [60].

Immune system responses are also controlled by the TGF β pathway, which, in general, exhibits immune-suppressive effects. Immune suppression by TGF β is a mechanism by which cancer cells may escape immune surveillance. TGF β disrupts the ability of immune cells to recognize and eliminate cancer cells by shifting the differentiation of naïve CD4⁺ cells towards T-regulatory cells, thereby restricting the production of effector T cells. In addition, it prevents natural killer cells from destroying tumor cells. A characteristic example for the role of TGF β in cancer progression, via modulation of the immune system, derives from a study using CCA cells. The findings of this study suggested that interference of the TGF β pathway by neutralizing antibodies against T β RII in dendritic cells (DC) caused increased activation of effector T cells and, in turn, enhanced targeting and lysis of co-cultured CCA cells by the immune system [61]. In addition, another cause of CCA progression is chronic inflammation, in response to injury or due to the presence of high levels of pro-inflammatory cytokines at the bile duct [62]. Interestingly, TGF β induces the expression of IL6 in iCCA cell lines, an effect that facilitates CCA growth [63]. Other major contributors of tumor progression are the tumor associated macrophages (TAM) [64]. These immune cells are recruited at the tumor site and secrete cytokines, such as TGF β , thereby creating a pro-inflammatory microenvironment that favors cancer progression. Also, TAM and other immune cells have been identified at tumor areas in patients with eCCA and are correlated with poor prognosis [65]. Interestingly, addition of supernatant from HuCCT1 CCA cells on TAM enforced the latter to express high levels of TGF β , IL10, and VEGF, suggesting that the interplay between TGF β and TAM is of high importance in CCA tumor microenvironment [66].

2.4. TGF β Functional Duality in Cancer

TGF β is a challenging target for cancer treatment, due to its ability to both inhibit and facilitate tumor progression. During tumor initiation, TGF β exhibits tumor-suppressing functions by halting proliferation and inducing programmed cell death. In contrast, in advanced malignancies, TGF β preferentially exerts tumor-promoting actions by affecting the behavior of the cancer cells themselves, or by creating a favorable microenvironment for tumor growth [67]. The unresponsiveness of advanced cancers to the tumor-restricting properties of TGF β is a consequence of either genetic mutations of downstream cytostatic genes that, otherwise, are induced by the pathway, or mutations in components of the core signaling pathway, such as *TGFBR2* and *SMAD4* [68]. For example, CCA cells do not respond to the growth inhibitory effects of TGF β , due to high expression levels of cyclin D1. On the other hand, normal biliary epithelial cells, expressing physiological levels of cyclin D1, undergo cell cycle arrest in response to TGF β [69]. In the case of liver cancer, TGF β induces cytostatic and pro-apoptotic factors at early stages of cancer, but later on it promotes EMT and also stimulates the generation of cancer associated fibroblasts (CAF) in the tumor stroma, which maintain active TGF β signaling and contribute to metastasis [70].

3. TGF β Pathway in Cholangiocarcinoma

3.1. Genomic Alterations

In many cancers, genes encoding members of the TGF β pathway are frequently subject to mutations, reflecting the importance of this pathway in tumor progression [71]. Indeed, studies involving whole genome or exome sequencing from human CCA tissues have recorded genomic aberrations in main signaling pathways, and among them, TGF β has a special place [27,28]. Using a large cohort of 103 iCCA patients, Zou and coworkers identified RAS/PI3K, P53, and TGF β pathways to be influenced by alterations in the exome. *SMAD4* was described as one of the 25 significantly mutated genes, with a mutational rate of 4% ($P < 0.01$) [28]. In another study, *SMAD4* was mutated in 3.6% of patients with iCCA ($n = 55$) and in 25% of patients with eCCA ($n = 20$) [72]. Although, this difference in the frequency of *SMAD4* mutations between iCCA and eCCA was not statistically significant ($P = 0.333$), probably as a result of a limited number of eCCA cases, it suggests that eCCA exhibits a molecular phenotype that resembles pancreatic cancer rather than iCCA [29,72]. The expression of *SMAD4* was also evaluated by immunohistochemistry in normal liver and iCCA tissues with different differentiation status and clinical stages. *SMAD4* inactivation was found in 22 out of 49 iCCA specimens (44.9%, $P = 0.029$), whereas all normal liver tissues ($n = 9$) expressed *SMAD4*. A significant negative correlation between *SMAD4* expression and advanced clinical stages was highlighted [73]. Indeed, loss of *SMAD4* expression was associated with lymph node and intrahepatic metastasis ($P < 0.001$), poorly to moderately differentiated histological grade ($P = 0.013$), and advanced TNM stage ($P = 0.018$). In addition, the expression of *SMAD4* and cell cycle regulators (P53, P16, P27, cyclin D1, and Rb) was measured in 42 resected iCCAs by immunohistochemistry. Loss of expression of *SMAD4* was reported in 45.2% of cases, as well as genes encoding cell cycle inhibitors (e.g., 35.7% for P16) [74]. Loss of *SMAD4* was positively correlated with advanced pTNM stage ($P = 0.039$). Taken together, these studies suggest that different mechanisms contribute to inactivate *SMAD4* and possibly the tumor suppressive arm of the TGF β pathway, both in eCCA, involving *SMAD4* inactivating point mutations, and in iCCA, involving a transcriptional down-regulation of *SMAD4*.

3.2. TGF β Regulates CCA Tumor Progression

TGF β is one of the main signaling pathways that promotes CCA progression. Many studies support the TGF β -mediated induction of EMT in CCA cell lines [75]. Stimulation of human CCKS-1 and TFK-1 cells with TGF β led to a significant induction of SNAIL, VIM, and S100A4 and the reduction of E-cadherin and cytokeratin 19, thereby promoting migration and invasion [76]. TGF β also exhibits pro-EMT functions in iCCA cell lines KKKU-M213 and HuCCA-1. Enhanced expression of VIM and SLUG and secretion of the metalloproteinase MMP9, accompanied by a concomitant increase in cell migration and invasion, were observed after TGF β treatment [77]. TGF β stimulation activated both SMAD-dependent and SMAD-independent pathways, exemplified by the induced phosphorylation of SMAD2/3 and ERK1/2, respectively. Interestingly, inhibition of the kinase activity of MEK diminished the induction of EMT but, in contrast, potentiated the anti-proliferative effects of TGF β [77]. This observation raises the possibility that selective blockade of the noncanonical TGF β signaling may suppress the pro-tumorigenic, while preserving the anti-tumorigenic effects of TGF β in CCA. Disruption of TGF β pathway in human CCA primary cell cultures, using the T β RI kinase inhibitor LY2157299 (galunisertib), compromised cell migration. Notably, the same inhibitor did not influence cell cycle progression or apoptosis, indicating a selective inhibitory effect towards EMT [78]. Moreover, inhibition of the kinase CK2, which is linked to TGF β signaling, attenuated proliferation and increased apoptosis of the primary CCA cells [78]. The observed effects propose a combinatorial use of T β RI and CK2 kinase inhibitors for the treatment of CCAs. In another report, treatment of Huh28 and RBE CCA cell lines with the 3-hydroxy-3-methylglutaryl-coenzyme-CoA (HMG-CoA) reductase inhibitor, lovastatin, resulted in reduced TGF β 1 expression as well as reduced tumor cell proliferation and migration [79]. An interesting link between TGF β signaling, EMT, and the generation

of CCA cells with stemness properties was provided by Shuang and co-workers [80]. According to this study, exposure of TFK-1 cells to TGF β resulted in the acquisition of a mesenchymal phenotype and an increase in the population of cells positive for the cancer stem cell marker aldehyde dehydrogenase1 (ALDH1). Moreover, decreased cell death rates of TFK-1 cells in response to the DNA-damaging agent 5-fluorouracil (5-FU) was observed as a consequence of TGF β treatment, suggesting that TGF β confers chemoresistance of CCA cells to anti-cancer drugs [80]. It is worth noting that BMP7, a ligand that belongs to the TGF β family of proteins but signals through a different combination of receptors and effector SMADs, is capable of antagonizing the effects of TGF β -induced EMT in CCAs. Indeed, combinatorial treatment of M213 and M139 CCA cells with TGF β 1 and BMP7 attenuated migration and the increased expression of the EMT markers TWIST and N-cadherin, observed after TGF β 1 stimulation [81].

The effects of hepatitis B (HBV) and C (HCV) viruses on the progression of iCCA were evaluated in vivo using zebrafish as an experimental model. Livers from animals overexpressing both hepatitis B virus X (HBx) and hepatitis C virus core (HCP) proteins developed fibrosis and iCCA characterized by activated noncanonical TGF β signaling, exemplified by enhanced MAPK and SMAD linker phosphorylation. In addition, in vivo disruption of TGF β 1 expression by morpholinos attenuated fibrosis and iCCA progression, suggesting the importance of TGF β signaling during hepatitis-induced iCCA [82]. Moreover, experiments in transgenic zebrafish, whereby the expression of *tgfb1* was specifically induced in hepatocytes, showed increased incidence of HCC and CCA as a consequence of *tgfb1* chronic expression. Interestingly, HCC tumors were characterized by a switch from Smad-dependent to Erk-dependent TGF β pathway, whereas CCA tumors exhibited activation of both canonical and Erk pathways [83].

On the other hand, there are very few reports about a tumor-suppressing role of the TGF β pathway in CCAs. TGF β -treated RBE human CCA cells exhibited elevated apoptosis, a process that was SMAD-dependent and augmented by inhibition of the c-Jun N-terminal kinase (JNK), using the chemical inhibitor SP600125 [84]. Blocking JNK activity not only resulted in enhanced TGF β -induced apoptosis, but also in increased levels of C-terminal phosphorylated SMAD2 and SMAD3 and a general induction of TGF β -dependent transcriptional responses. Nevertheless, JNK inhibition did not influence SMAD linker phosphorylation, implying alternative molecular mechanisms for the inhibitory role of JNK on SMAD activation, in this context [84]. In a recent report, PTPN3, a protein tyrosine phosphatase which acts as a tumor suppressor, was shown to enhance T β RI stability, independently of its catalytic activity. PTPN3 was described as an antagonist of SMURF2 by overlapping with its binding domain to T β RI. SMURF2 is a well-known ubiquitin E3 ligase, recruited by SMAD7 and targeting T β RI for proteasomal degradation [85]. Interestingly, PTPN3 point mutations (L232R) may disrupt its interaction with T β RI, allowing SMAD7/ubiquitin E3 ligase complexes to exert their inhibitory role towards T β RI and, thus, abolishing TGF β pro-cytostatic effects [86]. Mutant PTPN3 L232R is frequently found in iCCA and results in loss of its anti-tumorigenic function. This finding could be of clinical importance for the subset of patients that carry PTPN3 L232R mutations, although a direct correlation between mutant PTPN3 and inactivation of TGF β signaling in iCCA patients was not addressed in this study.

3.3. Noncoding RNAs as Emerging Effectors of TGF β Signaling during CCA Progression

Noncoding RNA (ncRNA) are transcripts that lack protein-coding potential. Instead, they are transcribed from genes and perform regulatory or structural roles in cells. Noncoding RNA longer than 200 nucleotides are defined as long ncRNA (lncRNA). The rest are collectively designated as short ncRNA and, among many types, include the microRNA (miRNA), whose functional role is to target messenger RNA (mRNA) for degradation or to inhibit mRNA translation, thereby limiting gene expression [87]. The role of ncRNA in cancer progression, downstream of TGF β signaling, has been appreciated during the last years. For instance, in HCC, the lncRNA activated by TGF β (lncRNA-ATB) is a target gene of TGF β signaling and contributes to metastasis by sponging miR-200,

thereby stabilizing the EMT transcription factors ZEB1 and ZEB2, and by stabilizing IL11 mRNA, thus favoring tumor cell dissemination [88].

The crosstalk between TGF β signaling and miRNA in modulating EMT has been investigated in samples from patients with CCA. After measuring the expression levels of epithelial and mesenchymal markers in tumor and adjacent nontumor tissues from 20 patients, Zhang and co-workers observed low expression of the epithelial markers E-cadherin and miR-200b, and high expression of the mesenchymal markers fibronectin and α -smooth muscle actin (α -SMA) in the tumor tissues [89]. In addition, higher levels of TGF β 1 were detected in the tumor tissues, as compared to nontumor adjacent tissues. In vitro experiments using human HCCC and RBE cell lines stimulated with TGF β confirmed the downregulation of miR-200b and the establishment of an EMT transcriptional program. The mRNA encoding AP2 α and MAPK7 proteins were identified as direct targets of miR-200b. Enforced expression of miR-200b led to decreased tumor formation and EMT in an in vivo mouse model, highlighting its tumor-suppressing role and the importance of its repression by TGF β to elicit the EMT program [89]. In a rat model of iCCA, increased tumor size and enhanced intrahepatic metastasis were detected upon overexpression of TGF β 1 [90]. In addition, TGF β 1 promoted colony formation in a rat bile duct epithelial cell line BDE-Neu. The authors also used RNA interference to deplete TGF β 1 and chemical inhibition of T β RI kinase activity by LY2157299 and SB431542 to further confirm that both manipulations negatively affected cell growth and migration of RBE and SSP25 CCA cell lines. At the molecular level, miR-34a was identified as a central effector downstream of TGF β signaling. The expression of miR-34a was reduced in response to TGF β , which was followed by an increase in the expression of the direct targets of miR-34a, including CDK6, cyclin D1, and c-Met. Stabilization of CDK6 and cyclin D1 promoted cell growth, and elevated c-Met levels reinforced migration [90]. In another study, the anti-migratory role of miR-34a in CCA was also established, although an alternative mechanism for the miR-34a-mediated suppression of EMT has been described. According to Qiao and co-workers, miR-34a targeted SMAD4 and suppressed TGF β -induced EMT in vitro. Moreover, the levels of miR-34a and SMAD4 were inversely correlated in human eCCA tissues, with decreased miR-34a and increased SMAD4 expression being observed [91]. The miRNA miR-29a was suppressed by TGF β in the CCA cell lines FRH-0201 and CCLP-1. In addition, CCA tissues expressed lower miR-29a levels compared to adjacent nontumor tissues. Overexpression of miR-29a dampened the TGF β -induced cell migration and invasion in CCA cells. The histone deacetylase HDAC4 was identified as a direct target of miR-29a and rescue experiments. Using simultaneous overexpression of both HDAC4 and miR-29a showed a reversion of the EMT phenotype, which was abolished, due to miR-29a overexpression alone. Furthermore, HDAC4 promoted EMT in CCA cells, although the direct molecular targets of HDAC4 were not deeply investigated [92].

A recent report highlighted the importance of intercellular transfer of miRNA through extracellular vesicles (EVs) during CCA progression. As an example, TGF β represses the expression of miR-30e, a negative regulator of SNAIL in HuCCT1 cells. Enforced expression of miR-30e results in a decreased expression of SNAIL1 and of several EMT markers. In addition, miR-30e was packaged in EVs and transported to recipient HuCCT1 cells, where it exhibited its anti-EMT functions. Interestingly, TGF β treatment reduced the abundance of miR-30e in EVs of CCA cells, suggesting a negative role of miR-30e in regulating EMT under physiological conditions [93].

Although several studies have identified lncRNA associated with CCA progression [94], the knowledge concerning lncRNA acting as regulators of CCA progression in response to TGF β signaling is currently limited. Nevertheless, the lncRNA TGF β -induced long noncoding RNA (TLINC) was reported to be highly expressed in response to TGF β stimulation and to facilitate a pro-inflammatory microenvironment by enhancing cytokines, such as IL8 [95]. This finding strengthens the idea that additional lncRNAs could be effectors of TGF β -regulated responses in CCA.

More recently, circular RNA (circRNA) emerged in the literature as a new class of ncRNA that may play a critical role in cancer. Numerous circRNA are generated from alternative back-splicing of coding and ncRNA, forming continuous loop without 3' and 5' extremities. Recent data highlighted

the deregulation of circRNA in several cancers [96]. At the functional level, evidence indicates that circRNA play a key regulatory role at transcriptional and translational levels by possibly acting as miRNA sponges, or as scaffolds for RNA binding proteins (RBP) to form RNA-protein complexes [97]. The singular circular structure of circRNA renders them extremely resistant to exoribonucleases. Therefore, circRNA exhibit an expanded half-life, as compared to their linear counterpart RNA. Moreover, studies demonstrated that circRNA may also be found circulating in body fluids, freely or in exosomes [98]. Altogether, these characteristics are promising regarding the capability of circRNA as reliable diagnostic and predictive biomarkers in cancer. Although there is currently no report about circRNA deregulation in CCA, this new class of ncRNA could act as key regulator in the TGF β network and thereby in cancer. Notably, Goodal et al. demonstrated that TGF β could promote RNA circularization by inducing the binding of the RBP Quaking on introns flanking circular junctions, which could foster the pro-oncogenic feature of some circRNA [99]. These data reinforce the idea that circRNA could also be effectors of TGF β -mediated responses in cancer, including CCAs.

3.4. Therapeutic Targeting of the TGF β Signaling in CCA

The importance of the TGF β signaling in CCA renders this pathway a promising target for developing anti-tumor therapies. Thus, drugs that specifically target the TGF β pathway at different levels (i.e., maturation of latent TGF β to active TGF β dimers, ligand binding to its receptors, T β RI kinase activity) have been or are currently tested for possible anti-tumor effects in different cancers, including liver cancers [100–104]. In CCA, anti-TGF β -based therapeutic strategies have been mainly evaluated in preclinical models. Thus, *in vivo* experiments using a rat model of induced liver fibrosis showed decreased levels of fibrosis and CCA in animals treated with a neutralizing monoclonal antibody against TGF β . In contrast, control animals exhibited extensive liver fibrosis, which, eventually, led to the development of larger tumors [105]. Other mouse xenograft models showed increased tumor dissemination and larger tumors at the site of injection when mice were injected with CCKS-1 cells and simultaneously treated with TGF β 1 as compared to vehicle control-treated mice. Interestingly, the pro-tumorigenic effects of TGF β were abolished upon administration of a soluble form of T β RII that competes for binding to TGF β ligands with the membrane-bound T β RII [76]. In addition to these preclinical models, ongoing clinical trials make use of the chimeric antibody M7824, which is composed of the extracellular domain of human T β RII and the C-terminus of human anti-PD-L1 heavy chain [106,107]. M7824 has a double anti-tumor function as it serves as a trap for TGF β ligand binding at the tumor microenvironment and restricts the immune checkpoint factor programmed cell death ligand-1 (PD-L1), thereby restoring immune responses against the tumor [106,107]. Thus, a recently started multicenter phase II clinical trial is evaluating M7824 monotherapy in locally advanced or metastatic second line biliary tract cancer, including CCAs and gallbladder cancer (NCT03833661).

4. Conclusions

This overview of TGF β functions in liver homeostasis underlined the critical role of its dysregulation in cancer onset and progression. The fact that CCA evolves in a desmoplastic microenvironment, whereby TGF β is extremely abundant and frequently associated with a poor prognosis, emphasizes the clinical relevance of TGF β -targeted-therapies. Unfortunately, the literature described TGF β as an elusive target considering that, depending on the cell transcriptional context, it exhibits a dichotomous action. Thus, any therapeutic strategy aiming at modulating the TGF β pathway must consider its potential repercussion, either by repressing the pro-apoptotic and tumor suppressor arm, or by improving the pro-oncogenic and pro-metastatic action. A better insight into the molecular mechanisms regulating the functional duality of TGF β could improve the efficiency of targeted therapies for a better management of patients with CCA (Figure 2).




Figure 2. TGFβ signaling in cholangiocarcinoma (CCA) progression. TGFβ activates SMAD-dependent (canonical) and SMAD-independent (non-canonical) pathways in order to evoke transcriptional programs that, ultimately, regulate physiological responses in CCA. Consistent with its dual role in cancer, TGFβ can either prevent CCA progression, by inducing apoptosis, or enhance CCA progression, by promoting EMT, migration, invasion, and suppression of the immune system. Many coding and noncoding TGFβ-target genes can mediate the effects of the pathway during CCA progression and a number of them could potentially serve as drugable targets and biomarkers of CCA. Current therapeutic approaches aim at targeting components of the core pathway, such as the TGFβ ligands and the type I TGFβ receptor (TβRI).

The latest evidence promises a bright future for ncRNAs including miRNA, lncRNA, and circRNA as innovative downstream TGFβ effectors, or as clinically relevant biomarkers in CCA (Figure 2). A better understanding of the intricate and coordinated network of coding and ncRNAs will certainly allow us to elucidate how the functional duality of TGFβ in CCA is regulated. Some of these ncRNAs are also found circulating in body fluids and could represent specific biomarkers for a better management of patients with CCA.

Funding: This research was funded by Inserm, Université de Rennes 1, INCa, and ITMO Cancer AVIESAN (Alliance Nationale pour les Sciences de la Vie et de la Santé) dans le cadre du Plan cancer (Noncoding RNA in cancerology: Fundamental to translational; [Grant C18007NS to C.C.]). P.P. is supported by a post-doctoral fellowship from ITMO Cancer AVIESAN. C.L. is supported by a PhD fellowship from Université de Rennes 1. The authors of this review article are members of the European Network for the Study of Cholangiocarcinoma and participate in the initiative COST Action EUROCHOLANGIO-NET granted by the COST Association (CA18122).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Rizvi, S.; Gores, G.J. Pathogenesis, Diagnosis, and Management of Cholangiocarcinoma. *Gastroenterology* **2013**, *145*, 1215–1229. [[CrossRef](#)] [[PubMed](#)]
2. Bertuccio, P.; Malvezzi, M.; Carioli, G.; Hashim, D.; Boffetta, P.; El-Serag, H.B.; La Vecchia, C.; Negri, E. Global trends in mortality from intrahepatic and extrahepatic cholangiocarcinoma. *J. Hepatol.* **2019**, *71*, 104–114. [[CrossRef](#)] [[PubMed](#)]
3. Sripa, B.; Pairojkul, C. Cholangiocarcinoma: Lessons from Thailand. *Curr. Opin. Gastroenterol.* **2008**, *24*, 349–356. [[CrossRef](#)] [[PubMed](#)]
4. Tyson, G.L.; El-Serag, H.B. Risk Factors of Cholangiocarcinoma. *Hepatology* **2011**, *54*, 173–184. [[CrossRef](#)] [[PubMed](#)]
5. Shaib, Y.H.; El-Serag, H.B.; Davila, J.A.; Morgan, R.; McGlynn, K.A. Risk factors of intrahepatic cholangiocarcinoma in the United States: A case-control study. *Gastroenterology* **2005**, *128*, 620–626. [[CrossRef](#)] [[PubMed](#)]
6. Lee, J.J.; Schindera, S.T.; Jang, H.-J.; Fung, S.; Kim, T.K. Cholangiocarcinoma and its mimickers in primary sclerosing cholangitis. *Abdom. Radiol.* **2017**, *42*, 2898–2908. [[CrossRef](#)] [[PubMed](#)]
7. Welzel, T.M.; Graubard, B.I.; Zeuzem, S.; El-Serag, H.B.; Davila, J.A.; McGlynn, K.A. Metabolic syndrome increases the risk of primary liver cancer in the United States: A study in the SEER-Medicare database. *Hepatology* **2011**, *54*, 463–471. [[CrossRef](#)] [[PubMed](#)]
8. Jarnagin, W.R.; Fong, Y.; Burke, E.C.; Bodniewicz, J.; Youssef, M.; Klimstra, D.; Blumgart, L.H.; DeMatteo, R.P.; Gonen, M. Staging, Resectability, and Outcome in 225 Patients with Hilar Cholangiocarcinoma. *Ann. Surg.* **2001**, *234*, 507–519. [[CrossRef](#)]
9. Bridgewater, J.; Galle, P.R.; Khan, S.A.; Llovet, J.M.; Park, J.-W.; Patel, T.; Pawlik, T.M.; Gores, G.J. Guidelines for the diagnosis and management of intrahepatic cholangiocarcinoma. *J. Hepatol.* **2014**, *60*, 1268–1289. [[CrossRef](#)]
10. Sulpice, L.; Desille, M.; Turlin, B.; Fautrel, A.; Boudjema, K.; Clément, B.; Coulouarn, C. Gene expression profiling of the tumor microenvironment in human intrahepatic cholangiocarcinoma. *Genom. Data* **2016**, *7*, 229–232. [[CrossRef](#)]
11. Sulpice, L.; Rayar, M.; Desille, M.; Turlin, B.; Fautrel, A.; Boucher, E.; Llamas-Gutierrez, F.; Meunier, B.; Boudjema, K.; Clement, B.; et al. Molecular profiling of stroma identifies osteopontin as an independent predictor of poor prognosis in intrahepatic cholangiocarcinoma. *Hepatology* **2013**, *58*, 1992–2000. [[CrossRef](#)] [[PubMed](#)]
12. Fabris, L.; Perugorria, M.J.; Mertens, J.; Björkström, N.K.; Cramer, T.; Lleo, A.; Solinas, A.; Sängler, H.; Lukacs-Kornek, V.; Moncsek, A.; et al. The tumour microenvironment and immune milieu of cholangiocarcinoma. *Liver Int.* **2019**, *39*, 63–78. [[CrossRef](#)] [[PubMed](#)]
13. Høgdall, D.; Lewińska, M.; Andersen, J.B. Desmoplastic Tumor Microenvironment and Immunotherapy in Cholangiocarcinoma. *Trends Cancer* **2018**, *4*, 239–255. [[CrossRef](#)] [[PubMed](#)]
14. Rimassa, L.; Personeni, N.; Aghemo, A.; Lleo, A. The immune milieu of cholangiocarcinoma: From molecular pathogenesis to precision medicine. *J. Autoimmun.* **2019**, *100*, 17–26. [[CrossRef](#)] [[PubMed](#)]
15. Huynh, J.; Etemadi, N.; Hollande, F.; Ernst, M.; Buchert, M.; Hollande, F. The JAK/STAT3 axis: A comprehensive drug target for solid malignancies. *Semin. Cancer Biol.* **2017**, *45*, 13–22. [[CrossRef](#)] [[PubMed](#)]
16. Yang, X.W.; Li, L.; Hou, G.J.; Yan, X.Z.; Xu, Q.G.; Chen, L.; Zhang, B.H.; Shen, F. STAT3 overexpression promotes metastasis in intrahepatic cholangiocarcinoma and correlates negatively with surgical outcome. *Oncotarget* **2017**, *8*, 7710–7721. [[CrossRef](#)] [[PubMed](#)]
17. Clapéron, A.; Mergey, M.; Ho-Bouldoires, T.H.N.; Vignjevic, D.M.; Wendum, D.; Chrétien, Y.; Merabtene, F.; Frazao, A.; Paradis, V.; Housset, C.; et al. EGF/EGFR axis contributes to the progression of cholangiocarcinoma through the induction of an epithelial-mesenchymal transition. *J. Hepatol.* **2014**, *61*, 325–332. [[CrossRef](#)]
18. Menakongka, A.; Suthiphongchai, T. Involvement of PI3K and ERK1/2 pathways in hepatocyte growth factor-induced cholangiocarcinoma cell invasion. *World J. Gastroenterol.* **2010**, *16*, 713–722. [[CrossRef](#)]
19. Geisler, F.; Strazzabosco, M. Emerging roles of Notch signaling in liver disease. *Hepatology* **2015**, *61*, 382–392. [[CrossRef](#)]

20. Sekiya, S.; Suzuki, A. Intrahepatic cholangiocarcinoma can arise from Notch-mediated conversion of hepatocytes. *J. Clin. Investig.* **2012**, *122*, 3914–3918. [[CrossRef](#)]
21. Omenetti, A.; Choi, S.; Michelotti, G.; Diehl, A.M. Hedgehog signaling in the liver. *J. Hepatol.* **2011**, *54*, 366–373. [[CrossRef](#)] [[PubMed](#)]
22. Gonnissen, A.; Isebaert, S.; Haustermans, K. Targeting the Hedgehog signaling pathway in cancer: Beyond Smoothed. *Oncotarget* **2015**, *6*, 13899–13913. [[CrossRef](#)] [[PubMed](#)]
23. Razumilava, N.; Gradilone, S.A.; Smoot, R.L.; Mertens, J.C.; Bronk, S.F.; Sirica, A.E.; Gores, G.J. Non-canonical Hedgehog signaling contributes to chemotaxis in cholangiocarcinoma. *J. Hepatol.* **2014**, *60*, 599–605. [[CrossRef](#)] [[PubMed](#)]
24. El Khatib, M.; Kalnytska, A.; Palagani, V.; Kossatz, U.; Manns, M.P.; Malek, N.P.; Wilkens, L.; Plentz, R.R. Inhibition of hedgehog signaling attenuates carcinogenesis in vitro and increases necrosis of cholangiocellular carcinoma. *Hepatology* **2013**, *57*, 1035–1045. [[CrossRef](#)] [[PubMed](#)]
25. Fujimoto, A.; Furuta, M.; Shiraiishi, Y.; Gotoh, K.; Kawakami, Y.; Arihiro, K.; Nakamura, T.; Ueno, M.; Ariizumi, S.; Nguyen, H.H.; et al. Whole-genome mutational landscape of liver cancers displaying biliary phenotype reveals hepatitis impact and molecular diversity. *Nat. Commun.* **2015**, *6*, 6120. [[CrossRef](#)] [[PubMed](#)]
26. Sia, D.; Losic, B.; Moeini, A.; Cabellos, L.; Hao, K.; Revill, K.; Bonal, D.; Miltiadous, O.; Zhang, Z.; Hoshida, Y.; et al. Massive parallel sequencing uncovers actionable FGFR2–PPHLN1 fusion and ARAF mutations in intrahepatic cholangiocarcinoma. *Nat. Commun.* **2015**, *6*, 6087. [[CrossRef](#)]
27. Nakamura, H.; Arai, Y.; Totoki, Y.; Shiota, T.; ElZawahry, A.; Kato, M.; Hama, N.; Hosoda, F.; Urushidate, T.; Ohashi, S.; et al. Genomic spectra of biliary tract cancer. *Nat. Genet.* **2015**, *47*, 1003–1010. [[CrossRef](#)] [[PubMed](#)]
28. Zou, S.; Li, J.; Zhou, H.; Frech, C.; Jiang, X.; Chu, J.S.C.; Zhao, X.; Li, Y.; Li, Q.; Wang, H.; et al. Mutational landscape of intrahepatic cholangiocarcinoma. *Nat. Commun.* **2014**, *5*, 5696. [[CrossRef](#)]
29. Ong, C.K.; Subimerb, C.; Pairojkul, C.; Wongkham, S.; Cutcutache, I.; Yu, W.; McPherson, J.R.; E Allen, G.; Ng, C.C.Y.; Wong, B.H.; et al. Exome sequencing of liver fluke-associated cholangiocarcinoma. *Nat. Genet.* **2012**, *44*, 690–693. [[CrossRef](#)]
30. Gao, Q.; Zhao, Y.; Wang, X.; Guo, W.; Gao, S.; Wei, L.; Shi, J.; Shi, G.; Wang, Z.; Zhang, Y.; et al. Activating Mutations in PTPN3 Promote Cholangiocarcinoma Cell Proliferation and Migration and Are Associated with Tumor Recurrence in Patients. *Gastroenterology* **2014**, *146*, 1397–1407. [[CrossRef](#)]
31. Bergeat, D.; Fautrel, A.; Turlin, B.; Merdrignac, A.; Rayar, M.; Boudjema, K.; Coulouarn, C.; Sulpice, L. Impact of stroma LOXL2 overexpression on the prognosis of intrahepatic cholangiocarcinoma. *J. Surg. Res.* **2016**, *203*, 441–450. [[CrossRef](#)] [[PubMed](#)]
32. Sulpice, L.; Rayar, M.; Turlin, B.; Boucher, E.; Bellaud, P.; Desille, M.; Meunier, B.; Clément, B.; Boudjema, K.; Coulouarn, C. Epithelial cell adhesion molecule is a prognosis marker for intrahepatic cholangiocarcinoma. *J. Surg. Res.* **2014**, *192*, 117–123. [[CrossRef](#)] [[PubMed](#)]
33. Borad, M.J.; Champion, M.D.; Egan, J.B.; Liang, W.S.; Fonseca, R.; Bryce, A.H.; McCullough, A.E.; Barrett, M.T.; Hunt, K.; Patel, M.D.; et al. Integrated Genomic Characterization Reveals Novel, Therapeutically Relevant Drug Targets in FGFR and EGFR Pathways in Sporadic Intrahepatic Cholangiocarcinoma. *PLoS Genet.* **2014**, *10*, e1004135. [[CrossRef](#)] [[PubMed](#)]
34. Golub, D.; Iyengar, N.; Dogra, S.; Wong, T.; Bready, D.; Tang, K.; Modrek, A.S.; Placantonakis, D.G. Mutant Isocitrate Dehydrogenase Inhibitors as Targeted Cancer Therapeutics. *Front. Oncol.* **2019**, *9*, 417. [[CrossRef](#)] [[PubMed](#)]
35. David, C.J.; Massagué, J. Contextual determinants of TGF β action in development, immunity and cancer. *Nat. Rev. Mol. Cell Biol.* **2018**, *19*, 419–435. [[CrossRef](#)] [[PubMed](#)]
36. Wang, J.; Shao, N.; Ding, X.; Tan, B.; Song, Q.; Wang, N.; Jia, Y.; Ling, H.; Cheng, Y. Crosstalk between transforming growth factor- β signaling pathway and long non-coding RNAs in cancer. *Cancer Lett.* **2016**, *370*, 296–301. [[CrossRef](#)] [[PubMed](#)]
37. Heldin, C.H.; Moustakas, A. Role of Smads in TGF β signaling. *Cell Tissue Res.* **2012**, *347*, 21–36. [[CrossRef](#)] [[PubMed](#)]
38. Chaikuad, A.; Bullock, A.N. Structural Basis of Intracellular TGF- β Signaling: Receptors and Smads. *Cold Spring Harb. Perspect. Biol.* **2016**, *8*, a022111. [[CrossRef](#)] [[PubMed](#)]

39. Derynck, R.; Budi, E.H. Specificity, versatility, and control of TGF- β family signaling. *Sci. Signal.* **2019**, *12*, eaav5183. [[CrossRef](#)]
40. Miyazono, K.; Katsuno, Y.; Koinuma, D.; Ehata, S.; Morikawa, M. Intracellular and extracellular TGF- β signaling in cancer: Some recent topics. *Front. Med.* **2018**, *12*, 387–411. [[CrossRef](#)]
41. Tang, J.; Gifford, C.C.; Samarakoon, R.; Higgins, P.J. Deregulation of Negative Controls on TGF- β 1 Signaling in Tumor Progression. *Cancers* **2018**, *10*, 159. [[CrossRef](#)] [[PubMed](#)]
42. Luo, K. Signaling Cross Talk between TGF- β /Smad and Other Signaling Pathways. *Cold Spring Harb. Perspect. Boil.* **2017**, *9*, a022137. [[CrossRef](#)] [[PubMed](#)]
43. Fouassier, L.; Marzioni, M.; Afonso, M.B.; Dooley, S.; Gaston, K.; Giannelli, G.; Rodrigues, C.M.P.; Lozano, E.; Mancarella, S.; Segatto, O.; et al. Signalling networks in cholangiocarcinoma: Molecular pathogenesis, targeted therapies and drug resistance. *Liver Int.* **2019**, *39* (Suppl. 1), 43–62. [[CrossRef](#)] [[PubMed](#)]
44. Nishio, M.; Sugimachi, K.; Goto, H.; Wang, J.; Morikawa, T.; Miyachi, Y.; Takano, Y.; Hikasa, H.; Itoh, T.; Suzuki, S.O.; et al. Dysregulated YAP1/TAZ and TGF-beta signaling mediate hepatocarcinogenesis in Mob1a/1b-deficient mice. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E71–E80. [[CrossRef](#)] [[PubMed](#)]
45. Yamada, D.; Kobayashi, S.; Wada, H.; Kawamoto, K.; Marubashi, S.; Eguchi, H.; Ishii, H.; Nagano, H.; Doki, Y.; Mori, M. Role of crosstalk between interleukin-6 and transforming growth factor-beta 1 in epithelial–mesenchymal transition and chemoresistance in biliary tract cancer. *Eur. J. Cancer* **2013**, *49*, 1725–1740. [[CrossRef](#)]
46. Bataller, R.; Brenner, D.A. Liver fibrosis. *J. Clin. Investig.* **2005**, *115*, 209–218. [[CrossRef](#)] [[PubMed](#)]
47. Coulouarn, C.; Clément, B. Stellate cells and the development of liver cancer: Therapeutic potential of targeting the stroma. *J. Hepatol.* **2014**, *60*, 1306–1309. [[CrossRef](#)]
48. Caja, L.; Dituri, F.; Mancarella, S.; Caballero-Diaz, D.; Moustakas, A.; Giannelli, G.; Fabregat, I. TGF- β and the Tissue Microenvironment: Relevance in Fibrosis and Cancer. *Int. J. Mol. Sci.* **2018**, *19*, 1294. [[CrossRef](#)]
49. Zhang, Y.; Alexander, P.B.; Wang, X.F. TGF-beta Family Signaling in the Control of Cell Proliferation and Survival. *Cold Spring Harb. Perspect. Biol.* **2017**, *9*. [[CrossRef](#)]
50. Schuster, N.; Kriegstein, K. Mechanisms of TGF- β -mediated apoptosis. *Cell and Tissue Research* **2002**, *307*, 1–14. [[CrossRef](#)]
51. Yamashita, M.; Fatyol, K.; Jin, C.; Wang, X.; Liu, Z.; Zhang, Y.E. TRAF6 mediates Smad-independent activation of JNK and p38 by TGF- β . *Mol. Cell* **2008**, *31*, 918–924. [[CrossRef](#)] [[PubMed](#)]
52. Bender, H.; Wang, Z.; Schuster, N.; Kriegstein, K. TIEG1 facilitates transforming growth factor- beta -mediated apoptosis in the oligodendroglial cell line OLI-neu. *J. Neurosci. Res.* **2004**, *75*, 344–352. [[CrossRef](#)] [[PubMed](#)]
53. Jang, C.W.; Chen, C.H.; Chen, C.C.; Chen, J.Y.; Su, Y.H.; Chen, R.H. TGF-beta induces apoptosis through Smad-mediated expression of DAP-kinase. *Nat. Cell Biol.* **2002**, *4*, 51–58. [[CrossRef](#)] [[PubMed](#)]
54. Tsubakihara, Y.; Moustakas, A. Epithelial-Mesenchymal Transition and Metastasis under the Control of Transforming Growth Factor β . *Int. J. Mol. Sci.* **2018**, *19*, 3672. [[CrossRef](#)] [[PubMed](#)]
55. Aiello, N.M.; Kang, Y. Context-dependent EMT programs in cancer metastasis. *J. Exp. Med.* **2019**, *216*, 1016–1026. [[CrossRef](#)] [[PubMed](#)]
56. Zheng, X.; Carstens, J.L.; Kim, J.; Scheible, M.; Kaye, J.; Sugimoto, H.; Wu, C.C.; LeBleu, V.S.; Kalluri, R. Epithelial-to-mesenchymal transition is dispensable for metastasis but induces chemoresistance in pancreatic cancer. *Nature* **2015**, *527*, 525–530. [[CrossRef](#)] [[PubMed](#)]
57. Fischer, K.R.; Durrans, A.; Lee, S.; Sheng, J.; Li, F.; Wong, S.T.; Choi, H.; El Rayes, T.; Ryu, S.; Troeger, J.; et al. Epithelial-to-mesenchymal transition is not required for lung metastasis but contributes to chemoresistance. *Nature* **2015**, *527*, 472–476. [[CrossRef](#)] [[PubMed](#)]
58. Singh, A.; Settleman, J. EMT, cancer stem cells and drug resistance: An emerging axis of evil in the war on cancer. *Oncogene* **2010**, *29*, 4741–4751. [[CrossRef](#)]
59. Malfettone, A.; Soukupova, J.; Bertran, E.; Crosas-Molist, E.; Lastra, R.; Fernando, J.; Koudelkova, P.; Rani, B.; Fabra, Á.; Serrano, T.; et al. Transforming growth factor- β -induced plasticity causes a migratory stemness phenotype in hepatocellular carcinoma. *Cancer Lett.* **2017**, *392*, 39–50. [[CrossRef](#)]
60. Moustakas, A.; Heldin, C.-H. Mechanisms of TGF β -Induced Epithelial–Mesenchymal Transition. *J. Clin. Med.* **2016**, *5*, 63. [[CrossRef](#)]
61. Thepmalee, C.; Panya, A.; Junking, M.; Chieochansin, T.; Yenchitsomanus, P.-T. Inhibition of IL-10 and TGF- β receptors on dendritic cells enhances activation of effector T-cells to kill cholangiocarcinoma cells. *Hum. Vaccines Immunother.* **2018**, *14*, 1423–1431. [[CrossRef](#)] [[PubMed](#)]

62. Landskron, G.; De La Fuente, M.; Thuwajit, P.; Thuwajit, C.; Hermoso, M.A. Chronic Inflammation and Cytokines in the Tumor Microenvironment. *J. Immunol. Res.* **2014**, *2014*, 1–19. [[CrossRef](#)] [[PubMed](#)]
63. Shimizu, T.; Yokomuro, S.; Mizuguchi, Y.; Kawahigashi, Y.; Arima, Y.; Taniai, N.; Mamada, Y.; Yoshida, H.; Akimaru, K.; Tajiri, T. Effect of transforming growth factor- β 1 on human intrahepatic cholangiocarcinoma cell growth. *World J. Gastroenterol.* **2006**, *12*, 6316–6324. [[CrossRef](#)] [[PubMed](#)]
64. Raggi, C.; Correnti, M.; Sica, A.; Andersen, J.B.; Cardinale, V.; Alvaro, D.; Chiorino, G.; Forti, E.; Glaser, S.; Alpini, G.; et al. Cholangiocarcinoma stem-like subset shapes tumor-initiating niche by educating associated macrophages. *J. Hepatol.* **2017**, *66*, 102–115. [[CrossRef](#)] [[PubMed](#)]
65. Kitano, Y.; Okabe, H.; Yamashita, Y.I.; Nakagawa, S.; Saito, Y.; Umezaki, N.; Tsukamoto, M.; Yamao, T.; Yamamura, K.; Arima, K.; et al. Tumour-infiltrating inflammatory and immune cells in patients with extrahepatic cholangiocarcinoma. *Br. J. Cancer* **2018**, *118*, 171–180. [[CrossRef](#)]
66. Hasita, H.; Komohara, Y.; Okabe, H.; Masuda, T.; Ohnishi, K.; Lei, X.F.; Beppu, T.; Baba, H.; Takeya, M. Significance of alternatively activated macrophages in patients with intrahepatic cholangiocarcinoma. *Cancer Sci.* **2010**, *101*, 1913–1919. [[CrossRef](#)]
67. Huang, J.J.; Blobel, G.C. Dichotomous Roles of TGF- β in Human Cancer. *Biochem. Soc. Trans.* **2016**, *44*, 1441–1454. [[CrossRef](#)]
68. Massague, J. TGFbeta in Cancer. *Cell* **2008**, *134*, 215–230. [[CrossRef](#)]
69. Zen, Y.; Harada, K.; Sasaki, M.; Chen, T.-C.; Chen, M.-F.; Yeh, T.-S.; Jan, Y.-Y.; Huang, S.-F.; Nimura, Y.; Nakanuma, Y. Intrahepatic cholangiocarcinoma escapes from growth inhibitory effect of transforming growth factor- β 1 by overexpression of cyclin D1. *Lab. Invest.* **2005**, *85*, 572–581. [[CrossRef](#)]
70. Fabregat, I.; Caballero-Díaz, D. Transforming Growth Factor- β -Induced Cell Plasticity in Liver Fibrosis and Hepatocarcinogenesis. *Front. Oncol.* **2018**, *8*, 357. [[CrossRef](#)]
71. Seoane, J.; Gomis, R.R. TGF- β Family Signaling in Tumor Suppression and Cancer Progression. *Cold Spring Harb. Perspect. Biol.* **2017**, *9*, a022277. [[CrossRef](#)] [[PubMed](#)]
72. Churi, C.R.; Shroff, R.; Wang, Y.; Rashid, A.; Kang, H.C.; Weatherly, J.; Zuo, M.; Zinner, R.; Hong, D.; Meric-Bernstam, F.; et al. Mutation Profiling in Cholangiocarcinoma: Prognostic and Therapeutic Implications. *PLoS ONE* **2014**, *9*, e115383. [[CrossRef](#)] [[PubMed](#)]
73. Yan, X.-Q.; Zhang, W.; Zhang, B.-X.; Liang, H.-F.; Zhang, W.-G.; Chen, X.-P. Inactivation of Smad4 is a prognostic factor in intrahepatic cholangiocarcinoma. *Chin. Med. J.* **2013**, *126*, 3039–3043.
74. Kang, Y.K.; Kim, W.H.; Jang, J.J. Expression of G1-S modulators (p53, p16, p27, cyclin D1, Rb) and Smad4/Dpc4 in intrahepatic cholangiocarcinoma. *Hum. Pathol.* **2002**, *33*, 877–883. [[CrossRef](#)] [[PubMed](#)]
75. Vaquero, J.; Guedj, N.; Clapéron, A.; Ho-Bouldoires, T.H.N.; Paradis, V.; Fouassier, L. Epithelial-mesenchymal transition in cholangiocarcinoma: From clinical evidence to regulatory networks. *J. Hepatol.* **2017**, *66*, 424–441. [[CrossRef](#)] [[PubMed](#)]
76. Sato, Y.; Harada, K.; Itatsu, K.; Ikeda, H.; Kakuda, Y.; Shimomura, S.; Ren, X.S.; Yoneda, N.; Sasaki, M.; Nakanuma, Y. Epithelial-Mesenchymal Transition Induced by Transforming Growth Factor- β 1/Snail Activation Aggravates Invasive Growth of Cholangiocarcinoma. *Am. J. Pathol.* **2010**, *177*, 141–152. [[CrossRef](#)] [[PubMed](#)]
77. Sritananuwat, P.; Sueangoen, N.; Thummarati, P.; Islam, K.; Suthiphongchai, T. Blocking ERK1/2 signaling impairs TGF- β 1 tumor promoting function but enhances its tumor suppressing role in intrahepatic cholangiocarcinoma cells. *Cancer Cell Int.* **2017**, *17*, 85. [[CrossRef](#)] [[PubMed](#)]
78. Lustrì, A.M.; Di Matteo, S.; Fraveto, A.; Costantini, D.; Cantafora, A.; Napoletano, C.; Bragazzi, M.C.; Giuliani, F.; De Rose, A.M.; Berloco, P.B.; et al. TGF- β signaling is an effective target to impair survival and induce apoptosis of human cholangiocarcinoma cells: A study on human primary cell cultures. *PLoS ONE* **2017**, *12*, e0183932. [[CrossRef](#)]
79. Yang, S.H.; Lin, H.Y.; Changou, C.A.; Chen, C.H.; Liu, Y.R.; Wang, J.; Jiang, X.; Luh, F.; Yen, Y. Integrin beta3 and LKB1 are independently involved in the inhibition of proliferation by lovastatin in human intrahepatic cholangiocarcinoma. *Oncotarget* **2016**, *7*, 362–373. [[CrossRef](#)]
80. Shuang, Z.-Y.; Wu, W.-C.; Xu, J.; Lin, G.; Liu, Y.-C.; Lao, X.-M.; Zheng, L.; Li, S. Transforming growth factor- β 1-induced epithelial-mesenchymal transition generates ALDH-positive cells with stem cell properties in cholangiocarcinoma. *Cancer Lett.* **2014**, *354*, 320–328. [[CrossRef](#)]
81. Duangkumpha, K.; Techasen, A.; Loilome, W.; Namwat, N.; Thanan, R.; Khuntikeo, N.; Yongvanit, P. BMP-7 blocks the effects of TGF- β -induced EMT in cholangiocarcinoma. *Tumor Biol.* **2014**, *35*, 9667–9676. [[CrossRef](#)] [[PubMed](#)]

82. Liu, W.; Chen, J.-R.; Hsu, C.-H.; Li, Y.-H.; Chen, Y.-M.; Lin, C.-Y.; Huang, S.-J.; Chang, Z.-K.; Chen, Y.-C.; Lin, C.-H.; et al. A zebrafish model of intrahepatic cholangiocarcinoma by dual expression of hepatitis B Virus X and hepatitis C virus core protein in liver. *Hepatology* **2012**, *56*, 2268–2276. [[CrossRef](#)] [[PubMed](#)]
83. Yan, C.; Yang, Q.; Shen, H.-M.; Spitsbergen, J.M.; Gong, Z. Chronically high level of *tgfb1a* induction causes both hepatocellular carcinoma and cholangiocarcinoma via a dominant Erk pathway in zebrafish. *Oncotarget* **2017**, *8*, 77096–77109. [[CrossRef](#)] [[PubMed](#)]
84. Lin, Y.; Zhang, B.; Lu, Y.; Liang, H.; Ai, X.; Zhang, B.; Chen, X. JNK inhibitor SP600125 enhances TGF- β -induced apoptosis of RBE human cholangiocarcinoma cells in a Smad-dependent manner. *Mol. Med. Rep.* **2013**, *8*, 1623–1629. [[CrossRef](#)] [[PubMed](#)]
85. Lin, X.; Liang, M.; Feng, X.-H. Smurf2 Is a Ubiquitin E3 Ligase Mediating Proteasome-dependent Degradation of Smad2 in Transforming Growth Factor- β Signaling. *J. Boil. Chem.* **2000**, *275*, 36818–36822. [[CrossRef](#)] [[PubMed](#)]
86. Yuan, B.; Liu, J.; Cao, J.; Yu, Y.; Zhang, H.; Wang, F.; Zhu, Y.; Xiao, M.; Liu, S.; Ye, Y.; et al. PTPN3 acts as a tumor suppressor and boosts TGF-beta signaling independent of its phosphatase activity. *EMBO J.* **2019**. [[CrossRef](#)] [[PubMed](#)]
87. Bunch, H. Gene regulation of mammalian long non-coding RNA. *Mol. Genet. Genomics* **2018**, *293*, 1–15. [[CrossRef](#)] [[PubMed](#)]
88. Yuan, J.-H.; Yang, F.; Wang, F.; Ma, J.-Z.; Guo, Y.-J.; Tao, Q.-F.; Liu, F.; Pan, W.; Wang, T.-T.; Zhou, C.-C.; et al. A Long Noncoding RNA Activated by TGF- β Promotes the Invasion-Metastasis Cascade in Hepatocellular Carcinoma. *Cancer Cell* **2014**, *25*, 666–681. [[CrossRef](#)]
89. Zhang, D.; Li, H.; Jiang, X.; Cao, L.; Wen, Z.; Yang, X.; Xue, P. Role of AP-2 α and MAPK7 in the regulation of autocrine TGF- β /miR-200b signals to maintain epithelial-mesenchymal transition in cholangiocarcinoma. *J. Hematol. Oncol.* **2017**, *10*, 170. [[CrossRef](#)]
90. Huang, C.-K.; Aihara, A.; Iwagami, Y.; Yu, T.; Carlson, R.; Koga, H.; Kim, M.; Zou, J.; Casulli, S.; Wands, J.R. Expression of Transforming Growth Factor β 1 Promotes Cholangiocarcinoma Development and Progression. *Cancer Lett.* **2016**, *380*, 153–162. [[CrossRef](#)]
91. Qiao, P.; Li, G.; Bi, W.; Yang, L.; Yao, L.; Wu, D. microRNA-34a inhibits epithelial mesenchymal transition in human cholangiocarcinoma by targeting Smad4 through transforming growth factor-beta/Smad pathway. *BMC Cancer* **2015**, *15*, 469. [[CrossRef](#)]
92. Wang, H.; Li, C.; Jian, Z.; Ou, Y.; Ou, J. TGF- β 1 Reduces miR-29a Expression to Promote Tumorigenicity and Metastasis of Cholangiocarcinoma by Targeting HDAC4. *PLoS ONE* **2015**, *10*, e0136703. [[CrossRef](#)] [[PubMed](#)]
93. Ota, Y.; Takahashi, K.; Otake, S.; Tamaki, Y.; Okada, M.; Aso, K.; Makino, Y.; Fujii, S.; Ota, T.; Haneda, M. Extracellular vesicle-encapsulated miR-30e suppresses cholangiocarcinoma cell invasion and migration via inhibiting epithelial-mesenchymal transition. *Oncotarget* **2018**, *9*, 16400–16417. [[CrossRef](#)] [[PubMed](#)]
94. Angenard, G.; Merdrignac, A.; Louis, C.; Edeline, J.; Coulouarn, C. Expression of long non-coding RNA ANRIL predicts a poor prognosis in intrahepatic cholangiocarcinoma. *Dig. Liver Dis.* **2019**, *51*, 1337–1343. [[CrossRef](#)] [[PubMed](#)]
95. Merdrignac, A.; Angenard, G.; Allain, C.; Petitjean, K.; Bergeat, D.; Bellaud, P.; Fautrel, A.; Turlin, B.; Clément, B.; Dooley, S.; et al. A novel transforming growth factor beta-induced long noncoding RNA promotes an inflammatory microenvironment in human intrahepatic cholangiocarcinoma. *Hepatol. Commun.* **2018**, *2*, 254–269. [[CrossRef](#)]
96. Arnaiz, E.; Sole, C.; Manterola, L.; Iparraguirre, L.; Otaegui, D.; Lawrie, C.H. CircRNAs and cancer: Biomarkers and master regulators. *Semin. Cancer Biol.* **2018**. [[CrossRef](#)]
97. Salzman, J. Circular RNA Expression: Its Potential Regulation and Function. *Trends Genet.* **2016**, *32*, 309–316. [[CrossRef](#)]
98. Zhang, Z.; Yang, T.; Xiao, J. Circular RNAs: Promising Biomarkers for Human Diseases. *EBioMedicine* **2018**, *34*, 267–274. [[CrossRef](#)]
99. Conn, S.J.; Pillman, K.A.; Toubia, J.; Conn, V.M.; Salmanidis, M.; Phillips, C.A.; Roslan, S.; Schreiber, A.W.; Gregory, P.A.; Goodall, G.J. The RNA Binding Protein Quaking Regulates Formation of circRNAs. *Cell* **2015**, *160*, 1125–1134. [[CrossRef](#)]
100. Akhurst, R.J.; Hata, A. Targeting the TGF β signalling pathway in disease. *Nat. Rev. Drug Discov.* **2012**, *11*, 790–811. [[CrossRef](#)]

101. Halder, S.K.; Beauchamp, R.D.; Datta, P.K. A Specific Inhibitor of TGF- β Receptor Kinase, SB-431542, as a Potent Antitumor Agent for Human Cancers. *Neoplasia* **2005**, *7*, 509–521. [[CrossRef](#)] [[PubMed](#)]
102. Giannelli, G.; Villa, E.; Lahn, M. Transforming Growth Factor- β as a Therapeutic Target in Hepatocellular Carcinoma. *Cancer Res.* **2014**, *74*, 1890–1894. [[CrossRef](#)] [[PubMed](#)]
103. Kelley, R.; Gane, E.; Assenat, E.; Siebler, J.; Galle, P.; Merle, P.; Hourmand, I.; Cleverly, A.; Zhao, Y.; Georguieva, I.; et al. A Phase 2 Study of Galunisertib (TGF- β 1 Receptor Type I Inhibitor) and Sorafenib in Patients With Advanced Hepatocellular Carcinoma. *Clin. Transl. Gastroenterol.* **2019**, *10*, e00056. [[CrossRef](#)] [[PubMed](#)]
104. Chen, Y.; Di, C.; Zhang, X.; Wang, J.; Wang, F.; Yan, J.F.; Xu, C.; Zhang, J.; Zhang, Q.; Li, H.; et al. Transforming growth factor beta signaling pathway: A promising therapeutic target for cancer. *J. Cell Physiol.* **2019**. [[CrossRef](#)] [[PubMed](#)]
105. Ling, H.; Roux, E.; Hempel, D.; Tao, J.; Smith, M.; Lonning, S.; Zuk, A.; Arbeeny, C.; Ledbetter, S. Transforming Growth Factor β Neutralization Ameliorates Pre-Existing Hepatic Fibrosis and Reduces Cholangiocarcinoma in Thioacetamide-Treated Rats. *PLoS ONE* **2013**, *8*, e54499. [[CrossRef](#)] [[PubMed](#)]
106. Batlle, E.; Massagué, J. Transforming Growth Factor- β Signaling in Immunity and Cancer. *Immunity* **2019**, *50*, 924–940. [[CrossRef](#)] [[PubMed](#)]
107. Lan, Y.; Zhang, D.; Xu, C.; Hance, K.W.; Marelli, B.; Qi, J.; Yu, H.; Qin, G.; Sircar, A.; Hernandez, V.M.; et al. Enhanced preclinical antitumor activity of M7824, a bifunctional fusion protein simultaneously targeting PD-L1 and TGF- β . *Sci. Transl. Med.* **2018**, *10*. [[CrossRef](#)] [[PubMed](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).