Corrigendum

Corrigendum to "CD36-Mediated Lipid Accumulation and Activation of NLRP3 Inflammasome Lead to Podocyte Injury in Obesity-Related Glomerulopathy"

Jing Zhao, Hong-liang Rui D, Min Yang, Li-jun Sun, Hong-rui Dong, and Hong Cheng D

Division of Nephrology, Beijing Anzhen Hospital, Capital Medical University, Beijing 100029, China

Correspondence should be addressed to Hong Cheng; drchengh@163.com

Received 11 July 2019; Accepted 15 July 2019; Published 23 September 2019

Copyright © 2019 Jing Zhao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled "CD36-Mediated Lipid Accumulation and Activation of NLRP3 Inflammasome Lead to Podocyte Injury in Obesity-Related Glomerulopathy" [1], there were some input errors in Table 3 as the unit of kidney weight "mg," Lee's index "g/cm," and urinary protein "mg/d" are wrong and should be corrected as kidney weight "g," Lee's index "g $^{1/3}$ /cm," and urinary protein " μ g/d." The corrected version of Table 3 is as follows:

Table 3: Biological parameters in the different groups at the 12th week $(\bar{x} \pm s)$.

Group	Control	Model
Body weight (g)	28.9 ± 1.8	31.9 ± 1.9*
Kidney weight (g)	0.34 ± 0.03	$0.37 \pm 0.02^*$
Lee's index (g ^{1/3} /cm)	16.86 ± 0.38	$17.64 \pm 0.56^*$
Visceral fat index	0.02 ± 0.01	$0.05 \pm 0.02^*$
Urinary protein (µg/d)	801.5 ± 178.9	$1001.7 \pm 230.1^*$
Serum creatinine (µmol/L)	11.3 ± 0.8	10.4 ± 1.1
Creatinine clearance rate (mL/min)	0.4 ± 0.1	$0.7 \pm 0.2^*$
Serum triglyceride (mmol/L)	0.5 ± 0.1	$0.9 \pm 0.4^*$
Serum cholesterol (mmol/L)	2.2 ± 0.4	$3.6 \pm 0.3^*$
Blood glucose (mmol/L)	9.5 ± 0.6	9.8 ± 1.3

^{*}P < 0.05 vs. control group.

The authors apologize for these errors and confirm that they do not affect the conclusions or the results of this article.

References

[1] J. Zhao, H.-l. Rui, M. Yang, L.-j. Sun, H.-r. Dong, and H. Cheng, "CD36-mediated lipid accumulation and activation of NLRP3 inflammasome lead to podocyte injury in obesity-related glomerulopathy," *Mediators of Inflammation*, vol. 2019, Article ID 3172647, 16 pages, 2019.