

Predicting weaning difficulty for planned extubation patients with an artificial neural network

Meng Hsuen Hsieh, BS^a, Meng Ju Hsieh, MD^b, Ai-Chin Cheng, RRT^{c,d}, Chin-Ming Chen, MD^{e,*}, Chia-Chang Hsieh, MD^f, Chien-Ming Chao, MD^g, Chih-Cheng Lai, MD^h, Kuo-Chen Cheng, MD^d, Willy Chou, MD^{i,j,*}

Abstract

This study aims to construct a neural network to predict weaning difficulty among planned extubation patients in intensive care units.

This observational cohort study was conducted in eight adult ICUs in a medical center about adult patients experiencing planned extubation.

The data of 3602 patients with planned extubation in ICUs of Chi-Mei Medical Center (from Dec. 2009 through Dec. 2011) was used to train and test an artificial neural network (ANN) model. The input features contain 47 clinical risk factors and the outputs are classified into three categories: simple, difficult, and prolonged weaning. A deep ANN model with four hidden layers of 30 neurons each was developed. The accuracy is 0.769 and the area under receiver operating characteristic curve for simple weaning, prolonged weaning, and difficult weaning are 0.910, 0.849, and 0.942 respectively.

The results revealed that the ANN model achieved a good performance in prediction the weaning difficulty in planned extubation patients. Such a model will be helpful for predicting ICU patients' successful planned extubation.

Abbreviations: ANN = artificial neural network, APACHE-II = acute physiology and chronic health evaluation II, AUROC = area under receiving operating characteristic, CI = confidence interval, DT = decision trees, ICU = intensive care unit, LDA = linear discriminant analysis, MEP = maximum expiratory pressure, MIP = maximum inspiratory pressure, NB = Naïve Bayes, PaCO₂ = arterial carbon dioxide tension, PEEP = positive end expiratory pressure, ROC = receiving operating characteristic, RSI = rapid shallow-breathing index, SBTs = spontaneous breathing trials, SE = standard error, SeLU = scaled exponential linear unit, SVM = support vector machines, TISS = therapeutic intervention scoring system.

Keywords: artificial neural network, planned extubation, prediction weaning difficulty

1. Introduction

Endotracheal intubation is a process commonly used in intensive care unit (ICU) patients. On average, 39% of ICU patients require endotracheal intubation with ventilatory support.^[1] Though required, prolonged ventilatory support can increase the risk of

certain complications, such as ventilation-associated pneumonia.^[2] The extubation of ventilated patients as early as possible is therefore desired through weaning.^[3] Weaning is the process of gradually removing ventilatory support in a patient by the process of extubation.^[4] The appropriate time to start the weaning process is determined by clinicians to avoid prolonged

Editor: Fu-Tsai Chung.

MHH, MJH, and ACC have the equal contribution.

The authors declare that there is no conflict of interest.

Supplemental Digital Content is available for this article.

^a Department of Electrical Engineering and Computer Science, University of California, Berkeley, Berkeley, CA, ^b Department of Medicine, Poznan University of Medical Science, Poznan, Poland, ^c Department of Medical Sciences Industry, Chang Jung Christian University, ^d Section of Respiratory Care, Department of Internal Medicine, Chi-Mei Medical Center, ^e Department of Intensive Care Medicine, Chi Mei Medical Center, Tainan, ^f China Medical University Children's Hospital, China Medical University, Taichung, ^g Department of Intensive Care Medicine, Chi Mei Medical Center, Liouying District, ^h Department of Internal Medicine, Kaohsiung Veterans General Hospital, Tainan Branch, Tainan, ⁱ Department of Physical Medicine and Rehabilitation, Chi Mei Medical Center, Chiali, ^j Department of Recreation and Health-Care Management, Chia Nan University of Pharmacy and Science, Tainan, Taiwan.

* Correspondence: Chin-Ming Chen, Department of Intensive Care Medicine, Chi-Mei Medical Center, 901 Chung Hwa Road, Yang Kang City, Tainan, Taiwan (e-mail: chencm3383@yahoo.com.tw); Willy Chou, Department of Physical Medicine and Rehabilitation, Chi Mei Medical Center, Chiali District, Tainan, Taiwan (e-mail: ufan0101@ms22.hinet.net).

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Hsieh MH, Hsieh MJ, Cheng AC, Chen CM, Hsieh CC, Chao CM, Lai CC, Cheng KC, Chou W. Predicting weaning difficulty for planned extubation patients with an artificial neural network. *Medicine* 2019;98:40(e17392).

Received: 3 January 2019 / Received in final form: 2 July 2019 / Accepted: 7 September 2019

<http://dx.doi.org/10.1097/MD.00000000000017392>

ventilatory support.^[5] Therefore, weaning profiles and extubation predictions are very important in assessing a patient undergoing endotracheal intubation.

In 2005, during the international consensus conference on weaning from mechanical ventilation, a patient classification system according to the weaning process was proposed. According to the duration of weaning and the number of spontaneous breathing trials (SBTs) preceding successful extubation, patients are classified into three groups: simple, difficult, and prolonged weaning.^[4] This weaning classification has been evaluated in clinical practice.^[6,7] In the studies, the prolonged weaning group was associated with increased mortality in the ICU.

In recent years, outcome prediction models using artificial neural network (ANN) and multivariable logistic regression analysis have been developed in many areas of health care research.^[8] There is a growing amount of publications regarding the use of machine learning algorithms in ICU subjects, particularly in the prediction of sepsis as well.^[9,10,11] ANNs are computer-based algorithms that mimic the habits and structures of neurons and can derive outcomes based on input data. With a clear classification system for patient weaning profiles, the aim of our study is to utilize ANNs to categorize patients into them to predict their individual weaning difficulty.

2. Materials and methods

2.1. Study design and setting

This study was a retrospective analysis using machine learning method based on the data collected based on a previous prospective study, which was conducted in eight adult ICUs of Chi-Mei Medical Center from December 2009 through December 2011. This is a 1288-bed tertiary medical center with 96 ICU beds: 48 medical ICU beds, 9 cardiac beds, and 39 surgical beds for adults. Every year, more than 5000 patients are admitted to the ICU in average. The ICU is covered by intensivists, senior residents, nurses, respiratory therapists, dietitians, physical therapists, and clinical pharmacists. The workload is the same in every shift and patient-to-nursing staff ratios of 2:1. There were no differences in nursing experience by shift. Each respiratory therapist was responsible for fewer than 10 patients at the same time on every shift. The ICU team made rounds at least once daily, and the physician decided the timing of initiating weaning process.

During the study period, a total of 3602 patients experiencing planned extubation were enrolled in this study. According to the weaning process classifications, all patients were separated into three groups: simple, difficult, and prolonged weaning. The definition of simple weaning is a successful extubation after the first SBT; an SBT trial is defined as a low-pressure support with ≤ 8 cm H₂O, or a T-piece trial.^[4] Difficult weaning is defined by a successful extubation after two or three SBTs, or a successful extubation within seven days of the first SBT. Prolonged weaning is classified by patients not weaned after more than three SBTs, or a weaning process greater than seven days. If the patients displayed unstable hemodynamics or desaturation during SBT trial, we would stop the trial. Either adjusting a higher support level or shifting to control mode would be performed for failed SBT. All patients' demographic and clinical information, laboratory results, comorbidities, severity scores, mortality,

and lengths of stays for both ICU and hospital were collected for analysis. The data were retrospectively collected before planned extubation after passed SBT and then analyzed. Therefore, informed consent was specifically waived and the study was approved by the Institutional Review Board of Chi Mei Medical Center (IRB: 10706-009).

2.1.1. Constructing the training data. All features are extracted from the original dataset. After normalizing and cleaning the data, there are 47 input features and three outputs, each representing a prediction of simple, difficult, and prolonged weaning. The 47 input features include subject age, gender, scoring systems as Acute Physiology and Chronic Health Evaluation II (APACHE-II), Therapeutic Intervention Scoring System (TISS) and Glasgow Coma Scales, comorbidities, etiology of intubation and respiratory failure, pre-extubation parameters, weaning methods and parameters, and pre-extubation data. The basis features are listed in Table 1 in the Supplemental Digital Content, <http://links.lww.com/MD/D264>. The data is then split into training and test sets at an approximately 9:1 ratio.^[12] Table 1 shows the data allocation between the test and train sets.

2.1.2. Algorithm and training. We used a multilayer perceptron deep neural network to train the data. To select the hyperparameters, optimizers, and loss function with the best performance, *k*-fold cross-validation with a *k* value of 10 is used over 10 epochs.^[12]

After the model selection process of comparing the performance of different models with *k*-fold cross-validation, the best-performing model consisted of one input layer of 47 dimensions, 4 hidden layers of 30 dimensions each, and an output layer of 3 dimensions. The network was trained using stochastic gradient descent and optimized using Adam with Nesterov Momentum.^[13] The input and hidden layers used the Scaled Exponential Linear Unit (SeLU) activation function, while the output layer used the Softmax activation function.^[14] Dropout of 20% was applied at the input layer and 50% at the output layer for regularization.^[15] The neuron weights were initialized using normalized He initialization.^[16] Since the ANN aims to solve a multi-class classification problem, the categorical cross-entropy function was used as the loss function. The model generates a probability for each category, and the patient is assigned to the category with the highest probability.

The software was implemented using Python (version 3.7.0) with the scikit-learn library (version 0.19.1).^[17] The ANN model was created and trained with the Tensorflow framework (version 1.9.0).^[18]

2.1.3. Statistical analyses. Mean values, standard deviations, and group sizes were used to summarize the results for continuous variables. Kruskal–Wallis ANOVA was used for comparison of continuous variables with Dunn's test for post hoc testing. The Chi-squared test for trends was used to compare categorical variables between the three weaning categories. A *P* value < .05 was considered statistically significant. Statistical analysis of the data was done with SPSS 21.0 for Windows (SPSS, Inc., IL).

The ANN performance was measured using the area under Receiving Operating Characteristic (ROC) curve. The area under ROC curve (AUROC) of the neural network was compared against the AUC of variables that had a significant difference in

Table 1**Baseline characteristics of the 3602 patients who started weaning, stratified by weaning category.**

Variable	Simple weaning n=2613 (72.6%)	Difficult weaning n=678 (18.8%)	Prolonged weaning n=311 (8.6%)	P value
Age (yr)	61.7±16.3	69.0±15.6	73.0±14.5	.001
≥65 years old	1186 (45.4%)	443 (65.3%)	231 (74.3%)	<.001
Males	1680 (64.3%)	443 (65.3%)	195 (62.7%)	.718
Internal Medicine	560 (21.4%)	430 (63.4%)	238 (76.5%)	<.001
APACHE II	14.5±6.7	20.6±6.8	22.8±6.9	*<.001
TISS Scale	26.9±7.9	28.2±7.6	27.2±7.5	#<.001
COMA Scale	12.7±3.2	10.5±3.6	9.5±3.4	<.001
Number of comorbidities	1.1±0.9	1.3±0.9	1.4±1.0	*<.001
Type of comorbidity				
Chronic heart failure	471 (18.0%)	139 (20.5%)	70 (22.5%)	.079
Chronic respiratory failure	162 (6.2%)	86 (12.7%)	37 (11.9%)	<.001
Chronic renal failure	291 (11.1%)	86 (12.7%)	42 (13.5%)	.299
Chronic liver disease	52 (2%)	17 (2.5%)	10 (3.2%)	.312
Diabetes	674 (25.8%)	263 (38.8%)	133 (42.8%)	<.001
Neurological disease	594 (22.7%)	226 (33.3%)	102 (32.8%)	<.001
Active cancer disease	647 (24.8%)	98 (14.5%)	38 (12.2%)	<.001
Immunocompromised	5 (0.2%)	1 (0.2%)	0 (0%)	.730
Etiology of Intubation				
Hypoventilation	218 (8.3%)	87 (12.8%)	37 (11.9%)	*.001
Airway obstruction	90 (3.4%)	41 (6.1%)	22 (7.1%)	*<.001
Pneumonia	141 (5.4%)	199 (29.4%)	120 (38.6%)	<.001
Cardiogenic pulmonary edema	163 (6.2%)	66 (9.7%)	27 (8.7%)	*.004
Septic shock	84 (3.2%)	80 (11.8%)	35 (11.3%)	*<.001
Chronic obstructive pulmonary disease	50 (1.9%)	30 (4.4%)	17 (5.5%)	*<.001
Post-operation	1866 (71.4%)	174 (25.7%)	54 (17.4%)	<.001
Etiology of Respiratory failure				
Pulmonary system	415 (15.9%)	259 (38.2%)	149 (47.9%)	<.001
Cardiovascular system	636 (24.3%)	92 (13.6%)	36 (11.6%)	*<.001
Neurological system	796 (30.5%)	125 (18.4%)	51 (16.4%)	*<.001
Renal system	196 (7.5%)	67 (9.9%)	20 (6.4%)	.075
Gastrointestinal system	357 (13.7%)	94 (13.9%)	42 (13.5%)	.984
Other system	213 (8.2%)	42 (6.2%)	13 (4.2%)	.016
Duration of mechanical ventilation (hours)	65.5±88.6	170.6±108.5	325.6±165.1	<.001
Pre-extubation parameter				
RSI	49.3±28.3	60.4±28.2	70.1±39.1	<.001
MIP (cmH ₂ O)	38.3±14.2	36.6±13.1	35.6±14.4	*<.017
MEP (cmH ₂ O)	63.0±28.5	56.2±31.1	50.5±28.7	<.016
Weaning methods				
T-piece	106 (4.1%)	488 (72.0%)	266 (85.5%)	<.001
Pressure support≤8cmH ₂ O	890 (34.1%)	549 (81.0%)	237 (76.2%)	<.001
SBT	994 (38.0%)	661 (97.5%)	307 (98.7%)	<.001
Number of attempts weaning				
T-piece	0±0.2	1±0.8	3.2±2.4	<.001
Pressure support≤8cmH ₂ O	0.3±0.5	0.9±0.5	0.9±0.7	*<.001
SBT	0.4±0.5	1.9±0.7	4.1±2.4	<.001
Pre-extubation data				
FI _O ₂	27.8±3.8	26.6±2.9	26.3±2.5	*<.001
Pressure level (cmH ₂ O)	9.5±1.6	8.4±1.1	8.3±1.2	*<.001
PEEP (cmH ₂ O)	5.1±0.5	5.1±0.5	5.1±0.4	ns
Minute ventilation (L/min)	7.8±2.7	7.9±2.4	8.0±2.4	ns
Heart rate	86.9±16.1	85.2±17.0	87.9±15.9	&.041
Mean arterial pressure	96.8±16.5	96.1±15.9	93.8±15.2	\$.006
Respiratory rate	16.2±5.0	17.8±5.3	19.5±5.1	<.001
pH	7.435±0.055	7.453±0.048	7.465±0.044	<.05
PaCO ₂ (mmHg)	36.9±5.8	38.9±6.5	39.8±6.9	*<.003
PaO ₂ (mmHg)	110.9±45.0	93.0±26.8	90.9±22.9	*<.001
PaO ₂ /FI _O ₂ (mmHg)	367.9±103.5	339.8±91.5	338.8±92.2	*<.001
Hemoglobin (g/dL)	11.6±2.0	10.6±1.6	10.4±1.5	*<.001
Hematocrit (%)	34.8±6.5	32.4±6.8	31.6±6.8	*<.001
BUN (mg/dL)	22.7±18.6	30.7±24.8	36.9±33.6	<.001
Cr (mg/dL)	1.66±2.09	1.77±2.00	1.72±2.02	ns

(continued)

Table 1
(continued).

Variable	Simple weaning n=2613 (72.6%)	Difficult weaning n=678 (18.8%)	Prolonged weaning n=311 (8.6%)	P value
Na (meq/L)	138.9±4.3	139.6±5.3	139.7±5.6	* <.017
K (meq/L)	3.8±0.5	3.9±0.5	4.0±0.5	<.022
Ca (mg/dL)	7.9±1.0	7.9±0.8	7.9±0.8	ns
P (mg/dL)	3.4±1.5	3.5±1.6	3.5±1.5	ns
Albumin (g/dL)	3.0±0.6	2.7±0.6	2.5±0.5	<.008

APACHE-II = acute physiology and chronic health evaluation II, MEP = maximum expiratory pressure, MIP = maximum inspiratory pressure, PEEP = positive end expiratory pressure, RSI = index of rapid shallow breathing, SBT = spontaneous breathing trial, TISS = therapeutic intervention scoring system.

* Simple vs difficult $P < .01$, simple vs prolonged $P < .001$.

Simple vs difficult $P < .01$.

§ Simple vs prolonged $P < .01$.

‡ Difficult vs prolonged $P < .01$.

terms of outcomes. The AUC was also compared against the ideal value of one.^[19]

3. Results

3.1. Results of clinical data

Of the 3602 patients included in the study, 50.9% were male and 49.1% were female. Patients were classified according to weaning classification: 2613 patients (72.6%) as simple weaning, 678 patients (18.8%) as difficult weaning, and 311 (8.6%) as prolonged weaning. The mean age of simple weaning group is 61.7 years, the mean age of difficult weaning group is 69.0 years, while the mean age of prolonged weaning group is 73.0 years. The mean APACHE II score in difficult group is 20.6 and in

prolonged group is 22.8, and both are significantly higher than that of simple weaning group, which is 14.5 ($P < .001$). The most common comorbidity is diabetes in the prolonged weaning group (42.8%) and in the difficult weaning group (38.8%). See Table 1 in the Supplemental Digital Content, <http://links.lww.com/MD/D264> for the full demographic and clinical characteristics ICU patients with planned extubation. The rate of extubation failure is 5.1% (185/3602).

3.2. Results of ANN

The accuracy of the ANN model is 0.769, and weighted k -fold cross-validation accuracy is 0.604. Figures 1 to 3 show the ROC curve of the artificial neural network, rapid shallow-breathing index (RSI), maximum expiratory pressure (MEP), and




Figure 1. Receiving operating characteristic (ROC) curve of artificial neural network (ANN), rapid shallow breathing (RSI), maximum expiratory pressure (MEP), and maximum inspiratory pressure (MIP) for simple weaning.




Figure 2. Receiving operating characteristic (ROC) curve of artificial neural network (ANN), rapid shallow breathing (RSI), maximum expiratory pressure (MEP), and maximum inspiratory pressure (MIP) for prolonged weaning.




Figure 3. Receiving operating characteristic (ROC) curve of artificial neural network (ANN), rapid shallow breathing (RSI), maximum expiratory pressure (MEP), and maximum inspiratory pressure (MIP) for difficult weaning.

Table 2
Area under ROC curve of ANN model in predicting the type of weaning across all data.

Weaning Type	AUROC	AUROC 95% CI	AUROC SE
Simple	0.910	0.901–0.920	0.005
Prolonged	0.849	0.836–0.863	0.007
Difficult	0.942	0.932–0.951	0.005

AUROC = area under ROC curve, CI = confidence interval, SE = standard error.

maximum inspiratory pressure (MIP) on all patient data for simple, prolonged, and difficult weaning respectively. Tables 2 and 3 show the AUROCs of all 3 weaning types for the ANN model and control predictors respectively. The AUROCs were calculated across all data.

4. Discussion

In this study, we demonstrated that a neural network model can be a good predictor for determining weaning classifications. As extubation failure remains prevalent in clinical practice, with reintubation rates reporting up to 19%,^[20] it is important to determine patients' weaning profiles. In common clinical practices, the extubation decision is based on a comprehensive assessment that considers a patient's clinical condition, arterial blood gases results, ventilator settings and weaning profiles.^[5] Despite this comprehensive assessment, extubation decisions carries risks of misjudgments that can be fatal. The model created in this study can aid in making decisions for patient extubation with laboratory data.

Additionally, the usage of ANNs in mortality prediction has been recorded since 2006, with its effectiveness noted. The ANN used in the 2006 study concluded that ANN mortality prediction outperformed traditional methods of prognosis assessment.^[21] With improved computing power since 2006, ANN models have significantly improved; as such, the ANN used in this study corroborates with the findings and can successfully categorize data into the preset categories for weaning prediction. We found that the collective predictive performance of ANN is better than several commonly used individual parameters in extubation assessment – Index of RSI, MIP, and MEP. This is consistent with a previous study^[8] where the proposed ANN model had better discrimination than existing predictors, such as the RSI and MIP,

Table 3
Area under ROC curve of RSI, MSI, and MEP in predicting the type of weaning across all data.

Weaning type	Variable	AUROC	AUROC 95% CI	AUROC SE
Simple	RSI	0.496	0.475–0.517	0.011
	MIP	0.551	0.530–0.572	0.011
	MEP	0.607	0.585–0.628	0.011
Prolonged	RSI	0.484	0.452–0.517	0.017
	MIP	0.566	0.532–0.601	0.018
	MEP	0.630	0.596–0.664	0.017
Difficult	RSI	0.503	0.479–0.527	0.012
	MIP	0.534	0.510–0.558	0.012
	MEP	0.573	0.548–0.598	0.013

AUROC = area under ROC curve, CI = confidence interval, SE = standard error.

in predicting successful extubation. This shows the widespread potential for ANN models in multiple scenarios wherever categorization and prediction is necessary.

Moreover, previous studies attempted to find appropriate predictors of weaning difficulty and presented different findings. These findings include older age, lower mean arterial pressure,^[22,23] arterial carbon dioxide tension (PaCO₂) under SBT and heart rate increase,^[7] lower BUN,^[24] MIP and PaCO₂,^[25] as well as the incorporation of respiratory rate, RSI, MIP, and APACHE II scores^[26] to aid in the prediction of weaning difficulty. In this study, all the aforementioned factors were included in the ANN model. Thus, the ANN model developed in this study can provide an accurate prediction based on the comprehensive information.

The usage of ANNs in the prediction of patient outcome in ventilator weaning has also been documented. However, the lack of patient data is noted to be a limitation of the studies. Arizmendi et al ran an ANN with 149 patients for the extubation process with a successful predictive capability but failed to categorize patient diagnosis criteria due to the low number of data points.^[27] For our study, using our ANN model with 3602 patient data points, the classification of patient weaning outcomes can be performed before the decision to extubate. The weaning classifications can even be used to further predict mortality.^[7] This allows for a tool to aid in considering extubation decisions for physicians that can ultimately prevent otherwise dangerous extubation procedures.

The ANN built in this study uses the open-source TensorFlow framework, which allows for easy reproduction of the study; further studies can be reproduced using differing data for a more comprehensive overview. Besides neural networks, other machine learning models such as Gaussian Naïve Bayes (NB), Decision Trees (DT), Linear Discriminant Analysis (LDA), and Support Vector Machines (SVM) can also be studied in the future.

5. Conclusions

Extubation strategies in ventilated ICU patients must be thoroughly planned. Previously, the clinical classifications of patient weaning difficulty are used as a characteristic of a patient, rather than assisting in the formulation of a strategy. The ANN used in this study showed that the patient classification can be accurately predicted before the weaning process. This allows the consideration of a patient weaning difficulty prior to the extubation procedure worthwhile.

Author contributions

Conceptualization: Meng Hsuen Hsieh, Ai-Chin Cheng, Chin-Ming Chen, Chih-Cheng Lai, Willy Chou.

Data curation: Ai-Chin Cheng, Chin-Ming Chen, Chien-Ming Chao, Kuo-Chen Cheng.

Formal analysis: Meng Hsuen Hsieh, Meng Ju Hsieh, Ai-Chin Cheng, Chin-Ming Chen, Chia-Chang Hsieh, Chien-Ming Chao, Kuo-Chen Cheng, Willy Chou.

Investigation: Meng Hsuen Hsieh, Meng Ju Hsieh, Ai-Chin Cheng, Chin-Ming Chen, Chia-Chang Hsieh, Chien-Ming Chao, Chih-Cheng Lai.

Methodology: Meng Hsuen Hsieh, Meng Ju Hsieh, Ai-Chin Cheng, Chin-Ming Chen, Chia-Chang Hsieh, Chien-Ming Chao, Chih-Cheng Lai, Willy Chou.

Project administration: Ai-Chin Cheng, Chin-Ming Chen, Chien-Ming Chao, Kuo-Chen Cheng.

Resources: Meng Hsuen Hsieh, Meng Ju Hsieh, Ai-Chin Cheng, Chin-Ming Chen, Chien-Ming Chao, Kuo-Chen Cheng.

Software: Meng Hsuen Hsieh, Meng Ju Hsieh, Chin-Ming Chen, Willy Chou.

Supervision: Ai-Chin Cheng, Chin-Ming Chen, Chia-Chang Hsieh, Chien-Ming Chao, Chih-Cheng Lai, Kuo-Chen Cheng, Willy Chou.

Validation: Meng Hsuen Hsieh, Meng Ju Hsieh, Willy Chou.

Visualization: Meng Hsuen Hsieh, Meng Ju Hsieh.

Writing – original draft: Meng Hsuen Hsieh, Meng Ju Hsieh, Chin-Ming Chen, Chia-Chang Hsieh, Chih-Cheng Lai.

Writing – review & editing: Meng Hsuen Hsieh, Meng Ju Hsieh, Chin-Ming Chen, Chia-Chang Hsieh, Chih-Cheng Lai, Willy Chou.

References

- [1] Esteban A, Anzueto A, Alia I, et al. How is mechanical ventilation employed in the intensive care unit? An international utilization review. *Am J Respir Crit Care Med* 2000;161:1450–8.
- [2] Thompson PJ, Greenough A, Hird MF, et al. Nosocomial bacterial infections in very low birth weight infants. *Eur J Pediatr* 1992;151:451–4.
- [3] Halliday HL. Towards earlier neonatal extubation. *Lancet* 2000;355:2091–2.
- [4] Boles JM, Bion J, Connors A, et al. Weaning from mechanical ventilation. *Eur Respir J* 2007;29:1033–56.
- [5] Fox WW, Schwartz JG, Shaffer TH. Successful extubation of neonates: clinical and physiological factors. *Crit Care Med* 1981;9:823–6.
- [6] Funk GC, Anders S, Breyer MK, et al. Incidence and outcome of weaning from mechanical ventilation according to new categories. *Eur Respir J* 2010;35:88–94.
- [7] Sellares J, Ferrer M, Cano E, et al. Predictors of prolonged weaning and survival during ventilator weaning in a respiratory ICU. *Intensive Care Med* 2011;37:775–84.
- [8] LeCun Y, Bengio Y, Hinton G, et al. *Nature* 2015;521:436–44. Kingma D, Adam JB. A method for stochastic optimization. *International Conference on Learning Representations (ICLR)*, 2015.
- [9] Dybowski R, Weller P, Chang R, et al. Prediction of outcome in critically ill patients using artificial neural network synthesised by genetic algorithm. *Lancet* 1996;347:1146–50.
- [10] 2008;Lukaszewski RA, Yates AM, Jackson MC, et al. Presymptomatic prediction of sepsis in intensive care unit patients[J] *Clin Vaccine Immunol.* 15:1089–94.
- [11] Brause R, Hanisch E, Paetz J, et al. Neural networks for sepsis prediction – the MEDAN-Project1. *J Intensive Care Med* 2004;11:40–3.
- [12] Hsieh MH, Hsieh MJ, Chen CM, et al. An artificial neural network model for predicting successful extubation in intensive care units. *J Clin Med* 2018;7:240.
- [13] Kingma D, Adam JB. A method for stochastic optimization. *International Conference on Learning Representations (ICLR)* 2015.
- [14] Klambauer G, Unterthiner T, Mayr A, et al. Self-normalizing neural networks. *Advances in Neural Information Processing Systems* 2017.
- [15] Srivastava N, Hinton G, Krizhevsky A, et al. A simple way to prevent neural networks from overfitting. *J Machine Learning Res* 2014;15:1929–58.
- [16] He H, Garcia EA. Learning from imbalanced data. *IEEE Transactions on knowledge and data engineering* 2009;21:1263–84.
- [17] Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: machine learning in python. *J Mach Learn Res* 2011;12:2825–30.
- [18] Martín Abadi , Paul Barham , Jianmin Chen , et al. TensorFlow: a system for large-scale machine learning. *OSDI* 2016;16:
- [19] Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* 1982;143:29–36.
- [20] Perren A, Previsdomini M, Llamas M, et al. Patients' prediction of extubation success. *Intensive Care Med* 2010;36:2045–52.
- [21] Silva A, Cortez P, Santos MF. Mortality assessment in intensive care units via adverse events using artificial neural networks. *Artif Intell Med* 2006;36:223–34.
- [22] Chen CM, Chan KS, Fong Y, et al. Age is an important predictor of failed unplanned extubation. *Int J Gerontol* 2010;4:120–9.
- [23] Cheng AC, Cheng KC, Chen CM, et al. The outcome and predictors of failed extubation in intensive care patients—the elderly is an important predictor. *Int J Gerontol* 2011;5:206–11.
- [24] Scheinhorn DJ, Hassenpflug M, Artinian BM, et al. Predictors of weaning after 6 weeks of mechanical ventilation. *Chest* 1995;107:500–5.
- [25] Nava S, Zanotti E, Rubini F. Weaning and outcome from mechanical ventilation. *Monaldi Arch Chest Dis* 1994;49:530–2.
- [26] Meade M, Guyatt G, Cook D, et al. Predicting success in weaning from mechanical ventilation. *Chest* 2001;120:400s–24s.
- [27] Arizmendi C, Romero E, Alquezar R, et al. Data mining of patients on weaning trials from mechanical ventilation using cluster analysis and neural networks. *Conf Proc IEEE Eng Med Biol Soc* 2009;2009:4343–6.