

Original article:

**THE EFFECTS OF VITAMIN D SUPPLEMENTATION ON MUSCLE
FUNCTION AMONG POSTMENOPAUSAL WOMEN:
A SYSTEMATIC REVIEW AND META-ANALYSIS OF RANDOMIZED
CONTROLLED TRIALS**

Reza Tabrizi^{1, 2}, Jamal Hallajzadeh³, Naghmeh Mirhosseini⁴, Kamran B Lankarani⁵,
Najmeh Maharlouei⁵, Maryam Akbari¹, Zatoollah Asemi^{*6}

¹ Health Policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran

² Clinical Neurology Research Center, Shiraz University of Medical Sciences Shiraz, Iran

³ Department of Biochemistry and Nutrition, Research Center for Evidence-Based Health Management, Maraghe University of Medical Science, Maraghe, Iran

⁴ School of Public Health, University of Saskatchewan, Saskatoon, Canada

⁵ Health Policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran

⁶ Research Center for Biochemistry and Nutrition in Metabolic Diseases, Kashan University of Medical Sciences, Kashan, Iran

* Corresponding author: Tel: +98-31-55463378; Fax: +98-31-55463377,
E-mail: asemi_r@yahoo.com

<http://dx.doi.org/10.17179/excli2019-1386>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

The loss of muscle mass and its strength is one of the most critical changes in aging which is associated with an increased risk of falls, osteoporotic fractures and mobility disability. Vitamin D, with its extra-skeletal benefits, might improve muscle function in elderly. The current systematic review and meta-analysis of randomized controlled trials (RCTs) was conducted to summarize available relevant data and determine the effect of vitamin D supplementation on muscle function among postmenopausal women. We reached databases including; Cochrane library, Embase, PubMed, and Web of Science database until the end of May 2018 to identify relevant published RCTs. Heterogeneity among included studies was assessed using Q-test and I² statistics. Random-effect model was applied to pool data and weighted mean difference (WMD) was calculated representing summary effect size. Outcomes of interest included the effects of vitamin D supplementation on hand grip strength (HGS), back muscle strength (BMS), and Timed Up and Go (TUG). Twelve RCTs out of 1739 potential reports were included in our meta-analysis. The pooled findings showed that vitamin D supplementation had no significant effect on HGS (WMD -0.03 kilogram (Kg); 95 % CI, -0.26, 0.20; P=0.78), BMS (WMD 7.21 newton (N); 95 % CI, -5.98, 20.40; P=0.28), and TUG (WMD 0.01 second (S); 95 % CI, -0.17, 0.18; P=0.93) in postmenopausal women. Overall, the current meta-analysis showed that taking vitamin D supplementation by postmenopausal women did not affect markers of muscle function. Further studies are required to confirm the effect of vitamin D supplementation on markers of muscle function.

Keywords: Vitamin D supplementation, muscle function, postmenopausal women, meta-analysis

INTRODUCTION

Muscle function is one of the main determinants of functional capacity among postmenopausal women. Lack of muscle mass and muscle strength occurs as important organic changes by aging and is remarkably associated with an elevated risk of falls and subsequent osteoporotic fractures, besides this severely influences the potential of movement (Orsatti et al., 2011; Muir and Montero-Odasso, 2011). Vitamin D deficiency is common among older populations worldwide, as more than 60 % of postmenopausal women are vitamin D deficient, regardless of other nutrient deficiencies (Rizzoli et al., 2006; Holick, 2007; Han et al., 2018). In observational studies, optimal serum 25(OH)D levels have been significantly associated with improved musculoskeletal function and muscle strength (Bischoff-Ferrari et al., 2004a; Visser et al., 2003). Further, postmenopausal women who were vitamin D deficient presented worse upper and lower limb muscle strength and physical performance compared to vitamin D sufficient control group (Iolascon et al., 2015). Skeletal muscle cells have vitamin D receptor and adequate protein synthesis in these cells, which might be directed through vitamin D function (Holick, 2007).

Current evidence evaluating the effects of vitamin D supplementation on musculoskeletal health has demonstrated promising results. In a meta-analysis conducted by Rosendahl-Riise et al. (2017), vitamin D supplementation alone or its co-administration with calcium supplements significantly improved muscle strength among community-dwelling older individuals. In another meta-analysis, consuming vitamin D by healthy people significantly increased upper and lower limb strength (Tomlinson et al., 2015). Moreover, Rejnmark (2011) reported the advantages of vitamin D supplementation on muscle strength and function in older adults. In another trial, taking vitamin D by postmenopausal women increased their muscle strength, controlled lean mass loss and protected them against the occurrence of sarcopenia (loss of skeletal muscle mass) (Cangussu et al.,

2015). One-year treatment with high-dose vitamin D did not show any significant effect on muscular strength, balance or quality of life in postmenopausal osteoporotic women (Grimnes et al., 2017).

Despite existing published RCTs evaluating the effect of vitamin D on muscle function, current evidence is inconclusive because of lack of trials conducted among younger populations and discrepant findings in older individuals. Discrepancies in current findings might be due to the differences in study design, study populations' characteristics, the dosage of vitamin D supplementation, the duration of intervention, co-supplementation with other nutrients and chronic comorbid conditions among elderly. This meta-analysis was conducted to summarize available data and to determine the effect of vitamin D supplementation on muscle function among postmenopausal women.

METHODS

Search strategy and study selection

Two independent authors systematically searched databases including PubMed, EMBASE, Scopus, Cochrane Library, and Web of Science databases to identify RCTs investigating the impact of vitamin D on muscle function outcomes. Included trials were restricted to human clinical trials published in English until May 2018. The following MeSH and text terms were used to determine related studies, using their title and abstract: participants ["post menopause" OR "postmenopausal" OR "PMP"], intervention ("ergocalciferol" OR "alphacalcidol" OR "cholecalciferol" OR "eldecalcitol" OR "alfacalcidol" OR "vitamin D" OR "vitamin D2" OR "vitamin D3" AND "intake" OR "supplementation"), and outcomes ["hand grip strength (HGS)" OR "back muscle strength (BMS)" OR "Timed Up and Go (TUG)"].

To increase the possibility of including more related studies, the reference list of selected studies and previously published relevant reviews were manually searched.

Inclusion and exclusion criteria

First, duplicate studies and those clearly were not relevant were removed from further investigation. Then, the full-texts of remaining studies were retrieved to assess whether these trials met the inclusion criteria for the current meta-analysis. The inclusion criteria were as followed: being a human RCTs (either parallel or cross-over method), study population were post-menopause women, being a placebo-controlled trial, mean difference and standard deviations (SDs), standard error of the mean (SEMs), or 95 % confidence intervals (CIs) were available for considered outcomes including HGS, BMS, and TUG at both baseline and end of the intervention for each group. Observational studies, case reports, animal, and *in vitro* studies, and RCTs did not meet the minimum requirement for the assessment of methodological quality were excluded from this meta-analysis.

Data extraction and quality assessment

The quality of selected RCTs methodology was assessed, using Cochrane Collaboration risk of bias tool and the data were extracted, using standardized excel sheet, by two independent authors (JH and MA). The following items were used to assess the quality of selected clinical RCTs and the probable risk of bias: "randomization generation, allocation concealment, blinding of participants and outcome assessment, incomplete outcome data, and selective outcome reporting, and other sources of bias". The following data were extracted from each study: first author's name, publication year, participants' age, study design and location, sample size in treatment and placebo groups, dosage of intervention, type of vitamin D supplement and placebo, and the mean (SD) changes for muscle outcomes in each group. If muscle outcomes were measured at multiple durations of intervention or using different dosages of vitamin D supplementation in selected studies, those measurements were considered as separate trials in meta-analysis. Any discrepancy in data extraction was resolved with consensus or by discussing with a third author (ZA).

Data synthesis and statistical analysis

Weighted mean differences (WMDs) were used to determine the impacts of vitamin D supplementation on muscle outcomes including HGS, BMS, and TUG for each clinical trial. The changed value approach was applied to estimate the pooled effect sizes for each outcome. Authors evaluated existing heterogeneity across included RCTs, using Cochrane's Q test and I-square (I^2). Heterogeneity was defined as the I^2 more than 50 percent with $P < 0.05$. Upon existing heterogeneity, random-effect model was applied, using DerSimonian and Laird method; otherwise a fixed-effect model with inverse variance method was used to calculate the pooled WMDs with 95 % confidence interval (CI). Further, sensitivity and subgroup analyses were conducted to calculate the effects of each study on pooled WMDs and to examine the source of heterogeneity, using the possible moderator variables, respectively. P value < 0.05 was considered as statistically significant level. Statistical analyses were performed using STATA software version 12.0 (Stata Corp., College Station, TX) and RevMan V.5.3 software (Cochrane Collaboration, Oxford, UK).

RESULTS

Description of the selected RCTs

Initially, literature search identified 1739 potential studies. After screening based on keywords, removing duplicates and assessing the quality of identified papers, twelve articles (or 16 trials) were included in the meta-analysis. The detailed flowchart of step-by-step identification and selection of the RCTs following literature review has been illustrated in Figure 1. The main characteristics of selected RCTs have been summarized in Table 1.

Overall in 16 trials, 7765 subjects (3788 individuals in intervention and 3977 in control groups) were included in the meta-analysis. Eleven RCTs reported the effects of vitamin D on HGS (kilogram (Kg)), three on BMS (newton (N)), and six on TUG (second (S)). This meta-analysis included RCTs pub-




Figure 1: Flowchart of literature search for selection of trials

lished from 2000 till 2018. Duration of intervention and the dosages of vitamin D supplement varied from 12 to 96 weeks and 10 to 50,000 IU/day, respectively.

Risk of bias and publication bias

Egger and Begg's statistics were used to assess the possibility of publication bias across included RCTs. Our findings indicated no significant evidence of publication bias for studies included in the meta-analyses evaluating the impact of vitamin D supplementation on HGS (P Egger's test=0.48, P Begg's test=0.58) and BMS (P_{Eg}=0.71, P_{Be}=0.60). We

found publication bias for TUG (P_{Eg}=0.02, P_{Be}=0.18), therefore non-parametric method (Duval and Tweedie) was applied to examine the censored study findings. However, the results showed that overall pooled WMD for TUG did not significantly vary between pre (WMD 0.01 S; 95 % CI, -0.17, 0.18) and post (WMD 0.01 S; 95 % CI, -0.17, 0.18) including censored RCTs., the methodological quality of the included RCTs was judged, following Cochrane Handbook for Systematic Reviews of Interventions, by authors and the findings for different items have been illustrated in Figure 2.

Table 1: Characteristics of included studies

Authors (Ref)	Publication year	Sample size (control/intervention)	Country/population	Intervention (name and daily dose)	Treatment duration	Dosage of vitamin D (IU/day)	Age (y) (control, intervention)	Presented data
Wood ^(a) et al.	2014	100/97	United Kingdom/PMP	400 IU vitamin D	48 weeks	400	60-70	HGS
Wood ^(b) et al.	2014	100/96	United Kingdom/PMP	1000 IU vitamin D	48 weeks	1000	60-70	HGS
Janssen et al.	2010	34/36	Netherlands/PMP	400 IU vitamin D+500 mg calcium	24 weeks	400	65 and older	HGS, TUG
Hansen ^(a) et al.	2015	73/74	Wisconsin/PMP	50,000 IU cholecalciferol	48 weeks	50000	75 years or younger	TUG
Hansen ^(b) et al.	2015	73/73	Wisconsin/PMP	800 IU cholecalciferol	48 weeks	800	75 years or younger	TUG
Saito et al.	2016	17/18	Japan/PMP	0.75 µg eldecalcitol+ alendronate at 35 mg/week	24 weeks	30	63–86	HGS, BMS, TUG
Zhu et al.	2010	132/129	Australia/PMP	1,000 IU vitamin D +calcium	48 weeks	1000	70-90	TUG
Cangussu et al.	2015	80/80	Brazil/PMP	1,000 IU vitamin D	36 weeks	1000	50-65	HGS
Gao ^(a) et al.	2015	262/107	China/PMP	800 IU vitamin D +600 mg calcium	96 weeks	800	63.44±5.04	HGS
Gao ^(b) et al.	2015	262/116	China/PMP	0.25 µg calcitriol+ 600 mg calcium	96 weeks	10	63.44±5.04	HGS
Anek et al.	2015	26/26	Thailand/PMP	20,000 IU/week vitamin D	4 weeks	2858	45-55	BMS
Glendenning et al.	2012	333/353	Australia/PMP	150,000 IU/every 3 months vitamin D	36 weeks	1667	77	HGS, TUG
Brunner ^(a) et al.	2008	1265/1334	USA/PMP	1,000 mg calcium carbonate plus 400 IU vitamin D3	24 weeks	400	50-79	HGS
Brunner ^(b) et al.	2008	1162/1185	USA/PMP	1,000 mg calcium carbonate plus 400 IU vitamin D3	96 weeks	400	50-79	HGS
Cavalcante et al.	2015	14/14	Brazil/PMP	943 IU vitamin D3	12 weeks	943	62	HGS
Hara et al.	2013	44/50	Japan/PMP	1.0 µg alfalcidol+ 200 mg calcium+ alendronate 35 mg/weeks	24 weeks	40	55-75	BMS

PMP, postmenopausal; HGS, hand grip strength; BMS, back muscle strength; TUG, Timed Up and Go

Number of RCTs used different types of dosage or duration for estimating the relevant outcomes, these articles were showed with subscript letters (a, b). We treated with these as a separate trial to be included in the current meta-analysis.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data addressed (attrition bias)	Selective reporting (reporting bias)	other sources of bias (e.g. bias of study design, trial stoppably, extreme baseline imbalance, and fraudulent trial)
Anek (2015)	+	+	-	-	?	+	+
Brunner (2008)	+	+	+	+	-	+	?
Cangussu (2015)	+	+	+	+	+	+	+
Cavalcante (2015)	+	+	+	+	?	+	+
Gao (2015)	+	+	-	-	?	+	-
Glendenning (2012)	+	+	+	+	?	+	-
Hansen (2015)	+	+	+	+	?	+	+
Hara (2013)	+	+	?	-	-	+	-
Janssen (2010)	+	+	+	+	?	+	?
Saito (2016)	+	+	-	-	+	+	+
Wood (2014)	+	+	+	+	+	+	+
Zhu (2010)	+	+	+	+	?	+	?

Figure 2: The methodological quality of included trials (risk of bias)

Main outcomes, muscle function

The pooled results of included RCTs indicated that vitamin D supplementation did not affect any significant effect on HGS (WMD -0.03 Kg; 95 % CI, -0.26, 0.20; P=0.78), BMS (WMD 7.21 N; 95 % CI, -5.98, 20.40; P=0.28), and TUG (WMD 0.01 S; 95 % CI, -0.17, 0.18; P=0.93) in postmenopausal women (Table 2 and Figure 3A-C).

Subgroups and sensitivity analyses

The findings of subgroup analyses for the effect of vitamin D supplementation on muscle outcomes have been summarized in Table 3.

Sensitivity analysis was applied to assess the impact of each study on the reliability of the association between vitamin D supple-

mentation and muscle function. After excluding each study from the analyses, we observed no considerable difference between the pre- and post-sensitivity pooled WMD for HGS, BMS, and TUG. However, the lower and upper pooled WMD for HGS in the sensitivity analysis were -0.09 Kg (95 % CI: -0.35, 0.17) after omitting Brunner^(b) et al. (2008) and 0.05 Kg (95 % CI: -0.20, 0.30) after omitting Glendenning et al. (2012), respectively. For BMS, the lower WMD in the sensitivity analysis was 5.15 N (95 % CI: -14.78, 25.10) after excluding Hara et al. (2013) and upper pooled WMD was 10.61 N (95 % CI: -5.42, 26.64) after excluding Anek et al. (2015). The lower WMD for TUG was -0.13 S (95 % CI: -0.41, 0.14) after omitting Glendenning et al. (2012) and upper pooled WMD was 0.02 S (95 % CI: -0.15, 0.20) after omitting Zhu et al. (2010).

Table 2: The effects of vitamin D supplementation on muscle strength in postmenopausal women

Variables	Number of study	Weighted mean difference	CI 95 %	P-value	Heterogeneity			
					I ² (%)	Q	P-value heterogeneity	
HGS	Placebo group (after vs. before)	8	-0.67	-1.13, -0.21	0.004	0.0	3.20	0.866
	Intervention group (after vs. before)	8	-0.35	-0.94, 0.25	0.250	40.7	11.80	0.107
	Change intervention group vs. placebo group	11	-0.03	-0.26, 0.20	0.788	0.0	9.51	0.482
BMS	Placebo group (after vs. before)	2	5.74	-12.50, 23.9	0.537	0.0	0.00	0.998
	Intervention group (after vs. before)	2	11.80	-9.39, 33.00	0.275	0.0	0.69	0.405
	Change intervention group vs. placebo group	3	7.21	-5.98, 20.40	0.284	0.0	0.78	0.678
TUG	Placebo group (after vs. before)	5	-0.43	-0.77, -0.09	0.012	0.0	3.74	0.443
	Intervention group (after vs. before)	5	-1.00	-1.83, -0.17	<0.001	76.1	16.74	0.002
	Change intervention group vs. placebo group	6	0.01	-0.17, 0.18	0.939	0.0	4.36	0.499


HGS, hand grip strength; BMS, back muscle strength; TUG, Timed Up and Go

Table 3: The association between vitamin D supplementation on muscle strength in postmenopausal women based on subgroup analysis


Variables	Number of trials	Subgroups	Pooled WMD	95 % CI	I ² (%)	P*	over-all I ² (%)
HGS	Dosage of vitamin D (IU/day)	<1000	0.11	-0.16, 0.37	0.0	0.05	0.0
		≥1000	-0.41	-0.85, 0.04	0.0		
	Duration of study (week)	>40 weeks	-0.00	-0.33, 0.33	29.9	0.80	
		≤40 weeks	-0.06	-0.38, 0.26	0.0		
	Treatment type	Vitamin D	-0.31	-0.69, 0.08	0.0	0.08	
		Vitamin D plus other nutrients	0.12	-0.17, 0.40	0.0		
BMS	Dosage of vitamin D (IU/day)	<1000	10.61	-5.43, 26.65	0.0	0.46	0.0
		≥1000	0.10	-23.08, 23.2	-		
	Duration of study (week)	>40 weeks	-	-	-	-	
		≤40 weeks	7.21	-5.98, 20.40	0.0		
	Treatment type	Vitamin D	0.10	-23.08, 23.2	-	0.46	
		Vitamin D plus other nutrients	10.61	-5.43, 26.65	0.0		
TUG	Dosage of vitamin D (IU/day)	<1000	-0.14	-0.52, 0.24	0.0	0.39	0.0
		≥1000	0.05	-0.15, 0.25	37.0		
	Duration of study (week)	>40 weeks	-0.11	-0.40, 0.18	11.2	0.31	
		≤40 weeks	0.07	-0.15, 0.29	0.0		
	Treatment type	Vitamin D	0.04	-0.14, 0.22	0.0	0.08	
		Vitamin D plus other nutrients	-0.67	-1.47, 0.13	0.0		

HGS, hand grip strength; BMS, back muscle strength; TUG, Timed Up and Go
 *Overall test for heterogeneity between sub-groups

3A



3B



3C




Figure 3A-C: Meta-analysis on muscle outcomes weighted mean differences estimates for (A) HGS, (B) for BMS, and (C) for TUG in vitamin D and placebo groups (CI=95 %)

DISCUSSION

To our best knowledge, this systematic review and meta-analysis was the first report determining the effect of vitamin D supplementation on muscle function among postmenopausal women. This meta-analysis showed that taking vitamin D supplementation by postmenopausal women did not affect markers of muscle function.

Current evidence demonstrating the impact of vitamin D supplementation on lower extremity muscle strength and function are controversial. While some trials have confirmed a significant effect of vitamin D administration on improving lower extremity muscle strength and function in older population (Gloth et al., 1995; Verhaar et al., 2000), others did not (Kenny et al., 2003; Latham et al, 2003). Most of these studies included males and females who were not vitamin D

deficient. In a study conducted by Zhu et al. (2010), 1,000 IU/day vitamin D supplementation for 1 year significantly improved muscle strength in subjects who had low baseline muscle strength and whose serum 25(OH)D levels were below 24 ng/mL. TUG represents basic mobility skills, it is an effective method and evaluates functional mobility in older population (Podsiadlo and Richardson, 1991). It is also a sensitive and specific measure for identifying the risk of fall among community-dwelling adults older than 65 years (Shumway-Cook et al., 2000). Further, vitamin D ingestion was shown to improve neuromuscular coordination in older adults who had a history of fall with serum 25(OH)D levels less than 30 nmol/L (12 ng/mL) (Dhesi et al., 2004). So, TUG improvement could be correlated with improved muscle strength and neuro-

muscular function. In a meta-analysis conducted by Beaudart et al. (2014), vitamin D supplementation had a modest positive impact on muscle strength. In another meta-analysis, vitamin D supplementation to healthy individuals increased upper and lower limb strength (Tomlinson et al., 2015). On the other hand, Rosendahl-Riise et al. (2017) did not observe any improvement in muscle strength following vitamin D supplementation (with or without calcium supplements) in community-dwelling older population in their meta-analysis. Epidemiological studies have proposed serum 25(OH)D levels greater than 20 ng/mL as a desirable level for optimal lower extremity strength (Wicherts et al., 2007; Bischoff-Ferrari et al., 2004b). The dosages of vitamin D used, type of vitamin D and the duration of intervention are some possible reasons which might explain the inconclusive results regarding the impacts of vitamin D on muscle function.

Low circulating levels of vitamin D is usually asymptomatic, however patient may present with loss of muscle mass and muscle weakness (Bergman et al., 2010). Muscle cells have vitamin D receptors (VDR) (Bischoff-Ferrari et al., 2009). Vitamin D metabolites can influence muscle cell metabolism through activating gene transcription, variation in the VDR allele, and rapid pathways not involved in DNA synthesis (Bischoff-Ferrari et al., 2009; Barr et al., 2010). Supplementation with vitamin D might enhance appendicular muscle strength and increase physical function, particularly in vitamin D deficient older population (Bischoff et al., 2003; Pfeifer et al., 2009).

The current study had a few limitations. We found a significant heterogeneity across included trials which might be explained through the large number of studies included and the variability observed between different protocols of supplementation. However, we considered this heterogeneity by applying a random effect model in meta-analyses. We were also unable to find any dose-response effect in this meta-analysis, which is probably

due to the variability of the different supplementation protocols across included studies.

CONCLUSIONS

Overall, the current meta-analysis showed that taking vitamin D supplementation by postmenopausal women did not affect markers of muscle function. Further studies are required to confirm the effect of vitamin D supplementation on markers of muscle function.

DECLARATIONS

Ethics approval and consent to participate

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments.

Consent for publication

Not applicable.

Availability of data and material

The primary data for this study is available from the authors on direct request.

Conflict of interest

Reza Tabrizi, Jamal Hallajzadeh, Naghmeh Mirhosseini, Kamran B Lankarani, Najmeh Maharlouei, Maryam Akbari and Zattollah Asemi declare that they have no conflict of interest.

Funding

The research grant (97-01-106-17275) was provided by Research Deputy of Shiraz University of Medical Sciences (SUMS).

Author contributions

ZA, JH, MA and RT contributed in conception, design, statistical analysis and drafting of the manuscript. NM, K-BL and NM contributed in conception, data collection and manuscript drafting. MM, NM and MC contributed in revised version. The final version was confirmed by all authors for submission.

Acknowledgements

The present study was supported by a grant from the Vice-chancellor for Research, SUMS, Shiraz, and Iran.

REFERENCES

- Anek A, Bunyaratavej N, Jittivilai T. Effects of short-term vitamin D supplementation on musculoskeletal and body balance for prevention of falling in postmenopausal women. *J Med Assoc Thai.* 2015;98 (Suppl 8):S26-31.
- Barr R, Macdonald H, Stewart A, McGuigan F, Rogers A, Eastell R, et al. Association between vitamin D receptor gene polymorphisms, falls, balance and muscle power: results from two independent studies (APOSS and OPUS). *Osteoporos Int.* 2010;21:457-66.
- Beaudart C, Buckinx F, Rabenda V, Gillain S, Cavalier E, Slomian J, et al. The effects of vitamin D on skeletal muscle strength, muscle mass, and muscle power: a systematic review and meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab.* 2014;99:4336-45.
- Bergman GJ, Fan T, McFetridge JT, Sen SS. Efficacy of vitamin D3 supplementation in preventing fractures in elderly women: a meta-analysis. *Curr Med Res Opin.* 2010;26:1193-201.
- Bischoff HA, Stahelin HB, Dick W, Akos R, Knecht M, Salis C, et al. Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. *J Bone Miner Res.* 2003;18:343-51.
- Bischoff-Ferrari HA, Borchers M, Gudat F, Durmuller U, Stahelin HB, Dick W. Vitamin D receptor expression in human muscle tissue decreases with age. *J Bone Miner Res.* 2004a;19:265-9.
- Bischoff-Ferrari HA, Dietrich T, Orav EJ, Hu FB, Zhang Y, Karlson EW, et al. Higher 25-hydroxyvitamin D concentrations are associated with better lower-extremity function in both active and inactive persons aged > or =60 y. *Am J Clin Nutr.* 2004b;80:752-8.
- Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck AE, Theiler R, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ.* 2009;339:b3692.
- Brunner RL, Cochrane B, Jackson RD, Larson J, Lewis C, Limacher M, et al. Calcium, vitamin D supplementation, and physical function in the Women's Health Initiative. *J Am Diet Assoc.* 2008;108:1472-9.
- Cangussu LM, Nahas-Neto J, Orsatti CL, Bueloni-Dias FN, Nahas EA. Effect of vitamin D supplementation alone on muscle function in postmenopausal women: a randomized, double-blind, placebo-controlled clinical trial. *Osteoporos Int.* 2015;26:2413-21.
- Cavalcante R, Maia J, Mesquita P, Henrique R, Griz L, Bandeira MP, et al. The effects of intermittent vitamin D3 supplementation on muscle strength and metabolic parameters in postmenopausal women with type 2 diabetes: a randomized controlled study. *Ther Adv Endocrinol Metab.* 2015;6:149-54.
- Dhesi JK, Jackson SH, Bearne LM, Moniz C, Hurley MV, Swift CG, et al. Vitamin D supplementation improves neuromuscular function in older people who fall. *Age Ageing.* 2004;33:589-95.
- Gao LH, Zhu WJ, Liu YJ, Gu JM, Zhang ZL, Wang O, et al. Physical performance and life quality in postmenopausal women supplemented with vitamin D: a two-year prospective study. *Acta Pharmacol Sin.* 2015;36:1065-73.
- Glendenning P, Zhu K, Inderjeeth C, Howat P, Lewis JR, Prince RL. Effects of three-monthly oral 150,000 IU cholecalciferol supplementation on falls, mobility, and muscle strength in older postmenopausal women: a randomized controlled trial. *J Bone Miner Res.* 2012;27:170-6.
- Gloth FM 3rd, Smith CE, Hollis BW, Tobin JD. Functional improvement with vitamin D replenishment in a cohort of frail, vitamin D-deficient older people. *J Am Geriatr Soc.* 1995;43:1269-71.
- Grimnes G, Emaus N, Cashman KD, Jorde R. The effect of high-dose vitamin D supplementation on muscular function and quality of life in postmenopausal women-A randomized controlled trial. *Clin Endocrinol (Oxf).* 2017;87:20-8.
- Han W, Fan Z, Zhou M, Guo X, Yan W, Lu X, et al. Low 25-hydroxyvitamin D levels in postmenopausal female patients with benign paroxysmal positional vertigo. *Acta Otolaryngol.* 2018;138:443-6.
- Hansen KE, Johnson RE, Chambers KR, Johnson MG, Lemon CC, Vo TN, et al. Treatment of vitamin D insufficiency in postmenopausal women: a randomized clinical trial. *JAMA Intern Med.* 2015;175:1612-21.
- Hara S, Kishimoto KN, Okuno H, Tanaka M, Saito H, Oizumi A, et al. Effects of alfacalcidol on back extensor strength gained through back extensor exercise in postmenopausal women with osteoporosis. *Am J Phys Med Rehabil.* 2013;92:101-10.
- Holick MF. Vitamin D deficiency. *N Engl J Med.* 2007;357:266-81.

- Iolascon G, de Sire A, Calafiore D, Moretti A, Gimigliano R, Gimigliano F. Hypovitaminosis D is associated with a reduction in upper and lower limb muscle strength and physical performance in postmenopausal women: a retrospective study. *Aging Clin Exp Res.* 2015;27(Suppl 1):S23-30.
- Janssen HC, Samson MM, Verhaar HJ. Muscle strength and mobility in vitamin D-insufficient female geriatric patients: a randomized controlled trial on vitamin D and calcium supplementation. *Aging Clin Exp Res.* 2010;22:78-84.
- Kenny AM, Biskup B, Robbins B, Marcella G, Burleson JA. Effects of vitamin D supplementation on strength, physical function, and health perception in older, community-dwelling men. *J Am Geriatr Soc.* 2003;51:1762-7.
- Latham NK, Anderson CS, Lee A, Bennett DA, Moseley A, Cameron ID. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: the Frailty Interventions Trial in Elderly Subjects (FITNESS). *J Am Geriatr Soc.* 2003; 51:291-9.
- Muir SW, Montero-Odasso M. Effect of vitamin D supplementation on muscle strength, gait and balance in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc.* 2011;59:2291-300.
- Orsatti FL, Nahas EA, Nahas-Neto J, Orsatti CL, Marocolo M, Barbosa-Neto O, et al. Low appendicular muscle mass is correlated with femoral neck bone mineral density loss in postmenopausal women. *BMC Musculoskel Disord.* 2011;12:225.
- Pfeifer M, Begerow B, Minne HW, Suppan K, Fahrleitner-Pammer A, Dobnig H. Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community-dwelling older individuals. *Osteoporos Int.* 2009;20: 315-22.
- Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc.* 1991;39:142-8.
- Rejnmark L. Effects of vitamin d on muscle function and performance: a review of evidence from randomized controlled trials. *Ther Adv Chronic Dis.* 2011;2:25-37.
- Rizzoli R, Eisman JA, Norquist J, Ljunggren O, Krishnarajah G, Lim SK, et al. Risk factors for vitamin D inadequacy among women with osteoporosis: an international epidemiological study. *Int J Clin Pract.* 2006;60:1013-9.
- Rosendahl-Riise H, Spielau U, Ranhoff AH, Gudbrandsen OA, Dierkes J. Vitamin D supplementation and its influence on muscle strength and mobility in community-dwelling older persons: a systematic review and meta-analysis. *J Hum Nutr Diet.* 2017;30:3-15.
- Saito K, Miyakoshi N, Matsunaga T, Hongo M, Kasukawa Y, Shimada Y. Eldecacitol improves muscle strength and dynamic balance in postmenopausal women with osteoporosis: an open-label randomized controlled study. *J Bone Miner Metab.* 2016;34:547-54.
- Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Phys Ther.* 2000;80:896-903.
- Tomlinson PB, Joseph C, Angioi M. Effects of vitamin D supplementation on upper and lower body muscle strength levels in healthy individuals. A systematic review with meta-analysis. *J Sci Med Sport.* 2015;18: 575-80.
- Verhaar HJ, Samson MM, Jansen PA, de Vreede PL, Manten JW, Duursma SA. Muscle strength, functional mobility and vitamin D in older women. *Aging (Milan, Italy).* 2000;12:455-60.
- Visser M, Deeg DJ, Lips P. Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the Longitudinal Aging Study Amsterdam. *J Clin Endocrinol Metab.* 2003;88:5766-72.
- Wicherts IS, van Schoor NM, Boeke AJ, Visser M, Deeg DJ, Smit J, et al. Vitamin D status predicts physical performance and its decline in older persons. *J Clin Endocrinol Metab.* 2007;92:2058-65.
- Wood AD, Secombes KR, Thies F, Aucott LS, Black AJ, Reid DM, et al. A parallel group double-blind RCT of vitamin D3 assessing physical function: is the biochemical response to treatment affected by overweight and obesity? *Osteoporos Int.* 2014;25:305-15.
- Zhu K, Austin N, Devine A, Bruce D, Prince RL. A randomized controlled trial of the effects of vitamin D on muscle strength and mobility in older women with vitamin D insufficiency. *J Am Geriatr Soc.* 2010;58: 2063-8.