

Characteristics and Clinical Outcomes of Critically Ill Cancer Patients Admitted to Korean Intensive Care Units

Soo Jin Na^{1*}, Tae Sun Ha^{2*}, Younsuck Koh³, Gee Young Suh^{1,4}, Shin Ok Koh⁵, Chae-Man Lim³, Won-Il Choi⁶, Young-Joo Lee⁷, Seok Chan Kim⁸, Gyu Rak Chon⁹, Je Hyeong Kim¹⁰, Jae Yeol Kim¹¹, Jaemin Lim¹², Sunghoon Park¹³, Ho Cheol Kim¹⁴, Jin Hwa Lee¹⁵, Ji Hyun Lee¹⁶, Jisook Park¹⁷, Juhee Cho¹⁸, Kyeongman Jeon^{1,4}; Validation of Simplified Acute Physiology Score 3 in Korean Intensive care unit (VSKI) study group, and the Korean Study Group On Respiratory Failure (KOSREF)

¹Department of Critical Care Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul; ²Department of Surgery, Soonchunhyang University Bucheon Hospital, Bucheon; ³Division of Pulmonary and Critical Care Medicine, Department of Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul; ⁴Division of Pulmonary and Critical Care Medicine, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul; ⁵Department of Anesthesiology and Pain Medicine, Anesthesia and Pain Research Institute, Yonsei University College of Medicine, Seoul; ⁶Division of Pulmonary and Critical Care Medicine, Department of Medicine, Dongsan Hospital, Keimyung University, Daegu; ⁷Department of Anesthesiology and Pain Medicine, Ewha Womans University Mokdong Hospital, Seoul; ⁸Division of Pulmonary and Critical Care Medicine, Department of Medicine, Seoul St. Mary's Hospital, College of Medicine, Catholic University of Korea, Seoul; ⁹Division of Pulmonary and Critical Care Medicine, Department of Medicine, Chungju Hospital, Konkuk University School of Medicine, Chungju; ¹⁰Department of Critical Care Medicine, Korea University Ansan Hospital, Ansan; ¹¹Division of Pulmonary and Critical Care Medicine, Department of Medicine, Chung-Ang University College of Medicine, Seoul; ¹²Division of Pulmonary and Critical Care Medicine, Department of Medicine, Gangneung Asan Hospital, University of Ulsan Medical College of Medicine, Gangneung; ¹³Department of Pulmonary, Allergy and Critical Care Medicine, Hallym University Sacred Heart Hospital, Anyang; ¹⁴Division of Pulmonary and Critical Care Medicine, Department of Medicine, College of Medicine, Gyeongsang Institute of Health Sciences, Gyeongsang National University, Jinju; ¹⁵Division of Pulmonary and Critical Care Medicine, Department of Medicine, Ewha Womans University School of Medicine, Seoul; ¹⁶Division of Pulmonary and Critical Care Medicine, Department of Medicine, CHA Bundang Hospital, CHA University, Seongnam; ¹⁷Department of Multimedia, Seoul Women's University, Seoul; ¹⁸Samsung Advanced Institute of Health Science and Technology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

Background: The objective of this study was to investigate the characteristics and clinical outcomes of critically ill cancer patients admitted to intensive care units (ICUs) in Korea.

Methods: This was a retrospective cohort study that analyzed prospective collected data from the Validation of Simplified Acute Physiology Score 3 (SAPS3) in Korean ICU (VSKI) study, which is a nationwide, multicenter, and prospective study that considered 5,063 patients from 22 ICUs in Korea over a period of 7 months. Among them, patients older than 18 years of age who were diagnosed with solid or hematologic malignancies prior to admission to the ICU were included in the present study.

Results: During the study period, a total of 1,762 cancer patients were admitted to the ICUs and 833 of them were deemed eligible for analysis. Six hundred fifty-eight (79%) had solid tumors and 175 (21%) had hematologic malignancies, respectively. Respiratory problems (30.1%) was the most common reason leading to ICU admission. Patients with hematologic malignancies had higher Sequential Organ Failure Assessment (12 vs. 8, $P < 0.001$) and SAPS3 (71 vs. 69, $P < 0.001$) values and were more likely to be associated with chemotherapy, steroid therapy, and immunocompromised status versus patients with solid tumors. The use of inotropes/vasopressors, mechanical ventilation, and/or continuous renal replacement therapy was more frequently required in hematologic malignancy patients. Mortality rates in the ICU (41.7% vs. 24.6%, $P < 0.001$) and hospital (53.1% vs. 38.6%, $P = 0.002$) were higher in hematologic malignancy patients than in solid tumor patients.

Conclusions: Cancer patients accounted for one-third of all patients admitted to the studied ICUs in Korea. Clinical characteristics were different according to the type of malignancy. Patients with hematologic malignancies had a worse prognosis than did patients with solid tumor.

Key Words: cancer; critical care; epidemiology; hematologic neoplasm; mortality

Original Article

Received: April 20, 2018

Revised: July 16, 2018

Accepted: July 17, 2018

Corresponding author
Kyeongman Jeon

Department of Critical Care Medicine
and Division of Pulmonary and Critical
Care Medicine, Department of Medicine,
Samsung Medical Center, Sungkyunkwan
University School of Medicine, 81 Irwon-
ro, Gangnam-gu, Seoul 06351, Korea
Tel: +82-2-3410-3429
Fax: +82-2-3410-6956
E-mail: kjeon@skku.edu

*The first two authors contributed equally
to this study.

Copyright © 2018 The Korean Society of
Critical Care Medicine

This is an Open Access article distributed
under the terms of Creative Attributions
Non-Commercial License ([http://
creativecommons.org/licenses/by-nc/4.0/](http://creativecommons.org/licenses/by-nc/4.0/))
which permits unrestricted noncommercial
use, distribution, and reproduction in any
medium, provided the original work is
properly cited.

INTRODUCTION

The increase in the number of individuals with cancer and their improved prognosis has led to a growing demand for intensive care unit (ICU) management [1]. Even with these changes, some physicians consider ICU management in cancer patients to be a futile effort, and metastatic cancer is a common reason for refusal of ICU admission in practice [2]. Considering the disappointing survival rates of critically ill cancer patients in studies published in the 1980s and 1990s, triaging physicians' refusal ICU admission to cancer patients seems reasonable [3,4].

However, recent investigations have demonstrated that advances in ICU management have led to an improvement of clinical outcomes in patients with and without cancer admitted to the ICU and suggested that critically ill cancer patients should not be excluded from the ICU only because they have cancer [5,6]. Domestic epidemiological data are needed to guide our practice in managing patients with cancer because there are geographic variations in the diagnosis of and type of cancer, but domestic data on clinical characteristics and outcomes in cancer patients are still limited [7]. Therefore, the present study was conducted to investigate the characteristics and clinical outcomes and predictors of mortality in critically ill cancer patients admitted to the ICU in Korea.

MATERIALS AND METHODS

Study Design and Population

This study used data from the Validation of Simplified Acute

KEY MESSAGES

- Cancer patients accounted for one-third of all patients admitted to the studied intensive care units in Korea.
- Clinical characteristics were different according to the type of malignancy.
- Patients with hematologic malignancies had a worse prognosis than did patients with solid tumor.

Physiology Score 3 (SAPS3) in Korean ICU (VSKI) study. The VSKI study, which is a nationwide, multicenter, and prospective study, was conducted by the Korean Study Group on Respiratory Failure between July 1, 2010 and January 31, 2011 to evaluate the prognostic performance of the SAPS3 in a large prospective cohort in Korea [8]. It included patients from 22 ICUs (14 medical, six surgical, and two multidisciplinary) in 15 tertiary or university-affiliated hospitals. Only the data from the index ICU admission were included.

A total of 5,063 patients were admitted to participating ICUs during the study period. Patients who were older than 18 years of age and diagnosed with solid or hematologic malignancies prior to admission to the ICU were eligible for inclusion in the present study. Among them, we excluded patients if they were electively admitted for postoperative care, transferred from other participating ICUs, or if they had no data available regarding hospital mortality (Figure 1). Patients were followed up until the time of death or hospital discharge. This study was approved by the relevant Institutional Review Board(s) of each participating hospital, and the requirement for informed




Figure 1. A schematic of study participant flow. ICU: intensive care unit.

consent was waived because of the noninterventional nature of the study.

Data Collection and Clinical Outcomes

Patient data were collected using a web-based database. The definitions of variables used in the original SAPS3 model were used for this study [9]. We used the most abnormal set of data from the one hour prior to or after ICU admission to calculate SAPS3. The following data were sourced from the medical charts of the participants: demographic data; location prior to ICU admission; reason(s) for ICU admission; infection and surgical status at the time of ICU admission; laboratory data and physiologic measurements, which were collected within 1 hour before and 24 hours after ICU admission; and treatment(s) received during ICU stay. The severity of illness was assessed by the SAPS3 and Sequential Organ Failure Assessment (SOFA) score. The primary outcome in this study was hospital mortality. Secondary outcomes were ICU mortality, length of hospital stay, and length of ICU stay.

Definition

Cancer was categorized as solid or hematologic malignancy. Solid tumors were defined as cancer with proven distant (not regional lymph node) metastasis by surgery, computed tomography scan, or any other appropriate method. Hematologic malignancies were defined as tumors of the hematopoietic and lymphoid tissue including lymphoma, acute leukemia, and multiple myeloma. Infection was defined as the presence of a pathogenic microorganism in a sterile site and/or clinically suspected infection, plus the administration of antibiotics. Severe sepsis and septic shock were defined as sepsis associated with acute organ dysfunction and sepsis with acute circulatory failure characterized by persistent arterial hypotension (i.e., systolic arterial pressure <90 mmHg, mean arterial pressure <60 mmHg, or a reduction in systolic blood pressure >40 mmHg from baseline) despite adequate volume resuscitation, respectively [10]. Acute lung injury and acute respiratory distress syndrome were defined as arterial oxygen pressure to inspiratory oxygen fraction ($\text{PaO}_2/\text{FiO}_2$) ratio <300 and <200, respectively, with the following criteria: acute onset, bilateral infiltrates on chest radiography, and pulmonary artery wedge pressure <18 mmHg or clinical evidence of left atrial hypertension. An unplanned ICU admission was defined as an admission planned less than 12 hours in advance or any unscheduled admission to the ICU that originated from the emergency department, general ward, operating room, or other location. Performance status was assigned a grade be-

tween 0 and 4 according to the Eastern Cooperative Oncology Group performance status classification [11]. We used the definition of variables employed in the original SAPS3 model to define variables in this study [12].

Statistical Analysis

Data are presented as median and interquartile range (25th and 75th percentiles) for continuous variables and as numbers (percentages) for categorical variables. Data were compared using the Mann-Whitney U-test for continuous variables and the chi-square or Fisher exact test for categorical variables, respectively, where applicable. Logistic regression analyses were performed to identify risk factors for the prediction of mortality. Variables that appeared to be related in the univariate analysis with a P-value of less than 0.2 were introduced into multivariate regression models using a forward stepwise method. For all analyses, a two-tailed test with a P-value of less than 0.05 was considered to be statistically significant. We used the SPSS version 20.0 (IBM SPSS Corp., Armonk, NY, USA) for statistical analysis.

RESULTS

Baseline Clinical Characteristics

As shown in Figure 1, a total of 1,762 cancer patients were admitted to the ICUs and 833 of them were deemed eligible for analysis. The baseline characteristics of the 833 eligible patients finally included in the present study are described in Table 1. Of these, 554 (66.5%) were male, and the median age was 63 years (range, 53 to 72 years). SOFA score and SAPS3 were 9 (range, 5 to 12) and 69 (range, 60 to 79), respectively. Unplanned admissions composed 74.4% of all ICU admissions. The most common reason for ICU admission was respiratory disease (30.1%), followed by surveillance (27.3%), cardiovascular disease (24.1%), and digestive disease (5.4%). About half of the patients included in this study had acute infection at the time of ICU admission, and most admissions were not related to surgery.

Comparison of Baseline Characteristics According to Type of Malignancy

Of the 833 patients, 658 (79%) had solid tumors and 175 (21%) had hematologic malignancies. The patients with solid tumors were typically older and more commonly male versus the patients with hematologic malignancies (Table 2). Performance status and comorbidities were similar between the two groups, except with regard to hypertension and liver cirrhosis. Patients

Table 1. Baseline characteristics of cancer patients admitted to the ICU

Characteristics	Value
Age (yr)	63 (53–72)
Male sex	554 (66.5)
ECOG performance status ^a	
0–1	349 (44.4)
2–4	438 (55.6)
Comorbidity	
Hypertension	219 (26.3)
Diabetes	153 (18.4)
Congestive heart failure	30 (3.6)
Stroke	35 (4.2)
Chronic lung disease	14 (1.7)
Liver cirrhosis	97 (11.6)
Chronic renal failure	42 (5.0)
Severity of illness	
SOFA	9 (5–12)
SAPS3	69 (60–79)
Location before ICU admission	
Emergency room	377 (45.3)
General ward	429 (51.5)
Other ICU	20 (2.4)
Other	7 (0.8)
Treatment history before ICU admission	
Chemotherapy	311 (37.3)
Radiotherapy	105 (12.6)
Steroid	29 (3.5)
Immune suppression status	79 (9.5)
CPR before ICU admission	44 (5.3)
Unplanned ICU admission	620 (74.4)
Reason for admission	
Surveillance	228 (27.3)
Cardiovascular	201 (24.1)
Respiratory	251 (30.1)
Neurological	29 (3.5)
Hepatic	29 (3.5)
Digestive	45 (5.4)
Renal	9 (1.1)
Metabolic	12 (1.4)
Hematological	5 (0.6)
Other ^b	24 (2.9)
Acute infection at ICU admission	427 (51.3)
Severe sepsis/septic shock	312 (37.5)
Surgical status at ICU admission	
No surgery	736 (88.4)
Emergent surgery	97 (11.6)

Values are presented as median (interquartile range) or number (%). ICU: intensive care unit; ECOG: Eastern Cooperative Oncology Group; SOFA: Sequential Organ Failure Assessment; SAPS3: Simplified Acute Physiology Score 3; CPR: cardiopulmonary resuscitation.

^aData were available for 787 patients; ^bIncludes patients with spinal stenosis, trauma, urinary incontinence, and mood depression disorder.

with hematologic malignancies had higher SOFA and SAPS3 values than did patients with solid tumors. The proportions of patients with a history of chemotherapy (57.1% vs. 32.1%, $P < 0.001$), steroid therapy (9.1% vs. 2.0%, $P < 0.001$), and compromised immune status (17.7% vs. 7.3%, $P < 0.001$) were significantly higher in the hematologic malignancies group, while, conversely, the proportion of patients who had undergone radiotherapy (13.8% vs. 8.0%, $P = 0.040$) was higher in the solid tumor group. Unplanned ICU admissions occurred more frequently among hematologic malignancy patients than among solid tumor patients. Respiratory (35.4% vs. 28.7%, $P = 0.095$) and cardiovascular (34.9% vs. 21.3%, $P < 0.001$) problems were common causes of ICU admission in both groups, with the exception of admission for surveillance. More patients with hematologic cancer showed an acute infection status at the time of ICU admission. In addition, rates of severe sepsis/septic shock were significantly higher in patients with hematologic cancer as compared with those with solid cancer (57.1% vs. 32.2%, $P < 0.001$).

ICU Treatments

The amount of total fluid administered within 24 hours (4,039 ml vs. 3,339 ml, $P < 0.001$) was higher and the use of an inotrope/vasopressor (54.8% vs. 44.7%, $P = 0.017$) was more prevalent, respectively, in patients with hematologic malignancies. In addition, hematologic malignancy patients more frequently required mechanical ventilation (46.9% vs. 33.3%, $P = 0.001$) and continuous renal replacement therapy (17.6% vs. 9.5%, $P = 0.004$) than did solid tumor patients.

Clinical Outcomes

Overall, 235 patients (28.2%) died in the ICU, while 112 patients (13.4%) died while hospitalized in other wards (Figure 2). Both ICU (41.7% vs. 24.6%, $P < 0.001$) and hospital (53.1% vs. 38.6%, $P = 0.002$) mortality rates were significantly higher in patients with hematologic malignancies than in patients with solid tumors. Multivariable logistic regression analysis revealed that age, demonstration of performance status 2 through 4, mechanical ventilation usage, and SAPS3 in solid tumor patients (Table 3) and mechanical ventilation usage and SAPS3 in hematologic malignancy patients were significant prognostic predictors for hospital mortality (Table 3).

In the total patient cohort, ICU length of stay and hospital length of stay were 5 days (range, 3 to 10 days) and 20 days (range, 11 to 39 days), respectively. Although the ICU length of stay (5 days vs. 6 days, $P = 0.160$) was similar between patients with solid tumor and those with hematologic malignancies,

Table 2. Baseline and treatment characteristics according to type of malignancy

Variable	Solid tumor (n=658)	Hematologic malignancies (n=175)	P-value
Age (yr)	65 (55-73)	55 (42-65)	0.005
Male sex	452 (70.2)	102 (58.6)	0.005
ECOG performance status ^a			0.928
0-1	271 (44.2)	78 (44.6)	
2-4	342 (55.8)	96 (54.9)	
Comorbidity (overlapped)			
Hypertension	191 (29.0)	28 (16.0)	0.001
Diabetes	121 (18.4)	32 (18.3)	>0.999
Congestive heart failure	21 (3.2)	9 (5.1)	0.251
Stroke	28 (4.3)	7 (4.0)	>0.999
Chronic lung disease	11 (1.7)	3 (1.7)	>0.999
Liver cirrhosis	93 (14.1)	4 (2.3)	<0.001
Chronic renal failure	32 (4.9)	10 (5.7)	0.703
Severity of illness			
SOFA	8 (4-12)	12 (8-15)	<0.001
SAPS3	69 (60-79)	71 (63-82)	<0.001
Location before ICU admission			<0.001
Emergency room	323 (49.1)	54 (30.9)	
General ward	314 (47.7)	115 (65.7)	
Other ICU	15 (2.3)	5 (2.9)	
Other ^b	6 (0.9)	1 (0.6)	
Hospital days before ICU admission	0 (0-2)	0 (0-9)	0.008
Treatment history before ICU admission			
Chemotherapy	211 (32.1)	100 (57.1)	<0.001
Radiotherapy	91 (13.8)	14 (8.0)	0.040
Steroid	13 (2.0)	16 (9.1)	<0.001
Immune suppression status	48 (7.3)	31 (17.7)	<0.001
CPR before ICU admission	26 (4.2)	18 (10.5)	0.001
Unplanned ICU admission	469 (71.3)	151 (86.3)	<0.001
Reason for ICU admission			
Surveillance	190 (28.9)	38 (21.7)	0.070
Cardiovascular	140 (21.3)	61 (34.9)	<0.001
Digestive	45 (6.8)	0	0.001
Hematological	3 (0.5)	2 (1.1)	0.283
Hepatic failure	29 (4.4)	0	0.004
Metabolic	12 (1.8)	0	0.081
Neurological	20 (3.0)	9 (5.1)	0.243
Renal	7 (1.1)	2 (1.1)	1.000
Respiratory	189 (28.7)	62 (35.4)	0.095
Other	20 (3.0)	1 (0.6)	0.098
Acute infection at ICU admission	298 (45.3)	129 (73.7)	<0.001
Severe sepsis/septic shock	212 (32.2)	100 (57.1)	<0.001

(Continued to the next page)

Table 2. Continued

Variable	Solid tumor (n=658)	Hematologic malignancies (n=175)	P-value
Surgical status at ICU admission			<0.001
No surgery	566 (86.0)	170 (97.1)	
Emergent surgery	92 (14.0)	5 (2.9)	
Initial laboratory variable			
White blood cell (mm ³)	9,750 (5,320–15,460)	3,640 (250–9,850)	<0.001
Total bilirubin (mg/dl)	0.9 (0.6–2.1)	1.1 (0.7–2.3)	0.297
Creatinine (mg/dl)	1.0 (0.7–1.6)	1.1 (0.7–1.8)	0.381
Arterial pH	7.39 (7.30–7.45)	7.40 (7.30–7.46)	0.159
Lactic acid (mmol/l)	2.90 (1.70–5.40)	2.65 (1.50–5.23)	0.468
C-reactive protein (mg/dl)	12.3 (4.3–21.3)	14.5 (6.3–24.3)	0.082
Procalcitonin (ng/ml)	3.64 (0.65–22.87)	8.12 (0.19–27.12)	0.988
NT-proBNP (pg/ml)	681 (161–3039)	965 (315–6070)	0.019
Treatment			
Total infused fluids within 24 hr (ml)	3,339 (2,275–4,775)	4,039 (3,001–5,246)	<0.001
Inotrope/vasopressor within 24 hr	293 (44.7)	96 (54.8)	0.017
Norepinephrine	251 (38.4)	84 (48.8)	0.012
Vasopressin	53 (8.3)	29 (17.2)	0.001
Dopamine	64 (10.0)	19 (11.3)	0.631
Dobutamine	51 (7.9)	20 (11.8)	0.108
Epinephrine	5 (0.8)	2 (1.2)	0.643
Mechanical ventilation	219 (33.3)	82 (46.9)	0.001
Continuous renal replacement therapy	60 (9.5)	30 (17.6)	0.004

Values are presented as median (interquartile range) or number (%).

ECOG: Eastern Cooperative Oncology Group; SOFA: Sequential Organ Failure Assessment; SAPS3: Simplified Acute Physiology Score 3; ICU: intensive care unit; CPR: cardiopulmonary resuscitation; NT-proBNP: N-terminal prohormone of brain natriuretic peptide.

^aData were available for 787 patients; ^bIncludes patients with spinal stenosis, trauma, urinary incontinence, and mood depression disorder.




Figure 2. A comparison of intensive care unit (ICU) and hospital mortality.

the hospital length of stay (28 days vs. 19 days, P=0.001) was noticeably longer in hematologic malignancy patients.

Table 3. Prognostic factors for hospital mortality

Variable	Adjusted OR	95% CI	P-value
Patient with solid tumor			
Age	0.96	0.93–0.99	0.019
ECOG performance status 2–4	2.63	1.29–5.37	0.008
Mechanical ventilation	2.85	1.35–6.04	0.006
SAPS3	1.07	1.04–1.10	<0.001
Patient with hematologic malignancies			
SAPS3	1.05	1.01–1.08	0.006
Mechanical ventilation	2.41	1.05–5.55	0.039

OR: odds ratio; CI: confidence interval; ECOG: Eastern Cooperative Oncology Group; SAPS3: Simplified Acute Physiology Score 3.

DISCUSSION

In this study, we investigated the epidemiology, characteristics, and clinical outcomes of critically ill cancer patients in

Korea. Within our cohort, patients with cancer accounted for 34.8% of the total number of patients admitted to participating ICUs during the study period, and respiratory and cardiovascular diseases were two of the common causes of ICU admission in patients with malignancies. Our percentage of cancer patients admitted was relatively higher than those seen in the results of other previous studies that were conducted involving a mixed population, which ranged from 13.5% to 21.5% of overall ICU admissions [5,13,14]. Although it is difficult to compare our results to those of other investigations directly because our study did not examine the details of cancer, geographic variations in incidence and type of cancer may have an effect on the gap in the proportion of cancer patients admitted to the ICU [7].

Acute respiratory failure was identified as one of the most common medical conditions causing cancer patients to enter the ICU in previous studies as well as in our study [13,15,16]. The incidence of acute respiratory failure in cancer patients ranges from 10% to 50%, and the risk of respiratory failure is higher in patients with hematologic malignancies, especially in patients who received allogeneic stem cell transplantation, as compared with in patients with solid tumor [6,17,18]. The most common cause of acute respiratory failure is pulmonary infection, but treatment-associated pulmonary toxicities, transfusion-related acute lung injury, lung involvement of underlying malignancy, and malignant airway obstruction also can cause a case of acute respiratory failure that requires ICU management [18].

Patients with hematologic malignancies have some clinical characteristics different from those of patients with solid tumor [13]. In our study, the proportion of patients who received chemotherapy and steroid therapy and that of patients with immunosuppression were higher, and observations of acute infection and severe sepsis/septic shock at the time of ICU admission were more frequent in patients with hematologic malignancies than in patients with solid tumor. In addition to a higher baseline SOFA score, the frequencies of vasopressor use, mechanical ventilation, and continuous renal replacement therapy during ICU treatment were all significantly higher in patients with hematologic malignancies. These findings are consistent with those of previous studies that showed that patients with hematologic malignancies are more likely to develop severe sepsis or septic shock in comparison with patients with solid tumor [13]. Neutropenia often occurs in hematologic malignancy patients due to an association with the underlying malignancy itself or chemotherapy; for example, in our study, the white blood cell counts in hematologic ma-

lignancy patients were significantly lower than those in solid tumor patients, and it is known that neutropenic patients are more vulnerable to bacterial or fungal infections that have the potential to cause sepsis [19,20].

Also, in regard to prognosis, solid tumor patients and hematologic malignancy patients showed different results. Although variations exist with regard to the type of cancer, patients with solid tumor have a global hospital mortality rate of 25% to 40%, and some studies have demonstrated that mortality in this population is similar to that in ICU patients without cancer [14,21]. On the other hand, the hospital mortality rate of patients with hematologic malignancies is about 40% to 70%, which is generally worse than that in solid tumor patients [6,22]. In particular, the mortality rate of patients with hematologic malignancies who receive hematopoietic stem cell transplantation is concerning, even in recent studies, at 63% to 75% [23,24]. Hospital mortality rates in solid tumor patients and hematologic malignancy patients in our study were 38.6% and 53.1%, respectively, which were similar to results reported in other countries.

SOFA score, multiple organ failure, the provision of organ support such as vasopressor use or mechanical ventilation, and allogeneic hematopoietic stem cell transplantation are known to be predictors of mortality in patients with cancer [21,22,25,26]. We also confirmed that higher SAPS3 value and mechanical ventilation support were associated with poor prognosis in both solid tumor and hematologic malignancy patients. Furthermore, age and pre-morbid performance status were independent prognostic factors for hospital mortality in patients with solid tumor. Some studies have demonstrated that advanced or recurrent cancer is a variable associated with poor outcomes, but others have suggested that cancer status generally does not affect the short-term outcomes of critically ill cancer patients [25,27,28]. Therefore, intensivists need to be careful when considering cancer status in deciding whether to pursue ICU admission for cancer patients.

Although our study investigated the clinical and treatment characteristics, outcomes, and variables associated with mortality in cancer patients admitted to the 22 medical and surgical ICUs in the Korea, there are several limitations that should be considered. First, our study did not record detailed data about the type, disease status, and treatment of cancer. Although some recent studies have suggested that the disease status of cancer does not adversely affect the short-term outcomes of cancer patients admitted to the ICU, progressive cancer or certain specific conditions, such as those that require mechanical ventilation for respiratory failure due to tumor involve-

ment, are associated with poor prognosis [25]. Therefore, further studies that include additional information on disease status should be conducted to elucidate more accurately the characteristics of and to assess the clinical outcomes in critically ill cancer patients. Also, data about the code status at the time of ICU admission and change(s) in code status during ICU management, which may influence therapeutic decisions and mortality, were not collected. Finally, analyses comparing the cancer patients admitted to the ICU with cancer patients not admitted to the ICU were not performed, and we could not determine whether ICU management is beneficial in cancer patients with acute deterioration.

In conclusion, the proportion of patients with cancer is higher in Korea than in other countries, accounting for about one-third of all patients admitted to the ICU. The most common reason for ICU admission in nonsurgical cancer patients was respiratory problems. Patients with hematologic malignancies had higher severity scores and rates of organ support treatment and mortality than did those with solid tumor. Mechanical ventilation and SAPS3 were independent predictors of hospital mortality in both solid tumor and hematologic malignancy patients.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

ORCID

Kyeongman Jeon <https://orcid.org/0000-0002-4822-1772>

REFERENCES

- van Vliet M, Verburg IW, van den Boogaard M, de Keizer NF, Peek N, Blijlevens NM, et al. Trends in admission prevalence, illness severity and survival of haematological patients treated in Dutch intensive care units. *Intensive Care Med* 2014;40:1275-84.
- Garrouste-Orgeas M, Montuclard L, Timsit JF, Reignier J, Desmettre T, Karoubi P, et al. Predictors of intensive care unit refusal in French intensive care units: a multiple-center study. *Crit Care Med* 2005;33:750-5.
- Schuster DP, Marion JM. Precedents for meaningful recovery during treatment in a medical intensive care unit: outcome in patients with hematologic malignancy. *Am J Med* 1983;75:402-8.
- Jackson SR, Tweeddale MG, Barnett MJ, Spinelli JJ, Sutherland HJ, Reece DE, et al. Admission of bone marrow transplant recipients to the intensive care unit: outcome, survival and prognostic factors. *Bone Marrow Transplant* 1998;21:697-704.
- Soares M, Caruso P, Silva E, Teles JM, Lobo SM, Friedman G, et al. Characteristics and outcomes of patients with cancer requiring admission to intensive care units: a prospective multicenter study. *Crit Care Med* 2010;38:9-15.
- Azoulay E, Mokart D, Pène F, Lambert J, Kouatchet A, Mayaux J, et al. Outcomes of critically ill patients with hematologic malignancies: prospective multicenter data from France and Belgium. A groupe de recherche respiratoire en réanimation onco-hématologique study. *J Clin Oncol* 2013;31:2810-8.
- Torre LA, Siegel RL, Ward EM, Jemal A. Global cancer incidence and mortality rates and trends: an update. *Cancer Epidemiol Biomarkers Prev* 2016;25:16-27.
- Lim SY, Koh SO, Jeon K, Na S, Lim CM, Choi WI, et al. Validation of SAPS3 admission score and its customization for use in Korean intensive care unit patients: a prospective multicentre study. *Respirology* 2013;18:989-95.
- Metnitz PG, Moreno RP, Almeida E, Jordan B, Bauer P, Campos RA, et al. SAPS 3: from evaluation of the patient to evaluation of the intensive care unit. Part 1: objectives, methods and cohort description. *Intensive Care Med* 2005;31:1336-44.
- Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003;31:1250-6.
- Oken MM, Creech RH, Tormey DC, Horton J, Davis TE, McFadden ET, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol* 1982;5:649-55.
- Moreno RP, Metnitz PG, Almeida E, Jordan B, Bauer P, Campos RA, et al. SAPS 3: from evaluation of the patient to evaluation of the intensive care unit. Part 2: development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med* 2005;31:1345-55.
- Staudinger T, Stoiser B, Müllner M, Locker GJ, Laczika K, Knapp S, et al. Outcome and prognostic factors in critically ill cancer patients admitted to the intensive care unit. *Crit Care Med* 2000;28:1322-8.
- Bos MM, de Keizer NF, Meynaar IA, Bakshi-Raiez F, de Jonge E. Outcomes of cancer patients after unplanned admission to general intensive care units. *Acta Oncol* 2012;51:897-905.
- Darmon M, Thiery G, Ciroldi M, de Miranda S, Galicier L, Raffoux E, et al. Intensive care in patients with newly diagnosed malignancies and a need for cancer chemotherapy. *Crit Care Med* 2005;33:2488-93.
- Hawari FI, Nazer LH, Addassi A, Rimawi D, Jamal K. Predic-

- tors of ICU admission in patients with cancer and the related characteristics and outcomes: a 5-year registry-based study. *Crit Care Med* 2016;44:548-53.
17. Chi AK, Soubani AO, White AC, Miller KB. An update on pulmonary complications of hematopoietic stem cell transplantation. *Chest* 2013;144:1913-22.
 18. Pastores SM, Voigt LP. Acute respiratory failure in the patient with cancer: diagnostic and management strategies. *Crit Care Clin* 2010;26:21-40.
 19. Bhatt V, Saleem A. Review: drug-induced neutropenia. Pathophysiology, clinical features, and management. *Ann Clin Lab Sci* 2004;34:131-7.
 20. Bodey GP, Buckley M, Sathe YS, Freireich EJ. Quantitative relationships between circulating leukocytes and infection in patients with acute leukemia. *Ann Intern Med* 1966;64:328-40.
 21. Puxty K, McLoone P, Quasim T, Sloan B, Kinsella J, Morrison DS. Risk of critical illness among patients with solid cancers: a population-based observational study. *JAMA Oncol* 2015; 1:1078-85.
 22. Depuydt PO, Benoit DD, Vandewoude KH, Decruyenaere JM, Colardyn FA. Outcome in noninvasively and invasively ventilated hematologic patients with acute respiratory failure. *Chest* 2004;126:1299-306.
 23. Platon L, Amigues L, Ceballos P, Fegueux N, Daubin D, Bernard N, et al. A reappraisal of ICU and long-term outcome of allogeneic hematopoietic stem cell transplantation patients and reassessment of prognosis factors: results of a 5-year cohort study (2009-2013). *Bone Marrow Transplant* 2016;51: 256-61.
 24. Depuydt P, Kerre T, Noens L, Nollet J, Offner F, Decruyenaere J, et al. Outcome in critically ill patients with allogeneic BM or peripheral haematopoietic SCT: a single-centre experience. *Bone Marrow Transplant* 2011;46:1186-91.
 25. Azevedo LC, Caruso P, Silva UV, Torelly AP, Silva E, Rezende E, et al. Outcomes for patients with cancer admitted to the ICU requiring ventilatory support: results from a prospective multicenter study. *Chest* 2014;146:257-66.
 26. Pène F, Aubron C, Azoulay E, Blot F, Thiéry G, Raynard B, et al. Outcome of critically ill allogeneic hematopoietic stem-cell transplantation recipients: a reappraisal of indications for organ failure supports. *J Clin Oncol* 2006;24:643-9.
 27. Soares M, Salluh JI, Spector N, Rocco JR. Characteristics and outcomes of cancer patients requiring mechanical ventilatory support for >24 hrs. *Crit Care Med* 2005;33:520-6.
 28. Peigne V, Rusinová K, Karlin L, Darmon M, Femand JP, Schlemmer B, et al. Continued survival gains in recent years among critically ill myeloma patients. *Intensive Care Med* 2009;35: 512-8.