



HHS Public Access

Author manuscript

Nat Rev Immunol. Author manuscript; available in PMC 2020 March 01.

Published in final edited form as:

Nat Rev Immunol. 2019 March ; 19(3): 154–169. doi:10.1038/s41577-018-0110-7.

Calcium signalling in T cells

Mohamed Trebak¹, Jean-Pierre Kinet²

¹Department of Cellular and Molecular Physiology, Pennsylvania State University College of Medicine, Hershey, PA, USA

²Department of Pathology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA.

Abstract

Calcium (Ca²⁺) signalling is of paramount importance to immunity. Regulated increases in cytosolic and organellar Ca²⁺ concentrations in lymphocytes control complex and crucial effector functions such as metabolism, proliferation, differentiation, antibody and cytokine secretion and cytotoxicity. Altered Ca²⁺ regulation in lymphocytes leads to various autoimmune, inflammatory and immunodeficiency syndromes. Several types of plasma membrane and organellar Ca²⁺-permeable channels are functional in T cells. They contribute highly localized spatial and temporal Ca²⁺ microdomains that are required for achieving functional specificity. While the mechanistic details of these Ca²⁺ microdomains are only beginning to emerge, it is evident that through crosstalk, synergy and feedback mechanisms, they fine tune T cell signalling to match complex immune responses. In this article, we review the expression and function of various Ca²⁺-permeable channels in the plasma membrane, endoplasmic reticulum, mitochondria and endolysosomes of T cells, and their role in shaping immunity and the pathogenesis of immune-mediated diseases.

Introduction

To mount effective immune responses, lymphocytes must transduce antigenic signals from the surface to their nuclei¹. Signal transduction requires intracellular messenger molecules, the function of which can be switched on and off with time. For Ca²⁺ ions to acquire the on/off switch of a signalling molecule, their concentration within cellular microdomains must vary with time. As such, Ca²⁺ that reaches, for instance, the cytosol when cells are stimulated must subsequently exit into the extracellular space or be sequestered within the endoplasmic reticulum (ER) when signalling is discontinued. Due to their charge however,

Correspondence to M.T. and J.-P.K. mtrebak@psu.edu; jkinet2@mac.com.

Author contributions

M.T. and J.-P.K. both wrote and edited the manuscript.

Competing interests

The authors declare no competing interests.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Reviewer information

Nature Reviews Immunology thanks S. Feske and other anonymous reviewer(s), for their contribution to the peer review of this work.

ions cannot diffuse freely across the lipid bilayer of biological membranes and instead require transmembrane channels and transporters to regulate Ca^{2+} concentrations within the cytosol and intracellular organelles².

T cells express several types of Ca^{2+} -permeable channels and transporters that control Ca^{2+} influx and efflux across the plasma membrane (PM)^{1,3–5}. Intracellular organelles such as the ER, mitochondria and lysosomes also express specific channels and transporters that control Ca^{2+} release into the cytosol and Ca^{2+} uptake into these organelles^{6–10} (FIG. 1). Ca^{2+} transport through channels is controlled by the concentration gradient of Ca^{2+} between each side of a biological membrane and by the electrical gradient or membrane potential [G] (V_m). The concentration gradient of Ca^{2+} across the PM favours a strong driving force of Ca^{2+} into the cytosol. T cells have a resting cytosolic Ca^{2+} concentration 50–100nM compared with the extracellular Ca^{2+} concentration which is 1–2mM¹¹. The PM resting V_m in T cells is –60mV to –50mV, which also favours Ca^{2+} influx into the cytosol. V_m is determined by ion channels [G] conducting sodium (Na^+), potassium (K^+) and chloride (Cl^-), which indirectly regulate Ca^{2+} transport by controlling membrane potential¹¹. These channels have been recently reviewed and are not discussed here⁴.

In T cells, Ca^{2+} influx across the PM is mediated mainly through receptor-activated Ca^{2+} channels, which are operational at the resting V_m . Although the expression of PM voltage-activated Ca^{2+} (Ca_v) channels [G] is generally associated with excitable cells such as muscle cells and neurons, several Ca_v channels have been associated with T cell function^{12–14}. An increase in cytosolic Ca^{2+} levels in T cells can also originate from the ER where Ca^{2+} concentrations are 300 μM –1mM¹⁵. ER Ca^{2+} release channels include inositol-1,4,5-trisphosphate receptors [G] (IP_3Rs)¹⁶ and ryanodine receptors [G] (RYRs)¹⁷. In turn, both Ca^{2+} entry and Ca^{2+} release channels couple receptor activation to metabolic activity and ATP production by providing cytosolic Ca^{2+} microdomains [G], in the order of 1–10 μM , for uptake by mitochondria through mitochondrial Ca^{2+} uniporter [G] (MCU)^{7,18,19} (FIG. 1). The activities of three dehydrogenases in the tricarboxylic acid (TCA) cycle are regulated by mitochondrial Ca^{2+} ^{20–23}, although this has not been shown specifically in T cells. Free mitochondrial Ca^{2+} concentration is estimated to vary between 0.1–10 μM and can, in some instances of receptor stimulation, reach sub-millimolar levels²⁴.

Ion pumps [G] transport Ca^{2+} ions against the electrochemical gradient, thus requiring energy in the form of ATP. These ion pumps include PM Ca^{2+} ATPases (PMCA)^{25,26} and sarcoplasmic/ER Ca^{2+} ATPases (SERCAs)²⁷, which pump Ca^{2+} from the cytosol to the extracellular space and the ER, respectively (FIG. 1). PMCA4b, SERCA2b and SERCA3 are the major isoforms in lymphocytes^{28,29}. Ca^{2+} efflux from mitochondria to the cytosol occurs via mitochondrial $\text{Na}^+/\text{Ca}^{2+}/\text{Li}^+$ exchanger (NCLX; also known as SLC24A6), a transporter that uses a Na^+ gradient to extrude mitochondrial Ca^{2+} ^{30–32} (FIG. 1). Ca^{2+} accumulation in endolysosomes and secretory vesicles occurs through the $\text{H}^+/\text{Ca}^{2+}$ exchanger, which uses the steep proton concentration gradient to drive uptake of Ca^{2+} into these organelles^{10,33}. Generation of high proton concentration in endolysosomes is a process that costs energy through the activity of vacuolar (V)-type H^+ ATPase³⁴. Measurements of Ca^{2+} from lysosomes of macrophages suggest a concentration of 400–600 μM , which varies depending on changes in pH within these organelles³⁵. Ca^{2+} release from endolysosomes and secretory

vesicles into the cytosol occurs through two pore channels [G] (TPCs)³⁶ and transient receptor potential mucolipin (TRPML) channels [G]³³ and is believed to regulate exocytosis and endolysosomal fusion, trafficking and function^{10,33,37}. Below, we discuss how Ca²⁺ microdomains generated by channels in the PM, ER, mitochondria and endolysosomes of T cells shape cellular Ca²⁺ signalling and immunity.

Plasma membrane Ca²⁺ channels

Antigen binding to the T cell receptor (TCR) initiates a cascade of protein phosphorylation that converges in the activation of phosphoinositide-specific phospholipase C γ 1 (PLC γ 1)³⁸ (FIG. 1). PLC γ 1 hydrolyzes, phosphatidylinositol-4,5-bisphosphate (PI(4,5)P₂) into the soluble head group inositol-1,4,5-trisphosphate (IP₃) and the PM-associated lipid diacylglycerol (DAG)⁶. Diffusible IP₃ binds IP₃R on the ER to achieve two interdependent events: first, cause Ca²⁺ release into the cytosol through IP₃R¹⁶; and second, empty ER Ca²⁺ stores and thus indirectly activate Ca²⁺ influx across the PM through store-operated Ca²⁺ entry [G] (SOCE)^{39,40} (FIG. 1). SOCE is mediated by PM channels activated as a consequence of emptying of ER Ca²⁺ stores. Inhibitors of SERCA pumps such as thapsigargin, which cause passive Ca²⁺ leak from the ER, activate SOCE to the same extent as receptor ligation⁴¹. Electrophysiological recordings of currents mediating SOCE identified an inwardly-rectifying Ca²⁺ selective current termed Ca²⁺ release-activated Ca²⁺ CRAC [G]⁴².

Store-operated Ca²⁺ channels: activators

The molecular mechanisms coupling Ca²⁺ store depletion to activation of CRAC channels were discovered in the past decade^{43,44}. The filling of ER stores is detected by the ER Ca²⁺ sensors stromal interaction molecule 1 (STIM1) and STIM2 in the ER membrane^{45,46} (FIG. 1). When ER stores are depleted, Ca²⁺ is released from the Ca²⁺-binding domain of STIMs, which triggers STIM aggregation and movement to junctional spaces where the ER and PM come into close contact, typically within 10–25nm. STIM2 is activated when modest store depletion occurs whereas STIM1 activation requires significant store depletion⁴⁷. ORAI1 is the molecular correlate of PM CRAC channels, which was identified almost simultaneously by three independent groups^{48–50}. ORAI1 and STIM1 are necessary and sufficient to support SOCE^{48–55}. ORAI1 coalesces with STIMs in ER–PM junctions at the immune synapse [G] on store depletion-cell activation⁵⁶.

ORAI1.—The role of CRAC channels in T cell function was recognized long before the discovery of STIMs and ORAI. CRAC channels in T cells are primarily required for the activation of Ca²⁺-dependent transcription factor isoforms of nuclear factor of activated T cells [G] (NFAT) (FIG. 2) and subsequent cytokine production, proliferation and immune competence^{1,4,5}. However, the role of SOCE and CRAC in T cell function is more complex since *in vitro* addition of IL-2 to STIM1-deficient and ORAI1-deficient T cells from either mice or patients only partially restores TCR-mediated proliferation^{57–60}, suggesting additional roles for SOCE beyond the production of proliferative cytokines. One of the mechanisms by which SOCE and NFAT control T cell proliferation involve *de novo* transcription of other transcription factors and metabolic genes (discussed below)⁶¹. Further, SOCE synergizes with other pathways and depending on the magnitude, duration and nature

of Ca^{2+} signals (sustained versus oscillatory), a plethora of different transcription factors are activated⁶² (FIG. 2).

T cells from patients suffering from a form of severe combined immunodeficiency (SCID) have normal Ca^{2+} release, whereas CRAC currents and SOCE were abrogated^{59,63}. These forms of primary SCID are associated with impaired proliferation in response to mitogens due to alterations or mutations in either *ORAI1* or *STIM1* genes, most notably a single amino acid autosomal recessive mutation in ORAI1 (R91W), which generates an inactive channel^{48,64}. Patients and mice deficient in either STIM1 or ORAI1 have increased susceptibility to bacterial, viral and fungal infections. In humans, immunodeficiency is accompanied by an autoimmune phenotype associated with lymphadenopathy, splenomegaly, thrombocytopenia and haemolytic anaemia that is presumably due to reduced numbers and altered suppressive function of FOXP3⁺ regulatory T (T_{reg}) cells which depend on NFAT-mediated transcription⁴. An increasing number of patients with abrogated SOCE due to autosomal recessive mutations in *STIM1* and *ORAI1* were identified with strong reduction in numbers of T_{reg} cells and invariant natural killer T (iNKT) cells^{64,65}.

Most *Orai1*^{-/-} and *Stim1*^{-/-} mice on the C57BL/6 background die perinatally within 24h postpartum⁶⁶⁻⁶⁸. In contrast, *Stim2*^{-/-}⁶⁸, *Orai2*^{-/-}⁶⁹ and *Orai3*^{-/-} (M. T., unpublished observations) mice are viable albeit *Stim2*^{-/-} mice show slight growth retardation and die within 4-5 weeks after birth of unknown reasons. Cytokine production by CD4⁺ and CD8⁺ T cells from STIM1- and ORAI1-deficient patients and mice is dramatically impaired, including IFN γ , tumour necrosis factor (TNF), IL-2, IL-4, IL-10 and IL-17^{4,5}. T cell-specific *Orai1* knockout mice show protection in the model experimental autoimmune encephalomyelitis (EAE) and have reduced T helper 1 (T_H1) and T_H17 cell-dependent cytokine production, including IL-17A and IFN γ ^{70,71}. Unlike the case in humans, *Orai1*^{-/-} mice have no defect in differentiation and function of induced T_{reg} cells, which is likely due to the fact that SOCE is partially inhibited but not completely abrogated, reflecting a potential role for ORAI2 in mice versus humans. Unlike human T cells, SOCE in T cells from either *Orai1*^{-/-} or ORAI1 R91W knock-in mice was only partially inhibited with a residual Ca^{2+} signal potentially reflecting ORAI2 involvement⁶⁷.

CD4⁺ T cell-specific *Stim1* knockout mice have dramatically reduced SOCE in response to anti-CD3 and undetectable CRAC currents. These cells failed to sustain NFAT nuclear translocation and produce IL-2 on anti-CD3 and anti-CD28 co-stimulation⁶⁸. STIM2 is a weaker ORAI1 activator compared with STIM1 and its activation occurs with slower kinetics^{47,72}. It is thus not surprising that STIM2 cannot fully compensate for the lack of STIM1⁴. CRAC currents and SOCE in STIM2-deficient CD4⁺ T cells was partially preserved but these cells failed to sustain higher plateaus of Ca^{2+} levels after 20min of TCR stimulation^{4,5,68}. While NFAT nuclear translocation occurred after stimulation of *Stim2*^{-/-} CD4⁺ T cells, it was not sustained and IL-2 production was inhibited⁶⁸.

The phenotype of *Stim1/2* CD4⁺ T cell-specific double knockout (*Stim1/2*-dKO) mice is more severe, involving splenomegaly, lymphadenopathy and inflammation. They completely lack SOCE and show almost complete abrogation of IL-2 production⁶⁸. *Stim1/2*-dKO mice develop humoral autoimmunity (at 6 months of age) including spontaneous

production of autoantibodies and altered differentiation of T follicular helper (T_{FH}) cells and T follicular regulatory (T_{FR}) cells. Indeed, STIM1 and STIM2 control the expression of the follicular T cell transcription factors IRF4, BATF and BCL-6 through SOCE activation of NFAT⁷³. They showed impaired germinal centre reactions within B cell follicles, failed to produce virus-specific antibodies⁷³ and had 80–90% reduction in T_{reg} cell numbers in lymphoid organs with the few remaining T_{reg} cells presenting markedly impaired suppressive function⁶⁸. Furthermore, differentiation of CD4⁺ T cells into inducible FOXP3⁺ T_{reg} cells and T_H17 cells are also impaired in *Stim1/2*-dKO mice⁴. The expression of glycolytic and oxidative phosphorylation enzymes as well as glucose transporters GLUT1 and GLUT3 were reduced in T cells from *Stim1/2*-dKO mice⁶¹. STIM1/2-mediated SOCE controls metabolic activity of T cells through calcineurin-mediated activation of NFAT and PI3K–mTOR signalling, which in turn regulate the expression of several transcription factors, including HIF1 α , IRF4 and MYC, that are critical for expression of glycolytic enzymes⁶¹. However, the molecular mechanisms connecting Ca²⁺ to the activation of PI3K–AKT–mTOR signalling remain unknown.

Whereas STIM1- and ORAI1-mediated Ca²⁺ signals are crucial for T cell activation, proliferation and metabolism, excessive or prolonged Ca²⁺ signalling can lead to cell death. ORAI1 deficiency renders T cells resistant to death upon long-term exposure to anti-CD3 and anti-CD28, which coincided with severe reduction in nuclear import of NFAT⁷⁴. Expression of either ORAI1 or constitutively active NFAT restored cell death in these cells. T cells from heterozygous *Orai1*^{+/-} mice are not resistant to cell death, suggesting that the levels of cytosolic Ca²⁺ attained differentially activate distinct isoforms of NFAT⁷⁵. NFAT2 is activated by moderate cytosolic Ca²⁺ levels and acts as a positive regulator of T cell proliferation, while NFAT1 activation requires high cytosolic Ca²⁺ and causes cell cycle arrest and death^{76,77}. Resistance to death in *Orai1*^{-/-} T cells correlated with reduced expression of death receptors and pro-apoptotic genes *Fasl* and *Pmaip1* (also known as *Noxa*)⁷⁴. Similarly, *Stim1*^{-/-} T cells were resistant to death on repeated TCR stimulation during chronic mice infection by *Mycobacterium tuberculosis*, with significantly decreased expression of the pro-apoptotic factors FAS, FASL and NOXA in the lungs⁷⁸. Collectively, these results show that SOCE is the major Ca²⁺ entry pathway in T cells required for T cell immunity. STIM1/STIM2- and ORAI1-mediated SOCE is the central activator, which generates critical Ca²⁺ microdomains required for nuclear translocation of NFAT isoforms and optimal cytokine gene expression, metabolism and proliferation.

ORAI2.—*Orai2*^{-/-} mice showed normal populations of thymocytes and peripheral lymphocytes and normal proportions of effector and memory T cells with largely maintained immune responses. ORAI2 was proposed as a negative modulator of ORAI1 that fine-tunes CRAC activity and the strength of T cell immunity⁶⁹. Deletion of *Orai2* either globally or in CD4⁺ T cells enhances SOCE only in naive T cells with no effect on effector T cells. By contrast, *Orai1*^{-/-} mice show strong SOCE reduction in effector T cells and moderate reduction in naive T cells. In fact, both ORAI1 and ORAI2 contribute to CRAC channels in naive T cells where ORAI2 negatively modulates CRAC function. However, effector T cells downregulate ORAI2 and upregulate ORAI1, resulting in a major reduction of SOCE in effector T cells from *Orai1*^{-/-} mice with marginal effects in the same cells from *Orai2*^{-/-}

mice⁶⁹. Increased ORAI2 in naive T cells is expected to yield heteromeric CRAC channels with enhanced Ca²⁺-dependent inactivation (CDI) that could control antigen-mediated signalling in cases of non-specific pathophysiological activation of naive T cells. In effector T cells however, strong ORAI1-mediated SOCE would allow for robust and immediate T cell activation in response to specific antigens. Consistent with a negative modulation of SOCE by ORAI2, an independent group reported that ORAI2 limits SOCE and degranulation in mast cells⁷⁹.

CD4⁺ T cell-specific *Orai1/Orai2* double knockout (*Orai1/2*-dKO) mice largely recapitulate the phenotype of *Stim1/2*-dKO mice. They show no developmental defects of conventional TCRαβ⁺ T cells while displaying splenomegaly and lymphadenopathy with increased proportions of immune cell subsets, including memory and effector T cells⁶⁹. Similar to *Stim1/2*-dKO mice, altered immune homeostasis in peripheral lymphoid organs of *Orai1/2*-dKO mice could be explained by reduced T_{reg} cell numbers in the thymus and peripheral lymphoid organs⁶⁹. Further, germinal centre formation, T_{FH} cell differentiation and antibody production in response to viral infection were significantly reduced in *Orai1/2*-dKO mice⁶⁹. T cells from *Orai1/2*-dKO mice show dramatically inhibited proliferation in response to anti-CD3 and anti-CD28 co-stimulation. IL-2 only partially restored proliferation, reflecting the IL-2-independent role of CRAC in T cell metabolism discussed above⁶⁹. Cytokine production was significantly hampered in T_{H1}, T_{H17} and T_{H2} cells from *Orai1/2*-dKO mice. In contrast, T cells from *Orai2*^{-/-} mice showed largely preserved cytokine production⁶⁹. Unlike wild-type mice and *Orai2*^{-/-} mice, *Orai1/2*-dKO mice are protected from T cell-mediated autoimmunity and alloimmunity in adoptive transfer models of inflammatory bowel disease (IBD) and graft-versus-host disease (GvHD)⁶⁹. Since ORAI2 is a negative regulator of SOCE, one might expect to see exacerbated immune responses in *Orai2*^{-/-} mice in the IBD and GvHD models, which was not the case. The authors speculated that it is likely due to low ORAI2 expression in effector T cells, which largely mediate immunopathology in the colitis and GvHD models⁶⁹.

ORAI3.—Although ORAI3 in heteromeric associations with ORAI1 forms a store-independent Ca²⁺ selective channel in other cell types^{80–84}, the role of ORAI3 in T cells remains largely obscure. Unlike ORAI1, ORAI3 channels are resistant to inhibitory oxidation by hydrogen peroxide (H₂O₂)⁸⁵. The equivalent of a reactive cysteine located in the second extracellular loop (Cys195) of ORAI1 is a glycine in ORAI3. Differentiation of CD4⁺ T cells into effector T cells was accompanied by progressive loss of sensitivity to H₂O₂ and increased expression of ORAI3, suggesting this switch could allow effector cells to continue to proliferate and secrete cytokines in an oxidant-rich inflammatory milieu⁸⁵. Knockdown of *Orai3* in a mouse model of collagen-induced arthritis slightly inhibited SOCE in splenocytes and thymocytes while reducing the production of inflammatory cytokines in joint protein extracts and serum and reducing the severity of arthritis⁸⁶.

Transient receptor potential channels: regulators

Transient receptor potential cation channel subfamily C, member 1 (TRPC1) and TRPC3 are expressed in T cells and knockdown of *TRPC3* caused a small reduction in SOCE and T cell proliferation⁸⁷. Altered TRPC3 expression was reported in several T cell mutants with

defective Ca^{2+} entry in response to TCR ligation and introduction of *TRPC3* cDNA restored TCR-mediated Ca^{2+} influx⁸⁸. The interpretation of these results is unclear and the involvement of TRPC3 in T cells remains highly contentious.

TRPM2 is expressed in T cells and is upregulated upon TCR ligation⁸⁹. Although there is no evidence that TCR ligation activates TRPM2 in T cells, native TRPM2-mediated currents in Jurkat T cells can be activated by cyclic ADP ribose (cADPR), ADPR and NAADP⁸⁹. TCR stimulation causes sustained increases in intracellular cADPR, and an antagonist of cADPR inhibited T cell activation and proliferation in response to TCR stimulation⁹⁰. These results however can be explained by an effect of cADPR on RYR (discussed below). *Trpm2*^{-/-} mice are partially protected in the EAE model and this correlates with reduced proliferation of CD4⁺ T cells and inhibited secretion of IL-17 and IFN γ in response to TCR stimulation⁹¹.

Increased cytosolic Ca^{2+} levels through SOCE upon TCR activation triggers Na^+ influx through TRPM4 (Kd ~400nM) to depolarize the PM and restrict further Ca^{2+} entry⁹². Ca^{2+} -induced activation of TRPM4 channels in T cells serves as a negative feedback mechanism, preventing excessive cytosolic Ca^{2+} -mediated toxicity and fine-tuning T cell responses^{1,92,93}. Indeed, in the absence of TRPM4 channels, SOCE-mediated intracellular Ca^{2+} oscillations in response to stimulation of T cells by mitogens became sustained, resulting in enhanced cytosolic Ca^{2+} and increased IL-2 production⁹³. The expression levels of TRPM4 channels seem to be higher in T_H2 cells than T_H1 cells, which could explain the reduced TCR-mediated Ca^{2+} flux in T_H2 cells⁹⁴, and might explain the relatively higher sensitivity to FAS-dependent apoptosis in T_H1 cells than T_H2 cells⁹⁵. In contrast to TRPM4, Ca^{2+} -activated and voltage-activated K^+ channels counteract TRPM4 and play a positive modulatory role in T cell Ca^{2+} signalling. K^+ efflux, which maintains a hyperpolarized PM Vm and enhances the driving force for Ca^{2+} entry, is a crucial regulator of T cell immunity⁹⁶.

TRPM7 is critical for T cell development as T cell-specific *Trpm7* knockout mice have reduced numbers of peripheral T cells due to halting of thymocyte development at the double negative CD4⁻CD8⁻ stage⁹⁷. Surprisingly, although these mice have a marked reduction in TRPM7 currents in thymocytes, acute Mg^{2+} uptake and total cellular Mg^{2+} levels were normal, suggesting that TRPM7 function in T cells likely occurs through either conduction of Ca^{2+} ions or its cytosolic serine/threonine kinase⁹⁷. Notwithstanding a potential direct regulation of cytosolic Ca^{2+} signalling, TRPM7 might indirectly regulate SOCE in lymphocytes⁹⁸ (FIG. 3). Deletion of TRPM7 or its pharmacological inhibition inhibits SOCE. The positive regulation of SOCE by TRPM7 seems to require the kinase activity, but not the channel domain of TRPM7⁹⁸. Thymocyte development was unaffected in TRPM7 kinase-dead (K1646R) knock-in mice, suggesting that the channel domain, but not the kinase domain, is crucial for T cell development⁹⁹. Splenic T cells from TRPM7-K1646R knock-in mice exhibited normal TRPM7 currents while showing reduced SOCE and reduced proliferation with mice exhibiting splenomegaly and altered blastogenesis in response to co-stimulation with anti-CD3 and anti-CD28 antibodies or phorbol myristate acetate (PMA) and ionomycin⁹⁹. An independent study showed that TRPM7-K1646R mice have normal T cell development, but TGF β -mediated expression of CD103, SMAD2 phosphorylation and nuclear translocation and T_H17 cell differentiation were altered¹⁰⁰.

TRPM7 kinase activity was required for gut colonization by alloreactive T cells during an acute model of GvHD¹⁰⁰. The TRPM7 channel domain, but not its kinase domain, regulates T cell homeostasis by mediating FAS-dependent T cell apoptosis through caspase activation¹⁰¹.

One group reported the expression of TRPV1 in CD4⁺ T cells¹⁰². These investigators reported a small inhibition of TCR-activated Ca²⁺ entry in TRPV1-deficient CD4⁺ T cells but only under low extracellular Ca²⁺ concentrations with no effect on SOCE activated by thapsigargin. These authors suggested that TRPV1 is activated downstream of TCR through TRPV1 phosphorylation by the tyrosine kinase LCK¹⁰². However, this study lacks thorough electrophysiological characterization supporting TRPV1 activation downstream of TCR. Mice with TRPV1-deficient CD4⁺ T cells showed reduced colitis severity during IBD¹⁰² and were protected from allergic rhinitis¹⁰³. The same group reported that TRPA1 inhibits TRPV1 through TRPA1–TRPV1 heteromultimeric assemblies to limit T cell activation and colitis in IBD¹⁰⁴. Nevertheless, in light of complete abrogation of TCR-mediated Ca²⁺ signals in *Stim1/2*-dKO, the involvement of TRPV1 and TRPA1 (or TRPC3) in TCR-mediated Ca²⁺ signalling is a highly contentious issue and stronger electrophysiological evidence is required to support a role for these channels in T cell Ca²⁺ signalling. In summary, TRP channel isoforms are important regulators of TCR signalling and can exert this function by controlling the driving force for SOCE through Na⁺ entry-mediated control of PM Vm (TRPM4), by regulating the activity of SOCE (TRPM7 kinase activity), or by potentially providing Ca²⁺ microdomains that either regulate T cell development or synergize with SOCE during T cell activation.

Purinergic ionotropic receptors: amplifiers

Purinergic ionotropic receptors (P2RX) represent a family of seven nonselective trimeric channels with a multitude of splice variants capable of homomerization and heteromerization¹⁰⁵. P2RX channels conduct Na⁺ and Ca²⁺ and are directly activated by extracellular ATP. Most P2RX isoforms are expressed in thymocytes and peripheral T cells¹⁰⁶. However, electrophysiological characterization of P2RX receptors in T cells is lacking and the precise ions (Ca²⁺ versus Na⁺ entry or K⁺ efflux) mediating P2RX receptor effects are unclear. While knockout mice for either *P2rx1*, *P2rx4* or *P2rx7* have normal T cell development, one group reported that *P2rx2/P2rx3* double knockout mice exhibit splenomegaly¹⁰⁷, with a different group reporting reduced cellular numbers in the thymus and bone marrow and altered lymph follicles in spleen and peripheral lymph nodes¹⁰⁸, suggesting isoform redundancy. Very little is known about P2RX5 and P2RX6, except that the expression of P2RX5 is increased in activated T cells¹⁰⁹. Based on small interfering RNA-mediated knockdown of *P2RX1*, *P2RX4* and *P2RX7* in primary T cells and Jurkat T cells, these channels seem to couple to activation of NFAT-dependent IL-2 production and T cell proliferation in response to TCR ligation¹⁰⁶. TCR-mediated activation of P2RX receptors could presumably be initiated by IP₃R-dependent and SOCE-dependent mitochondrial Ca²⁺ uptake leading to the production of ATP and its extracellular export to activate P2RX receptors in an autocrine and paracrine fashion (FIG. 4). In support of this, Pannexin1 channels, which can export ATP to the extracellular space, translocate to the immune synapse where they colocalize with STIM1, ORAI1, P2RX1 and P2RX4¹¹⁰. In this

regard, P2RX receptors serve as amplifiers of TCR-mediated SOCE and T cell effector functions. TCR ligation is known to promote mitochondrial ATP production, which would fuel P2RX receptor-mediated Ca^{2+} entry to further enhance ATP production, thus amplifying weak TCR signals to produce calcineurin–NFAT activation and mount adequate T cell responses (FIG. 4).

P2RX7 was initially proposed to mediate lymphocyte cytotoxicity and cell death in response to enhanced extracellular ATP levels. However, under moderate conditions of stimulation, P2RX7 can promote T cell activation and growth by coupling to mitochondrial bioenergetics and ATP production (presumably by providing cytosolic Ca^{2+} for mitochondrial uptake) and by enhancing glycolysis through transcriptional upregulation of glycolytic enzymes¹⁰⁶. Excessive P2RX7 stimulation causes mitochondrial Ca^{2+} overload, uncoupling of oxidative phosphorylation and cell death¹⁰⁶, which is analogous to TCR-mediated activation of SOCE where the strength of stimulation and the expression levels of STIM and ORAI can promote either T cell activation or death. Increasing evidence points towards a role for P2RX1, P2RX4 and P2RX7 in promoting T cell-mediated immunity and inflammatory responses in vivo, with P2RX7 being the most studied in this context^{105,106}. A homeostatic role for P2RX1 was proposed to be required for T cell activation. This consists of a positive feedback loop involving basal mitochondrial ATP production, ATP release and activation of P2RX1¹¹¹. Stimulated T cells in lymph nodes secrete ATP and activate P2RX4 and P2RX7 to mediate paracrine calcium waves in neighbouring resting lymphocytes to reduce their motility and help antigenic scanning of resident dendritic cells¹¹². P2RX4 and P2RX7 seem to play an important role in the amplification of TCR-mediated Ca^{2+} signalling in non-conventional $\gamma\delta$ T cell differentiation and cytokine production^{113,114}.

Pan inhibition of P2RX receptors in mice using oxidized ATP prevented the development of colitis in adoptive T cell transfer models and was associated with increased T_{reg} cell numbers in peripheral lymph nodes and reduced production of IL-17. The activation of the ubiquitous P2RX7 inhibits the immunosuppressive functions of T_{reg} cell and facilitates their conversion into $T_{\text{H}}17$ cells, which is accompanied by reduced expression of FOXP3 and enhanced expression of the $T_{\text{H}}17$ cell-specific transcription factor ROR γ t¹¹⁵ (FIG. 4). Consistent with these findings, inactivation of P2RX7 with oxidized ATP showed preserved pancreatic islet allografts in mice and reduced $T_{\text{H}}1$ and $T_{\text{H}}17$ cells in this model¹¹⁶. Inflammatory sites contain high concentrations of ATP, in the order of hundred micromolar¹⁰⁶. P2RX7 expression was specifically upregulated in graft-infiltrating lymphocytes in humans and mice with cardiac transplants and P2RX7 inhibition delayed coronaropathy and enhanced survival of cardiac allografts in a mouse model of chronic rejection¹¹⁷. In ischaemia-reperfusion injury, P2RX receptor inhibition by oxidized ATP promoted expansion of T_{reg} cells, and blunted acute renal injury and improved renal recovery¹¹⁸. *P2rx7*^{-/-} mice are protected in the EAE model¹¹⁹ and in a model of experimental autoimmune uveoretinitis¹²⁰. *P2rx7*^{-/-} mice have enhanced susceptibility to *Plasmodium chabaudi* malaria infection, which correlated with decreased IL-2 and IFN γ secretion by CD4⁺ T cells, T-bet expression and impaired $T_{\text{H}}1$ cell differentiation. The requirement for P2RX7 in $T_{\text{H}}1$ cell differentiation during infection was accompanied by a reduction in the splenic T_{FH} cell population of infected mice¹²¹. Of note, the beneficial effects of P2RX7 inhibition on reducing inflammation through promotion of T_{reg} cells in the

digestive tract in a colitis model were offset by anti-apoptotic effects of P2RX7 blockade on intestinal epithelial cells, which caused increased proliferation of these cells to promote colitis-associated cancer¹²². Deletion or pharmacological inhibition of P2RX7 is known to enhance tumour metastasis and to drastically alter immunogenic cell death of cancers, presumably because P2RX7 is required for dendritic cells to release IL-1 β (for review, see¹⁰⁶).

Recent work showed that P2RX7 is required for generation and function of durable long-lived memory CD8⁺ T cells but not short-lived effector CD8⁺ T cells. P2RX7 achieves this role by promoting metabolic fitness of these cells through induction of AMP-activated protein kinase and maintenance of mitochondrial homeostasis and metabolism¹²³. While overall data supports P2RX7 activity as pro-inflammatory, an earlier report showed that *P2rx7*^{-/-} mice have exacerbated EAE due to reduced apoptosis of lymphocytes¹²⁴. This could be explained by the *P2rx7*^{-/-} strain used that lacks P2RX7 in macrophages and dendritic cells while unexpectedly expressing higher levels in T cells¹²⁵. Nevertheless, there are conflicting reports showing that exogenous ATP suppresses T cell activation and IL-2 production¹²⁶ and ATP production by dying acute myeloid leukemia cells during chemotherapy causes immune suppression¹²⁷. The overall data suggests the P2RX receptors play a homeostatic (P2RX1) and amplifying (P2RX4 and P2RX7) role of TCR signalling. P2RX receptors seem to synergize with SOCE through a positive feedback loop involving mitochondrial Ca²⁺ uptake and enhanced ATP production. SOCE-mediated mitochondrial ATP production, ATP export by pannexin1 channels and ATP action on P2RX receptors can cause further Ca²⁺ entry to amplify weak TCR signals and help in antigen scanning.

Ca_v channels: modulators

L-type Ca²⁺ channels.—Transcripts and proteins of all four members of the L-type (long-lasting) Ca_v channel family and regulatory β 3 and β 4 subunits are expressed in mouse and human T cells and their expression is increased upon anti-CD3 and anti-CD28 co-stimulation¹²⁸. Yet, the involvement of voltage-activated Ca²⁺ channels in T cell function is a highly contentious issue. T cells rely on a negative membrane potential for Ca²⁺ entry and in which depolarization is expected to inhibit Ca²⁺ transport across the PM. Nevertheless, several studies reported that T cells express truncated or alternatively spliced forms of L-type Ca_v channels, which either lack one voltage sensor (e.g. Ca_v1.4) or a linker adjacent to voltage sensor (Ca_v1.1)¹²⁹, suggesting that Ca_v channels in T cells might not be voltage-gated and can be activated by alternative mechanisms. Such mechanisms may involve phosphorylation of Ca_v channels by protein kinase C (PKC) and protein kinase A (PKA); both kinases are activated downstream SOCE and are Ca_v regulators in different cell types¹³⁰ (FIG. 5). Of significance, constitutively active Ca_v1.2 channels, which required PKC, were reported in arterial smooth muscle cells^{131–134}. Quantal Ca²⁺ sparklets and single Ca_v1.2 channel currents were recorded at membrane potentials as low as -90 mV¹³⁵. Similar to its non-channel function in skeletal muscle¹³⁶, the Ca_v1.1 splice variant expressed in T cells¹³⁷ could be involved in a voltage-independent Ca²⁺ entry-independent conformational coupling with RYR at ER-PM junctions to induce ER Ca²⁺ release via RYR (discussed below) (FIG. 5). However, experimental data in support of conformational

coupling of Ca_v1.1 to RYR or Ca_v1.2 channel clustering and activation in lymphocytes is lacking.

One major limitation is the lack of thorough biophysical characterization of L-type currents in T cells by patch clamp electrophysiology [G] as any alteration of Ca²⁺ signalling measured with fluorescent dyes can be equally explained by changes in membrane potential. Although a small Ca_v-like current activated by depolarizing voltage ramps was reported in wild-type T cells and was lost in T cells from Ca_v1.4-deficient mice, these currents, which were recorded in non-physiological extracellular solutions containing high Ba²⁺ concentrations (100 mM) as the charge carrier, remain controversial¹³. Independent recordings from activated human T cells readily revealed CRAC currents but failed to detect voltage-activated Ca²⁺ currents, using the same conditions that produced voltage-activated Ca²⁺ currents in neuronal PC12 cells⁵⁶. Ca²⁺ measurements in response to TCR stimulation or thapsigargin showed reduction in Ca²⁺ release and Ca²⁺ entry in Ca_v1.4-deficient T cells, leading the authors to propose that Ca_v1.4 mediates basal Ca²⁺ entry important for ER refilling¹³ (FIG. 5). Whether loss of Ca_v1.4 in T cells alters the expression of ER and PM Ca²⁺ transport proteins or membrane potential was not explored. Ca_v1.4-deficient mice showed a reduction in mature thymocytes and in peripheral CD4⁺ and CD8⁺ T cells, suggesting a role for Ca_v1.4 in promoting T cell survival and expansion¹³. Similarly, knockout of the auxiliary subunits β3 and β4 in mice led to reduction in Ca_v1.1 expression and in partial inhibition of TCR-activated Ca²⁺ signals and NFAT1 and NFAT2 activation in CD4⁺ T cells¹². The β subunit-deficient CD4⁺ T cells displayed defects in cytokine production (IL-2, IL-4 and IFNγ) but proliferation in response to TCR stimulation was unaffected¹². Deletion of the β3 subunit led to loss of Ca_v1.4 expression, impaired TCR-activated Ca²⁺ influx and inhibited survival of CD8⁺ T cells¹³⁸. T cell-specific deletion of the β2 subunit reduced the expression of Ca_v1.2 and Ca_v1.3 and altered T cell development, causing a dramatic decrease in thymocytes and peripheral T cell numbers¹³⁹. Ca_v1.2 and Ca_v1.3 were proposed to be crucial for T_H2 cells (but not for T_H1 cells); their inhibition was protective in experimental models of asthma¹²⁹.

T-type Ca²⁺ channels.—The α1 pore-forming subunit of Ca_v3.1 is expressed at the PM of CD4⁺ T cells¹⁴. Notwithstanding a potential activation of T-type (transient) Ca²⁺ channels through voltage-independent mechanisms, robust T-type currents were readily apparent in T cells at resting V_m and had a range of window currents from -65mV to -25mV, which is compatible with T-type channels being functional under resting membrane potential in T cells¹⁴ (FIG. 5). Knockout of the Ca_v3.1 α1 subunit in mice led to a ~50% reduction of T-type currents in T cells, with the remaining current likely reflecting compensatory upregulation of Ca_v3.2 and Ca_v3.3 channels¹⁴. T cells from Ca_v3.1-deficient mice showed normal TCR-activated Ca²⁺ entry, suggesting that Ca_v3.1 mediates constitutive Ca²⁺ entry independently of SOCE. Naive CD4⁺ T cells from Ca_v3.1-deficient mice as well as T cells with acute deletion of Ca_v3.1 and polarized in vitro into T_H17 cells showed decreased IL-17A, IL-17F and IL-21 production¹⁴. Ca_v3.1-deficient mice were protected from EAE and showed inhibited production of granulocyte-macrophage colony-stimulating factor (GM-CSF), a cytokine implicated in susceptibility to EAE, by brain-infiltrating T_H1 and T_H17 cells¹⁴. Nuclear translocation of NFAT1 and NFAT2 after anti-CD3 and anti-CD28 co-

stimulation was reduced in $\text{Ca}_v3.1$ -deficient CD4^+ T cells, suggesting that basal $\text{Ca}_v3.1$ -mediated Ca^{2+} increase is crucial for calcineurin–NFAT activation to drive transcription of cytokines and other transcription factors, including GM-CSF and $\text{ROR}\gamma\text{t}^{14}$ (FIG. 5).

Regardless of how L-type and T-type channel isoforms are activated in T cells, they likely provide discrete Ca^{2+} microdomains that fine-tune TCR-mediated Ca^{2+} signals. This can be achieved through coupling of Ca^{2+} microdomains to SERCA for ER Ca^{2+} store refilling and optimal SOCE function, by directly sustaining NFAT activation and production of cytokines during T cell activation, or by coupling to downstream pathways that control differentiation of specific T cell populations (such as $\text{T}_\text{H}17$ cells).

ER Ca^{2+} release channels: initiators and orchestrators

ER Ca^{2+} release channels provide initial cytosolic Ca^{2+} microdomains in response to TCR ligation. They function through this primary mode to stimulate further ER Ca^{2+} release and Ca^{2+} entry across the PM required for proliferation and secretion of cytokines and provide Ca^{2+} microdomains for uptake by mitochondria to enhance T cell metabolism and support activation. In the long term, ER Ca^{2+} release channels can have a sustaining and synergistic effect on cellular Ca^{2+} signals emanating from PM and organellar Ca^{2+} channels.

IP₃Rs.—The initial step in response to TCR stimulation, preceding Ca^{2+} entry across the PM, is Ca^{2+} release from the ER by soluble second messengers. The most extensively studied Ca^{2+} -releasing second messenger is IP_3 , which causes Ca^{2+} release through IP_3R^6 . T cells express all three isoforms of IP_3R , which have distinct sensitivity to IP_3 , allosteric regulation by Ca^{2+} and Ca^{2+} release patterns⁸. Heteromerization of different IP_3R isoforms can further enhance the diversity of Ca^{2+} release signals¹⁴⁰. Type I IP_3R ($\text{IP}_3\text{R1}$) is essential for IP_3 -mediated Ca^{2+} release in response to TCR stimulation¹⁴¹. T cells deficient in $\text{IP}_3\text{R1}$ are resistant to apoptosis induced by TCR ligation, dexamethasone and FAS stimulation¹⁴². Recently, a Tie2-Cre mouse was used to conditionally delete all three genes encoding IP_3R isoforms in mice. In IP_3R triple knockout mice, thymocyte development was arrested at the double negative 4 (DN4 ; $\text{CD44}^-\text{CD25}^-$) and immature single positive (ISP) stages, which coincided with the development of T cell acute lymphoblastic leukemia (T-ALL) in knockout mice¹⁴³. Thymocyte development was normal up to the $\text{CD4}^+\text{CD8}^+$ double positive (DP) stage when one or two IP_3R types were deleted, highlighting IP_3R redundancy in T cell function¹⁴³. Mechanistically, IP_3R triple knockout leads to maintained expression of SOX13 past the DN3 stage. SOX13 is an antagonist of the WNT transcription factor TCF7 (also known as TCF1). The continued expression of SOX13 inhibits TCF7 and halts thymocyte development at the DN to DP transition¹⁴³. IP_3R -mediated Ca^{2+} signals were also proposed to inhibit oncogenic Notch1 transcriptional target HES1 to promote tumorigenesis¹⁴³.

RYRs.—Guse and co-workers proposed a role for the second messengers NAADP and cADPR in T cell Ca^{2+} signalling. NAADP and cADPR, which act on RYR Ca^{2+} release channels, are produced after TCR ligation^{17,90}. NAADP, which is produced early after TCR stimulation, provides a trigger by targeting RYR1 through binding to an unknown accessory molecule¹⁴⁴. Early (within 20 milliseconds) Ca^{2+} microdomains near the PM following

activation by either TCR ligation or NAADP injection were absent or markedly reduced in T cells from *Ryr1*^{-/-} mice¹⁴⁵. This early Ca²⁺ signal facilitates the activation of IP₃R by IP₃ within seconds of TCR stimulation through the process of Ca²⁺-induced Ca²⁺ release (FIG. 6). Later, within tens of minutes, production of cADPR would act on RYR2 or RYR3 to sustain Ca²⁺ signalling for an extended period of time^{17,144}. In T cells, production of IP₃ and cADPR occur within tens of seconds and minutes respectively, and were proposed to sustain late phases of Ca²⁺ signalling¹⁴⁶ (FIG. 6). However, although the role of IP₃ in sustaining Ca²⁺ signalling is well established, the involvement of cADPR is less certain.

A pharmacological compound that blocks the interaction of NAADP with RYR1 inhibited NAADP-mediated Ca²⁺ release, formation of the immune synapse, nuclear translocation of NFAT, IL-2 production and proliferation of CD4⁺ T cells¹⁴⁷. However, pharmacological inhibition of RYR by ryanodine suggested that Ca²⁺ release via RYR was not the direct result of TCR activation but rather depended on RYR activation by Ca²⁺ originating from SOCE¹⁴⁸. In turn, Ca²⁺ release via RYR maintains proximal ER store depletion to sustain SOCE¹⁴⁹ (FIG. 6). Inhibition of RYR reduced T cell proliferation and IL-2 production^{148,149}. However, an independent study showed that T cells lacking RYR3, which is the main isoform in T cells, have normal proliferation to mitogens¹⁵⁰. Furthermore, depending on the cell type considered, the channel (or channels) targeted by NAADP remain uncertain. Some studies suggested that NAADP activates the endolysosomal channels, TPCs¹⁵¹, with potential significance for the function of cytotoxic T cells (discussed below). Other studies showed Ca²⁺ release from the ER in response to NAADP but failed to observe activation of Ca²⁺ release from acidic organelles¹⁵², whereas independent studies showed that NAADP released Ca²⁺ from both ER and acidic organelles¹⁵³.

Mitochondrial Ca²⁺ signalling

Mitochondria control biosynthesis, thermogenesis, ATP and reactive oxygen species (ROS) production and cell death through apoptosis¹⁵⁴. Mitochondria are active participants in Ca²⁺ signalling^{7,155–158}. They uptake cytosolic Ca²⁺ to enhance bioenergetics necessary for T cell activation and proliferation but can also act as buffers and thus shape the spatial distribution of cytosolic Ca²⁺ signals^{7,155–158} (FIG. 6). In T cells, mitochondria move towards the immune synapse, during interactions between T cells and antigen-presenting cells^{159,160}. ORAI1 channels in T cells are also located at the immune synapse and Ca²⁺ buffering by mitochondria within close vicinity of ORAI1 relieves the channel's CDI enabling cells to maintain sustained levels of Ca²⁺ necessary for efficient NFAT activation^{161,162}. While mitochondria clearly cannot fit within the small space of the ER-PM junctions *per se*, they locate as close as ~200nm from the PM¹⁶³. Intriguingly, at the immune synapse mitochondria regulate PMCA¹⁶⁴. Typically, PMCA is inhibited at resting Ca²⁺ levels but as cytosolic Ca²⁺ increases, its activity slowly increases due to binding of Ca²⁺-calmodulin to its C-terminus, a process known as modulation, which allows fast Ca²⁺ transients to proceed relatively unaffected while sustained Ca²⁺ signals that might be toxic to cells are curtailed. During immune synapse formation, PMCA4b, the most common isoform of PMCA in T cells, redistributes into areas that are juxtaposed with mitochondria¹⁶⁴. This close apposition allows mitochondria to buffer Ca²⁺ away from PMCA4b, preventing its modulation and reducing Ca²⁺ extrusion outside the cell (FIG. 6), thus maximizing ORAI1-mediated Ca²⁺

signalling for robust T cell activation¹⁶⁴. Further, STIM1 physically interacts with PMCA at the IS to inhibit PMCA-mediated Ca^{2+} clearance during T cell activation¹⁶⁵.

Mitochondrial Ca^{2+} uptake through the inner membrane channel MCU^{18,19} is driven by the steep mitochondrial membrane potential (~ -180 mV) whereas mitochondrial Ca^{2+} extrusion is achieved mainly by NCLX^{30,158,166} and potentially by $\text{H}^+/\text{Ca}^{2+}$ exchange activity¹⁶⁷. Mitochondrial Ca^{2+} extrusion can shuttle Ca^{2+} to different areas of the cytosol and support efficient ER refilling by dumping Ca^{2+} at the vicinity of SERCA pumps¹⁶⁸. Sustained elevation of cytosolic Ca^{2+} in response to TCR stimulation requires not only mitochondrial Ca^{2+} uptake but also depends on mitochondrial Ca^{2+} extrusion. Inhibition of mitochondrial Ca^{2+} extrusion in T cells through Na^+ depletion inhibits TCR- and store depletion-activated Ca^{2+} entry¹⁶¹. Na^+ depletion or NCLX deficiency prevents mitochondrial Ca^{2+} extrusion, cause mitochondrial Ca^{2+} overload and enhance mitochondrial ROS, which cause ORAI1 oxidation at Cys195 and its inhibition³¹. The role of MCU and NCLX in T cell Ca^{2+} signalling and function remain essentially unknown.

Vesicular Ca^{2+} channels

Trafficking by means of endosomes, recycling endosomes, lysosomes and secretory lysosomes is essential for T cell function^{169,170}. Ca^{2+} release from these organelles has emerged as crucial for T cell activation¹⁶⁹. Recycling endosomes can sustain TCR signalling at the immune synapse through endocytosis of exhausted signalling molecules and replenishment of endocytosed molecules, including TCR-CD3 complexes¹⁷¹. Guided secretion of cytokines and lytic enzymes at the immune synapse also occurs through secretory vesicles, exosomes and secretory lysosomes. The best studied are lysosomes or lysosome-like acidic organelles where Ca^{2+} release was proposed to occur through NAADP action on TPCs¹⁷²⁻¹⁷⁴. TPC2 is found in lysosomes whereas TPC1 is predominantly expressed in endosomes¹⁷⁵. Whether TPCs are Ca^{2+} permeable channels is a contentious issue since whole-endolysosome patch clamp measurements suggested that TPCs that are activated by $\text{PI}(3,5)\text{P}_2$ are Na^+ -selective^{176,177}. Unlike $\text{PI}(4,5)\text{P}_2$, which is associated with PM, $\text{PI}(3,5)\text{P}_2$ is enriched in late endosomes and lysosomes¹⁷⁸. TPCs play a crucial role in vesicular trafficking associated with endocytosis and autophagy¹⁰. Upon cytotoxic T cell activation, NAADP-mediated activation of TPCs cause Ca^{2+} release from cytolytic granules, migration of these granules to the immune synapse and exocytosis of perforin and granzymes in the synaptic space to induce target cell killing¹⁷⁹. These TPC functions could not be recapitulated by increasing global cytosolic Ca^{2+} with either a cell permeant IP_3 or ionomycin, suggesting that IP_3 -mediated Ca^{2+} release and SOCE (which are crucial for exocytosis) are not sufficient and additional Ca^{2+} microdomains at the outer surface of secretory lysosomes are required¹⁷⁹.

Mucolipin TRPs (TRPML1-TRPML3) are non-selective Ca^{2+} -permeable channels found both at the PM and in endolysosomes³³. TRPML1 is ubiquitously expressed throughout the body whereas TRPML2 and TRPML3 are more tissue-restricted with TRPML2 prominently expressed in immune cells^{180,181}. TRPML1 is enriched on the lysosomal surface, whereas TRPML2 is localized mainly to endosomes¹⁸². TRPML3 is found on lysosomes and on the PM^{181,183}. Similar to TPCs, the activation of mucolipin TRPs depends on $\text{PI}(3,5)\text{P}_2$ ¹⁸⁴.

TRPML1 is required for vesicular fusion, trafficking of endocytic vesicles, lysosomal exocytosis and phagocytosis^{185,186}.

Intriguingly, TRPML channels are inhibited by PM-localized PI(4,5)P₂, suggesting that TRPMLs function is restricted to endolysosomes^{187,188}. Adenosine in the lysosomal lumen was also reported to be a TRPML1 inhibitor. Abnormally increased adenosine within lysosomes (normally generated from nucleic acids and ATP) impairs the function of TRPML1 channels leading to dysfunctional, enlarged and alkaline lysosomes and altered lymphocyte function¹⁸⁹. Increased lysosomal adenosine in patients deficient in adenosine deaminase have impaired immune function reminiscent of SCID¹⁹⁰.

Summary and perspectives

T cells express a plethora of Ca²⁺ permeable channels at various locations, with unique activation mechanisms that are necessary for T cell activation, maturation and secretion of cytokines. Ca²⁺ signalling in lymphocytes is tightly coupled to the regulation of adaptive immunity, differentiation and metabolism. Future research should tackle this increasingly diverse array of Ca²⁺ channels, their regulation, temporal expression, unique functions and how they synergize in controlling specific aspects of immunity. Why do cells require this large arsenal of channels to conduct the same ion? Distinct channels positioned in discrete locations within the PM, mitochondria and other organelles sculpt a dynamic cytosolic “Ca²⁺ signature” in the form of Ca²⁺ microdomains at the vicinity of key Ca²⁺-activated effector proteins. Studies showed that Ca²⁺ concentrations reach hundreds of μM nanometers away from a channel pore, while few μm away from the pore Ca²⁺ concentrations precipitously fall to the nM range¹⁹¹. Because Ca²⁺ signals can poorly diffuse through cellular buffers, highly localized spatial and temporal Ca²⁺ microdomains for Ca²⁺ signalling and Ca²⁺ transfer between different organelles are exquisitely specific regulators of T cell function. Ca²⁺ is transferred between organelles through highly-specialized and tight contact sites, epitomized by the mitochondria-associated membranes (MAMs), which are intimate ER-mitochondria contact regions. Therefore, the source and location of a Ca²⁺ signal is what determines which downstream signalling pathways and gene programs are activated, and it is the major determinant by which the ubiquitous Ca²⁺ signal achieves specificity.

Depending on the nature of stimuli, their concentration and affinity for receptors, Ca²⁺ signals can take the shape of either sustained plateaus or regenerative oscillations. While it is appreciated that the strength and duration of Ca²⁺ plateaus can determine whether a T cell proliferates or dies, different levels and durations of Ca²⁺ plateaus likely activate different transcription programs in a T cell subset-dependent manner. The frequency and amplitude of Ca²⁺ oscillations, which are controlled by factors such as the strength of stimulation and levels of ion channels (e.g. TRPM4), provide a bandwidth of digitally distinct signals that are decoded by downstream cytosolic and nuclear proteins. It is therefore not surprising that a large number of channel isoforms are required to mount effective immune responses. To ensure a high degree of specificity, T cells probably require few “initiating” Ca²⁺ signals (for example, through IP₃R) and “activating” Ca²⁺ signals (for example, through ORAI1), as well as many modulatory Ca²⁺ signals (for example, through Ca_v, P2RX and TRP), for fine

tuning T cell activation. This incredibly complex network of channels provides the necessary diversity and flexibility to match the staggering complexity of immune responses.

While evidence exists for Ca^{2+} transfer from ER and PM to mitochondria, potential sites for Ca^{2+} transfer between endolysosomes and mitochondria are unknown. The field of Ca^{2+} signalling by intracellular vesicles is at its infancy with even scarcer knowledge as it relates to immune cells. The contribution of Ca^{2+} release from organelles to T cell activation and the potential differences in the patterns of expression of specific vesicular Ca^{2+} channels in different T cell subsets await future investigations. The advent of super-resolution imaging is beginning to allow for dynamic measurements of Ca^{2+} microdomains in living T cells. The development of cells and mice with tissue-specific expression of genetically-encoded Ca^{2+} indicators with custom-made dissociation constants for Ca^{2+} that can be targeted to specific PM areas, different sides of organellar membranes or even tethered to channels will likely speed up our understanding of the contributions of different channels to T cell function. In any case, the era of studying Ca^{2+} signalling in a specific cellular compartment in isolation is coming to an end. Ca^{2+} signalling operates as a network and regulates what was long considered housekeeping functions, such as metabolism. Future studies should consider simultaneous measurements of Ca^{2+} in different organelles, something that is now possible. As we toil to better understand Ca^{2+} signalling in T cells, we ought to adapt inclusive approaches that addresses not only signalling and gene transcription, but also metabolism, organellar trafficking and secretion for all these converge to exert crucial control over T cell activation and immune function. This understanding will lead to specific channel inhibitors for therapy of transplant rejection, immune disease and inflammatory disease.

Acknowledgements

Research in the authors' laboratories is supported by grants R01HL123364, R01HL097111 and R21AG050072 from the National Institutes of Health, and grant NPRP8-110-3-021 from the Qatar National Research Fund (QNRF) to M.T.

Glossary

Ca^{2+} microdomains

Discrete sites in the cytosol of cells localized near the mouth (within few nm) of Ca^{2+} channels of either plasma membrane or organellar membranes. These regions, which contain high Ca^{2+} concentrations, are the sites where specific Ca^{2+} -activated effector proteins are located. Ca^{2+} microdomains near specific Ca^{2+} channels are the major means by which the ubiquitous Ca^{2+} ion ensures specificity of signal transduction

Membrane potential

The difference in electric potential between the interior and the exterior of a biological membrane. In resting T cells, the plasma membrane potential is typically between -60 and -50 mV

Ion channels

Transmembrane proteins that form oligomers around a central pore, which allows specific ions to flow across biological membranes. Channels conduct ions according to the

electrochemical gradient of this membrane and therefore this process does not consume energy in the form of ATP

Voltage-activated Ca^{2+} (Ca_V) channels

Ca^{2+} selective channels located at the plasma membrane (PM) of excitable cells, such as muscle cells and neurons, and activated in response to PM depolarization. In T cells Ca_V channels might be activated by voltage-dependent or voltage-independent means

Inositol-1,4,5-trisphosphate receptors

Ca^{2+} release channels present in the endoplasmic reticulum (ER) membrane that release Ca^{2+} from the ER lumen to the cytosol in response to allosteric binding of Ca^{2+} and inositol-1,4,5-trisphosphate

Ryanodine receptors

Ca^{2+} release channels present in the endoplasmic reticulum (ER) membrane that mediate release of Ca^{2+} from the ER lumen to the cytosol on activation by Ca^{2+} , nicotinic acid adenine dinucleotide phosphate (NAADP) and cyclic ADP-ribose (cADPR)

Mitochondrial Ca^{2+} uniporter

A mitochondrial Ca^{2+} selective channel complex located in the inner mitochondrial membrane that conducts Ca^{2+} from the cytosol to the mitochondrial matrix

Ion pumps

Transmembrane proteins that transport ions against the electrochemical gradient of a membrane and this function requires energy in the form of ATP hydrolysis. Examples include: sarcoplasmic/endoplasmic reticulum Ca^{2+} ATPase (SERCA), which pumps Ca^{2+} from the cytosol into the ER; and plasma membrane Ca^{2+} ATPase (PMCA), which pumps Ca^{2+} from the cytosol to the extracellular space

Two pore channels

Ion channels located in the membrane of endolysosomes and proposed to release Ca^{2+} and Na^+ from endolysosomes to the cytosol and are activated by NAADP and by phosphoinositide species localized in the endolysosomal membrane, such as phosphatidylinositol-3,5-bisphosphate

Transient receptor potential mucolipin (TRPML) channels

Non-selective cation channels located on the surface of endolysosomes that release Ca^{2+} and Na^+ from these organelles into the cytosol. TRPML channels are activated by phosphoinositide species localized in the endolysosomal membrane such as phosphatidylinositol-3,5-bisphosphate

Store-operated Ca^{2+} entry

The most ubiquitous Ca^{2+} influx pathway in non-excitable cells, which is activated when endoplasmic reticulum Ca^{2+} stores are depleted. It is mediated by plasma membrane ORAI Ca^{2+} channels activated by direct binding of stromal interaction molecule (STIM) proteins

Ca^{2+} release-activated Ca^{2+} (CRAC)

The biophysical manifestation of store-operated Ca^{2+} entry and ORAI channels measured by whole-cell patch clamp electrophysiology. CRAC currents are highly Ca^{2+} -selective

Nuclear factor of activated T cells (NFAT)

An important family of transcription factors that are Ca^{2+} -activated. Ca^{2+} -calmodulin activates the phosphatase calcineurin, which then dephosphorylates NFAT causing its import into the nucleus to mediate gene transcription of many cytokines, transcription factors and metabolic genes

Immune synapse

The nanoscale interface of interaction between a lymphocyte and an antigen presenting cell

Patch clamp electrophysiology

A laboratory technique used to measure ionic currents through specific channels from single living cells or from a patch of cell membrane. Under the voltage clamp configuration, controlled (clamped) voltage values are applied to the cell membrane by the experimenter and the resulting currents are measured

References

1. Vig M. & Kinet JP Calcium signaling in immune cells. *Nat Immunol* 10, 21–27 (2009). [PubMed: 19088738]
2. Cai X., Wang X., Patel S. & Clapham DE Insights into the early evolution of animal calcium signaling machinery: a unicellular point of view. *Cell Calcium* 57, 166–173 (2015). [PubMed: 25498309]
3. Hogan PG, Lewis RS & Rao A. Molecular basis of calcium signaling in lymphocytes: STIM and ORAI. *Annu Rev Immunol* 28, 491–533 (2010). [PubMed: 20307213]
4. Feske S., Wulff H. & Skolnik EY Ion channels in innate and adaptive immunity. *Annu Rev Immunol* 33, 291–353 (2015).
These are detailed review articles that provide excellent overview of different ions and ion channels and their role in controlling innate and adaptive immunity
[PubMed: 25861976]
5. Feske S., Skolnik EY & Prakriya M. Ion channels and transporters in lymphocyte function and immunity. *Nat Rev Immunol* 12, 532–547 (2012). [PubMed: 22699833]
6. Berridge MJ The Inositol Trisphosphate/Calcium Signaling Pathway in Health and Disease. *Physiol Rev* 96, 1261–1296 (2016).
An excellent and comprehensive review on IP3 signaling pathways in health and disease
[PubMed: 27512009]
7. De Stefani D., Rizzuto R. & Pozzan T. Enjoy the Trip: Calcium in Mitochondria Back and Forth. *Annu Rev Biochem* 85, 161–192 (2016).
An excellent review on mitochondrial Ca^{2+} signaling and its role in shaping cellular signaling and cell function.
[PubMed: 27145841]
8. Foskett JK, White C., Cheung KH & Mak DO Inositol trisphosphate receptor Ca^{2+} release channels. *Physiol Rev* 87, 593–658 (2007). [PubMed: 17429043]
9. Raffaello A., Mammucari C., Gherardi G. & Rizzuto R. Calcium at the Center of Cell Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes. *Trends Biochem Sci* 41, 1035–1049 (2016). [PubMed: 27692849]

10. Xiong J. & Zhu MX Regulation of lysosomal ion homeostasis by channels and transporters. *Sci China Life Sci* 59, 777–791 (2016).

This review provides an excellent introduction to ion channel networks in lysosomes and their role in lysosomal function

[PubMed: 27430889]

11. Cahalan MD & Chandy KG The functional network of ion channels in T lymphocytes. *Immunol Rev* 231, 59–87 (2009).

These are detailed review articles that provide excellent overview of different ions and ion channels and their role in controlling innate and adaptive immunity

[PubMed: 19754890]

12. Badou A. et al. Critical role for the beta regulatory subunits of Cav channels in T lymphocyte function. *Proc Natl Acad Sci U S A* 103, 15529–15534 (2006).
13. Omilusk K. et al. The Ca(v)1.4 calcium channel is a critical regulator of T cell receptor signaling and naive T cell homeostasis. *Immunity* 35, 349–360 (2011). [PubMed: 21835646]
14. Wang H. et al. Low-Voltage-Activated CaV3.1 Calcium Channels Shape T Helper Cell Cytokine Profiles. *Immunity* 44, 782–794 (2016).

This article provides the first patch clamp evidence of T-type Ca²⁺ channels in T cells and their role in cytokine production and immune function

[PubMed: 27037192]

15. Parker I. & Smith IF Recording single-channel activity of inositol trisphosphate receptors in intact cells with a microscope, not a patch clamp. *J Gen Physiol* 136, 119–127 (2010). [PubMed: 20660654]
16. Mikoshiba K. IP3 receptor/Ca²⁺ channel: from discovery to new signaling concepts. *J Neurochem* 102, 1426–1446 (2007). [PubMed: 17697045]
17. Guse AH & Wolf IM Ca(2+) microdomains, NAADP and type 1 ryanodine receptor in cell activation. *Biochim Biophys Acta* 1863, 1379–1384 (2016). [PubMed: 26804481]
18. Baughman JM et al. Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter. *Nature* 476, 341–345 (2011).

These two papers were first to identify the molecular identity of the mitochondrial Ca²⁺ uniporter (MCU) largely responsible for Ca²⁺ uptake by mitochondria.

[PubMed: 21685886]

19. De Stefani D., Raffaello A., Teardo E., Szabo I. & Rizzuto R. A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature* 476, 336–340 (2011).

These two papers were first to identify the molecular identity of the mitochondrial Ca²⁺ uniporter (MCU) largely responsible for Ca²⁺ uptake by mitochondria.

[PubMed: 21685888]

20. McCormack JG & Denton RM The effects of calcium ions and adenine nucleotides on the activity of pig heart 2-oxoglutarate dehydrogenase complex. *Biochem J* 180, 533–544 (1979). [PubMed: 39549]
21. Denton RM & McCormack JG Ca²⁺ as a second messenger within mitochondria of the heart and other tissues. *Annu Rev Physiol* 52, 451–466 (1990). [PubMed: 2184763]
22. McCormack JG, Halestrap AP & Denton RM Role of calcium ions in regulation of mammalian intramitochondrial metabolism. *Physiol Rev* 70, 391–425 (1990). [PubMed: 2157230]
23. Hansford RG Physiological role of mitochondrial Ca²⁺ transport. *J Bioenerg Biomembr* 26, 495–508 (1994). [PubMed: 7896765]
24. Montero M. et al. Chromaffin-cell stimulation triggers fast millimolar mitochondrial Ca²⁺ transients that modulate secretion. *Nat Cell Biol* 2, 57–61 (2000). [PubMed: 10655583]

25. Cali T., Brini M. & Carafoli E. Regulation of Cell Calcium and Role of Plasma Membrane Calcium ATPases. *Int Rev Cell Mol Biol* 332, 259–296 (2017). [PubMed: 28526135]
26. Stafford N., Wilson C., Oceandy D., Neyses L. & Cartwright EJ The Plasma Membrane Calcium ATPases and Their Role as Major New Players in Human Disease. *Physiol Rev* 97, 1089–1125 (2017). [PubMed: 28566538]
27. Chemaly ER, Troncone L. & Lebeche D. SERCA control of cell death and survival. *Cell Calcium* 69, 46–61 (2018). [PubMed: 28747251]
28. Wu KD, Lee WS, Wey J., Bungard D. & Lytton J. Localization and quantification of endoplasmic reticulum Ca(2+)-ATPase isoform transcripts. *Am J Physiol* 269, C775–784 (1995). [PubMed: 7573409]
29. Chen J. et al. CD22 attenuates calcium signaling by potentiating plasma membrane calcium-ATPase activity. *Nat Immunol* 5, 651–657 (2004). [PubMed: 15133509]
30. Palty R. et al. NCLX is an essential component of mitochondrial Na⁺/Ca²⁺ exchange. *Proc Natl Acad Sci U S A* 107, 436–441 (2010).
This paper is the first to identify the molecular identity of the Na⁺/Ca²⁺ exchanger in mitochondria (NCLX), responsible for Ca²⁺ extrusion from mitochondria.
[PubMed: 20018762]
31. Ben-Kasus Nissim T. et al. Mitochondria control store-operated Ca(2+) entry through Na(+) and redox signals. *EMBO J* 36, 797–815 (2017). [PubMed: 28219928]
32. Sekler I. Standing of giants shoulders the story of the mitochondrial Na(+)Ca(2+) exchanger. *Biochem Biophys Res Commun* 460, 50–52 (2015). [PubMed: 25998733]
33. Xu H. & Ren D. Lysosomal physiology. *Annu Rev Physiol* 77, 57–80 (2015). [PubMed: 25668017]
34. Paroutis P., Touret N. & Grinstein S. The pH of the secretory pathway: measurement, determinants, and regulation. *Physiology (Bethesda)* 19, 207–215 (2004). [PubMed: 15304635]
35. Christensen KA, Myers JT & Swanson JA pH-dependent regulation of lysosomal calcium in macrophages. *J Cell Sci* 115, 599–607 (2002). [PubMed: 11861766]
36. Morgan AJ, Davis LC, Ruas M. & Galione A. TPC: the NAADP discovery channel? *Biochem Soc Trans* 43, 384–389 (2015). [PubMed: 26009180]
37. Morgan AJ, Platt FM, Lloyd-Evans E. & Galione A. Molecular mechanisms of endolysosomal Ca²⁺ signalling in health and disease. *Biochem J* 439, 349–374 (2011). [PubMed: 21992097]
38. Courtney AH, Lo WL & Weiss A. TCR Signaling: Mechanisms of Initiation and Propagation. *Trends Biochem Sci* 43, 108–123 (2018). [PubMed: 29269020]
39. Putney JW Jr. A model for receptor-regulated calcium entry. *Cell Calcium* 7, 1–12 (1986). [PubMed: 2420465]
40. Putney JW Jr. Capacitative calcium entry revisited. *Cell Calcium* 11, 611–624 (1990). [PubMed: 1965707]
41. Takemura H., Hughes AR, Thastrup O. & Putney JW Jr. Activation of calcium entry by the tumor promoter thapsigargin in parotid acinar cells. Evidence that an intracellular calcium pool and not an inositol phosphate regulates calcium fluxes at the plasma membrane. *J Biol Chem* 264, 12266–12271 (1989).
42. Hoth M. & Penner R. Depletion of intracellular calcium stores activates a calcium current in mast cells. *Nature* 355, 353–356 (1992). [PubMed: 1309940]
43. Prakriya M. & Lewis RS Store-Operated Calcium Channels. *Physiol Rev* 95, 1383–1436 (2015).
An exhaustive review that provides an outstanding overview of store-operated Ca²⁺ channels, including their biophysical properties, activation, regulation, and cellular function.
[PubMed: 26400989]
44. Trebak M. & Putney JW Jr. ORAI Calcium Channels. *Physiology (Bethesda)* 32, 332–342 (2017).
This review provides a brief historical overview of store-operated Ca²⁺ channels and a summary of the various native Ca²⁺ channels encoded by ORAI proteins.
[PubMed: 28615316]

45. Liou J. et al. STIM is a Ca²⁺ sensor essential for Ca²⁺-store-depletion-triggered Ca²⁺ influx. *Curr Biol* 15, 1235–1241 (2005). [PubMed: 16005298]
46. Roos J. et al. STIM1, an essential and conserved component of store-operated Ca²⁺ channel function. *J Cell Biol* 169, 435–445 (2005). [PubMed: 15866891]
47. Brandman O., Liou J., Park WS & Meyer T. STIM2 is a feedback regulator that stabilizes basal cytosolic and endoplasmic reticulum Ca²⁺ levels. *Cell* 131, 1327–1339 (2007). [PubMed: 18160041]
48. Feske S. et al. A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* 441, 179–185 (2006). [PubMed: 16582901]
49. Vig M. et al. CRACM1 is a plasma membrane protein essential for store-operated Ca²⁺ entry. *Science* 312, 1220–1223 (2006). [PubMed: 16645049]
50. Zhang SL et al. Genome-wide RNAi screen of Ca(2+) influx identifies genes that regulate Ca(2+) release-activated Ca(2+) channel activity. *Proc Natl Acad Sci U S A* 103, 9357–9362 (2006). [PubMed: 16751269]
51. Mercer JC et al. Large store-operated calcium selective currents due to co-expression of Orai1 or Orai2 with the intracellular calcium sensor, Stim1. *J Biol Chem* 281, 24979–24990 (2006).
52. Peinelt C. et al. Amplification of CRAC current by STIM1 and CRACM1 (Orai1). *Nat Cell Biol* 8, 771–773 (2006). [PubMed: 16733527]
53. Prakriya M. et al. Orai1 is an essential pore subunit of the CRAC channel. *Nature* 443, 230–233 (2006). [PubMed: 16921383]
54. Soboloff J. et al. Orai1 and STIM reconstitute store-operated calcium channel function. *J Biol Chem* 281, 20661–20665 (2006).
55. Vig M. et al. CRACM1 multimers form the ion-selective pore of the CRAC channel. *Curr Biol* 16, 2073–2079 (2006). [PubMed: 16978865]
56. Lioudyno MI et al. Orai1 and STIM1 move to the immunological synapse and are up-regulated during T cell activation. *Proc Natl Acad Sci U S A* 105, 2011–2016 (2008). [PubMed: 18250319]
57. Feske S. et al. Severe combined immunodeficiency due to defective binding of the nuclear factor of activated T cells in T lymphocytes of two male siblings. *Eur J Immunol* 26, 2119–2126 (1996). [PubMed: 8814256]
58. Fuchs S. et al. Antiviral and regulatory T cell immunity in a patient with stromal interaction molecule 1 deficiency. *J Immunol* 188, 1523–1533 (2012). [PubMed: 22190180]
59. Le Deist F. et al. A primary T-cell immunodeficiency associated with defective transmembrane calcium influx. *Blood* 85, 1053–1062 (1995). [PubMed: 7531512]
60. Picard C. et al. STIM1 mutation associated with a syndrome of immunodeficiency and autoimmunity. *N Engl J Med* 360, 1971–1980 (2009). [PubMed: 19420366]
61. Vaeth M. et al. Store-Operated Ca(2+) Entry Controls Clonal Expansion of T Cells through Metabolic Reprogramming. *Immunity* 47, 664–679 e666 (2017).
An outstanding paper providing evidence for the role of store-operated Ca²⁺ entry in T cell metabolism and immunity through control of glycolysis and oxidative phosphorylation genes. [PubMed: 29030115]
62. Dolmetsch RE, Lewis RS, Goodnow CC & Healy JI Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration. *Nature* 386, 855–858 (1997). [PubMed: 9126747]
63. Partiseti M. et al. The calcium current activated by T cell receptor and store depletion in human lymphocytes is absent in a primary immunodeficiency. *J Biol Chem* 269, 32327–32335 (1994).
64. Lacruz RS & Feske S. Diseases caused by mutations in ORAI1 and STIM1. *Ann N Y Acad Sci* 1356, 45–79 (2015). [PubMed: 26469693]
65. Vaeth M. & Feske S. Ion channelopathies of the immune system. *Curr Opin Immunol* 52, 39–50 (2018). [PubMed: 29635109]
66. Oh-Hora M. et al. Agonist-selected T cell development requires strong T cell receptor signaling and store-operated calcium entry. *Immunity* 38, 881–895 (2013). [PubMed: 23499491]

67. Vig M. et al. Defective mast cell effector functions in mice lacking the CRACM1 pore subunit of store-operated calcium release-activated calcium channels. *Nat Immunol* 9, 89–96 (2008). [PubMed: 18059270]
68. Oh-Hora M. et al. Dual functions for the endoplasmic reticulum calcium sensors STIM1 and STIM2 in T cell activation and tolerance. *Nat Immunol* 9, 432–443 (2008). [PubMed: 18327260]
69. Vaeth M. et al. ORAI2 modulates store-operated calcium entry and T cell-mediated immunity. *Nat Commun* 8, 14714 (2017).
This paper provides the first evidence that ORAI2 channels act as negative modulators of store-operated Ca²⁺ entry and T cell immunity.
70. Kaufmann U. et al. Selective ORAI1 Inhibition Ameliorates Autoimmune Central Nervous System Inflammation by Suppressing Effector but Not Regulatory T Cell Function. *J Immunol* 196, 573–585 (2016). [PubMed: 26673135]
71. Kim KD et al. Calcium signaling via Orai1 is essential for induction of the nuclear orphan receptor pathway to drive Th17 differentiation. *J Immunol* 192, 110–122 (2014). [PubMed: 24307733]
72. Wang X. et al. Distinct Orai-coupling domains in STIM1 and STIM2 define the Orai-activating site. *Nat Commun* 5, 3183 (2014). [PubMed: 24492416]
73. Vaeth M. et al. Store-Operated Ca(2+) Entry in Follicular T Cells Controls Humoral Immune Responses and Autoimmunity. *Immunity* 44, 1350–1364 (2016). [PubMed: 27261277]
74. Kim KD et al. ORAI1 deficiency impairs activated T cell death and enhances T cell survival. *J Immunol* 187, 3620–3630 (2011). [PubMed: 21873530]
75. Srikanth S., Woo JS, Sun Z. & Gwack Y. Immunological Disorders: Regulation of Ca(2+) Signaling in T Lymphocytes. *Adv Exp Med Biol* 993, 397–424 (2017). [PubMed: 28900926]
76. Robbs BK, Cruz AL, Werneck MB, Mognol GP & Viola JP Dual roles for NFAT transcription factor genes as oncogenes and tumor suppressors. *Mol Cell Biol* 28, 7168–7181 (2008). [PubMed: 18809576]
77. Srinivasan M. & Frauwirth KA Reciprocal NFAT1 and NFAT2 nuclear localization in CD8+ anergic T cells is regulated by suboptimal calcium signaling. *J Immunol* 179, 3734–3741 (2007). [PubMed: 17785810]
78. Desvignes L. et al. STIM1 controls T cell-mediated immune regulation and inflammation in chronic infection. *J Clin Invest* 125, 2347–2362 (2015). [PubMed: 25938788]
79. Tsvilovskyy V. et al. Deletion of Orai2 augments endogenous CRAC currents and degranulation in mast cells leading to enhanced anaphylaxis. *Cell Calcium* 71, 24–33 (2018). [PubMed: 29604961]
80. Shuttleworth TJ Selective activation of distinct Orai channels by STIM1. *Cell Calcium* 63, 40–42 (2017). [PubMed: 27847114]
81. Desai PN et al. Multiple types of calcium channels arising from alternative translation initiation of the Orai1 message. *Sci Signal* 8, ra74 (2015).
82. Gonzalez-Cobos JC et al. Store-independent Orai1/3 channels activated by intracrine leukotriene C4: role in neointimal hyperplasia. *Circ Res* 112, 1013–1025 (2013). [PubMed: 23349245]
83. Zhang X. et al. Mechanisms of STIM1 activation of store-independent leukotriene C4-regulated Ca²⁺ channels. *Mol Cell Biol* 33, 3715–3723 (2013). [PubMed: 23878392]
84. Zhang X. et al. Complex role of STIM1 in the activation of store-independent Orai1/3 channels. *J Gen Physiol* 143, 345–359 (2014). [PubMed: 24567509]
85. Bogeski I. et al. Differential redox regulation of ORAI ion channels: a mechanism to tune cellular calcium signaling. *Sci Signal* 3, ra24 (2010).
86. Liu S., Kiyoi T., Takemasa E. & Maeyama K. Systemic lentivirus-mediated delivery of short hairpin RNA targeting calcium release-activated calcium channel 3 as gene therapy for collagen-induced arthritis. *J Immunol* 194, 76–83 (2015). [PubMed: 25422506]
87. Wenning AS et al. TRP expression pattern and the functional importance of TRPC3 in primary human T-cells. *Biochim Biophys Acta* 1813, 412–423 (2011). [PubMed: 21215279]
88. Philipp S. et al. TRPC3 mediates T-cell receptor-dependent calcium entry in human T-lymphocytes. *J Biol Chem* 278, 26629–26638 (2003).

89. Beck A., Kolisek M., Bagley LA, Fleig A. & Penner R. Nicotinic acid adenine dinucleotide phosphate and cyclic ADP-ribose regulate TRPM2 channels in T lymphocytes. *FASEB J* 20, 962–964 (2006). [PubMed: 16585058]
90. Guse AH et al. Regulation of calcium signalling in T lymphocytes by the second messenger cyclic ADP-ribose. *Nature* 398, 70–73 (1999). [PubMed: 10078531]
91. Melzer N., Hicking G., Gobel K. & Wiendl H. TRPM2 cation channels modulate T cell effector functions and contribute to autoimmune CNS inflammation. *PLoS One* 7, e47617 (2012).
92. Launay P. et al. TRPM4 is a Ca²⁺-activated nonselective cation channel mediating cell membrane depolarization. *Cell* 109, 397–407 (2002). [PubMed: 12015988]
93. Launay P. et al. TRPM4 regulates calcium oscillations after T cell activation. *Science* 306, 1374–1377 (2004). [PubMed: 15550671]
94. Weber KS, Hildner K., Murphy KM & Allen PM Trpm4 differentially regulates Th1 and Th2 function by altering calcium signaling and NFAT localization. *J Immunol* 185, 2836–2846 (2010). [PubMed: 20656926]
95. Zhang X. et al. Unequal death in T helper cell (Th)1 and Th2 effectors: Th1, but not Th2, effectors undergo rapid Fas/FasL-mediated apoptosis. *J Exp Med* 185, 1837–1849 (1997). [PubMed: 9151709]
96. Chandy KG et al. K⁺ channels as targets for specific immunomodulation. *Trends Pharmacol Sci* 25, 280–289 (2004). [PubMed: 15120495]
97. Jin J. et al. Deletion of Trpm7 disrupts embryonic development and thymopoiesis without altering Mg²⁺ homeostasis. *Science* 322, 756–760 (2008). [PubMed: 18974357]
98. Faouzi M., Kilch T., Horgen FD, Fleig A. & Penner R. The TRPM7 channel kinase regulates store-operated calcium entry. *J Physiol* 595, 3165–3180 (2017). [PubMed: 28130783]
99. Beesetty P. et al. Inactivation of TRPM7 kinase in mice results in enlarged spleens, reduced T-cell proliferation and diminished store-operated calcium entry. *Sci Rep* 8, 3023 (2018). [PubMed: 29445164]
100. Romagnani A. et al. TRPM7 kinase activity is essential for T cell colonization and alloreactivity in the gut. *Nat Commun* 8, 1917 (2017). [PubMed: 29203869]
101. Desai BN et al. Cleavage of TRPM7 releases the kinase domain from the ion channel and regulates its participation in Fas-induced apoptosis. *Dev Cell* 22, 1149–1162 (2012). [PubMed: 22698280]
102. Bertin S. et al. The ion channel TRPV1 regulates the activation and proinflammatory properties of CD4(+) T cells. *Nat Immunol* 15, 1055–1063 (2014). [PubMed: 25282159]
103. Samivel R. et al. The role of TRPV1 in the CD4+ T cell-mediated inflammatory response of allergic rhinitis. *Oncotarget* 7, 148–160 (2016). [PubMed: 26700618]
104. Bertin S. et al. The TRPA1 ion channel is expressed in CD4+ T cells and restrains T-cell-mediated colitis through inhibition of TRPV1. *Gut* 66, 1584–1596 (2017). [PubMed: 27325418]
105. Di Virgilio F., Sarti AC & Grassi F. Modulation of innate and adaptive immunity by P2X ion channels. *Curr Opin Immunol* 52, 51–59 (2018). [PubMed: 29631184]
106. Di Virgilio F., Dal Ben D., Sarti AC, Giuliani AL & Falzoni S. The P2X7 Receptor in Infection and Inflammation. *Immunity* 47, 15–31 (2017).
- Provides an excellent overview on the role of P2XR7 in regulating innate and adaptive immunity. [PubMed: 28723547]
107. Coutinho-Silva R., Knight GE & Burnstock G. Impairment of the splenic immune system in P2X(2)/P2X(3) knockout mice. *Immunobiology* 209, 661–668 (2005). [PubMed: 15804044]
108. Cockayne DA et al. P2X2 knockout mice and P2X2/P2X3 double knockout mice reveal a role for the P2X2 receptor subunit in mediating multiple sensory effects of ATP. *J Physiol* 567, 621–639 (2005). [PubMed: 15961431]
109. Abramowski P., Ogradowczyk C., Martin R. & Pongs O. A truncation variant of the cation channel P2RX5 is upregulated during T cell activation. *PLoS One* 9, e104692 (2014).

110. Woehrle T. et al. Pannexin-1 hemichannel-mediated ATP release together with P2X1 and P2X4 receptors regulate T-cell activation at the immune synapse. *Blood* 116, 3475–3484 (2010). [PubMed: 20660288]
111. Ledderose C. et al. Mitochondrial Dysfunction, Depleted Purinergic Signaling, and Defective T Cell Vigilance and Immune Defense. *J Infect Dis* 213, 456–464 (2016). [PubMed: 26150546]
112. Wang CM, Ploia C., Anselmi F., Sarukhan A. & Viola A. Adenosine triphosphate acts as a paracrine signaling molecule to reduce the motility of T cells. *EMBO J* 33, 1354–1364 (2014). [PubMed: 24843045]
113. Manohar M. et al. ATP release and autocrine signaling through P2X4 receptors regulate gammadelta T cell activation. *J Leukoc Biol* 92, 787–794 (2012). [PubMed: 22753954]
114. Frascoli M., Marcandalli J., Schenk U. & Grassi F. Purinergic P2X7 receptor drives T cell lineage choice and shapes peripheral gammadelta cells. *J Immunol* 189, 174–180 (2012). [PubMed: 22649196]
115. Schenk U. et al. ATP inhibits the generation and function of regulatory T cells through the activation of purinergic P2X receptors. *Sci Signal* 4, ra12 (2011).
116. Vergani A. et al. Effect of the purinergic inhibitor oxidized ATP in a model of islet allograft rejection. *Diabetes* 62, 1665–1675 (2013). [PubMed: 23315496]
117. Vergani A. et al. Long-term heart transplant survival by targeting the ionotropic purinergic receptor P2X7. *Circulation* 127, 463–475 (2013). [PubMed: 23250993]
118. Koo TY et al. The P2X7 receptor antagonist, oxidized adenosine triphosphate, ameliorates renal ischemia-reperfusion injury by expansion of regulatory T cells. *Kidney Int* 92, 415–431 (2017). [PubMed: 28396117]
119. Sharp AJ et al. P2 \times 7 deficiency suppresses development of experimental autoimmune encephalomyelitis. *J Neuroinflammation* 5, 33 (2008). [PubMed: 18691411]
120. Takeda A. et al. Crucial role of P2X7 receptor for effector T cell activation in experimental autoimmune uveitis. *Jpn J Ophthalmol* 62, 398–406 (2018). [PubMed: 29572578]
121. Salles EM et al. P2X7 receptor drives Th1 cell differentiation and controls the follicular helper T cell population to protect against *Plasmodium chabaudi* malaria. *PLoS Pathog* 13, e1006595 (2017).
122. Hofman P. et al. Genetic and pharmacological inactivation of the purinergic P2RX7 receptor dampens inflammation but increases tumor incidence in a mouse model of colitis-associated cancer. *Cancer Res* 75, 835–845 (2015). [PubMed: 25564520]
123. Borges da Silva H. et al. The purinergic receptor P2RX7 directs metabolic fitness of long-lived memory CD8(+) T cells. *Nature* 559, 264–268 (2018). [PubMed: 29973721]
124. Chen L. & Brosnan CF Exacerbation of experimental autoimmune encephalomyelitis in P2X7R $-/-$ mice: evidence for loss of apoptotic activity in lymphocytes. *J Immunol* 176, 3115–3126 (2006). [PubMed: 16493071]
125. Taylor SR et al. Lymphocytes from P2X7-deficient mice exhibit enhanced P2X7 responses. *J Leukoc Biol* 85, 978–986 (2009). [PubMed: 19276178]
126. Shinohara Y. & Tsukimoto M. Guanine and inosine nucleotides/nucleosides suppress murine T cell activation. *Biochem Biophys Res Commun* 498, 764–768 (2018). [PubMed: 29524424]
127. Lecciso M. et al. ATP Release from Chemotherapy-Treated Dying Leukemia Cells Elicits an Immune Suppressive Effect by Increasing Regulatory T Cells and Tolerogenic Dendritic Cells. *Front Immunol* 8, 1918 (2017). [PubMed: 29312358]
128. Badou A., Jha MK, Matza D. & Flavell RA Emerging roles of L-type voltage-gated and other calcium channels in T lymphocytes. *Front Immunol* 4, 243 (2013). [PubMed: 24009608]
129. Robert V., Triffaux E., Savignac M. & Pelletier L. Singularities of calcium signaling in effector T-lymphocytes. *Biochim Biophys Acta* 1833, 1595–1602 (2013). [PubMed: 23266355]
130. Navedo MF & Santana LF CaV1.2 sparklets in heart and vascular smooth muscle. *J Mol Cell Cardiol* 58, 67–76 (2013). [PubMed: 23220157]
131. Navedo MF, Amberg GC, Nieves M., Molkentin JD & Santana LF Mechanisms underlying heterogeneous Ca $^{2+}$ sparklet activity in arterial smooth muscle. *J Gen Physiol* 127, 611–622 (2006). [PubMed: 16702354]

132. Nieves-Cintrón M., Amberg GC, Navedo MF, Molkentin JD & Santana LF The control of Ca²⁺ influx and NFATc3 signaling in arterial smooth muscle during hypertension. *Proc Natl Acad Sci U S A* 105, 15623–15628 (2008).
133. Dixon RE et al. Graded Ca²⁺(+)/calmodulin-dependent coupling of voltage-gated Ca_v1.2 channels. *Elife* 4 (2015).
134. Dixon RE, Yuan C., Cheng EP, Navedo MF & Santana LF Ca²⁺ signaling amplification by oligomerization of L-type Cav1.2 channels. *Proc Natl Acad Sci U S A* 109, 1749–1754 (2012). [PubMed: 22307641]
135. Navedo MF, Amberg GC, Votaw VS & Santana LF Constitutively active L-type Ca²⁺ channels. *Proc Natl Acad Sci U S A* 102, 11112–11117 (2005).
136. Bannister RA & Beam KG Ca_v(V)1.1: The atypical prototypical voltage-gated Ca²⁺(+) channel. *Biochim Biophys Acta* 1828, 1587–1597 (2013). [PubMed: 22982493]
137. Matza D. et al. T Cell Receptor Mediated Calcium Entry Requires Alternatively Spliced Cav1.1 Channels. *PLoS One* 11, e0147379 (2016).
138. Jha MK et al. Defective survival of naive CD8⁺ T lymphocytes in the absence of the beta3 regulatory subunit of voltage-gated calcium channels. *Nat Immunol* 10, 1275–1282 (2009). [PubMed: 19838200]
139. Jha A. et al. Essential roles for Cavbeta2 and Cav1 channels in thymocyte development and T cell homeostasis. *Sci Signal* 8, ra103 (2015).
140. Chandrasekhar R., Alzayady KJ, Wagner LE 2nd & Yule DI Unique Regulatory Properties of Heterotetrameric Inositol 1,4,5-Trisphosphate Receptors Revealed by Studying Concatenated Receptor Constructs. *J Biol Chem* 291, 4846–4860 (2016). [PubMed: 26755721]
141. Jayaraman T., Ondriasova E., Ondrias K., Harnick DJ & Marks AR The inositol 1,4,5-trisphosphate receptor is essential for T-cell receptor signaling. *Proc Natl Acad Sci U S A* 92, 6007–6011 (1995). [PubMed: 7597070]
142. Jayaraman T. & Marks AR Calcineurin is downstream of the inositol 1,4,5-trisphosphate receptor in the apoptotic and cell growth pathways. *J Biol Chem* 275, 6417–6420 (2000). [PubMed: 10692444]
143. Ouyang K. et al. Loss of IP3R-dependent Ca²⁺ signalling in thymocytes leads to aberrant development and acute lymphoblastic leukemia. *Nat Commun* 5, 4814 (2014). [PubMed: 25215520]
144. Wolf IMA & Guse AH Ca²⁺ Microdomains in T-Lymphocytes. *Front Oncol* 7, 73 (2017). [PubMed: 28512623]
145. Wolf IM et al. Frontrunners of T cell activation: Initial, localized Ca²⁺ signals mediated by NAADP and the type 1 ryanodine receptor. *Sci Signal* 8, ra102 (2015).
146. Kunerth S. et al. Amplification and propagation of pacemaker Ca²⁺ signals by cyclic ADP-ribose and the type 3 ryanodine receptor in T cells. *J Cell Sci* 117, 2141–2149 (2004). [PubMed: 15054112]
147. Dammermann W. et al. NAADP-mediated Ca²⁺ signaling via type 1 ryanodine receptor in T cells revealed by a synthetic NAADP antagonist. *Proc Natl Acad Sci U S A* 106, 10678–10683 (2009).
148. Dadsetan S., Zakharova L., Molinski TF & Fomina AF Store-operated Ca²⁺ influx causes Ca²⁺ release from the intracellular Ca²⁺ channels that is required for T cell activation. *J Biol Chem* 283, 12512–12519 (2008).
149. Thakur P., Dadsetan S. & Fomina AF Bidirectional coupling between ryanodine receptors and Ca²⁺ release-activated Ca²⁺ (CRAC) channel machinery sustains store-operated Ca²⁺ entry in human T lymphocytes. *J Biol Chem* 287, 37233–37244 (2012).
150. Takeshima H. et al. Generation and characterization of mutant mice lacking ryanodine receptor type 3. *J Biol Chem* 271, 19649–19652 (1996).
151. Davis LC, Platt FM & Galione A. Preferential Coupling of the NAADP Pathway to Exocytosis in T-Cells. *Messenger (Los Angel)* 4, 53–66 (2015). [PubMed: 27330870]
152. Steen M., Kirchberger T. & Guse AH NAADP mobilizes calcium from the endoplasmic reticular Ca²⁺ store in T-lymphocytes. *J Biol Chem* 282, 18864–18871 (2007).

153. Gerasimenko JV, Sherwood M., Tepikin AV, Petersen OH & Gerasimenko OV NAADP, cADPR and IP3 all release Ca²⁺ from the endoplasmic reticulum and an acidic store in the secretory granule area. *J Cell Sci* 119, 226–238 (2006). [PubMed: 16410548]
154. Mehta MM, Weinberg SE & Chandel NS Mitochondrial control of immunity: beyond ATP. *Nat Rev Immunol* 17, 608–620 (2017). [PubMed: 28669986]
155. Bagur R. & Hajnoczky G. Intracellular Ca(2+) Sensing: Its Role in Calcium Homeostasis and Signaling. *Mol Cell* 66, 780–788 (2017). [PubMed: 28622523]
156. Csordas G., Weaver D. & Hajnoczky G. Endoplasmic Reticular-Mitochondrial Contactology: Structure and Signaling Functions. *Trends Cell Biol* (2018).
An excellent review on the intimate crosstalk between ER and mitochondrial Ca²⁺ signaling networks, their structure, organization, protein composition and novel methods to study them.
157. Santo-Domingo J. & Demaurex N. Calcium uptake mechanisms of mitochondria. *Biochim Biophys Acta* 1797, 907–912 (2010). [PubMed: 20079335]
158. Pathak T. & Trebak M. Mitochondrial Ca(2+) signaling. *Pharmacol Ther* (2018).
159. Quintana A. & Hoth M. Mitochondrial dynamics and their impact on T cell function. *Cell Calcium* 52, 57–63 (2012). [PubMed: 22425631]
160. Junker C. & Hoth M. Immune synapses: mitochondrial morphology matters. *EMBO J* 30, 1187–1189 (2011). [PubMed: 21468096]
161. Hoth M., Fanger CM & Lewis RS Mitochondrial regulation of store-operated calcium signaling in T lymphocytes. *J Cell Biol* 137, 633–648 (1997). [PubMed: 9151670]
162. Hoth M., Button DC & Lewis RS Mitochondrial control of calcium-channel gating: a mechanism for sustained signaling and transcriptional activation in T lymphocytes. *Proc Natl Acad Sci U S A* 97, 10607–10612 (2000).
163. Quintana A. et al. T cell activation requires mitochondrial translocation to the immunological synapse. *Proc Natl Acad Sci U S A* 104, 14418–14423 (2007).
164. Quintana A. et al. Calcium microdomains at the immunological synapse: how ORAI channels, mitochondria and calcium pumps generate local calcium signals for efficient T-cell activation. *EMBO J* 30, 3895–3912 (2011). [PubMed: 21847095]
165. Ritchie MF, Samakai E. & Soboloff J. STIM1 is required for attenuation of PMCA-mediated Ca²⁺ clearance during T-cell activation. *EMBO J* 31, 1123–1133 (2012). [PubMed: 22246182]
166. Verkhatsky A., Trebak M., Perocchi F., Khananshvil D. & Sekler I. Crosslink between calcium and sodium signalling. *Exp Physiol* 103, 157–169 (2018). [PubMed: 29210126]
167. Jiang D., Zhao L. & Clapham DE Genome-wide RNAi screen identifies Letm1 as a mitochondrial Ca²⁺/H⁺ antiporter. *Science* 326, 144–147 (2009). [PubMed: 19797662]
168. Kim B., Takeuchi A., Koga O., Hikida M. & Matsuoka S. Pivotal role of mitochondrial Na(+) (–)Ca(2)(+) exchange in antigen receptor mediated Ca(2)(+) signalling in DT40 and A20 B lymphocytes. *J Physiol* 590, 459–474 (2012). [PubMed: 22155933]
169. Finetti F., Onnis A. & Baldari CT Regulation of vesicular traffic at the T cell immune synapse: lessons from the primary cilium. *Traffic* 16, 241–249 (2015). [PubMed: 25393976]
170. Voskoboinik I., Whisstock JC & Trapani JA Perforin and granzymes: function, dysfunction and human pathology. *Nat Rev Immunol* 15, 388–400 (2015). [PubMed: 25998963]
171. Benzing C., Rossy J. & Gaus K. Do signalling endosomes play a role in T cell activation? *FEBS J* 280, 5164–5176 (2013). [PubMed: 23834225]
172. Brailoiu E. et al. Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *J Cell Biol* 186, 201–209 (2009). [PubMed: 19620632]
173. Calcraft PJ et al. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* 459, 596–600 (2009). [PubMed: 19387438]
174. Zong X. et al. The two-pore channel TPCN2 mediates NAADP-dependent Ca(2+)-release from lysosomal stores. *Pflugers Arch* 458, 891–899 (2009). [PubMed: 19557428]
175. Galione A. A primer of NAADP-mediated Ca(2+) signalling: From sea urchin eggs to mammalian cells. *Cell Calcium* 58, 27–47 (2015). [PubMed: 25449298]
176. Wang X. et al. TPC proteins are phosphoinositide- activated sodium-selective ion channels in endosomes and lysosomes. *Cell* 151, 372–383 (2012). [PubMed: 23063126]

177. Cang C. et al. mTOR regulates lysosomal ATP-sensitive two-pore Na⁽⁺⁾ channels to adapt to metabolic state. *Cell* 152, 778–790 (2013). [PubMed: 23394946]
178. Li X. et al. Genetically encoded fluorescent probe to visualize intracellular phosphatidylinositol 3,5-bisphosphate localization and dynamics. *Proc Natl Acad Sci U S A* 110, 21165–21170 (2013).
179. Davis LC et al. NAADP activates two-pore channels on T cell cytolytic granules to stimulate exocytosis and killing. *Curr Biol* 22, 2331–2337 (2012). [PubMed: 23177477]
180. Cuajungco MP, Silva J., Habibi A. & Valadez JA The mucolipin-2 (TRPML2) ion channel: a tissue-specific protein crucial to normal cell function. *Pflugers Arch* 468, 177–192 (2016). [PubMed: 26336837]
181. Cheng X., Shen D., Samie M. & Xu H. Mucolipins: Intracellular TRPML1–3 channels. *FEBS Lett* 584, 2013–2021 (2010). [PubMed: 20074572]
182. Sun L., Hua Y., Vergarajauregui S., Diab HI & Puertollano R. Novel Role of TRPML2 in the Regulation of the Innate Immune Response. *J Immunol* 195, 4922–4932 (2015). [PubMed: 26432893]
183. Venkatachalam K., Hofmann T. & Montell C. Lysosomal localization of TRPML3 depends on TRPML2 and the mucopolipidosis-associated protein TRPML1. *J Biol Chem* 281, 17517–17527 (2006).
184. Dong XP et al. PI(3,5)P₂ controls membrane trafficking by direct activation of mucolipin Ca⁽²⁺⁾ release channels in the endolysosome. *Nat Commun* 1, 38 (2010). [PubMed: 20802798]
185. Samie M. et al. A TRP channel in the lysosome regulates large particle phagocytosis via focal exocytosis. *Dev Cell* 26, 511–524 (2013). [PubMed: 23993788]
186. LaPlante JM et al. Lysosomal exocytosis is impaired in mucopolipidosis type IV. *Mol Genet Metab* 89, 339–348 (2006). [PubMed: 16914343]
187. Zhang X., Li X. & Xu H. Phosphoinositide isoforms determine compartment-specific ion channel activity. *Proc Natl Acad Sci U S A* 109, 11384–11389 (2012).
188. Feng X. et al. Drosophila TRPML forms PI(3,5)P₂-activated cation channels in both endolysosomes and plasma membrane. *J Biol Chem* 289, 4262–4272 (2014). [PubMed: 24375408]
189. Zhong XZ et al. Inhibition of Transient Receptor Potential Channel Mucolipin-1 (TRPML1) by Lysosomal Adenosine Involved in Severe Combined Immunodeficiency Diseases. *J Biol Chem* 292, 3445–3455 (2017). [PubMed: 28087698]
190. Whitmore KV & Gaspar HB Adenosine Deaminase Deficiency - More Than Just an Immunodeficiency. *Front Immunol* 7, 314 (2016). [PubMed: 27579027]
191. Shuai J. & Parker I. Optical single-channel recording by imaging Ca²⁺ flux through individual ion channels: theoretical considerations and limits to resolution. *Cell Calcium* 37, 283–299 (2005). [PubMed: 15755490]
192. Potier M. & Trebak M. New developments in the signaling mechanisms of the store-operated calcium entry pathway. *Pflugers Arch* 457, 405–415 (2008). [PubMed: 18536932]
193. Gwack Y., Feske S., Srikanth S., Hogan PG & Rao A. Signalling to transcription: store-operated Ca²⁺ entry and NFAT activation in lymphocytes. *Cell Calcium* 42, 145–156 (2007). [PubMed: 17572487]
194. Fukushima M., Tomita T., Janoshazi A. & Putney JW Alternative translation initiation gives rise to two isoforms of Orai1 with distinct plasma membrane mobilities. *J Cell Sci* 125, 4354–4361 (2012). [PubMed: 22641696]
195. Ruhle B. & Trebak M. Emerging roles for native Orai Ca²⁺ channels in cardiovascular disease. *Curr Top Membr* 71, 209–235 (2013). [PubMed: 23890117]
196. Willoughby D. et al. Direct binding between Orai1 and AC8 mediates dynamic interplay between Ca²⁺ and cAMP signaling. *Sci Signal* 5, ra29 (2012). An article identifying the physical interaction between ORAI1 and Adenylyl Cyclase 8 as a determinant for Ca²⁺/cAMP signaling crosstalk.
197. Zhang W. et al. Leukotriene-C₄ synthase, a critical enzyme in the activation of store-independent Orai1/Orai3 channels, is required for neointimal hyperplasia. *J Biol Chem* 290, 5015–5027 (2015). [PubMed: 25540197]

198. Krishnamoorthy M. et al. The channel-kinase TRPM7 regulates antigen gathering and internalization in B cells. *Sci Signal* 11 (2018).
199. Mammucari C. et al. Mitochondrial calcium uptake in organ physiology: from molecular mechanism to animal models. *Pflugers Arch* (2018).
200. Ghosh D. et al. Calcium Channels in Vascular Smooth Muscle. *Adv Pharmacol* 78, 49–87 (2017). [PubMed: 28212803]
201. Guse AH & Diercks BP Integration of nicotinic acid adenine dinucleotide phosphate (NAADP)-dependent calcium signalling. *J Physiol* (2018).
202. Huang P. et al. P2X4 forms functional ATP-activated cation channels on lysosomal membranes regulated by luminal pH. *J Biol Chem* 289, 17658–17667 (2014).




Figure 1 | Calcium signalling in T cells.

Stimulation of the T cell receptor (TCR) by specific antigens leads to activation of phospholipase C γ 1 (PLC γ 1), the production of inositol-1,4,5-trisphosphate (IP₃) and Ca²⁺ release from endoplasmic reticulum (ER) Ca²⁺ stores via IP₃ receptor (IP₃R) channels. The decrease in Ca²⁺ levels within the ER lumen is sensed by low affinity EF-hands of stromal interaction molecule 1 (STIM1) and STIM2. STIM proteins gain an extended conformation to trap and activate ORAI1 proteins at the plasma membrane (PM) and induce store operated Ca²⁺ entry (SOCE)¹⁹². SOCE activates Ca²⁺-calmodulin and its target enzymes and transcription factors, most notably nuclear factor for activated T cells (NFAT) isoforms¹⁹³. Other PM channels are involved in mediating Ca²⁺ signals during T cell activation and include non-selective transient receptor potential (TRP) channels, purinergic ionotropic receptors (P2RX) and Ca_v channels. Ca²⁺ release by IP₃R (and Ca²⁺ entry through PM channels) is transferred into mitochondria through the mitochondrial Ca²⁺ uniporter (MCU) at highly specialized membrane contact sites termed mitochondria-associated membranes (MAMs), which effectively couple TCR ligation to enhanced bioenergetics and ATP production required for clonal expansion and secretion of cytokines. Sources of Ca²⁺ uptake into endolysosomes remain incompletely understood but contributions from the ER and PM are likely. Endolysosomal Ca²⁺ release controls vesicular fusion, trafficking and secretion of cargo and replenishment of exhausted signalling molecules at the PM of activated T cells¹⁶⁹. Cytoplasmic, ER and mitochondrial Ca²⁺ homeostasis are maintained by the actions of transporters and pumps, including the PM Ca²⁺ ATPase (PMCA), the sarcoplasmic/ER Ca²⁺ ATPase (SERCA) and the mitochondrial Na⁺/Ca²⁺/Li⁺ exchanger (NCLX). LAT, linker for activation of T cells; ZAP70, ζ -chain-associated protein kinase of 70 kDa.




Figure 2 | ORAI channels: major players in T cell activation.

ORAI proteins form highly Ca^{2+} selective homo-hexameric and hetero-hexameric channels in the plasma membrane (PM). On T cell receptor (TCR) ligation and subsequent inositol-1,4,5-trisphosphate (IP_3)-mediated endoplasmic reticulum (ER) Ca^{2+} store depletion, stromal interaction molecule (STIM) proteins move to ER–PM junctions and physically interact with ORAI channels, causing their activation. Mammals express three ORAI proteins (ORAI1–ORAI3) encoded by three separate genes and the major isoform mediating store operated Ca^{2+} entry (SOCE) in most cells including T cells is ORAI1. ORAI1 exists in two isoforms ORAI1 α (long) and ORAI1 β (short) due to alternative translation-initiation of *Orai1* mRNA^{81,194}. ORAI1 α and ORAI1 β do not seem to form hetero-hexamers. ORAI1 β has not been studied in T cells. ORAI2 can mediate residual SOCE in T cells from *Orai1*^{-/-} mice and was proposed to negatively modulate ORAI1 activity through formation of ORAI1–ORAI2 heteromeric associations. ORAI3, which unlike ORAI1, is resistant to inhibitory oxidation and is upregulated in effector T cells. The exclusively mammalian ORAI1 α and ORAI3 proteins were shown to form heteromeric channels that are activated independently of store depletion in other cell types¹⁹⁵, but their existence in T cells is unknown. Ca^{2+} entry through ORAI channels activates calcineurin and subsequent nuclear translocation of isoforms of nuclear factor for activated T cells (NFAT)¹⁹³. ORAI1 α physically associates through its N-terminus with the Ca^{2+} -activated adenylyl cyclase 8 (AC8)¹⁹⁶, thus connecting Ca^{2+} microdomains through ORAI1 α to cAMP production, activation of protein kinase A (PKA) and the transcription factor cAMP-responsive-element-binding protein (CREB). Ca^{2+} entry through heteromeric ORAI1 α –ORAI3 channels was proposed to regulate the AKT pathway¹⁹⁷, although the exact mechanisms remain unknown. AP-1, activator protein-1; CaM, calmodulin; CaMK-II, Ca^{2+} -calmodulin-dependent kinase II; MEF2, myocyte-specific enhancer factor 2; mTOR, mammalian target of rapamycin; NF- κ B, nuclear factor- κ B; PI3K, phosphoinositide 3-kinase.




Figure 3 | TRP channels: regulators of T cell Ca^{2+} signalling.

The transient receptor potential melastatin 4 (TRPM4) is a Ca^{2+} -activated Na^+ selective channel that is a potent regulator of ORAI-mediated Ca^{2+} entry in T cells. TRPM4 mediates its action through Na^+ entry, causing plasma membrane depolarization, which limits the driving force for Ca^{2+} and inhibits Ca^{2+} entry through ORAI channels. As such, during Ca^{2+} responses in T cells, Ca^{2+} activation by TRPM4 can shape the magnitude of sustained Ca^{2+} signals (plateaus) as well as the frequency and amplitude of oscillatory Ca^{2+} responses and therefore determine which downstream transcription factors and gene programmes are activated. An opposing role is played by Ca^{2+} - and voltage-dependent K^+ channels (KCa and Kv), which maintain hyperpolarized V_m to support Ca^{2+} entry. The Mg^{2+} - and Ca^{2+} -permeable channel-enzyme TRPM7 is one of few reported Ca^{2+} channels with a crucial role in T cell development and homeostasis and this function appears to be mediated by Ca^{2+} entry through the channel domain. The kinase domain of TRPM7 regulates ORAI1 signalling and coordinates antigen receptor signalling termination in lymphocytes, likely through phosphorylation of phospholipase $\text{C}\gamma$ (PLC γ) isoforms¹⁹⁸. TRPM2 channels are non-selective Ca^{2+} -conducting channels which are activated by hydrogen peroxide (H_2O_2) and through the cytosolic second messengers nicotinic acid adenine dinucleotide phosphate (NAADP) and cyclic ADP ribose (cADPR). TRPM2 has a more established role in neutrophils and was proposed to support T cell Ca^{2+} signalling in high oxidant inflammatory milieu.




Figure 4 | P2RX receptors: amplifiers of T cell receptor-mediated Ca^{2+} signalling through paracrine and autocrine ATP.

Purinergic ionotropic receptors (P2RX), including P2RX1, P2RX4 and P2RX7 are trimeric ionotropic non-selective Ca^{2+} -conducting channels that are activated by direct binding of extracellular ATP. Mitochondrial ATP production is enhanced in activated T cells by Ca^{2+} transfer from ORAI- and IP_3R -generated Ca^{2+} microdomains to the mitochondrial matrix via the mitochondrial Ca^{2+} uniporter (MCU)¹⁹⁹. ATP is then exported outside T cells by the pannexin 1 hemichannels and activates P2RX receptors to cause further Ca^{2+} entry and mitochondrial ATP production. In this regard, P2RX receptor signalling acts as a Ca^{2+} signalling enhancer for nuclear factor for activated T cells (NFAT)-mediated transcription. P2RX7 was shown to promote T helper 17 (TH17) cell differentiation and inflammation by promoting retinoic acid receptor-related orphan receptor γt (ROR γt) while inhibiting forkhead box P3 (FOXP3) transcription to suppress regulatory T (Treg) cells. However, Ca^{2+} signals through ORAI1 are required for Treg cell development (for both thymic Treg cells and induced Treg cells in peripheral lymphoid organs).




Figure 5 | CaV channels: modulators of T cell Ca²⁺ signalling.

Based on knockout of the $\beta 3$ and $\beta 4$ regulatory subunits of L-type Ca²⁺ channels which causes decreased expression of CaV1.2 and CaV1.1 at the plasma membrane and decreased T cell receptor (TCR)-activated Ca²⁺ entry and nuclear factor for activated T cells (NFAT) activity, CaV1.2 and CaV1.1 were proposed to play a role in TCR-mediated Ca²⁺ signalling. Although no CaV1.2 currents were reported in T cells, there is evidence of spontaneous CaV1.2 currents from channel clusters at hyperpolarized membrane potentials, as low as -90 mV, in smooth muscle cells^{135,200}. In this case, CaV1.2 activity can be sensitized by protein kinase A (PKA) and protein kinase C (PKC)-mediated phosphorylation; both enzymes can be activated downstream of store operated Ca²⁺ entry (SOCE). One intriguing hypothesis in T cells is the existence of a conformational coupling between CaV1.1 and ryanodine receptors (RYRs), similar to skeletal muscle. Voltage-activated CaV1.4 was proposed to support SOCE by mediating constitutive Ca²⁺ activity to help refill the ER Ca²⁺ stores and sustain CRAC channel activity, but these studies remain highly controversial. Voltage-activated whole-cell T-type Ca²⁺ currents mediated by CaV3.1 were recorded in T cells, generating window currents between -65 and -25mV, which are within T cell resting membrane potential. CaV3.1-mediated Ca²⁺ entry operates independently of SOCE and provides Ca²⁺ microdomains that synergize with SOCE to support NFAT activity. CaV3.1 activity is required for the expression of GM-CSF and RORγt.




Figure 6 | Organellar Ca^{2+} channels: initiators and master orchestrators of Ca^{2+} signalling microdomains during T cell activation.

The role of inositol-1,4,5-trisphosphate (IP_3) in initiating and sustaining T cell Ca^{2+} signalling is well established. Nicotinic acid adenine dinucleotide phosphate (NAADP) and cyclic ADP ribose (cADPR) were proposed to synergize with IP_3 during T cell Ca^{2+} signalling, but their role requires further clarification. For instance, the identity of the NAADP-producing enzyme downstream TCR ligation and the accessory molecule required for NAADP action on type 1 ryanodine receptors (RYR1) remain obscure. Nevertheless, according to this model, NAADP produced during the first few seconds of T cell receptor (TCR) stimulation was proposed to represent the crucial initial trigger for T cell Ca^{2+} signalling, preceding IP_3 -mediated ER Ca^{2+} release. NAADP would activate RYR1 and cause endoplasmic reticulum (ER) Ca^{2+} release, which would synergize with IP_3 to cause further Ca^{2+} release via IP_3 receptor (IP_3R). Subsequent production of cADPR and activation of RYR2 and RYR3 by cADPR would serve to sustain Ca^{2+} signalling for extended periods of time²⁰¹. Through a positive feedback loop, Ca^{2+} entry through ORAI channels might activate RYR isoforms to maintain store depletion and store operated Ca^{2+} entry (SOCE). Mitochondrial Ca^{2+} uptake through the mitochondrial Ca^{2+} uniporter (MCU) at the vicinity of ORAI channels maintains CRAC channel activity by relieving its Ca^{2+} -dependent inhibition (CDI). Mitochondria can also buffer Ca^{2+} near plasma membrane Ca^{2+} ATPase (PMCA) pumps, thus preventing PMCA modulation, inhibiting Ca^{2+} extrusion to the outside and maintaining cytosolic Ca^{2+} signalling during T cell activation. Ca^{2+} shuttling to the outside of mitochondria by the activity of the $\text{Na}^+/\text{Ca}^{2+}/\text{Li}^+$ exchanger (NCLX) provides Ca^{2+} microdomains to the sarcoplasmic/endoplasmic Ca^{2+} ATPase (SERCA) pump and serves to replenish ER Ca^{2+} levels by these mobile organelles, thus sustaining SOCE and lymphocyte activation. NAADP-mediated activation of lysosomal two pore channels (TPC) is important for secretion of cytolytic granules by cytotoxic T cells. The endolysosomal TRP mucolipin (TRPML) channels are crucial to endolysosomal function, such as endocytosis, exocytosis and autophagy, and are likely important, during T cell activation, for endocytosis of exhausted membrane proteins and replenishment of these

proteins through exocytosis at the immune synapse. Purinergic ionotropic receptor P2RX4 located in endolysosomal membranes was proposed to be important for vesicular fusion²⁰², but its role in T cell endolysosomal function remains unknown. CICR, Ca²⁺-induced Ca²⁺ release.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript