



Published in final edited form as:

J Am Chem Soc. 2016 September 28; 138(38): 12316–12319. doi:10.1021/jacs.6b06847.

A Catalytic Intermolecular Formal Ene Reaction between Ketone-Derived Silyl Enol Ethers and Alkynes

Stephen D. Holmbo[#], Nicole A. Godfrey[#], Joshua J. Hirner[†], Sergey V. Pronin^{*}

Department of Chemistry, University of California, Irvine, California 92697-2025, United States

[#] These authors contributed equally to this work.

Abstract

A catalytic formal ene reaction between ketone-derived silyl enol ethers and terminal alkynes is described. This transformation is uniquely capable of bimolecular assembly of 2-siloxy-1,4-dienes and can be used to access β,γ -unsaturated ketones containing quaternary carbons in the α -position.

Intermolecular alkenylation of enolates or their surrogates offers direct and convenient access to β,γ -unsaturated ketones, versatile building blocks that contain two orthogonal functionalization sites. Over 30 years ago, Migita reported pioneering examples of such sp^2 - sp^3 coupling processes, which relied on a Pd(0)-catalyzed alkenylation of *in situ*-generated tri-*n*-butyltin enolates with substituted vinyl bromides.¹ Subsequent works from other laboratories further demonstrated the power of Pd(0)- and Ni(0)-catalyzed alkenylations in the synthesis of β,γ -unsaturated ketones,^{2–8} although these reactions are typically limited to aryl alkyl or alkenyl alkyl ketones. Intermolecular alkenylations of dialkyl ketones bearing two unactivated enolizable positions are much less represented.^{9,10} In this context, selective formation of quaternary centers poses a particular challenge, and only isolated examples can be found in the literature.^{7,11} It is noteworthy that analogous intramolecular alkenylations are much more common and have found a number of applications in synthesis, likely owing to high levels of regiocontrol enforced by kinetic selectivity during cyclization.¹² Here we demonstrate a new catalytic formal ene reaction between silyl enol ethers and terminal alkynes that is uniquely capable of bimolecular assembly of 2-siloxy-1,4-dienes and can be used to access β,γ -unsaturated ketones containing quaternary carbons in the α -position.

During our work on the synthesis of paxilline indole diterpenes, we identified an indium(III) bromide-mediated alkenylation of a silyl enol ether with a terminal alkyne as suitable means for assembly of the requisite β,γ -unsaturated ketone.¹³ We became intrigued by the idea of performing intermolecular alkenylations in a catalytic fashion with retention of the silyl enol ether functionality in the product.¹⁴ Electrophilic activation of alkynes toward nucleophilic attack by silyl enol ethers has found extensive application in α alkenylation of ketones. In

^{*}Corresponding Author spronin@uci.edu.

[†]Present Address J.J.H.: Honeywell UOP, Des Plaines, IL 60016

Notes

The authors declare no competing financial interest.

his pioneering work, Conia described the first, to our knowledge, examples of such reactivity.¹⁵ A number of reports of related Al(III)-mediated¹⁶ and transition-metal-catalyzed^{17–22} intramolecular alkenylations of ketone-derived enoxysilanes followed in subsequent years, including some asymmetric variants.²³ However, relevant intermolecular alkenylations of silyl enol ethers find very little precedent,^{13,24,25} and the only available catalytic intermolecular process requires the use of 1,3-dicarbonyls as activated precursors²⁶ (Scheme 1). In the latter example, facile enolization of the substrates enables the ene-like reaction with alkynes. A similar process between ketone-derived silyl enol ethers and alkynes is considerably more challenging, as it requires a net proton transfer from the allylic position of the enoxysilane to the former alkyne fragment that has never been observed in a bimolecular setting.²⁷ Furthermore, the propensity of silyl enol ethers derived from unsymmetrical dialkyl ketones to isomerize under acidic conditions^{28,29} poses a potential issue of site selectivity. Finally, the product of the alkenylation, a 2-siloxy-1,4-diene, would itself represent a substrate for the reaction, posing an inherent issue of chemoselectivity. In particular, application of such a reaction to the selective construction of quaternary centers would require preferential alkenylation of fully substituted enoxysilane starting materials in the presence of less substituted enoxysilane products.

With these considerations in mind, we set out to investigate the catalytic formal ene reaction between silyl enol ether **1** and 1-octyne (Table 1; see Supporting Information (SI) for additional details). Gold(I) complexes¹⁹ proved to be poor catalysts for the desired transformation with the best-performing combination of a NHC-derived precatalyst and a halidescavenging additive providing only small amounts of diene **2** and the hydrolyzed product **3** (entry 1). Application of zinc(II) bromide²² was unsuccessful (entry 2). Conditions similar to those previously employed by Nakamura²⁶ afforded only traces of enoxysilane **2** (entry 3). However, changing the counter-anion in the indium(III)-based catalyst produced instructive trends. While chloride (entry 4) offered only marginal improvement over triflate, bromide (entry 5) allowed for the efficient alkenylation of substrate **1**, and only small amounts of ketone **3**, allene **4**,³⁰ and regioisomer **5** were observed. Application of indium(III) iodide and introduction of halide-scavenging additives were less efficient (entries 6 and 7). Catalytic alkenylation in the presence of indium(III) bromide could be performed at temperatures as low as 50 °C, which decreased the content of allene **4** and regioisomer **5** (entry 8). Attempted alkenylation in the presence of catalytic amounts of hydrogen bromide (generated *in situ*, entry 9) as well as triflic acid (entry 10) did not lead to product formation, confirming the catalytic role of the metal derivative. The robust TIPS-enol ethers³¹ performed better than more labile TMS and TBS derivatives. Interestingly, attempted alkenylation of isomeric silyl enol ether **6** afforded siloxydiene **2** as a major product, suggesting isomerization of **6** to **1** under the reaction conditions.

It is important to note that the product of double alkenylation was not observed during these studies. Alkenylation of the fully substituted enoxysilane **1** must therefore proceed

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/jacs.6b06847](https://doi.org/10.1021/jacs.6b06847).
Detailed experimental procedures and characterization data for new compounds (PDF)

considerably faster under our conditions than alkenylation of product 2. Although speculatively, we attribute the observed selectivity to the difference in the conformational preferences of the bulky siloxy group.³² To demonstrate the application of such unusual selectivity, the synthesis of diketone **9** was undertaken (Scheme 2). Starting diketone **8** contains three reactive α -positions, C1, C8, and C10, and selective alkenylation at C8 represents a considerable challenge. To our delight, subjection of the bis-silyl enol ether derived from diketone **8** to our alkenylation conditions resulted in a selective reaction of the fully substituted siloxyalkene fragment, delivering monoalkenylated product **9** after a mild hydrolytic workup.

With a functional set of conditions in hand, we investigated the preliminary scope of this catalytic formal ene reaction. We found that alkyl acetylenes with aromatic, halide, and ether functionalities as well as conjugated alkynes successfully participate in the alkenylation process (dienes **10–14** in Table 2). The reaction also tolerated a Lewis basic ester group (diene **15**), although more forcing conditions were required in this case.³³

Trimethylsilylacetylene and internal alkynes did not participate in the alkenylation. The formal ene reaction was effective at forming quaternary centers containing four non-methyl substituents (dienes **16** and **17**), and various substitution patterns of the cyclopentene ring were tolerated (dienes **18–21**). Synthesis of dienes **19–21** demonstrates a convenient approach to diastereoselective vicinal difunctionalization of a simple alkenone³⁴ and can be useful in the context of natural product synthesis.¹³ Cyclohexanone-derived silyl enol ethers also underwent successful alkenylation, and corresponding ketones **22** and **23** could be readily isolated after a mild hydrolytic workup.³⁵ Furthermore, acyclic silyl enol ethers were suitable substrates for this formal ene reaction (e.g., see diene **24**). Because the resulting 2-siloxy-1,4-dienes often proved hydrolytically unstable relative to their cyclic counterparts, the reaction mixtures were subjected to a mild hydrolytic workup, delivering the corresponding β , γ -unsaturated ketones (products **25–30**). In almost all cases, the desired 1,1-disubstituted alkenes were produced in a highly regioselective manner, and only small quantities of corresponding 1,2-disubstituted isomers were observed.

We propose that the mechanism of this catalytic formal ene reaction involves initial nucleophilic attack of enoxysilane **1** on the alkyne–indium(III) bromide complex³⁶ to form alkenylindium **31** (or the corresponding divinylindium derivative; Scheme 3).³⁷ The formation and stereochemistry of intermediate **31** or related species are supported by observation of deuterated ketone **34** upon treatment of the reaction mixture with CD₃OD.³⁴ Similar alkenylindium derivatives have been previously isolated and characterized by Baba.²⁴ Intermediate **31** can undergo protodemetalation with indium(III) bromide–ketone complex **32** to produce ketone **3** and enolate **33**, respectively.^{26,38} Participation of ketone **3**, which is formed early and whose concentration remains constant during the reaction,³⁹ is supported by the observation that an additive of ketone **35** undergoes silylation in situ to form enoxysilane **36**.³⁴ Subsequent silylation of enolate **33** is expected to afford the desired siloxydiene **2**.

In summary, we disclose a new catalytic intermolecular ene reaction between ketone-derived silyl enol ethers and terminal alkynes. Among the features of this alkenylation are selective formation of highly sterically encumbered quaternary centers and direct access to 2-

siloxo-1,4-dienes and corresponding β,γ -unsaturated ketones with synthetically challenging substitution patterns. The reaction has its limitations and future work will focus on expanding the substrate scope and further improving the efficiency of the process. However, this transformation has no alternatives among the current methods and can be expected to find broad application in natural product synthesis.

ACKNOWLEDGMENTS

Funding from the School of Physical Sciences at the University of California Irvine, Chao Family Comprehensive Cancer Center, and the National Science Foundation (DGE-1321846 to N.A.G.) is gratefully acknowledged. We thank Profs. Larry Overman, Chris Vanderwal, and Suzanne Blum for providing access to their instrumentation and helpful discussions.

REFERENCES


- (1). (a) Kosugi M; Hagiwara I; Migita T *Chem. Lett* 1983, 12, 839–841. For application in synthesis see: (b) Gracia J; Thomas EJJ *Chem. Soc., Perkin Trans. 1* 1998, 1, 2865–2871. (c) Almendros P; Rae A; Thomas EJ *Tetrahedron Lett* 2000, 41, 9565–9568.
- (2). (a) Chieffi A; Kamikawa K; Åhman J; Fox JM; Buchwald SL *Org. Lett* 2001, 3, 1897–1900. [PubMed: 11405739] (b) Hamada T; Buchwald SL *Org. Lett* 2002, 4, 999–1001. For application in synthesis see: [PubMed: 11893206] (c) Riou M; Barriault LJ *Org. Chem* 2008, 73, 7436–7439.
- (3). Huang J; Bunel E; Faul MM *Org. Lett* 2007, 9, 4343–4346. [PubMed: 17887766]
- (4). Lou S; Fu GC *J. Am. Chem. Soc* 2010, 132, 5010–5011. [PubMed: 20302338]
- (5). (a) Cosner CC; Helquist P *Org. Lett* 2011, 13, 3564–3567. [PubMed: 21688856] (b) Grigalunas M; Ankner T; Norrby P-O; Wiest O; Helquist P *Org. Lett* 2014, 16, 3970–3973. [PubMed: 25032503] (c) Grigalunas M; Ankner T; Norrby P-O; Wiest O; Helquist PJ *Am. Chem. Soc* 2015, 137, 7019–7022. (d) Grigalunas M; Norrby P-O; Wiest O; Helquist P *Angew. Chem., Int. Ed* 2015, 54, 11822–11825.
- (6). Su W; Raders S; Verkade JG; Liao X; Hartwig JF *Angew. Chem., Int. Ed* 2006, 45, 5852–5855.
- (7). For a relevant Heck reaction of alkyl vinyl ethers see: Datta GK; Larhed M *Org. Biomol. Chem* 2008, 6, 674–676. [PubMed: 18264566]
- (8). For a recent review of alkenylation of enolates see: Ankner T; Cosner CC; Helquist P *Chem. - Eur. J* 2013, 19, 1858–1871. [PubMed: 23325616]
- (9). See refs 1a, 5c, and 6. For an elegant solution see ref 4.
- (10). For relevant examples of alkenylation see: (a) Ooi T; Goto R; Maruoka K *J. Am. Chem. Soc* 2003, 125, 10494–10495. [PubMed: 12940712] (b) Nandi RK; Takeda N; Ueda M; Miyata O *Tetrahedron Lett* 2016, 57, 2269–2272.
- (11). See refs 1a and 1b for relevant discussions.
- (12). (a) Piers E; Marais PC *J. Org. Chem* 1990, 55, 3454–3455. (b) Piers E; Renaud JJ *Org. Chem* 1993, 58, 11–13. (c) Wang T; Cook JM *Org. Lett* 2000, 2, 2057–2059. [PubMed: 10891229] (d) Sole D; Peidro E; Bonjoch J *Org. Lett* 2000, 2, 2225–2228. [PubMed: 10930249] (e) Zhao S; Liao X; Cook JM *Org. Lett* 2002, 4, 687–690. [PubMed: 11869102] (f) Yu J; Wang T; Liu X; Deschamps J; Flippen-Anderson J; Liao X; Cook JM *J. Org. Chem* 2003, 68, 7565–7581. [PubMed: 14510528] (g) Sole D; Diaba F; Bonjoch J *J. Org. Chem* 2003, 68, 5746–5749. [PubMed: 12839475] (h) Cao H; Yu J; Wearing XZ; Zhang C; Liu X; Deschamps J; Cook JM *Tetrahedron Lett* 2003, 44, 8013–8017. (i) Zhou H; Liao X; Cook JM *Org. Lett* 2004, 6, 249–252. [PubMed: 14723540] (j) Yu J; Wearing XZ; Cook JM *J. Org. Chem* 2005, 70, 3963–3979. [PubMed: 15876085] (k) Sole D; Urbaneja X; Bonjoch J *Org. Lett* 2005, 7, 5461–5464. [PubMed: 16288531] (l) Dounay AB; Humphreys PG; Overman LE; Wroblewski AD *J. Am. Chem. Soc* 2008, 130, 5368–5377. [PubMed: 18303837] (m) Shen L; Zhang M; Wu Y; Qin Y *Angew. Chem., Int. Ed* 2008, 47, 3618–3621. (n) Yao Y; Liang G *Org. Lett* 2012, 14, 5499–5501. [PubMed: 23095081]
- (13). George DT; Kuenstner EJ; Pronin SV *J. Am. Chem. Soc* 2015, 137, 15410–15413. [PubMed: 26593869]

- (14). The utility of silyl enol ethers in modification of corresponding ketones has been appreciated for a long time: Brownbridge P *Synthesis* 1983, 1983, 1–28.
- (15). (a) Drouin J; Boaventura M-A; Conia J-M. *J. Am. Chem. Soc* 1985, 107, 1726–1729. Also see: (b)Drouin J; Boaventura MA *Tetrahedron Lett* 1987, 28, 3923–3926. For application in synthesis see:(c)Forsyth CJ; Clardy JJ *Am. Chem. Soc* 1990, 112, 3497–3505.(d)Huang H; Forsyth CJ *J. Org. Chem* 1995, 60, 2773–2779.(e)Huang H; Forsyth CJ *J. Org. Chem* 1995, 60, 5746–5747.
- (16). Imamura K; Yoshikawa E; Gevorgyan V; Yamamoto Y *Tetrahedron Lett* 1999, 40, 4081–4084.
- (17). (a) Maeyama K; Iwasawa N *J. Am. Chem. Soc* 1998, 120, 1928–1929.(b)Iwasawa N; Maeyama K; Kusama H *Org. Lett* 2001, 3, 3871–3873. [PubMed: 11720557] (c)Kusama H; Yamabe H; Iwasawa N *Org. Lett* 2002, 4, 2569–2571. [PubMed: 12123378] (d)Iwasawa N; Miura T; Kiyota K; Kusama H; Lee K; Lee PH *Org. Lett* 2002, 4, 4463–4466. [PubMed: 12465913] (e)Grandmarre A; Kusama H; Iwasawa N *Chem. Lett* 2007, 36, 66–67.
- (18). Nevado C; Cardenas DJ; Echavarren AM *Chem. - Eur. J* 2003, 9, 2627–2635. [PubMed: 12794906]
- (19). (a) Staben ST; Kennedy-Smith JJ; Huang D; Corkey BK; LaLonde RL; Toste FD *Angew. Chem., Int. Ed* 2006, 45, 5991 For selected applications in synthesis see:(b)Linghu X; Kennedy-Smith JJ; Toste FD *Angew. Chem., Int. Ed* 2007, 46, 7671–7673.(c)Nicolaou KC; Tria GS; Edmonds DJ; Kar MJ *Am. Chem. Soc* 2009, 131, 15909–15917.(d)Huwlyer N; Carreira EM *Angew. Chem., Int. Ed* 2012, 51, 13066–13069.(e)Lu Z; Li Y; Deng J; Li A *Nat. Chem* 2013, 5, 679–684. [PubMed: 23881499] (f)Moreno J; Picazo E; Morrill LA; Smith JM; Garg NK *J. Am. Chem. Soc* 2016, 138, 1162–1165. [PubMed: 26783944] (g)Li Y; Zhu S; Li J; Li AJ *Am. Chem. Soc* 2016, 138, 3982–3985.
- (20). (a) Barabe F; Betournay G; Bellavance G; Barriault L *Org. Lett* 2009, 11, 4236–4238. [PubMed: 19739690] (b)Sow B; Bellavance G; Barabe F; Barriault L *Beilstein J. Org. Chem* 2011, 7, 1007–1013. [PubMed: 21915201] (c)Barabe F; Levesque P; Korobkov I; Barriault L *Org. Lett* 2011, 13, 5580–5583. [PubMed: 21916520]
- (21). (a) Ito H; Ohmiya H; Sawamura M *Org. Lett* 2010, 12, 4380–4383. [PubMed: 20822100] (b)Iwai T; Okochi H; Ito H; Sawamura M *Angew. Chem., Int. Ed* 2013, 52, 4239–4242.
- (22). Han Y; Zhu L; Gao Y; Lee C-S *Org. Lett* 2011, 13, 588–591. [PubMed: 21218775]
- (23). (a) Corkey BK; Toste FD *J. Am. Chem. Soc* 2007, 129, 2764–2765. [PubMed: 17305344] (b)Brazeau J-F; Zhang S; Colomer I; Corkey BK; Toste FD *J. Am. Chem. Soc* 2012, 134, 2742–2749. [PubMed: 22296571]
- (24). Nishimoto Y; Moritoh R; Yasuda M; Baba A *Angew. Chem., Int. Ed* 2009, 48, 4577–4580.
- (25). (a) Yamaguchi M; Tsukagoshi T; Arisawa M *J. Am. Chem. Soc* 1999, 121, 4074–4075. (b)Arisawa M; Miyagawa C; Yamaguchi M *Synthesis* 2002, 2002, 138–145. Also see:(c)Jones IL; Moore FK; Chai CLL *Org. Lett* 2009, 11, 5526–5529. [PubMed: 19874042]
- (26). Nakamura M; Endo K; Nakamura EJ *Am. Chem. Soc* 2003, 125, 13002 See also:Itoh Y; Tsuji H; Yamagata K.-i.; Endo K; Tanaka I; Nakamura M; Nakamura E *J. Am. Chem. Soc* 2008, 130, 17161–17167 and references therein. [PubMed: 19053468]
- (27). For intramolecular examples see refs 17c, 17d, and 18.
- (28). Stork G; Hudrlik PF *J. Am. Chem. Soc* 1968, 90, 4462–4464.
- (29). Formation of trace quantities of strong Brønsted acids can be associated with processes catalyzed by metal salts: Wabnitz TC; Yu J-Q; Spencer JB *Chem. - Eur. J* 2004, 10, 484–493. [PubMed: 14735517]
- (30). Formation of 4 likely involves a regioisomeric carbometalation of alkyne followed by an intramolecular hydride migration. For relevant observations see: Harrak Y; Simonneau A; Malacria M; Gandon V; Fensterbank L *Chem. Commun* 2010, 46, 865–867.
- (31). TIPS-enol ether precursors are readily available from corresponding ketones. See SI for preparation.
- (32). Such selectivity for fully substituted silyl enol ethers is not exclusive. See SI for a detailed discussion.
- (33). Application to aldehyde-, ketone-, nitrile-, and amide-containing alkynes was unsuccessful thus far.
- (34). See SI for details.


- (35). Attempted purification of corresponding siloxydienes was unsuccessful.
- (36). Surendra K; Corey EJ J. Am. Chem. Soc 2014, 136, 10918–10920. [PubMed: 25095905]
- (37). For a review of organoindium compounds see: Shen Z-L; Wang S-Y; Chok Y-K; Xu Y-H; Loh T-P Chem. Rev 2013, 113, 271–401. [PubMed: 23110495]
- (38). Indium(III) bromide–ketone complex 32 can be expected to be a strong Brønsted acid: Ren J; Cramer CJ; Squires RR J. Am. Chem. Soc 1999, 121, 2633–2634.
- (39). Protodemetalation of intermediate 10 or related species with alkyne–indium(III) bromide complex can be responsible for initial generation of ketone 3, which cannot be accounted for by the presence of residual TIPSOH or adventitious moisture.

Previous work:


- Alkenylation of silyl ketene acetals²⁴ (stoichiometric in metal; also see ref. 13, 25)




- Alkenylation of β -ketoesters and 1,3-diketones²⁶ (catalytic in metal)

**This work:**


- A catalytic formal ene reaction between enoxysilanes and terminal alkynes

**Scheme 1.**

Comparison of Relevant Intermolecular Alkenylations with Current Work




Scheme 2.
Synthesis of Diketone 9




Scheme 3.
Proposed Mechanism of the Alkenylation

Table 1.

Development of the Catalytic Formal Ene Reaction



Observed by-products:



Entry	Conditions ^a	2(%) ^b	3(%) ^b	4(%) ^{b,c}	5(%) ^b
1	5 mol% IPrAuCl, 5 mol% NaBARF, 65°C	11	8	10	<1
2	10 mol% ZnBr ₂ , 65°C	4	5	1	<1
3	5 mol% In(OTf) ₃ , 65°C	<1	1	<1	<1
4	5 mol% InCl ₃ , 65°C	5	9	<1	<1
5	5 mol% InBr₃, 65°C	72	5	6	2
6	5 mol% InI ₃ , 65°C	36	6	7	<1
7	5 mol% InBr ₃ , 5 mol% NaBARF, 65°C	40	11	3	<1
8 ^d	5 mol% InBr₃, 50°C	65	9	3	<1
9	5 mol% MeOH, 8 mol% TMSBr, 65°C	<1	<1	<1	<1
10	5 mol% TfOH, 65°C	<1	<1	<1	<1

Attempted alkenylation of silyl enol ether 6:



^a0.25–0.5 mmol of 1, 1.5 equiv of alkyne, 0.25–0.5 mL of (CH₂Cl)₂, 20–28 h.^bBased on internal standard and determined by ¹H NMR.^cdr = 1:1.^d65% isolated yield of 2.

Table 2.

Preliminary Substrate Scope of the Alkenylation^a

^aTypically, starting silyl enol ethers were mixtures of isomers; see SI for details. 1 equiv of silyl enol ether, 1.5 equiv of alkyne, 10 mol% of InBr₃, (CH₂Cl)₂ (1 M in enoxysilane), 65 °C, ca. 24 h; rr 10:1 (except **15**, rr 9:1; **24**, rr 7:1). Siloxydienes (except **13**, **14**, and **24**) contained ca. 5 mol% of inseparable allenes.

^bHeated to 50 °C.

^cHeated to 80 °C.

^d5 mol% of InBr₃.

^eNeat. ^f5 M in silyl enol ether.

^gHeated for ca. 72 h.

^hHeated for ca. 48 h.

ⁱAfter hydrolysis.