



HHS Public Access

Author manuscript

Inj Prev. Author manuscript; available in PMC 2020 June 01.

Published in final edited form as:

Inj Prev. 2019 June ; 25(3): 191–198. doi:10.1136/injuryprev-2017-042676.

Impacts of an opioid overdose prevention intervention delivered subsequent to acute care

Caleb Banta-Green,

University of Washington, Alcohol and Drug Abuse Institute; University of Washington School of Public Health, Health Services

Phillip O. Coffin,

San Francisco Department of Public Health, San Francisco, California, USA; Department of Medicine, University of California San Francisco, San Francisco, California, USA

Joseph O. Merrill,

Department of Medicine, University of Washington, Seattle, Washington, USA

Jeanne M. Sears,

Department of Health Services, School of Public Health and Harborview Injury Prevention and Research Center, University of Washington, Seattle, Washington, USA; Institute for Work and Health, Toronto, Ontario, Canada

Chris Dunn,

Department of Psychiatry & Behavioral Sciences, University of Washington, Seattle, Washington, USA

Anthony S. Floyd,

Alcohol & Drug Abuse Institute and Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington, USA

Lauren K. Whiteside,

Department of Emergency Medicine, University of Washington, Seattle Washington, USA

N. David Yanez,

Oregon Health & Science University/Portland State University School of Public Health, Oregon Health & Science University, Portland, Oregon, USA

Dennis M. Donovan

Alcohol & Drug Abuse Institute and Department of Psychiatry & Behavioral Sciences, University of Washington, Seattle, Washington, USA

Abstract

Background—Opioid overdose is a major and increasing cause of injury and death. There is an urgent need for interventions to reduce overdose events among high-risk persons.

Methods—Adults at elevated risk for opioid overdose involving heroin or pharmaceutical opioids who had been cared for in an emergency department (ED) were randomized to overdose education combined with a brief behavioral intervention and take-home naloxone or usual-care. Outcomes included: 1) time to first opioid overdose-related event resulting in medical attention or death using competing risks survival analysis; and 2) ED visit and hospitalization rates, using negative binomial regression and adjusting for time at risk.

Results—During the follow-up period, 24% of the 241 participants had at least one overdose event, 85% had one or more ED visits, and 55% had at least one hospitalization, with no significant differences between intervention and comparison groups. The instantaneous risk of an overdose event was not significantly lower for the intervention group (subhazard ratio: 0.83; 95% CI: 0.49–1.40).

Discussion—These null findings may be due in part to the severity of the population in terms of housing insecurity (70% impermanently housed), drug use, unemployment, and acute health care issues. Given the high overdose and health care utilization rates, more intensive interventions, such as direct referral and provision of housing and opioid agonist treatment medications may be necessary to have a substantial impact on opioid overdoses for this high acuity population in acute care settings.

Keywords

opioid; overdose; behavioral intervention

INTRODUCTION

Opioid overdose deaths continue to increase and are a major cause of preventable death.[1] According to the Centers for Disease Control and Prevention, opioid-involved overdose deaths quadrupled from 1999 to 2015 when there were 33,091 such deaths in the United States.[2] Additionally, opioid-related Emergency Department (ED) visits and inpatient admissions have increased dramatically over the years reflecting the increase in non-fatal overdose.[3] Many overdoses are amenable to intervention due to biological and social circumstances.[4] Opioid overdoses rarely lead to sudden death, with death usually occurring several hours after consumption,[5] though this may be changing as illicit synthetic opioids with their rapid rates of onset and high potency emerge as major causes of death.[6] Moreover, most overdoses occur in the presence of another person,[7,8] providing an opportunity for bystander intervention.

Brief behavior change counseling is based upon motivational interviewing[9], has been found to help reduce drug use frequency[10], and to significantly improve health behaviors such as alcohol use and injury, to increase entry into drug abuse treatment, and reduce costs in emergency departments (ED). [11,12] In pharmaceutical opioid using patients in the ED with elevated risk for overdose, patients receiving a brief behavioral intervention had decreased overdose risk behaviors.[13] Additionally, brief intervention has been used to decrease drug use among patients in the ED.[14] However, these studies did not specifically target illicit opioid use and did not combine brief behavior change counseling with take-home naloxone. We combined these interventions based on the information-motivation-

behavior model, positing that overdose education, combined with self-identified motivating factors and the behavioral skills to utilize naloxone might impact overdose occurrence.[15]

Naloxone is an opioid-antagonist prescription medication that reverses opioid overdoses by preferentially binding to opioid receptors and displacing opioids such as heroin, morphine, oxycodone, and fentanyl and reversing respiratory depression and sedation. Naloxone cannot be abused, has no psycho-active effects and has been found to be extremely safe.[16,17] Since the 1990s, naloxone has increasingly been provided to people who use drugs through low-threshold service programs – such that by 2014 community based programs were distributing naloxone to laypersons at 644 sites in the U.S.[18] Take-home-naloxone for lay people has been recommended by organizations such as the World Health Organization and in the Substance Abuse and Mental Health Services Administration opioid overdose toolkit. [19,20]

Research on take-home naloxone provided to people at risk for having or witnessing an overdose indicates that: 1) naloxone administration has not resulted in dangerous health consequences;[21] 2) lay persons can be trained to recognize an overdose and evaluate whether administration of naloxone is warranted as well as medical experts;[22] 3) illicit drug users are willing to administer naloxone to each other;[23] 4) naloxone availability does not increase drug use;[24] 5) many opioid overdoses have been reversed with naloxone as a result of overdose prevention and recognition training combined with the distribution of take-home naloxone;[25] and 6) provision of naloxone is associated with reduced mortality in communities that implement the program compared to communities that do not and reduced opioid overdose mortality after release from prison.[25,26] However, there are no trials that aim to assess the impact of pairing naloxone provision with a brief behavioral intervention in health care settings on subsequent overdose events and health care utilization.

Emergency departments (EDs) and acute care settings are potentially advantageous settings to reach populations vulnerable to opioid overdose that may not access health care in other settings such as substance use disorder treatment centers or primary care. In this study, we tested an intervention for opioid users at elevated risk for overdose that was delivered during or after an acute care episode that combined opioid overdose education, a take-home-naloxone kit, and brief behavior change counseling to determine the impact on participants' subsequent opioid overdoses, ED visits, and hospitalizations.

METHODS

Setting

Participants were enrolled subsequent to a visit at one of two EDs in Seattle: Harborview Medical Center (HMC) and the University of Washington Medical Center (UWMC). HMC is a large urban academic medical center dedicated to an underserved population including those with mental health and substance use problems. UWMC is an academic tertiary care hospital providing extensive specialty care. Potential participants were identified through medical records review by study staff or health care staff referrals. Recruitment and enrollment generally occurred between the hours of 12pm and 8pm weekdays in the ED, in other hospital units during subsequent inpatient admission, or at respite care (a recuperative

care facility adjacent to HMC for homeless people who require medical assistance and shelter and do not require inpatient treatment).

Participants

Potential participants were identified either by study staff reviewing electronic medical records or by medical staff referral with eligibility confirmed via a screening questionnaire. Eligibility criteria included being at elevated risk of opioid overdose based upon: (1) reason for visit was opioid overdose; (2) use of pharmaceutical opioids not prescribed 2 or more times in the prior month; (3) use of other opioids, alcohol, sedatives, or stimulants within two hours of using opioids 2 or more times in the prior month; (4) average daily dose of prescribed opioids greater than 10mg morphine equivalent dose or higher for 15 or more of the last 30 days; or (5) enrolled in an opioid agonist therapy (OAT) program and receiving methadone or buprenorphine. Opioids needed to be used at least twice in the last 30 days (or if institutionalized recently, in the most recent month they were not institutionalized) with pharmaceutical users also needing to have other risks present. Subjects were not excluded if pregnant and were offered naloxone if in the intervention arm and informed during consent about potential risks to a fetus due to precipitated withdrawal and the need to seek emergency medical care.

Exclusion criteria included: (1) refusing access to follow up medical or drug treatment records; (2) inability to communicate in English; (3) current suicidal ideation; (4) significant cognitive or psychiatric impairment; (5) inability to provide adequate contact information to assist with follow-up (the number of required contacts was reduced after a month of recruitment from three to one as most homeless people were being excluded which would have negatively impacted the generalizability of the findings); (6) under age 18 or over age 70; (7) not living in Washington State or planning to move from Washington State within a year; (8) receiving treatment for sexual assault; or (9) currently having non-expired naloxone.

Potential participants provided informed consent for eligibility screening. If eligible and interested in the study, consent was obtained for study participation. Eligibility screening and study participation were remunerated with \$5 and \$20 store gift cards respectively. Follow up surveys were conducted at 3, 6 and 12 months (with \$10, \$10 and \$20 gift card remunerations respectively), these data are not presented here as follow up rates were below 50% at each time point. Releases of information and HIPAA authorizations were obtained to access medical records and drug treatment data. Baseline data were collected by interventionists in clinical settings, with an attempt to maximize privacy, and participants were randomized; this process took approximately 30 to 45 minutes. An unrestricted or “fair-coin” randomization process was utilized to generate a study assignment table based upon study identification numbers and implemented automatically via REDCap with interventionists learning study assignment at the same time as the participant.

Study data were collected and managed using REDCap (Research Electronic Data Capture) electronic data capture, secure web-based tools hosted at the Institute of Translational Health Sciences at the University of Washington.

Intervention

The intervention consisted of 1) overdose education, 2) a brief behavioral change counseling component to assist participants in identifying their overdose risks and the steps they were interested in taking to reduce those risks and 3) a naloxone kit. The intervention was provided by two interventionists who had master's degrees and at least basic training in motivational interviewing.

Overdose education included watching an 8-minute video and reviewing an informational flier with the interventionist, which addressed risk factors for an opioid overdose, overdose recognition, recommendation to call 911, how to administer naloxone, and guidance to remain with the overdose victim for several hours. The flier also provided specific information about locations where naloxone could be obtained either free at area syringe exchange programs or for purchase at a local pharmacy. The flier included a link to www.stopoverdose.org which has online overdose educational materials, including the training video utilized at the time created by the New York City Department of Health as well as a naloxone locator for Washington State.

Naloxone administration training included hands-on practice assembling the kit which included a luer lock syringe, 2mg/2ml naloxone (Amphastar NDC#76329-3369-1), and a mucosal atomization device. Intranasal administration was an off-label route of administration and a Food and Drug Administration (FDA) Investigational New Drug (IND) application was required (#112,043). The kit included two doses of naloxone, two mucosal atomizers, a disposable rescue breathing mask, a wallet card with information about Washington State's Good Samaritan Overdose and naloxone access law (RCW 69.50.315), and the educational flier all packaged in a nylon pouch. Participants were directly handed the kit by study staff, however they did not need to accept the offer of the kit to be considered a study participant.

Participants assigned to the comparison group were provided the informational flier.

Intervention fidelity

Sessions were audio recorded and initial training and regular supervision with a doctoral level psychologist (CD) occurred throughout the study. A sample of intervention recordings were reviewed for fidelity. The intervention was motivational interviewing (MI)-inspired in order to facilitate rapport building and participant engagement. Despite not being a full MI intervention, given the didactic and interactive educational components, fidelity was measured using scales from the Motivational Interviewing Treatment Integrity (MITI) 3.1.1. Interventionist behavioral counts were measured along with MI Spirit, an average measure of the quality of MI delivery.[27]

Data sources

The baseline survey included demographic information, housing status, and relationship status. Opioid use in the prior 30 days was categorized as: only pharmaceutical opioids whether prescribed to the participant or not, OAT from an approved provider (licit source), heroin and OAT, and OAT no heroin. Route of administration was coded into whether or not

a person had smoked, snorted or injected any opioids (96% reported injecting). Days of opioid use in the prior 30 days was recorded. Protective factors, including whether “anyone you have regular contact with” had “overdose education” or “regularly carry or have quick access to naloxone”, were documented. Overdose risk factors included: overdose history; using opioids when no one else was around or behind a locked door; and using opioids within two hours of alcohol, sedatives/downers (specific brand names were provided), using another kind of opioid, or stimulants including cocaine, methamphetamine or pharmaceutical stimulants.

Health care utilization data from UW Medicine included encounter and billing data from HMC and UWMC, their EDs, and affiliated onsite and offsite clinics in the Seattle, Washington metropolitan area. The Comprehensive Hospital Abstract Reporting System (CHARS) maintained by the Washington State Department of Health was used for statewide capture of opioid overdose events resulting in inpatient or hospital observation stays. CHARS contains hospital discharge information for inpatient and observation stays derived from billing records for essentially all Washington State community hospitals. Statewide death certificate data were obtained from the Washington State Department of Health.

Outcome measures

ED visits were defined as an encounter that had at least one charge originating from the ED, regardless of whether it resulted in inpatient admission. Some encounters may have been counted as both an ED visit and an inpatient admission, but each metric was analyzed separately. Index encounters were defined as ED visits or inpatient admissions (1) beginning on or before the randomization date and (2) concluding on or after the randomization date. ED visits and inpatient admissions were counted separately, and included all encounters for any principal diagnosis with admission/visit dates occurring after the discharge date of the index visit through December 31, 2015. Index encounters were excluded from outcome encounter counts and excluded from consideration as the first opioid overdose event (described below), because the need for these encounters was evidenced prior to the intervention and hence not properly considered an outcome.

All three administrative data sources (i.e., UW Medicine, CHARS, death certificate data) were used jointly to identify the first opioid overdose event occurring after randomization and discharge from the index encounter, and censored at December 31, 2015. Time to the first opioid overdose event was measured from randomization to the date of the first-occurring qualifying event: (1) UW Medicine ED, inpatient, or outpatient encounter for opioid overdose, (2) CHARS inpatient admission or observation stay for opioid overdose, or (3) death from opioid overdose. The definitions for an opioid overdose based upon ICD 9 and 10 codes across datasets are detailed in the supplement.

Sample size

Power calculations were based on the estimated annual overdose rate of 20% for heroin users and 10% for pharmaceutical opioid users (seen in the ED), and reduction in opioids overdoses of 50% due to the intervention. The sample size for heroin users to meet these parameters was 219 with one year of follow up for overdose. For pharmaceutical users with

an estimated annual overdose rate of 10%, we would require roughly twice the number of subjects or double the follow up time to have the same number of overdose events.

Data analysis

Health care utilization outcomes (i.e., number of ED visits, number of inpatient admissions) were analyzed using negative binomial regression with robust variance estimates and a time at-risk exposure adjustment for available follow-up time. Mean rates per person-year were calculated using these models.

Kaplan-Meier survival function curves were used to depict time from randomization to the first opioid overdose event, with days of follow-up as the time scale. Death due to causes other than opioid overdose was treated as a censoring event, but cannot be considered independent of randomization assignment. We therefore treated death due to causes other than opioid overdose as a competing risk, using competing risk survival analysis models to analyze time from randomization to the first opioid overdose event.[28] The STATA command -stcrreg- (based on the Fine and Gray semiparametric method) was used to produce subhazard ratios (SHR).[29,30]

All statistical tests were two-tailed, with statistical significance defined as $p < 0.05$. Analyses were performed using Stata/MP 13.1 for Windows. This study was approved by the Washington State Institutional Review Board and the University of Washington Human Subjects Division. The study was registered at clinicaltrials.gov (NCT01788306).

RESULTS

Study enrollment

Enrollment occurred between January 31, 2013 through April 3, 2015, allowing for at least 272 and up to 1,064 days of follow-up, which ended on December 31, 2015. Participant enrollment, allocation and analysis are outlined in the Consort diagram in Figure 1, as are reasons for study exclusion. Among the 430 assessed for study eligibility, 256 were enrolled and randomized, with 125 allocated to the intervention and 131 to the comparison group.

The study under-enrolled compared to the original study design of 500 heroin and 500 pharmaceutical opioid users. The randomization process appears to have achieved sufficient balance; baseline characteristics in Table 1 are comparable (all p -values > 0.05). For these analyses only those who provided consent to access secondary data were included, 115 in the intervention arm and 126 in the comparison group, a combined total of 241. The care setting in which participants were recruited included ED ($n=149$), respite care ($n=37$), and hospital inpatient ($n=55$) and did not differ significantly by study assignment. We identified potential subjects as quickly as possible and among those enrolled in respite care 89% had their baseline assessment completed within three days of starting the assessment (maximum 5 days) and among inpatients 98% had the assessment completed within 2 days (maximum 3).

Characteristics of Intervention and Comparison Groups

The majority of participants were male, White, non-Hispanic, homeless, not in a relationship, unemployed, used heroin and not in OAT, and used opioids by routes of ingestion including smoking, snorting or injecting (Table 1). The mean number of days that opioids were used in the past 30 was 24.7 (8.6 s.d.) with a median of 30. Protective factors for overdose were reported by a minority of participants including others in their life having overdose education or possessing naloxone. The majority had a history of an opioid overdose, 19% in the prior three months, 38% sometime prior to the most recent three months. The vast majority, 78%, reported using opioids alone sometimes or always in the prior three months and many reported using a range of other substances within two hours of using opioids. A substantial minority reported having had prior overdose education and knowing others who had naloxone.

Motivational interviewing fidelity

MI Spirit was calculated on 61 intervention sessions (out of 76 recordings), with an average score of 4.23 (out of 5). Recorded length across the 76 sessions for the overdose education and brief behavior change counseling content averaged 27 minutes, with length ranging from 10–55 minutes.

Overdose events by intervention assignment

Opioid overdose events and censoring by randomization status are presented in Table 2 and indicate that 23.7% of participants had at least one overdose event of some type, 6.2% had a non-overdose fatality and 70.1% had no observed event prior to censoring at the end of study follow up.

Health care utilization by intervention assignment

The majority (55%) of all participants had a hospital admission during the follow-up period with an average annual rate of 1.17 (95% CI 0.95–1.44) visits, and with no significant difference between the intervention and comparison groups (Table 3). A substantial majority (85%) of all participants had a subsequent emergency department visit during the follow-up period, with an average annual rate of 4.90 (95% CI 4.25–5.66) visits, and with no significant difference by intervention assignment.

Time to first overdose by intervention assignment

In the competing risk regression analysis, the difference in the time to first overdose event was not significantly lower for the intervention group relative to the comparison group (subhazard ratio: 0.83; 95% CI: 0.49–1.40). These data are presented as a survival curve in Figure 2.

DISCUSSION

In this study of patients at high risk for opioid overdose presenting for or soon after emergency care, an overdose prevention intervention was found to have no statistically significant impact on subsequent overdoses, either positive or negative. This null finding is

perhaps not surprising given the medical and social acuity of the population in terms of homelessness, drug use, and other health and social issues. A brief, one-time intervention in acute care settings or subsequent to receiving acute care may not be sufficient to reduce serious overdose events. The multi-part intervention was likely more intensive and time consuming than most overdose education and naloxone distribution program in community or medical settings, although there is great heterogeneity in these interventions.

Population-based studies have found decreased mortality rates associated with distributing naloxone to illicit drug users.[25] Estimates of the lifetime impact of naloxone distribution to individual heroin users are modest, a 6% mortality reduction.[31] Naloxone distribution programs provide clear life-saving benefits, however they are also insufficient to substantially address opioid overdose alone. Brief interventions in the ED have shown modest benefits in decreasing opioid-related risk behaviors and drug use, however these studies were of those with recent prescription opioid misuse and utilized a much broader definition of overdose that was not limited to opioids. [13,14] We did not use self-report of overdose as an outcome as follow up rates at each time point were less than 50%, perhaps due to the high levels of housing impermanence. We chose to enroll a high acuity population, despite knowing that this might lessen the impact of the intervention and lower follow up rates as we felt it was important to enroll a population representative of that seen in the care settings. It is possible that the informational flyer provided to the comparison group combined with the substantial increases in take-home naloxone within the community were sufficient to reduce any differential effect of the intervention.

The statistical power to detect differences was limited by the sample size and given the modest, though significant, impact of naloxone distribution found in other studies likely was an important limitation in our ability to find any potential impact of our intervention. We attempted to address this by adding measures of non-fatal overdose as an outcome because studies with an outcome of fatal overdose require sample sizes of many thousands,[32] but we could only capture overdose events that resulted in an outpatient visit, ED or hospital admission. Naloxone distribution studies are further limited in that naloxone distributed to one person is often administered to another, potentially under-estimating the total impact of the intervention. We did not account for heroin users being on OAT in our sampling design, which is a complicating issue in that research indicates being on OAT reduces fatal overdose occurrence, potentially reducing the overdose event rate and the statistical power to detect any differences.[33] A sub-analysis of the time to overdose event adjusting for opioid-use type did not significantly impact results (data not shown).

An important limitation was that study recruitment was challenging given the care settings and the acuity of the population, acute care settings are very hectic and complicated the logistics identifying, approaching and enrolling subjects in a study. Medical providers were re-oriented to the study multiple times in order to reinforce that pharmaceutical opioid users, including those prescribed the medications were indeed eligible for the study. There were also challenges in identifying pharmaceutical users due to very limited use of and access to Washington State's prescription drug monitoring program (PDMP) by medical providers at the time of the study. Study staff were precluded from accessing the PDMP for research purposes per state law.

In addition to overdose education, counseling, and naloxone, other more robust interventions such as direct referral and provision of housing and OAT medications may be necessary to have a clinically-significant impact on opioid overdoses for this high-acuity population served in acute care settings. The ED is a challenging setting for delivering an intervention logistically, in terms of timing and space constraints, and due to the medical state of patients. The most common reasons for refusing the screening interview (n=510), among those approached (n=936) were “not interested” (49%), “not feeling well” (39%) and “no time” (8%). Only a small proportion of those enrolled in the study were seen for an opioid overdose (12%). Recent findings suggest that brief overdose and naloxone training is sufficient and it appears that a population level mortality benefit is associated with higher rates of naloxone distribution in a community. [25,34–41] Therefore, ED overdose prevention interventions might reasonably be limited to brief education with the direct provision of take-home-naloxone. Future research on the impact of a more modest intervention with a larger number of acute care participants may be worthwhile.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

Research reported in this publication was supported by the National Institute on Drug Abuse of the National Institutes of Health under award number 1R01A030351. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Study data were collected and managed using REDCap electronic data capture tools hosted at the Institute of Translational Health Sciences. REDCap at ITHS is supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number UL1 TR002319.

We would like to thank:

Barbara Burke, LeslieENZian, Shira Hasson-Schiff, Sheree Miller, Susan Stern, Steve Mitchell, Melissa Phares, Jeff Purcell, Harborview Emergency Department, Harborview Adult Medicine Clinic, Harborview Respite Program at Jefferson Terrace (Edward Thomas House), Data Safety Monitoring Oversight Committee, and Administrative Support staff at the UW Alcohol and Drug Abuse Institute.

REFERENCES

1. Jones CM, Mack KA, Paulozzi LJ. Pharmaceutical overdose deaths, United States, 2010. *JAMA* 2013;309:657–9. doi:10.1001/jama.2013.272 [PubMed: 23423407]
2. Rudd RA, Seth P, David F, et al. Increases in Drug and Opioid-Involved Overdose Deaths — United States, 2010–2015. *MMWR Morb Mortal Wkly Rep* 2016;65:1445–52. doi:10.15585/mmwr.mm6505051e1 [PubMed: 28033313]
3. Weiss AJ, Elixhauser A, Barrett ML, et al. Opioid-Related Inpatient Stays and Emergency Department Visits by State, 2009–2014. *HCUP Stat Br* #219 2014;;1–15.<http://www.hcupus.ahrq.gov/reports/statbriefs/sb219-Opioid-Hospital-Stays-ED-Visits-by-State.pdf>
4. McGregor C, Darke S, Ali R, et al. Experience of non-fatal overdose among heroin users in Adelaide, Australia: circumstances and risk perceptions. *Addiction* 1998;93:701–11.<http://www.ncbi.nlm.nih.gov/pubmed/9692269> (accessed 25 Oct 2017). [PubMed: 9692269]
5. Baca CT, Grant KJ. Take-home naloxone to reduce heroin death. *Addiction* 2005;100:1823–31. doi: 10.1111/j.1360-0443.2005.01259.x [PubMed: 16367983]
6. Fairbairn N, Coffin PO, Walley AY. Naloxone for heroin, prescription opioid, and illicitly made fentanyl overdoses: Challenges and innovations responding to a dynamic epidemic. *Int J Drug Policy* 2017;46:172–9. doi:10.1016/j.drugpo.2017.06.005 [PubMed: 28687187]

7. Davidson PJ, Ochoa KC, Hahn JA, et al. Witnessing heroin-related overdoses: the experiences of young injectors in San Francisco. *Addiction* 2002;97:1511–6. <http://www.ncbi.nlm.nih.gov/pubmed/12472634> (accessed 25 Oct 2017). [PubMed: 12472634]
8. Strang J, Powis B, Best D, et al. Preventing opiate overdose fatalities with take-home naloxone: pre-launch study of possible impact and acceptability. *Addiction* 1999;94:199–204. <http://www.ncbi.nlm.nih.gov/pubmed/10396785> (accessed 25 Oct 2017). [PubMed: 10396785]
9. Miller WR (William R, Rollnick S. *Motivational interviewing: preparing people for change*. Guilford Press 2002.
10. Whitlock EP, Polen MR, Green CA, et al. Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2004;140:557–68. <http://www.ncbi.nlm.nih.gov/pubmed/15068985> (accessed 25 Oct 2017). [PubMed: 15068985]
11. Estee S, Wickizer T, He L, et al. Evaluation of the Washington State Screening, Brief Intervention, and Referral to Treatment Project. *Med Care* 2010;48:18–24. doi:10.1097/MLR.0b013e3181bd498f [PubMed: 19927016]
12. Gentilello LM, Rivara FP, Donovan DM, et al. Alcohol interventions in a trauma center as a means of reducing the risk of injury recurrence. *Ann Surg* 1999;230:473–80; discussion 480–3. <http://www.ncbi.nlm.nih.gov/pubmed/10522717> (accessed 25 Oct 2017). [PubMed: 10522717]
13. Bohnert ASB, Bonar EE, Cunningham R, et al. A pilot randomized clinical trial of an intervention to reduce overdose risk behaviors among emergency department patients at risk for prescription opioid overdose. *Drug Alcohol Depend* 2016;163:40–7. doi:10.1016/j.drugalcdep.2016.03.018 [PubMed: 27062245]
14. Blow FC, Walton MA, Bohnert ASB, et al. A randomized controlled trial of brief interventions to reduce drug use among adults in a low-income urban emergency department: the Healthi ER You study. *Addiction* 2017;112:1395–405. doi:10.1111/add.13773 [PubMed: 28127808]
15. Fisher JD, Fisher WA, Amico KR, et al. An information-motivation-behavioral skills model of adherence to antiretroviral therapy. *Health Psychol* 2006;25:462–73. doi:10.1037/0278-6133.25.4.462
16. Maxwell S, Bigg D, Stanczykiewicz K, et al. Prescribing Naloxone to Actively Injecting Heroin Users. *J Addict Dis* 2006;25:89–96. doi:10.1300/J069v25n03_11
17. Sporer KA, Kral AH. Prescription Naloxone: A Novel Approach to Heroin Overdose Prevention. *Ann Emerg Med* 2007;49:172–7. doi:10.1016/j.annemergmed.2006.05.025 [PubMed: 17141138]
18. Wheeler E, Jones TS, Gilbert MK, et al. Opioid Overdose Prevention Programs Providing Naloxone to Laypersons - United States, 2014. *Morb Mortal Wkly Report Mortal Wkly Rep* 2015;64:631–5. <http://www.ncbi.nlm.nih.gov/pubmed/26086633>
19. WHO. Community management of opioid overdose In: SpringerReference. 2014. doi:10.1007/SpringerReference_44514
20. SAMHSA. Opioid Overdose TOOLKIT Facts for Community Members Five Essential Steps for First Responders Information for Prescribers Safety Advice for Patients & Family Members Recovering from Opioid Overdose. HHS Publ No 2013;:13–4742. <http://www.store.samhsa.gov>.
21. Doe-Simkins M, Walley AY, Epstein A, et al. Saved by the Nose: Bystander-Administered Intranasal Naloxone Hydrochloride for Opioid Overdose. *Am J Public Health* 2009;99:788–91. doi:10.2105/AJPH.2008.146647 [PubMed: 19363214]
22. Green TC, Heimer R, Grau LE. Distinguishing signs of opioid overdose and indication for naloxone: an evaluation of six overdose training and naloxone distribution programs in the United States. *Addiction* 2008;103:979–89. doi:10.1111/j.1360-0443.2008.02182.x [PubMed: 18422830]
23. Lagu T, Anderson BJ, Stein M. Overdoses among friends: Drug users are willing to administer naloxone to others. *J Subst Abuse Treat* 2006;30:129–33. doi:10.1016/j.jsat.2005.05.010 [PubMed: 16490676]
24. Seal KH, Downing M, Kral AH, et al. Attitudes About Prescribing Take-Home Naloxone to Injection Drug Users for the Management of Heroin Overdose: a Survey of Street-Recruited Injectors in the San Francisco Bay Area. *J Urban Heal Bull New York Acad Med* 2003;80:291–301. doi:10.1093/jurban/jtg032

25. Walley AY, Xuan Z, Hackman HH, et al. Opioid overdose rates and implementation of overdose education and nasal naloxone distribution in Massachusetts: interrupted time series analysis. *BMJ* 2013;346:f1744<http://www.ncbi.nlm.nih.gov/pubmed/23372174> (accessed 9 Dec 2015). [PubMed: 23372174]
26. Strang J, Bird SM, Parmar MKB. Take-Home Emergency Naloxone to Prevent Heroin Overdose Deaths after Prison Release: Rationale and Practicalities for the N-ALIVE Randomized Trial. *J Urban Heal* 2013;90:983–96. doi:10.1007/s11524-013-9803-1
27. Moyers TB, Martin T, Manuel JK, et al. Revised Global Scales: Motivational Interviewing Treatment Integrity 3.1.1 (MITI 3.1.1). Published Online First: 2010https://casaa.unm.edu/download/miti3_1.pdf (accessed 25 Oct 2017).
28. Pintilie M *Competing risks: a practical perspective*. West Sussex, England: John Wiley & Sons 2006 https://books.google.com/books?hl=en&lr=&id=ZRh2gl-U6isC&oi=fnd&pg=PR7&dq=Pintilie+M.+Competing+Risks:+A+Practical+Perspective.+West+Sussex,+England:+Wiley%3B+2006.&ots=NdLT548oNj&sig=Jm47omSXqSKRT8K0-Ejb0KEbq_0#v=onepage&q&f=false (accessed 25 Oct 2017).
29. Cleves MA, Gould WW, Gutierrez RG, et al. *An introduction to survival analysis using Stata*. 2008.
30. Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk. *J Am Stat Assoc* 1999;94:496–509. doi:10.1080/01621459.1999.10474144
31. Coffin PO, Sullivan SD. Cost-effectiveness of distributing naloxone to heroin users for lay overdose reversal. *Ann Intern Med* 2013;158:1–9. doi:10.7326/0003-4819-158-1-201301010-00003 [PubMed: 23277895]
32. Strang J, Bird SM, Parmar MKB. Take-home emergency naloxone to prevent heroin overdose deaths after prison release: rationale and practicalities for the N-ALIVE randomized trial. *J Urban Health* 2013;90:983–96. doi:10.1007/s11524-013-9803-1 [PubMed: 23633090]
33. Pierce M, Bird SM, Hickman M, et al. Impact of treatment for opioid dependence on fatal drug-related poisoning: a national cohort study in England. *Addiction* 2016;111:298–308. doi:10.1111/add.13193 [PubMed: 26452239]
34. Larimer ME, Malone DK, Garner MD, et al. Health care and public service use and costs before and after provision of housing for chronically homeless persons with severe alcohol problems. *JAMA* 2009;301:1349–57. doi:10.1001/jama.2009.414 [PubMed: 19336710]
35. Banta-Green CJ, Coffin PO. Commentary on Pierce et al. (2016): Raising the bar of addiction treatment—first do no harm. *Addiction* 2016;111. doi:10.1111/add.13228
36. Polsky D, Glick HA, Yang J, et al. Cost-effectiveness of extended buprenorphine-naloxone treatment for opioid-dependent youth: Data from a randomized trial. *Addiction* 2010;105:1616–24. doi:10.1111/j.1360-0443.2010.03001.x [PubMed: 20626379]
37. Hser YI, Evans E, Huang D, et al. Long-term outcomes after randomization to buprenorphine/naloxone versus methadone in a multi-site trial. *Addiction* 2016;111:695–705. doi:10.1111/add.13238 [PubMed: 26599131]
38. D’Onofrio G, O’Connor PG, Pantalon MV., et al. Emergency Department–Initiated Buprenorphine/Naloxone Treatment for Opioid Dependence. *JAMA* 2015;313:1636. doi:10.1001/jama.2015.3474 [PubMed: 25919527]
39. D’Onofrio G, Chawarski MC, O’Connor PG, et al. Emergency Department-Initiated Buprenorphine for Opioid Dependence with Continuation in Primary Care: Outcomes During and After Intervention. *J Gen Intern Med* 2017;32:660–6. doi:10.1007/s11606-017-3993-2 [PubMed: 28194688]
40. Kourounis G, Richards BDW, Kyprianou E, et al. Opioid substitution therapy: Lowering the treatment thresholds. *Drug Alcohol Depend* 2016;161:1–8. doi:10.1016/j.drugalcdep.2015.12.021 [PubMed: 26832931]
41. Behar E, Santos G-M, Wheeler E, et al. Brief overdose education is sufficient for naloxone distribution to opioid users. *Drug Alcohol Depend* 2015;148:209–12. doi:10.1016/j.drugalcdep.2014.12.009 [PubMed: 25595053]

What is already known on this subject

- Opioid overdoses are increasing rapidly
- Overdose education and take-home-naloxone decrease population rates of overdose

What this study adds

- A brief behavioral-educational intervention combined with an offer of naloxone did not reduce opioid overdose events or health care utilization among a high acuity population seen post-acute care
- Patients seen in acute care settings at elevated risk for overdose had very high rates of subsequent ED visits and hospitalizations and warrant more intensive interventions
- Clinical trials of emergent issues may be impacted by rapid changes in the public health and health care environments




Figure 1.
CONSORT Diagram, Study enrollment, allocation and analysis




Figure 2.
Time to the first opioid overdose event (encounter or death) (n=241)

Table 1.

Participant demographic characteristics, opioid use patterns, overdose history and risk and protective factors

	Intervention		Comparison		Total	
	n=115		n=126		n=241	
Age mean sd	40.2	11.5	42.3	11.5	41.3	11.5
Female	32	28%	37	29%	69	29%
Hispanic	14	12%	21	17%	35	15%
Race						
White	60	53%	67	54%	127	53%
Black	17	15%	14	11%	31	13%
Asian	0	0%	2	2%	2	1%
Hawaiian/Pacific Islander	0	0%	2	2%	2	1%
American Indian/Native Alaskan	6	5%	6	5%	12	5%
Other	13	11%	14	11%	27	11%
More than one race	18	16%	19	15%	37	16%
Housing						
Permanent	29	25%	44	35%	73	30%
Impermanent	19	17%	22	18%	41	17%
Homeless	67	58%	59	47%	126	53%
Education						
<HS	38	33%	35	28%	73	30%
HS	37	32%	45	36%	82	34%
>HS	40	35%	46	37%	86	36%
Relationship						
Not in a relationship	86	75%	83	66%	169	71%
In a relationship	28	25%	42	34%	70	29%
Employment						
Employed	9	8%	9	7%	18	8%
Unemployed	88	77%	88	70%	176	73%
Retired	6	5%	4	3%	10	4%
Unable to work	7	6%	22	18%	29	12%
Other*	5	4%	2	2%	7	3%
Opioid type						
Pharmaceutical only	18	16%	13	10%	31	13%
Heroin & No Opioid agonist therapy	67	58%	74	59%	141	59%
Heroin & Opioid agonist therapy	24	21%	24	19%	48	20%
Opioid agonist therapy no Heroin	6	5%	15	12%	21	9%
Smoke/snort/inject opioids # of days used opioids past 30 days mean sd	24.3	8.4	25.1	8.8	24.7	8.6
Know others with OD education						
Yes	45	39%	62	49%	107	44%
No/Do not know	69	60%	64	51%	133	55%

	Intervention		Comparison		Total	
Know others who have naloxone						
Yes	33	29%	41	33%	74	31%
No/Do not know	82	71%	85	67%	167	69%
Opioid overdose history						
Never overdosed	48	42%	56	44%	104	43%
Overdosed, not past 3 months	48	42%	43	34%	91	38%
Overdosed past 3 months	19	17%	27	21%	46	19%
Used alone past 3 months Always/sometimes use ___ within 2 hours of opioids						
Alcohol	36	31%	54	43%	90	37%
Sedatives/downers	48	42%	61	48%	109	45%
More than one kind of opioid	53	46%	54	43%	107	44%
Uppers- cocaine, methamphetamine, pharmaceutical	57	50%	72	57%	129	54%

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

Opioid overdose events and censoring by randomization status

Group	Total N	Opioid overdose event status by data source: n (percent)					Mean days to first overdose event ^{*****}
		First opioid overdose in UW Medicine [*]	First opioid overdose in CHARS ^{**}	First opioid overdose fatality in DOH ^{***}	Nonoverdose fatality in DOH ^{***}	No event prior to censoring at study end ^{****}	
Comparison group	126	21 (16.7%)	6 (4.8%)	6 (4.8%)	7 (5.6%)	86 (68.3%)	836
Intervention group	115	18 (15.7%)	4 (3.5%)	2 (1.7%)	8 (7.0%)	83 (72.2%)	870
Combined	241	39 (16.2%)	10 (4.2%)	8 (3.3%)	15 (6.2%)	169 (70.1%)	852

^{*} Local inpatient admission or emergency department visit

^{**} Statewide hospital discharge for inpatient admission or observation visit

^{***} Death certificate data

^{****} Available follow-up time: 272 to 1064 days

^{*****} Mean is underestimated because the largest observed analysis time is censored.

Table 3.

Annual local health care utilization (all-cause) after study enrollment by randomization status

Group	Emergency department visits*				Inpatient admissions			
	Median annual rate	IQR	Mean annual rate**	95% CI	Median annual rate	IQR	Mean annual rate**	95% CI
Comparison group	2.72	5.49	4.85	3.96–5.93	0.43	1.50	1.05	0.80–1.39
Intervention group	2.42	7.35	4.96	4.04–6.10	0.39	1.80	1.29	0.95–1.76
Combined	2.57	6.93	4.90	4.25–5.66	0.41	1.52	1.17	0.95–1.44

* Regardless of discharge status (may have resulted in inpatient admission)

** Mean rate per person-year calculated using negative binomial regression with time at-risk exposure variable for available follow-up time (272 to 1064 days, censored at death or December 31, 2015). Note: Rate differences between the intervention and comparison groups were not statistically significant.