



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Mobile phone text messaging and app-based interventions for smoking cessation (Review)

Whittaker R, McRobbie H, Bullen C, Rodgers A, Gu Y, Dobson R

Whittaker R, McRobbie H, Bullen C, Rodgers A, Gu Y, Dobson R.  
Mobile phone text messaging and app-based interventions for smoking cessation.  
*Cochrane Database of Systematic Reviews* 2019, Issue 10. Art. No.: CD006611.  
DOI: [10.1002/14651858.CD006611.pub5](https://doi.org/10.1002/14651858.CD006611.pub5).

[www.cochranelibrary.com](http://www.cochranelibrary.com)

## TABLE OF CONTENTS

HEADER .....	1
ABSTRACT .....	1
PLAIN LANGUAGE SUMMARY .....	2
SUMMARY OF FINDINGS .....	3
BACKGROUND .....	6
OBJECTIVES .....	6
METHODS .....	6
RESULTS .....	8
Figure 1. ....	9
Figure 2. ....	13
DISCUSSION .....	15
Figure 3. ....	17
AUTHORS' CONCLUSIONS .....	17
ACKNOWLEDGEMENTS .....	17
REFERENCES .....	19
CHARACTERISTICS OF STUDIES .....	28
DATA AND ANALYSES .....	73
Analysis 1.1. Comparison 1 Text messaging versus minimal smoking cessation support, Outcome 1 Long-term abstinence (all randomised)). ....	74
Analysis 1.2. Comparison 1 Text messaging versus minimal smoking cessation support, Outcome 2 Long-term abstinence (complete case). ....	74
Analysis 2.1. Comparison 2 Text messaging versus other smoking cessation intervention, Outcome 1 Long-term abstinence (all randomised). ....	75
Analysis 2.2. Comparison 2 Text messaging versus other smoking cessation intervention, Outcome 2 Long-term abstinence (complete case). ....	76
Analysis 3.1. Comparison 3 Text messaging + other smoking cessation support versus other smoking cessation support alone, Outcome 1 Long-term abstinence (all randomised). ....	76
Analysis 3.2. Comparison 3 Text messaging + other smoking cessation support versus other smoking cessation support alone, Outcome 2 Long-term abstinence (complete case). ....	77
Analysis 4.1. Comparison 4 High-frequency versus low-frequency text messaging, Outcome 1 Long-term abstinence (all randomised). ....	77
Analysis 4.2. Comparison 4 High-frequency versus low-frequency text messaging, Outcome 2 Long-term abstinence (complete case). ....	78
Analysis 5.1. Comparison 5 Smartphone app versus lower-intensity smoking cessation support, Outcome 1 Long-term abstinence (all randomised). ....	78
Analysis 5.2. Comparison 5 Smartphone app versus lower-intensity smoking cessation support, Outcome 2 Long-term abstinence (complete case). ....	79
Analysis 6.1. Comparison 6 CO monitoring + contingency management versus smoking cessation support, Outcome 1 Long-term abstinence (all randomised). ....	80
Analysis 6.2. Comparison 6 CO monitoring + contingency management versus smoking cessation support, Outcome 2 Long-term abstinence (complete case). ....	80
Analysis 7.1. Comparison 7 Smartphone app + text messaging versus web-based intervention, Outcome 1 Long-term abstinence (all randomised). ....	81
Analysis 7.2. Comparison 7 Smartphone app + text messaging versus web-based intervention, Outcome 2 Long-term abstinence (complete case). ....	81
APPENDICES .....	81
WHAT'S NEW .....	81
HISTORY .....	82
CONTRIBUTIONS OF AUTHORS .....	82
DECLARATIONS OF INTEREST .....	82
SOURCES OF SUPPORT .....	83
DIFFERENCES BETWEEN PROTOCOL AND REVIEW .....	83
INDEX TERMS .....	83

[Intervention Review]

# Mobile phone text messaging and app-based interventions for smoking cessation

Robyn Whittaker<sup>1</sup>, Hayden McRobbie<sup>2</sup>, Chris Bullen<sup>1</sup>, Anthony Rodgers<sup>3</sup>, Yulong Gu<sup>4</sup>, Rosie Dobson<sup>1</sup>

<sup>1</sup>National Institute for Health Innovation, University of Auckland, Auckland, New Zealand. <sup>2</sup>National Drug and Alcohol Research Centre, University of New South Wales, Sydney, Australia. <sup>3</sup>The George Institute for Public Health, Sydney, Australia. <sup>4</sup>School of Health Sciences, Stockton University, Galloway, New Jersey, USA

**Contact address:** Robyn Whittaker, National Institute for Health Innovation, University of Auckland, Tamaki Campus, Private Bag 92019, Auckland, 1142, New Zealand. [r.whittaker@auckland.ac.nz](mailto:r.whittaker@auckland.ac.nz).

**Editorial group:** Cochrane Tobacco Addiction Group.

**Publication status and date:** New search for studies and content updated (no change to conclusions), published in Issue 10, 2019.

**Citation:** Whittaker R, McRobbie H, Bullen C, Rodgers A, Gu Y, Dobson R. Mobile phone text messaging and app-based interventions for smoking cessation. *Cochrane Database of Systematic Reviews* 2019, Issue 10. Art. No.: CD006611. DOI: [10.1002/14651858.CD006611.pub5](https://doi.org/10.1002/14651858.CD006611.pub5).

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Mobile phone-based smoking cessation support (mCessation) offers the opportunity to provide behavioural support to those who cannot or do not want face-to-face support. In addition, mCessation can be automated and therefore provided affordably even in resource-poor settings. This is an update of a Cochrane Review first published in 2006, and previously updated in 2009 and 2012.

### Objectives

To determine whether mobile phone-based smoking cessation interventions increase smoking cessation rates in people who smoke.

### Search methods

For this update, we searched the Cochrane Tobacco Addiction Group's Specialised Register, along with [clinicaltrials.gov](http://clinicaltrials.gov) and the ICTRP. The date of the most recent searches was 29 October 2018.

### Selection criteria

Participants were smokers of any age. Eligible interventions were those testing any type of predominantly mobile phone-based programme (such as text messages (or smartphone app) for smoking cessation. We included randomised controlled trials with smoking cessation outcomes reported at at least six-month follow-up.

### Data collection and analysis

We used standard methodological procedures described in the *Cochrane Handbook for Systematic Reviews of Interventions*. We performed both study eligibility checks and data extraction in duplicate. We performed meta-analyses of the most stringent measures of abstinence at six months' follow-up or longer, using a Mantel-Haenszel random-effects method, pooling studies with similar interventions and similar comparators to calculate risk ratios (RR) and their corresponding 95% confidence intervals (CI). We conducted analyses including all randomised (with dropouts counted as still smoking) and complete cases only.

### Main results

This review includes 26 studies (33,849 participants). Overall, we judged 13 studies to be at low risk of bias, three at high risk, and the remainder at unclear risk. Settings and recruitment procedures varied across studies, but most studies were conducted in high-income countries. There was moderate-certainty evidence, limited by inconsistency, that automated text messaging interventions were more effective than minimal smoking cessation support (RR 1.54, 95% CI 1.19 to 2.00;  $I^2 = 71%$ ; 13 studies, 14,133 participants). There was

also moderate-certainty evidence, limited by imprecision, that text messaging added to other smoking cessation interventions was more effective than the other smoking cessation interventions alone (RR 1.59, 95% CI 1.09 to 2.33;  $I^2 = 0\%$ , 4 studies, 997 participants). Two studies comparing text messaging with other smoking cessation interventions, and three studies comparing high- and low-intensity messaging, did not show significant differences between groups (RR 0.92 95% CI 0.61 to 1.40;  $I^2 = 27\%$ ; 2 studies, 2238 participants; and RR 1.00, 95% CI 0.95 to 1.06;  $I^2 = 0\%$ , 3 studies, 12,985 participants, respectively) but confidence intervals were wide in the former comparison. Five studies compared a smoking cessation smartphone app with lower-intensity smoking cessation support (either a lower-intensity app or non-app minimal support). We pooled the evidence and deemed it to be of very low certainty due to inconsistency and serious imprecision. It provided no evidence that smartphone apps improved the likelihood of smoking cessation (RR 1.00, 95% CI 0.66 to 1.52;  $I^2 = 59\%$ ; 5 studies, 3079 participants). Other smartphone apps tested differed from the apps included in the analysis, as two used contingency management and one combined text messaging with an app, and so we did not pool them. Using complete case data as opposed to using data from all participants randomised did not substantially alter the findings.

### Authors' conclusions

There is moderate-certainty evidence that automated text message-based smoking cessation interventions result in greater quit rates than minimal smoking cessation support. There is moderate-certainty evidence of the benefit of text messaging interventions in addition to other smoking cessation support in comparison with that smoking cessation support alone. The evidence comparing smartphone apps with less intensive support was of very low certainty, and more randomised controlled trials are needed to test these interventions.

## PLAIN LANGUAGE SUMMARY

### Can programmes delivered by mobile phones help people to stop smoking?

#### Background

Tobacco smoking is a leading cause of preventable death. Mobile phones can be used to support people who want to quit smoking. In this review, we have focused on programmes that use text messages or smartphone apps to do so.

#### Search date

We searched for published and unpublished studies in October 2018.

#### Study characteristics

We included 26 randomised controlled studies (involving over 33,000 people) that compared smoking quit rates in people who received text messages or smartphone apps to help them quit, with people who did not receive these programmes. We were interested in studies that measured smoking for six months or longer.

#### Key results

We found that text messaging programmes may be effective in supporting people to quit, increasing quit rates by 50% to 60%. This was the case when they were compared to minimal support or were tested as an addition to other forms of stop-smoking support. There was not enough evidence to determine the effect of smartphone apps.

#### Quality and completeness of the evidence

Most of the studies were of high quality, although three studies had high drop out rates. We are moderately confident in the results of the text messaging interventions, but there were some issues with unexplained differences between study findings and for some comparisons there was not much data. We have low confidence in the results concerning smartphone apps, and more studies are needed in this field.

## SUMMARY OF FINDINGS

### Summary of findings for the main comparison. Text messaging compared to minimal support for smoking cessation

#### Text messaging compared to minimal support for smoking cessation

**Patient or population:** people who smoke  
**Setting:** community  
**Intervention:** text messaging  
**Comparison:** minimal smoking cessation support

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with minimal SC	Risk with text messaging				
<b>Long-term abstinence</b> (all randomised)	Study population		RR 1.54 (1.19 to 2.00)	14,133 (13 RCTs)	⊕⊕⊕⊖ <b>Moderate</b> <sup>a</sup>	
Measured with self-report and biochemical validation at 6 to 12 months	6 per 100	9 per 100 (7 to 11)				

\***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**CI:** confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio; **SC:** smoking cessation

#### GRADE Working Group grades of evidence

**High certainty:** we are very confident that the true effect lies close to that of the estimate of the effect.

**Moderate certainty:** we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

**Low certainty:** our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

**Very low certainty:** we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

<sup>a</sup>Downgraded one level due to inconsistency: substantial unexplained heterogeneity ( $I^2 = 71\%$ ).

### Summary of findings 2. Text messaging in addition to other smoking cessation support

#### Text messaging in addition to other smoking cessation support compared to other smoking cessation support alone for smoking cessation

**Patient or population:** people who smoke  
**Setting:** community  
**Intervention:** text messaging + other smoking cessation support

**Comparison:** other smoking cessation support alone

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with other SC support alone	Risk with text messaging + other SC support				
<b>Long-term abstinence</b> (all randomised)	Study population		RR 1.59 (1.09 to 2.33)	997 (4 RCTs)	⊕⊕⊕⊖ <b>Moderate</b> <sup>a</sup>	
Measured as self-reported and biochemical validation at 6 to 12 months	8 per 100	12 per 100 (9 to 18)				

\***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**CI:** confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio; **SC:** smoking cessation

#### GRADE Working Group grades of evidence

**High certainty:** we are very confident that the true effect lies close to that of the estimate of the effect.

**Moderate certainty:** we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

**Low certainty:** our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

**Very low certainty:** we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

<sup>a</sup>Downgraded one level due to imprecision: fewer than 300 events overall.

### Summary of findings 3. Smartphone app compared to lower-intensity support for smoking cessation

#### Smartphone app compared to lower-intensity support for smoking cessation

**Patient or population:** people who smoke

**Setting:** community

**Intervention:** smartphone app

**Comparison:** lower-intensity smoking cessation support

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with lower intensity SC support	Risk with Smartphone app SC support				

<b>Long-term abstinence</b> (all randomised)  Measured with self-report and biochemical validation at 6 months	Study population		RR 1.00 (0.66 to 1.52)	3079 (5 RCTs)	⊕○○○ <b>Very low</b> <sup>a,b</sup>
	8 per 100	8 per 100 (5 to 12)			

\***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**CI:** confidence interval; **RCT:** randomised controlled trial; **RR:** risk ratio; **SC:** smoking cessation

#### GRADE Working Group grades of evidence

**High certainty:** we are very confident that the true effect lies close to that of the estimate of the effect.

**Moderate certainty:** we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

**Low certainty:** our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

**Very low certainty:** we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

<sup>a</sup>Downgraded one level due to inconsistency: considerable unexplained statistical heterogeneity ( $I^2 = 59\%$ ).

<sup>b</sup>Downgraded two levels due to imprecision: confidence intervals encompass both clinically significant harm and clinically significant benefit.

## BACKGROUND

### Description of the condition

Tobacco remains one of the most important risk factors for poor health across the globe (IHME 2018). Many countries are looking for sustainable options for the provision of smoking cessation support on a large scale.

### Description of the intervention

'mHealth' describes the use of mobile communications technologies and mobile phones to support health care. In this review, we are specifically interested in the use of text messaging and smartphone applications (apps) to support smoking cessation.

### How the intervention might work

The benefits of mobile phone-based smoking cessation support (mCessation) interventions are: the ease of use anywhere at any time; cost-effective delivery and scalability to large populations, regardless of location; the ability to tailor messages to key user characteristics (such as age, sex, ethnicity); the ability to send time-sensitive messages with an 'always on' device; the provision of content that can distract the user from cravings; and the ability to link the user with others for social support.

A key benefit of the use of mobile phones for health programmes is their widespread uptake in those areas where health services are not easily accessible or used. In 2018, the number of mobile phone subscriptions globally topped 8 billion, with the developing world now having more mobile phone subscriptions than population (population penetration of 102%; ITU 2018). There is evidence to suggest that people from lower socioeconomic groups may prefer mCessation interventions due to the greater feeling of control associated with the ability to decide when and where they engage with messages, and the perception of around-the-clock support (Boland 2017). Focusing mCessation efforts on the populations in greatest need, could help to address the health inequalities that come about from high use of tobacco and lack of accessible health promotion and prevention services in low-resource settings globally.

Furthermore, initial research suggests that the use of text messaging for smoking cessation is cost effective. Guerriero 2013 found that the cost of text message-based support was GBP 278 per quitter. When the future health service costs saved (as a result of smoking cessation) were included, with 0.5 quality-adjusted life years (QALYs) gained per quitter, text-based support was considered to be cost saving.

### Why it is important to do this review

Smartphones (mobile phones with a computer operating system) are fast becoming the computer of choice, or at least the most accessible computer, in many countries. According to the International Telecommunications Union only 36.3% of low- and middle-income countries has a computer in the household, but 61% have mobile broadband subscriptions (allowing mobile phones to access to the Internet; ITU 2018). Therefore, it was important to update this review to include studies on the effectiveness of smartphone apps, as well as text messaging interventions, for smoking cessation.

## OBJECTIVES

To determine whether mobile phone-based smoking cessation interventions increase smoking cessation rates in people who smoke.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

Randomised or quasi-randomised trials. Cluster-randomised trials were eligible for inclusion.

#### Types of participants

People who smoked at study enrolment.

#### Types of interventions

We included studies that examined any intervention that could be considered predominantly a mobile phone-based programme (such as text messaging or smartphone apps) for smoking cessation. We excluded interventions where mobile phones were seen as an adjunct to a predominantly face-to-face or Internet programme, such as to remind participants of appointments, or where the effects of the various components of a multi-faceted programme could not be separated. We also excluded interventions that could be performed via any type of telephone such as telephone counselling. We did not exclude any studies based on comparator, but instead grouped studies by comparators in the analyses.

#### Types of outcome measures

The primary outcome was smoking abstinence at longest follow-up, and at least six months from baseline. Where multiple measures were available, we preferred sustained abstinence to point prevalence abstinence, and biochemically validated results to self-report.

There is no obvious risk of adverse events for text messaging or smartphone app interventions, and so we have not included this as an outcome in this review.

### Search methods for identification of studies

For the present update of the review, we searched the Cochrane Tobacco Addiction Group's Specialised Register on 29 October 2018 using the terms 'mobile phone', 'cell phone', 'txt', 'pvt', 'sms', or 'mms' in the title, abstract or keyword fields. The Specialised Register includes reports of possible controlled trials of smoking cessation interventions identified from sensitive searches of databases. At the time of the search, the Register included the following results of searches

- Cochrane Central Register of Controlled trials (CENTRAL; 2018, Issue 1)
- MEDLINE (via Ovid, to 26 October 2018)
- Embase (via Ovid, to 28 October 2018)
- PsycINFO (via Ovid; to 22 October 2018)

See the [Cochrane Tobacco Addiction Group website](#) for full search strategies and a list of other resources searched. We also searched the World Health Organization International Clinical Trials



Registry Platform (WHO ICTRP; [apps.who.int/trialsearch/](https://apps.who.int/trialsearch/)) and [ClinicalTrials.gov](https://clinicaltrials.gov) trials registers for ongoing or recently completed studies. We searched through the reference lists of identified studies for any additional eligible studies and attempted to contact the authors of ongoing studies.

We placed no restrictions on publication language or date.

## Data collection and analysis

### Selection of studies

The Cochrane Tobacco Addiction Group's Information Specialist ran the searches and provided the results. Two review authors (YG, HM) independently pre-screened the titles and abstracts of records identified in duplicate to exclude reports that had no relevance to the topic and to provide a list of potentially relevant citations. A third reviewer (CB) resolved any differences in initial screening. Two review authors (from RW, YG, CB, RD) independently reviewed full-text manuscripts in duplicate for the final eligibility screen.

We resolved any disagreements by discussion or by obtaining further information through contacting study authors. We recorded reasons for exclusion of studies in the [Characteristics of excluded studies](#) table. We contacted authors of unpublished, registered studies, which could potentially have been completed, to determine ongoing status or to request unpublished data.

### Data extraction and management

We extracted the following methodological details from the included study reports and presented them in the [Characteristics of included studies](#) table. Two review authors (from RW, YG, RD, CB, HM) independently extracted data using the standardised [Covidence](#) data extraction form. A third review author provided a review of the quality assessment and a consensus check.

- Funding source
- Authors' declarations of interest
- Country and context of the study
- Study design
- Number of participants
- Age and other relevant recorded characteristics of study participants
- Inclusion criteria
- Exclusion criteria
- Intervention details
- Control details
- Definition of abstinence outcome
- Smoking cessation rates at six months (self-reported abstinence or biochemically verified abstinence, or both)
- Smoking cessation rates at final follow-up (if follow-up greater than six months and where these data were available)

### Assessment of risk of bias in included studies

Two review authors (from RW, YG, RD, CB, HM) independently assessed the risk of bias for included studies, based on the guidance of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2017](#)), and the Cochrane Tobacco Addiction Group. For each study, we assessed the following domains.

- Random sequence generation

- Allocation concealment
- Blinding of outcome assessment
- Incomplete outcome data
- Other sources of bias

Specific 'Risk of bias' guidance developed by the Cochrane Tobacco Addiction Group to assess smoking cessation studies states that performance bias (relating to the blinding of participants and providers) should not be assessed for behavioural interventions, as it is impossible to blind people to these types of interventions. We graded detection bias as low where there was biochemical verification of abstinence, or where abstinence was self-reported with no difference in face-to-face contact between control and intervention arms. We considered bias due to incomplete outcome as low risk where numbers lost to follow-up were clearly reported for each group, the overall loss was not greater than 50%, and the difference between groups was not greater than 20%, or sensitivity analysis showed that the direction of effect was not sensitive to different imputation methods for loss to follow-up.

Each review author recorded information in study reports relevant to each relevant domain and then judged each domain as either at low, high, or unclear risk of bias. We resolved disagreements through discussion with a third review author.

### Measures of treatment effect

We recorded the information below.

- Smoking cessation rates at six months or longer using the most stringent measure available
- Biochemically verified abstinence, where available

We calculated risk ratios (RR) and 95% confidence intervals (CI) for the smoking cessation outcome for each included study. We calculated outcomes on an intention-to-treat basis, including all participants randomised to a trial arm and assuming that participants lost to follow-up had continued to smoke or relapsed.

### Dealing with missing data

If we found any important study characteristics or outcome data to be missing, we followed up with study authors where possible.

### Assessment of heterogeneity

In order to assess whether it was appropriate to pool studies and conduct meta-analyses we assessed the characteristics of included studies to identify any clinical or methodological heterogeneity. Where we deemed studies homogeneous enough to be combined meaningfully, we conducted a meta-analysis, and we assessed statistical heterogeneity using the  $I^2$  statistic; we deemed an  $I^2$  value greater than 50% to indicate substantial heterogeneity ([Higgins 2003](#)).

### Assessment of reporting biases

We planned to use funnel plots to assess reporting bias for any comparisons where we identified and analysed abstinence rates from at least 10 studies. Only the 'text messaging versus minimal smoking cessation support' comparison met this criteria in this review; therefore a funnel plot was generated for this comparison only. Funnel plots illustrate the relationship between the effect estimates from individual studies against their size or precision. The

greater the degree of asymmetry, the greater the risk of reporting bias.

### Data synthesis

We conducted meta-analyses of the included studies, using the Mantel-Haenszel random-effects method to pool RRs and 95% CIs calculated for the smoking abstinence outcome, across the following comparisons.

- Text messaging versus minimal smoking cessation support (including standard self-help materials, as is standard practice in the Cochrane Tobacco Addiction Group)
- Text messaging in addition to another form of smoking cessation support
- Text messaging versus other smoking cessation support
- Higher- versus lower-frequency text messaging
- Smartphone app versus less intensive smoking cessation support

Where studies had multiple intervention arms relevant to a single meta-analysis, we split control arm data to avoid double-counting.

### Subgroup analysis and investigation of heterogeneity

We carried out the following subgroup analyses.

- We split the 'smartphone app versus less intensive smoking cessation support' comparison into two subgroups to reflect the different comparators used across studies; either minimal non-app smoking cessation support (e.g. self-help materials, information on existing stop-smoking services) or a less intensive smartphone app.

### Sensitivity analysis

We conducted the following sensitivity analyses.

- We calculated pooled RRs and 95% CIs for all analyses using complete case data to calculate quit rates. People may drop out of studies for reasons other than still smoking, and these reasons may differ between groups. For example, people who successfully stop smoking may withdraw from receiving an intervention if the text messages remind them of smoking. Therefore, this analysis tests whether assuming that all people lost to follow-up are smoking (as in our primary analyses of all participants randomised) is potentially biasing our results.
- Removing any studies judged to be at high risk of bias from all comparisons
- Removing the only cluster-RCT (Haug 2013), as information was not available to adjust for any potential clustering effect
- Removing the two studies carried out in a pregnant (Abroms 2017), or postnatal population (Yu 2017), as these populations differ substantially from those recruited in the other studies.

### 'Summary of findings' tables

Following standard Cochrane methods (Schünemann 2017), we created a 'Summary of findings' table for the primary outcome (smoking abstinence), for the following comparisons.

- Text messaging versus minimal smoking cessation support
- Text messaging in addition to other smoking cessation support
- Smartphone app versus less intensive smoking cessation support

Also following standard Cochrane methodology (Schünemann 2017), we used the five GRADE considerations (risk of bias, inconsistency, imprecision, indirectness and publication bias) to assess the certainty of the body of evidence for the abstinence outcome for each comparison, and to draw conclusions about the certainty of evidence within the text of the review.

## RESULTS

### Description of studies


See [Characteristics of included studies](#); [Characteristics of excluded studies](#); and [Characteristics of ongoing studies](#) for further details.

### Results of the search


The previous version of this review (Whittaker 2016), included 12 studies (Abroms 2014; Bock 2013; Borland 2013; Ferguson 2015; Free 2009; Free 2011; Gritz 2013; Haug 2013; Naughton 2014; Rodgers 2005; Tseng 2017; Whittaker 2011). Gritz 2013 was excluded at this update, as their intervention (telephone counselling and help line) was significantly different from the other interventions included in this review. A telephone help line intervention does not need to be carried out using a mobile phone specifically. Therefore, 11 of the previously included studies were included at this update, as well as one previously 'ongoing' study that was changed to 'included' as the study is now complete and data was available (Danaher 2019).

For this update of our review, the new literature search identified 370 studies (Figure 1). Many were duplicates, or unrelated and were immediately excluded at the title and abstract screening phase. We screened the full-text of 71 reports of 62 studies, excluding 16 studies, and leaving 14 new studies eligible for inclusion at this update (Abroms 2017; Alessi 2017; Augustson 2017; Baskerville 2018; BinDhim 2018; Chan 2015; Cobos-Campos 2017; Garrison 2018; Herbec 2019; Liao 2018; Peiris 2019; Squiers 2017; Wilson 2016; Yu 2017). Data were supplied by the authors for two studies (Danaher 2019; Herbec 2019). Reasons for excluding studies included: intervention that was not predominantly a mobile phone programme; not a randomised controlled trial; relapse prevention only; or no abstinence outcome measured at  $\geq 6$  months follow-up (see [Characteristics of excluded studies](#) table for further details).

**Figure 1. Study flow diagram for this update**



**Figure 1. (Continued)**



We also identified 32 ongoing studies at this update. When added to the previously identified ongoing studies there was a total of 34 ongoing studies (for further details see the [Characteristics of ongoing studies](#) table).

**Included studies**

**Context and participants**

The settings and recruitment methods, and therefore the participants, varied considerably across studies. Where previously this review had included only studies from a small range of high-income countries, the new studies included in this update provided greater variation in settings, including China ([Augustson 2017](#); [Chan 2015](#); [Liao 2018](#)).

[Bock 2013](#) (USA) found usual in-person recruitment methods slow and shifted to online recruitment methods during the study. [Baskerville 2018](#) (Canada), [Borland 2013](#) (Australia), [Danaher 2019](#) (USA), [Garrison 2018](#) (USA), [Squiers 2017](#) (USA), [Herbec 2019](#) (UK), and [Abroms 2014](#) (USA) also used online recruitment via Internet advertisements. In [Abroms 2014](#) this initially led to some fraudulent participants who were discovered and disqualified, and extra procedures were put in place to prevent this from happening again. [Free 2009](#) and [Free 2011](#) recruited via advertisements at UK primary care centres, smoking cessation clinics, pharmacies, newspapers, websites, bus billboards and on the radio in the UK, and [Liao 2018](#) used similar advertising methods in China. [Rodgers 2005](#) also used direct advertising via websites, email, and posters at tertiary institutions across New Zealand. Similarly [Whittaker 2011](#) (New Zealand) used a wide range of advertising media, including Māori-specific media, and targeted young people. [Alessi 2017](#) recruited through email, flyers, and print advertisements and [Ferguson 2015](#) (Australia) used advertisements in papers, radio, and Facebook. [Abroms 2017](#) was embedded in the Text4Baby text message (three messages a week) health information programme for pregnant women in the USA. Women who had smoked at least one puff in the past two weeks were eligible to also receive Quit4Baby text messages (between one to eight messages a day) to support smoking cessation. [Augustson 2017](#) recruited through Nokia Life Tools, a service providing more than 100 million users with tools pre-installed on their Nokia mobile phones, in urban and rural areas of China's Zhejiang, Heilongjiang and Shaanxi provinces. [BinDhim 2018](#) recruited through the Apple App Store in several countries (Australia, Singapore, UK, USA). Participants were advised that by downloading the app they would be participating in a study. [Chan 2015](#) recruited through a Quit and Win competition in Hong Kong that was promoted in shopping malls and other public areas. [Wilson 2016](#) mailed letters to potential participants

in the US Veterans Administration health system. [Naughton 2014](#) was set in primary care practices in the UK with trained smoking cessation advisors providing smoking cessation advice; [Cobos-Campos 2017](#) in two health clinics in Spain with health advice provided by a doctor or nurse; and [Tseng 2017](#) in large urban HIV clinics. [Haug 2013](#) recruited in vocational schools and differed from the other studies by allowing the inclusion of occasional smokers (at least four cigarettes in the past month or at least one in the preceding week). [Peiris 2019](#) (Australia) recruited via an Aboriginal Community Controlled Health Service, a regional community event, and the New South Wales Government telephone coaching service. [Yu 2017](#) recruited in maternal-child health centres in China after asking mothers about household second-hand smoke exposure. The intervention included messages on both the harms of second-hand smoke (to the mother and her husband) and additional messages to the husband to encourage quitting.

Four studies deliberately targeted young adults ([Baskerville 2018](#) in Canada, [Haug 2013](#) in Switzerland; [Squiers 2017](#) in USA; [Whittaker 2011](#) in New Zealand). Most studies had similar proportions of men and women or slightly more women than men. The exceptions were [Abroms 2017](#), as the intervention was targeted at pregnant women (100% women), [Wilson 2016](#), which recruited 89% male veterans, and the studies in China, where the rates of smoking in women are low ([Chan 2015](#) > 80% men, [Liao 2018](#) 94.6% men, [Yu 2017](#) 100% men).

**Intervention programmes**

**Text messaging**

All studies tested automated text messaging interventions. Eighteen of the included studies used text messaging (SMS) as a central component of the intervention ([Abroms 2014](#); [Abroms 2017](#); [Augustson 2017](#); [Bock 2013](#); [Borland 2013](#); [Chan 2015](#); [Cobos-Campos 2017](#); [Ferguson 2015](#); [Free 2009](#); [Free 2011](#); [Haug 2013](#); [Liao 2018](#); [Naughton 2014](#); [Rodgers 2005](#); [Tseng 2017](#); [Squiers 2017](#); [Whittaker 2011](#); [Yu 2017](#)). [Whittaker 2011](#) sent text messages containing links to theoretically driven video messages from 'ordinary' role models coping with quitting. Several studies paired text messages with in-person visits or assessments ([Bock 2013](#); [Cobos-Campos 2017](#); [Haug 2013](#); [Naughton 2014](#)).

The text message interventions varied in length from one week ([Chan 2015](#)), to five weeks ([Ferguson 2015](#)), six weeks ([Augustson 2017](#); [Yu 2017](#)), eight weeks ([Bock 2013](#); [Squiers 2017](#)), three months ([Abroms 2017](#); [Haug 2013](#); [Naughton 2014](#); [Tseng 2017](#)), and six months ([Abroms 2014](#); [Cobos-Campos 2017](#); [Free 2009](#); [Free](#)

2011; Liao 2018; Rodgers 2005; Whittaker 2011), or were variable (Borland 2013).

Eight studies did not state that text messages were tailored to the individual (Abroms 2017; Augustson 2017; Chan 2015; Cobos-Campos 2017; Liao 2018; Tseng 2017; Squiers 2017; Yu 2017). In other studies using text messages, the degree of individual tailoring varied:

- Abroms 2014 tailored messages to include first name, quit date, top three reasons for quitting, money saved by quitting, and use of quit-smoking medications;
- Bock 2013 and Haug 2013 tailored messages to the stage of readiness to quit;
- Borland 2013's programme could be interacted with by reporting changes in smoking behavior (e.g. a quit attempt, relapse), so that appropriate stage-specific messages could be sent;
- Ferguson 2015 tailored their intervention text messages to contain advice and encouragement tailored to participants' current quit status (preparing to quit, first week of the quit attempt, second week of attempt etc.)
- Free 2009 and Free 2011 tailored the messages to information collected at baseline about the individual;
- Naughton 2014 individually tailored messages using 24 items from the iQuit questionnaire and information on smoking status at three and seven weeks;
- Rodgers 2005 matched participant characteristics to messages by keyword to create an individualised programme;
- Whittaker 2011's participants selected the role model from whom they wished to receive messages.

A number of text messaging interventions included interactive components such as:

- the ability to text for more support in the instance of cravings or lapses (Abroms 2014; Bock 2013; Free 2011; Liao 2018; Naughton 2014; Rodgers 2005);
- an optional Quit Buddy in Rodgers 2005 and Free 2011;
- a Quit support network in Bock 2013;
- polls and quizzes (Rodgers 2005);
- regular checking in on smoking status (Haug 2013).

Borland 2013 was the only study to include some degree of choice. Participants received offers of support via a personalised tailored Internet programme, a text message programme, both programmes, a choice of all three, or a minimal control. For the purposes of meta-analyses, we compared the text message group with the control group.

Some of the included interventions were somewhat related to each other. The text messaging intervention in Rodgers 2005 was developed in New Zealand, and later adapted and tested in a UK pilot study (Free 2009), and then a large randomised controlled study (Free 2011). The intervention in Abroms 2017 was developed for pregnant women from the same group's previous intervention for adult smokers in Abroms 2014. The Augustson 2017 intervention in China was adapted from the smoke-free text programme that was evaluated in Squiers 2017. For further details of the messaging interventions across individual studies see the [Characteristics of included studies](#) table.

The control conditions used in the text message studies could be categorised into four groups.

- Minimal smoking cessation support (13 studies): the control programmes across the studies in this category varied from no smoking cessation support (Haug 2013; Yu 2017), to non-smoking-related text messages sent two-weekly (Free 2009; Free 2011; Rodgers 2005; Whittaker 2011), or weekly (Liao 2018), to written or Internet untailored materials (Abroms 2014; Chan 2015; Ferguson 2015), to links to smoking cessation support (Borland 2013; Rodgers 2005), or regular general health advice provided by a clinician (Cobos-Campos 2017). Abroms 2017's control group participants received standard non-smoking-related Text4Baby text messages (three a week) without the extra smoking cessation-related Quit4Baby text messages.
- Another form of smoking cessation support (matched to support received by the intervention group, but without the text messaging intervention; four studies): support varied across studies and included a single session of smoking cessation counselling plus non-smoking-related text messages (Bock 2013); smoking cessation behavioural support and pharmacotherapy (Naughton 2014), and behavioural support and pharmacotherapy (Tseng 2017). Participants in the comparison arm of Yu 2017 received in-person counselling and materials on establishing a smoke-free home.
- Another form of smoking cessation support (not matched in the intervention arm; two studies): an Internet-based interactive smoking cessation programme (Borland 2013), and a five-minute smoking cessation counselling session (Chan 2015).
- Higher- versus lower-frequency text messaging. Three studies examined the effect of higher- versus lower-frequency text messages (Augustson 2017; Liao 2018; Squiers 2017). In Augustson 2017 this was comparing 91 messages over six weeks (three a day initially, followed by two a day, then one a day), with one text message a week for six weeks. In Liao 2018 this was three to five messages per day compared with three to five messages per week. Squiers 2017 compared smoking assessment and quit date messages only, with those messages plus motivational preparatory messages for two weeks prior to quitting, and with all of those messages plus six weeks of follow-up post-quit messages.

#### Smartphone apps

Five studies tested the effectiveness of smoking cessation smartphone apps alone (Baskerville 2018; BinDhim 2018; Garrison 2018; Herbec 2019; Peiris 2019). These apps varied considerably in intervention content and components. The app in Baskerville 2018 was described as comprehensive and evidence-informed, including components such as a quit plan, contingency reinforcement, a link to an online Facebook community, supportive messages through the app, web-based distraction, information and performance feedback, access to evidence-based cessation services. BinDhim 2018 described their app as a decision aid (based on the Ottawa Decision Support Framework drawing from a number of psychological and behavioural theories) with additional support with push notifications, messages, diary and benefits tracker. The Garrison 2018 app training modules taught mindfulness for smoking cessation and how to work through cravings. Herbec 2019 included craving management tools within an app that supported smokers to be smoke free for 28 days.



The control conditions used in these smoking cessation app studies could be categorised into two groups:

- minimal non-app smoking cessation support that included: a printed self-help guide ([Baskerville 2018](#)), and encouragement to access available smoking cessation services ([Peiris 2019](#));
- less intensive app support that included an app that provided only basic information. In [BinDhim 2018](#) this included information only on quitting (no structured process or support). [Garrison 2018](#) delivered experience sampling to query smoking, craving, and mindfulness in real time, and the control app in [Herbec 2019](#) was designed to be a minimally credible intervention that resembled the intervention but without key intervention components.

#### Carbon monoxide (CO) monitoring and contingency management

[Alessi 2017](#) and [Wilson 2016](#) used mobile phone technology slightly differently to the above studies, by specifically using mobile phones to monitor the concentration of carbon monoxide in end-expiratory air (CO levels). In [Alessi 2017](#) interactive voice response calls would prompt the participant to conduct a CO test using a CO monitor. This was video recorded on the mobile phone and submitted using multimedia messaging. The CO result was provided via interactive voice response call. In the reinforcement arm of the trial, this was supplemented by negative CO test results (not smoking), which were rewarded with chances to win prizes. Therefore, the study had two arms that received mHealth CO monitoring as well as counselling and nicotine replacement therapy (NRT), with one of the arms also receiving rewards for smoking abstinence. [Wilson 2016](#) combined cognitive behavioural telephone counselling and access to NRT with a mobile app for CO monitoring and contingency management in one study arm, and compared this to the same intervention without the CO monitoring and contingency management app. Participants provided CO readings twice a day by video through the app and received payment for abstinence in the intervention arm.

#### Smartphone app plus text messaging

[Danaher 2019](#) tested an intervention that used both an integrated mobile web app and text messaging. Text messages were prompts and motivations to visit parts of the web programme as well as information, motivation and smoking questions (290 messages

over six months). The control group received a PC-based web intervention with interactive and multimedia features based on phases of quitting, the main difference to the intervention app being that it was not adapted for the small screen and did not include text messaging. Emails were sent as prompts if there were periods of inactivity.

#### Outcome

The included studies provided a range of abstinence outcome measures. Five studies ([Cobos-Campos 2017](#); [Free 2009](#); [Free 2011](#); [Liao 2018](#); [Peiris 2019](#)), reported the strictest outcome as biochemically verified sustained/continuous abstinence, and [Abrams 2014](#) and [Alessi 2017](#) defined abstinence as biochemically confirmed repeat point prevalence at six months.

Seven additional studies reported self-reported continuous abstinence at six months, without biochemical verification ([Baskerville 2018](#); [BinDhim 2018](#); [Borland 2013](#); [Herbec 2019](#); [Naughton 2014](#); [Rodgers 2005](#); [Whittaker 2011](#)).

Two studies used self-reported four-week or 30-day point prevalence abstinence at six-month follow-up ([Abrams 2017](#); [Haug 2013](#)), three studies used self-reported seven-day point prevalence at six months ([Augustson 2017](#); [Bock 2013](#); [Danaher 2019](#)), one used self-reported point prevalence abstinence at 32 weeks ([Squiers 2017](#)), one at 12 months ([Yu 2017](#)), and an additional four studies used six-month biochemically verified measures of seven-day point prevalence ([Chan 2015](#); [Ferguson 2015](#); [Garrison 2018](#); [Tseng 2017](#)). [Chan 2015](#) also provided biochemically verified seven-day point prevalence abstinence rates at 12-month follow-up. [Wilson 2016](#) reported six month follow-up data in their trial registry entry; however they do not specify whether these rates were validated or not.

#### Risk of bias in included studies

The [Characteristics of included studies](#) table provides details of 'Risk of bias' judgements for each domain of each included study. [Figure 2](#) illustrates judgements for each included study. Overall, we judged 13 studies to be at low risk of bias (judged at low risk for all domains), and three to be at high risk (judged to be at high risk in at least one domain). We judged the remaining studies to be at unclear risk (judged to be at unclear risk of bias for at least one domain, but with no judgements of high risk).

**Figure 2. 'Risk of bias' summary: review authors' judgements about each 'Risk of bias' item for each included study**

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Other bias
Abroms 2014	+	+	+	+	?
Abroms 2017	+	+	+	+	
Alessi 2017	+	?	+	+	
Augustson 2017	?	?	+	-	
Baskerville 2018	+	+	+	+	
BinDhim 2018	+	+	+	+	
Bock 2013	+	+	+	+	
Borland 2013	+	+	+	+	
Chan 2015	+	?	+	+	
Cobos-Campos 2017	+	+	+	-	
Danaher 2019	+	+	+	+	
Ferguson 2015	?	?	+	?	
Free 2009	+	+	+	+	
Free 2011	+	+	+	+	
Garrison 2018	+	?	+	+	
Haug 2013	+	?	+	+	?
Herbec 2019	+	+	+	-	
Liao 2018	+	+	+	+	
Naughton 2014	+	+	+	+	
Peiris 2019	+	+	+	+	
Rodgers 2005	+	+	+	+	?
Rodgers 2017	?	+	+	+	

**Figure 2. (Continued)**

Rodgers 2005	+	+	+	+	?
Squiers 2017	?	+	+	+	
Tseng 2017	+	+	+	+	
Whittaker 2011	+	+	+	+	
Wilson 2016	+	?	+	+	
Yu 2017	+	?	+	+	

### Selection bias

The majority of studies (17 of 26) appeared to have adequate procedures for random sequence generation and allocation concealment, so we judged them to be at low risk of bias for these domains; however [Alessi 2017](#); [Augustson 2017](#); [Chan 2015](#); [Ferguson 2015](#); [Garrison 2018](#); [Haug 2013](#); [Squiers 2017](#); [Wilson 2016](#) and [Yu 2017](#) did not provide sufficient description of either randomisation, concealment procedures, or both. Therefore, it is impossible to know whether the lack of information is due to actual bias or simply because it has not been reported, and we judged them to be at unclear risk of bias for at least random sequence generation or allocation concealment.

### Detection bias

Blinding of participants is not possible in studies of behavioural interventions. In this case participants knew if they were receiving text messages or using an app. Therefore, we did not assess performance bias, and instead judged the likelihood of detection bias. We did not deem a study to be high risk for this domain where there was biochemical verification of abstinence, or where both arms received the same amount of face-to-face contact (or none).

In most cases, studies collected outcomes electronically and remotely ([Abroms 2014](#); [Augustson 2017](#); [Baskerville 2018](#); [BinDhim 2018](#); [Bock 2013](#); [Danaher 2019](#); [Free 2009](#); [Free 2011](#); [Garrison 2018](#); [Squiers 2017](#)). [Chan 2015](#); [Herbec 2019](#); [Liao 2018](#); [Haug 2013](#) and [Wilson 2016](#) all collected outcomes by phone, and [Naughton 2014](#) by mailed questionnaire or in person. [Cobos-Campos 2017](#) collected outcomes in person in the clinic and this was not blinded, however this was mitigated by biochemical verification of quitting.

A number of the trials sought biochemical verification of long-term abstinence with salivary cotinine ([Abroms 2014](#); [Free 2009](#); [Free 2011](#)), urinary cotinine ([Liao 2018](#)), or expired CO ([Cobos-Campos 2017](#); [Garrison 2018](#)). [Chan 2015](#) assessed both CO and cotinine concentrations. [Abroms 2017](#) biochemically validated their primary outcome at three months, but not at six months, and [Rodgers 2005](#) validated abstinence at six weeks but not long-term follow-up. Similarly, [Naughton 2014](#) used verification at four weeks only. [Wilson 2016](#) stated that they planned to verify abstinence at all follow-up points using salivary cotinine; however it is not stated whether the abstinence rates reported in their trial registry entry were the validated rates or not. However, as data was collected remotely this study was still deemed to be at low risk of bias for this domain. In fact, we deemed all studies to be at low risk of detection bias.

### Attrition bias

We judged three studies to be at high risk of bias due to greater than 50% of participants lost to follow-up at six months ([Augustson 2017](#); [Cobos-Campos 2017](#); [Herbec 2019](#)). Several other studies had moderately high loss to follow-up but the numbers were clearly reported. The difference between groups was not greater than 20%, and overall loss was not greater than 50%. [Ferguson 2015](#) did not report loss to follow-up and so we judged it to be at unclear risk of attrition bias.

### Other

In [Abroms 2014](#) there were some issues with fraudulent enrolment at the outset of the study, although this was corrected once detected. In [Haug 2013](#), although clustering is adjusted for in this study's analysis the authors do not report the clustering effect, making it impossible to adjust for this in our analysis. Therefore, it is not clear how much the clustering adjustment influences the result from this study and our meta-analyses. [Rodgers 2005](#) suggested that some participants in their control group may have thought their incentive at follow-up (a month of free text messaging) depended on reporting quitting. This could account for an unexpected increase in control group participants reporting quitting from six weeks (109 participants) to six months (202 participants reporting no smoking in the past seven days), which could have led to an underestimation of the effect of the intervention.

### Effects of interventions

See: [Summary of findings for the main comparison Text messaging compared to minimal support for smoking cessation](#); [Summary of findings 2 Text messaging in addition to other smoking cessation support](#); [Summary of findings 3 Smartphone app compared to lower-intensity support for smoking cessation](#)

#### Text messaging versus minimal smoking cessation support

We pooled those studies that compared a text messaging intervention with minimal smoking cessation support. This included 13 studies ([Abroms 2014](#); [Abroms 2017](#); [Borland 2013](#); [Chan 2015](#); [Cobos-Campos 2017](#); [Ferguson 2015](#); [Free 2009](#); [Free 2011](#); [Haug 2013](#); [Liao 2018](#); [Rodgers 2005](#); [Whittaker 2011](#); [Yu 2017](#)). The analysis of all randomised participants, with those lost to follow-up classified as smokers resulted in a RR of 1.54 (95% CI 1.19 to 2.00;  $I^2 = 71%$ ; 14,133 participants; [Analysis 1.1](#)) with minimal difference found in the result when we carried out a complete case analysis (RR 1.56, 95% CI 1.21 to 2.02;  $I^2 = 72%$ ; 11,969 participants; [Analysis 1.2](#)).



We conducted the following sensitivity analyses:

- removing studies with very different populations from the main analysis (i.e. [Analysis 1.1](#)), pregnant women only in [Abroms 2017](#) and postnatal families only in [Yu 2017](#). This made very little difference to the overall result (RR 1.57, 95% CI 1.18 to 2.07;  $I^2 = 68%$ ; 13,408 participants);
- removing the only cluster-randomised trial, which we were unable to adjust for ([Haug 2013](#)). That again had minimal impact on the result (RR 1.57, 95% CI 1.19 to 2.07;  $I^2 = 73%$ ; 13,378 participants);
- removing the only study judged to be at high risk of bias ([Cobos-Campos 2017](#)), which again had minimal impact on the result (RR 1.49, 95% CI 1.13 to 1.96;  $I^2 = 72%$ ; 13,813 participants).

### Text messaging versus other smoking cessation intervention

Only two studies ([Borland 2013](#); [Chan 2015](#); 2238 participants), compared text messaging with another smoking cessation intervention. When pooled these did not show a superior effect of either text message support to quit or the other forms of smoking cessation intervention in either an analysis including all randomised participants (RR 0.92, 95% CI 0.61 to 1.40;  $I^2 = 27%$ ; 2238 participants; [Analysis 2.1](#)) or a complete case analysis (RR 0.93, 95% CI 0.63 to 1.36;  $I^2 = 20%$ ; 1813 participants; [Analysis 2.2](#)).

### Text messaging plus other smoking cessation support versus other smoking cessation support alone

Four studies ([Bock 2013](#); [Naughton 2014](#); [Tseng 2017](#); [Yu 2017](#); 997 participants), compared those who received both text messaging and another form of smoking cessation support with those only receiving the other form of smoking cessation support. The analysis of all randomised participants, assuming those lost to follow-up were smoking, showed a benefit of adding the text messaging with RR of 1.59 (95% CI 1.09 to 2.33;  $I^2 = 0%$ ; 997 participants; [Analysis 3.1](#)). The result was comparable when we carried out a complete case analysis (RR 1.63; 1.12 to 2.37;  $I^2 = 0%$ ; 796 participants; [Analysis 3.2](#)).

We carried out a sensitivity analysis on [Analysis 3.1](#) removing [Yu 2017](#), as it had a substantially different population (postnatal families). The interpretation of the effect remained the same (RR 1.87, 95% CI 1.13 to 3.09;  $I^2 = 0%$ ; 769 participants).

### High-frequency versus low-frequency text messaging

Three studies ([Augustson 2017](#); [Liao 2018](#); [Squiers 2017](#); 12,985 participants), compared high-frequency text messaging interventions with low-frequency text messaging interventions. The pooled effect indicated no difference in cessation rates between groups in either the analysis of all participants randomised (RR 1.00, 95% CI 0.95 to 1.06;  $I^2 = 0%$ ; [Analysis 4.1](#)) or the complete case analysis (RR 1.04, 95% CI 1.00 to 1.09;  $I^2 = 0%$ ; 6798 participants; [Analysis 4.2](#)). A sensitivity analysis removing the one study judged to be at high risk of bias ([Augustson 2017](#)), led to no difference in the interpretation of the effect (RR 1.02, 95% CI 0.92 to 1.12;  $I^2 = 0%$ ; 4985 participants).

### Smartphone app versus lower-intensity smoking cessation support

We divided studies of smartphone apps according to the type of control. Two studies ([Baskerville 2018](#); [Peiris 2019](#); 1645

participants), compared a smartphone app with minimal non-app smoking cessation support. There was no evidence of a favourable effect of smartphone apps in comparison with minimal non-app smoking cessation support (RR 0.82, 95% CI 0.56 to 1.18;  $I^2 = n/a$  as [Peiris 2019](#) had no events; [Analysis 5.1](#)). Interpretation remained the same when we carried out a complete case analysis (RR 0.87, 95% CI 0.62 to 1.23;  $I^2 = n/a$ ; 771 participants; [Analysis 5.2](#)). Three studies ([BinDhim 2018](#); [Garrison 2018](#); [Herbec 2019](#); 2175 participants), compared a smoking cessation smartphone app with a less intensive smoking cessation smartphone app. The analysis including all randomised participants resulted in an RR of 1.12 (95% CI 0.60 to 2.09;  $I^2 = 68%$ ; [Analysis 5.1](#)) with a very similar result in the complete case analysis (RR 1.18, 95% CI 0.67 to 2.09;  $I^2 = 65%$ ; 1003 participants). When we pooled all five studies, the resulting RR for all randomised participants was 1.00 (95% CI 0.66 to 1.52;  $I^2 = 59%$ ; 3079 participants; [Analysis 5.1](#)), providing no clear evidence of an increase in quit rates as a result of smart phone smoking cessation apps when compared to smoking cessation support of lower intensity. A sensitivity analysis removing the only study judged to be at high risk of bias ([Herbec 2019](#)), led to no difference in the interpretation of the effect (RR 1.10, 95% CI 0.60 to 2.00;  $I^2 = 71%$ ; 2654 participants).

### Carbon monoxide monitoring + contingency management versus smoking cessation support

Neither of the studies that used mobile phones to monitor CO and provide contingency management provided evidence that these strategies were more effective than standard smoking cessation support.

[Alessi 2017](#) compared messages prompting CO monitoring via video alone with the same CO monitoring plus reinforcement (with the chance to win prizes) for negative readings, and resulted in a RR of 0.88 (95% CI 0.35 to 2.21; 90 participants; [Analysis 6.1](#)).

[Wilson 2016](#) compared CO monitoring and contingency management combined with smoking cessation telephone counselling and NRT, with the counselling and NRT alone, and resulted in an RR of 0.94 (95% CI 0.64 to 1.38; 310 participants; [Analysis 6.1](#)).

In both cases carrying out a complete case analysis resulted in a change in the direction of the effect estimate; however CIs still incorporated evidence of both considerable benefit and harm ([Alessi 2017](#): RR 1.13, 95% CI 0.44 to 2.93; 81 participants; [Wilson 2016](#): RR 1.06, 95% CI 0.74 to 1.54; 250 participants; [Analysis 6.2](#)).

### Smartphone app + text messaging versus web-based interventions

[Danaher 2019](#) compared a smartphone app plus text messaging with a web-based smoking cessation intervention and found evidence for a benefit of the app plus text messaging (RR 1.80, 95% CI 1.32 to 2.45; 1271 participants; [Analysis 7.1](#)). Complete case analysis resulted in a similar point estimate (RR 1.56, 95% CI 1.19 to 2.05; 463 participants; [Analysis 7.2](#)).

## DISCUSSION

### Summary of main results

We found 26 randomised controlled trials of mobile phone smoking cessation interventions that met our inclusion criteria.

Whilst text messaging interventions tend to be very similar in design and content, the choice of control varied considerably. In this update, we separated out comparisons ensuring that only similar interventions and similar controls were pooled in meta-analyses.

Our analyses found moderate-certainty evidence ([Summary of findings for the main comparison](#)), that text messaging interventions are more effective than minimal smoking cessation support ([Abroms 2014](#); [Abroms 2017](#); [Borland 2013](#); [Chan 2015](#); [Cobos-Campos 2017](#); [Ferguson 2015](#); [Free 2009](#); [Free 2011](#); [Haug 2013](#); [Liao 2018](#); [Rodgers 2005](#); [Whittaker 2011](#); [Yu 2017](#)). Text messaging added to other smoking cessation interventions also appeared more effective than the other smoking cessation interventions alone ([Bock 2013](#); [Naughton 2014](#); [Tseng 2017](#); [Yu 2017](#); [Summary of findings 2](#)).

However, when text messaging was compared with other smoking cessation interventions, the analysis did not find evidence that either the text messaging intervention or the other smoking cessation interventions resulted in superior quit rates. It is important to highlight that there were just two studies in this analysis and they each had slightly different contexts: [Borland 2013](#) included people not seeking cessation support and participants were given 'suggestions about resources to use'; [Chan 2015](#) was in the context of a Quit & Win contest.

We were also able to assess the effect of higher- versus lower-intensity text messages on long-term abstinence rates, using data pooled from three studies providing direct comparisons ([Augustson 2017](#); [Liao 2018](#); [Squiers 2017](#)). The frequency of messaging did differ somewhat between studies (e.g. [Augustson 2017](#) used on average 15 versus 1 text message per week; [Liao 2018](#) used 21 to 35 versus 3 to 5 messages per week; and [Squiers 2017](#) used, on average, 16 versus 5 versus 1 text message per week), but overall, this analysis did not provide evidence that the intensity of the text messaging intervention impacted on abstinence rates. On average high intensity interventions resulted in abstinence rates of 26.6% versus 27.1% in low intensity interventions.

Studies of smartphone apps also included various control programmes. We found no evidence for a benefit of high intensity smartphone apps when compared with lower-intensity smoking cessation apps ([BinDhim 2018](#); [Garrison 2018](#); [Herbec 2019](#)), or minimal non-app smoking cessation support ([Baskerville 2018](#); [Peiris 2019](#)), but we judged the evidence to be of very low certainty, meaning we have very little confidence in the effect estimate ([Summary of findings 3](#)).

[Danaher 2019](#) was the only intervention that used both text messaging and a smartphone app and found that this combination resulted in higher quit rates than a web-based smoking cessation intervention.

### Overall completeness and applicability of evidence

Our review includes 26 studies with 33,849 participants. In comparison with previous reviews, there is now a much greater number of eligible studies, with increased sample sizes and including a greater diversity of settings and countries. We also found a large number of ongoing studies ( $n = 34$ ), the results of which are likely to increase the diversity of contexts even further.

This is the first update of this review where there were randomised controlled trials of smartphone apps eligible to be included. In 2011, a review of available smoking cessation apps found them to be lacking in adherence to cessation guidelines or theory ([Abroms 2011](#)). In this review the included smartphone apps, although few in number, tended to be based on evidence or theory and were tested in high-quality randomised controlled trials.

There has been criticism that smartphone apps may not be widely accessible to all, as they may rely on a certain degree of digital literacy and technology access that may not be widely dispersed in the population. It is important to note that in the included studies of smartphone apps there were reasonably high levels of education: 84% of participants in [Garrison 2018](#) had greater than high school education; in [BinDhim 2018](#), 53.7% had graduate level or higher education; in [Baskerville 2018](#), 55.5% had post-secondary education or higher; [Danaher 2019](#) included 70% with a high school graduate education and higher; and in [Herbec 2019](#), 68.7% had a post-16 years qualification.

A common criticism of randomised controlled trials is that whilst they might provide evidence of effectiveness in a clinical trial setting, these data are not applicable to 'real-world' settings. We are aware that many countries are implementing mCessation interventions and encourage routine monitoring and evaluation of these programmes, which will provide important 'real-world' evidence for consideration alongside the research evidence.

### Certainty of the evidence

There was moderate-certainty evidence that text messaging increases quit rates by approximately 50% when compared to minimal support for smoking cessation ([Summary of findings for the main comparison](#)). We downgraded the evidence by one level due to inconsistency as there was substantial unexplained statistical heterogeneity. This means the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

There was also moderate-certainty evidence that text messages increase quit rates by approximately 60% when tested as an addition to other smoking cessation support ([Summary of findings 2](#)). We downgraded results by one level due to imprecision: there were fewer than 300 events overall, and confidence intervals encompassed minimal benefit and substantial benefit.

There was very low-certainty evidence regarding the effect of smartphone apps compared to lower-intensity support ([Summary of findings 3](#)). This is due to inconsistency (considerable unexplained statistical heterogeneity) and very serious imprecision, with confidence intervals encompassing both clinically significant harm and clinically significant benefit.

### Potential biases in the review process


A wide variability of control group programmes is potentially important in ensuring that the studies can provide the best information for decision makers who may want to compare mCessation with what already exists in their context. However, it could also lead to difficulties in the interpretation of the results. In some cases control groups received substantial smoking cessation support and the details of this were not always clear. This is supported by the fact that in some cases high quit rates, over what might have been expected, were observed in control

groups (Rodgers 2005; Squiers 2017), with high rates in both the intervention and control groups in another study (Augustson 2017). This could indicate some degree of trial effect (everyone does better just through being involved in a research study), social desirability bias, or that minimal mobile phone interventions (just for reminders, prompts or data collection) may also be effective in producing behaviour change. High-intensity control groups leading

to high quit rates could have underestimated the relative effect of mobile phone interventions.

Though we searched trial registries, there remains a risk that there were eligible but unpublished studies we failed to identify. Reassuringly, a funnel plot (Figure 3), showed no evidence of asymmetry.

**Figure 3. Funnel plot of comparison 1. Text messaging versus minimal smoking cessation support, outcome: 1.1 long-term abstinence (all randomised))**



**Agreements and disagreements with other studies or reviews**

This review agrees with other reviews of the benefits of text messaging to support healthy behaviour change (Armanasco 2017; Thakkar 2016; Scott-Sheldon 2016). Several reviews have shown mixed results with respect to the effectiveness of smartphone apps for behaviour change, with significant issues relating to the size and quality of studies (Byambasuren 2018; Dirieto 2016; Lunde 2018; Schoeppe 2016; Zhao 2016).

**AUTHORS' CONCLUSIONS**

**Implications for practice**

There is moderate-certainty evidence that text-message-based interventions improve smoking cessation rates, either delivered on their own or as an add-on to other treatments. There is insufficient evidence with which to evaluate the effect of mobile app interventions, but there are many ongoing studies, so evidence on these interventions will continue to evolve over time.

**Implications for research**

Research in diverse populations and contexts is still required in order to understand what types of mCessation might be effective for particular groups and those most in need of support. The heterogeneity in text message programmes and the variation in functionality within the apps means further research is also needed to understand the effective elements, components and durations of these types of interventions. The variety of control programmes in the studies reviewed, and the often unexpectedly high abstinence rates in control groups, is an issue that may require further research in order to determine the actual size of the effect of interventions and potentially the 'minimal' effective mCessation intervention. More large-scale randomised controlled trials are needed in order to establish whether mobile app interventions are effective for smoking cessation.

**ACKNOWLEDGEMENTS**

We acknowledge the assistance of Cochrane Tobacco Addiction Editorial base in the preparation of this review, whose time on the

review was supported by the National Institute for Health Research (NIHR), via Cochrane Infrastructure and Cochrane Programme Grant funding to Cochrane Tobacco Addiction. The views and opinions expressed therein are those of the authors and do not

necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health and Social Care. We also gratefully acknowledge the consumer review by Lee Bromhead.

## REFERENCES

### References to studies included in this review

#### Abroms 2014 {published data only}

Abroms L, Ahuja M, Kodl Y, Thaweethai L, Sims J, Winickoff J, et al. Text2Quit: results from a pilot test of a personalised, interactive mobile health smoking cessation program. *Journal of Health Communication* 2012;**17**(Suppl 1):44-53.

\* Abroms LC, Boal AL, Simmens SJ, Mendel JA, Windsor RA. A randomized trial of Text2Quit: a text messaging program for smoking cessation. *American Journal of Preventive Medicine* 2014;**47**(3):242-50. [DOI: [10.1016/j.amepre.2014.04.010](https://doi.org/10.1016/j.amepre.2014.04.010)]

Heminger CL, Boal AL, Zumer M, Abroms LC. Text2Quit: an analysis of participant engagement in the mobile smoking cessation program. *American Journal of Drug and Alcohol Abuse* 2016;**42**(4):450-8. [DOI: [10.3109/00952990.2016.1149591](https://doi.org/10.3109/00952990.2016.1149591)]

Hoepfner B, Hoepfner S, Abroms L. How do text-messaging smoking cessation interventions confer benefit? A multiple mediation analysis of Text2Quit. *Addiction* 2017;**112**(4):673-82. [DOI: [10.1111/add.13685](https://doi.org/10.1111/add.13685)]

#### Abroms 2017 {published data only}

Abroms L, Johnson P, Leavitt L, Cleary S, Bushar J, Brandon T, et al. A randomized trial of text messaging for smoking cessation in pregnant women. *American Journal of Preventive Medicine* 2017;**53**(6):781-90. [DOI: [10.1016/j.amepre.2017.08.002](https://doi.org/10.1016/j.amepre.2017.08.002)]

#### Alessi 2017 {published data only}

Alessi SM, Rash CJ. Treatment satisfaction in a randomized clinical trial of mHealth smoking abstinence reinforcement. *Journal of Substance Abuse Treatment* 2017;**72**:103-10. [DOI: [10.1016/j.jsat.2016.06.013](https://doi.org/10.1016/j.jsat.2016.06.013)]

\* Alessi SM, Rash CJ, Petry NM. A randomized trial of adjunct mHealth abstinence reinforcement with transdermal nicotine and counseling for smoking cessation. *Nicotine & Tobacco Research* 2017;**19**(3):290-8. [DOI: [10.1093/ntr/ntw155](https://doi.org/10.1093/ntr/ntw155)]

#### Augustson 2017 {published data only}

Augustson E, Engelgau M, Zhang S, Cai Y, Cher W, Li R, et al. Text to quit China: an mHealth smoking cessation trial. *American Journal of Health Promotion* 2017;**31**(3):217-25. [DOI: [10.4278/ajhp.140812-QUAN-399](https://doi.org/10.4278/ajhp.140812-QUAN-399)]

#### Baskerville 2018 {published data only}

Baskerville N, Struik L, Dash D. Crush the Crave: development and formative evaluation of a smartphone app for smoking cessation. *JMIR Mhealth Uhealth* 2018;**6**(3):e52. [DOI: [10.2196/mhealth.9011](https://doi.org/10.2196/mhealth.9011)]

\* Baskerville N, Struik L, Guindon G, Norman C, Whittaker R, Burns C, et al. Effect of a mobile phone intervention on quitting smoking in a young adult population of smokers: randomized controlled trial. *JMIR Mhealth and Uhealth* 2018;**6**(10):e10893.

Baskerville N, Struik L, Hammond D, Guindon G, Norman C, Whittaker R, et al. Effect of a mobile phone intervention on quitting smoking in a young adult population of smokers:

randomized controlled trial study protocol. *JMIR Research Protocols* 2015;**4**(1):e10. [DOI: [10.2196/resprot.3823](https://doi.org/10.2196/resprot.3823)]

NCT01983150. A randomized controlled trial to test the effect of a smartphone quit smoking intervention on young adult smokers. [clinicaltrials.gov/ct2/show/NCT01983150](https://clinicaltrials.gov/ct2/show/NCT01983150) (first received 13 November 2013).

#### BinDhim 2018 {published data only}

ACTRN12613000833763. Assessing the effect of an interactive decision-aid smartphone smoking cessation application (app) on quit rates: a double-blind randomized control trial [In adult smokers, does an interactive smartphone smoking cessation application, compared to standard information, increase the quit rates?]. [anzctr.org.au/Trial/Registration/TrialReview.aspx?id=364613](http://anzctr.org.au/Trial/Registration/TrialReview.aspx?id=364613) (first received 17 July 2013).

\* BinDhim N, McGeechan K, Revena L. Smartphone smoking cessation application (SSC App) trial: a multicountry double-blind automated randomised controlled trial of a smoking cessation decision-aid 'app'. *BMJ Open* 2018;**8**(1):e017105. [DOI: [10.1136/bmjopen-2017-017105](https://doi.org/10.1136/bmjopen-2017-017105)]

BinDhim N, McGeechan K, Trevena L. Assessing the effect of an interactive decision-aid smartphone smoking cessation application (app) on quit rates: a double-blind automated randomised control trial protocol. *BMJ Open* 2014;**4**(7):e005371. [DOI: <http://dx.doi.org/10.1136/bmjopen-2014-005371>]

#### Bock 2013 {published data only}

\* Bock B, Heron K, Jennings E, Morrow K, Cobb V, Magee J, et al. A text message delivered smoking cessation intervention: the initial trial of TXT-2-Quit: randomized controlled trial. *JMIR MHealth and UHealth* 2013;**1**(2):e17. [CENTRAL: 1015646; CRS: 940012900003441]

#### Borland 2013 {published and unpublished data}

Balmford J, Borland R. How do smokers use a smoking cessation text messaging intervention?. *Nicotine & Tobacco Research* 2014;**16**(12):1586-92. [DOI: [10.1093/ntr/ntu111](https://doi.org/10.1093/ntr/ntu111)]

Balmford J, Borland R, Benda P, Howard S. Factors associated with use of automated smoking cessation interventions: findings from the eQuit study. *Health Education Research* 2013;**28**(2):288-99. [CENTRAL: 921007; CRS: 9400107000001543; PUBMED: 23107931]

\* Borland R, Balmford J, Benda P. Population-level effects of automated smoking cessation help programs: a randomized controlled trial. *Addiction* 2013;**108**(3):618-28. [CENTRAL: 921001; CRS: 9400107000000767; PUBMED: 22994457]

#### Chan 2015 {published data only}

Chan SS, Wong DC, Cheung YT, Leung DY, Lau L, Lai V, et al. A block randomized controlled trial of a brief smoking cessation counselling and advice through short message service on participants who joined the Quit to Win Contest in Hong Kong. *Health Education Research* 2015;**30**(4):609-21. [DOI: [10.1093/her/cyv023](https://doi.org/10.1093/her/cyv023)]



**Cobos-Campos 2017** {published data only}

Cobos-Campos R, de Larrinoa A, Morinigo A, Parraza N, Barandiaran F. Effectiveness of text messaging as an adjuvant to health advice in smoking cessation programs in primary care. A randomized clinical trial. *Nicotine & Tobacco Research* 2017;**19**(8):901-7. [DOI: [10.1093/ntr/ntw300](https://doi.org/10.1093/ntr/ntw300)]

**Danaher 2019** {published data only}

\* Danaher B, Tyler M, Crowley R, Brendryen H, Seeley J. Outcomes and device usage for fully automated Internet interventions designed for a smartphone or personal computer: the MobileQuit smoking cessation randomized controlled trial. *JMIR* 2019;**21**(6):e13290.

NCT01952236. Web and mobile smoking cessation (MobileQuit). [clinicaltrials.gov/show/nct01952236](https://clinicaltrials.gov/show/nct01952236) (first received 27 September 2013).

**Ferguson 2015** {published and unpublished data}

\* Ferguson SG, Walters JA. The effect of mobile phone text messages on short and long term quitting in motivated smokers: a randomised controlled trial. Society for Research on Nicotine and Tobacco 21st Annual Meeting; 2015 Feb 25-28; Philadelphia. 2015. [CRS: 9400131000001072; POS3-80]

Schuz N, Walters JA, Frandsen M, Bower J, Ferguson SG. Compliance with an EMA monitoring protocol and its relationship with participant and smoking characteristics. *Nicotine & Tobacco Research* 2014;**16**(S2):S88-92. [CRS: 9400129000001511; EMBASE: 2014262309]

**Free 2009** {published and unpublished data}

Free C, Whittaker R, Knight R, Abramsky T, Rodgers A, Roberts IG. Txt2stop: a pilot randomised controlled trial of mobile phone-based smoking cessation support. *Tobacco Control* 2009;**18**:88-91.

**Free 2011** {published and unpublished data}

Douglas N, Free C. 'Someone battling in my corner': experiences of smoking-cessation support via text message. *British Journal of General Practice* 2013;**63**(616):e768-76. [CENTRAL: 1000333; CRS: 9400129000002807; PUBMED: 24267860]

Free C, Hoile E, Robertson S, Knight R. Three controlled trials of interventions to increase recruitment to a randomized controlled trial of mobile phone based smoking cessation support. *Clinical Trials* 2010;**7**(3):265-73.

\* Free C, Knight R, Robertson S, Whittaker R, Edwards P, Zhou W, et al. Smoking cessation support delivered via mobile phone text messaging (Txt2stop): a single-blind, randomised trial. *Lancet* 2011;**378**:49-55.

Guerriero C, Cairns J, Roberts I, Rodgers A, Whittaker R, Free C. The cost-effectiveness of smoking cessation support delivered by mobile phone text messaging: Txt2stop. *European Journal of Health Economics* 2013;**14**(5):789-97. [CENTRAL: 986154; CRS: 9400129000001252; PUBMED: 22961230]

Michie S, Free C, West R. Characterising the 'Txt2Stop' smoking cessation text messaging intervention in terms of

behaviour change techniques. *Journal of Smoking Cessation* 2012;**7**(1):55-60. [CRS: 9400123000018452]

Severi E, Free C, Knight R, Robertson S, Edwards P, Hoile E. Two controlled trials to increase participant retention in a randomized controlled trial of mobile phone-based smoking cessation support in the United Kingdom. *Clinical Trials* 2011;**8**(5):654-60.

**Garrison 2018** {published data only}

\* Garrison K, Pal P, O'Malley S, Pittman B, Gueorguieva R, Rojiani R, et al. Craving to Quit: a randomised controlled trial of smartphone app-based mindfulness training for smoking cessation. *Nicotine & Tobacco Research* 2018 [Epub ahead of print]:1-8. [DOI: [10.1093/ntr/nty126](https://doi.org/10.1093/ntr/nty126)]

Garrison KA, Pal P, Rojiani R, Dallery J, O'Malley SS, Brewer JA. A randomized controlled trial of smartphone-based mindfulness training for smoking cessation: a study protocol. *BMC Psychiatry* 201;**15**(1):83. [DOI: [10.1186/s12888-015-0468-z](https://doi.org/10.1186/s12888-015-0468-z)]

**Haug 2013** {published and unpublished data}

Haug S, Meyer C, Dymalski A, Lippke S, John U. Efficacy of a text messaging (SMS) based smoking cessation intervention for adolescents and young adults: study protocol of a cluster randomised controlled trial. *BMC Public Health* 2012;**12**:51. [CENTRAL: 814353; CRS: 9400123000011961; EMBASE: 22260736; PUBMED: 22260736]

\* Haug S, Schaub MP, Venzin V, Meyer C, John U. Efficacy of a text message-based smoking cessation intervention for young people: a cluster randomized controlled trial. *Journal of Medical Internet Research* 2013;**15**(8):142-55. [CENTRAL: 980800; CRS: 9400129000000569; PUBMED: 23956024]

Haug S, Schaub MP, Venzin V, Meyer C, John U. Moderators of outcome in a text messaging (SMS)-based smoking cessation intervention for young people [Differenzielle Wirksamkeit eines Short Message Service (SMS)-basierten Programms zur Forderung des Rauchstopps bei Jugendlichen]. *Psychiatrische Praxis* 2013;**40**(6):339-46. [CENTRAL: 875575; CRS: 9400126000000210; EMBASE: 2013576012; PUBMED: 24008683]

Haug S, Schaub Michael P, Schmid H. Predictors of adolescent smoking cessation and smoking reduction. *Patient Education and Counseling* 2014;**95**(3):378-83. [CRS: 9400129000001807; PUBMED: 24674150]

**Herbec 2019** {unpublished data only}

\* Herbec A, Brown J, Shahab L, West R. Lessons learned from unsuccessful use of personal carbon monoxide monitors to remotely assess abstinence in a pragmatic trial of a smartphone stop smoking app – a secondary analysis. *Addictive Behaviors Reports* 2019;**9**:100122.

Herbec A, Shahab L, Matei A, Brown J, Kaur Ubhi H, et al. Does inclusion of craving management tools increase effectiveness and usage of a stop smoking app? Results from BupaQuit trial. 3rd UCL Centre for Behaviour Change Digital Health Conference 2017: Harnessing digital technology for behaviour change; 2017 Feb 22-23; London, UK. 2017.

ISRCTN10548241. Study of effectiveness of a smartphone application for quitting smoking. [isrctn.com/ISRCTN10548241](http://isrctn.com/ISRCTN10548241) (first received 10 February 2015).

**Liao 2018** {published data only}

\* Liao Y, Wu Q, Kelly B, Zhang F, Tang YY, Want Q, et al. Effectiveness of a text-messaging-based smoking cessation intervention ("Happy Quit") for smoking cessation in China: a randomized controlled trial. *PLOS Medicine* 2018;**15**(12):e1002713.

Liao Y, Wu Q, Tang J, Zhang F, Wang X, Qi C, et al. The efficacy of mobile phone-based text message interventions ('Happy Quit') for smoking cessation in China. *BMC Public Health* 2016;**16**(1):833. [DOI: [10.1186/s12889-016-3528-5](https://doi.org/10.1186/s12889-016-3528-5)]

**Naughton 2014** {published data only}

Faulkner K, Sutton S, Jamison J, Sloan M, Boase S, Naughton F. Are nurses and auxiliary healthcare workers equally effective in delivering smoking cessation support in primary care?. *Nicotine & Tobacco Research* 2016;**18**(5):1054-60. [DOI: [10.1093/ntr/ntv206](https://doi.org/10.1093/ntr/ntv206)]

\* Naughton F, Jamison J, Boase S, Sloan M, Gilbert H, Prevost AT, et al. Randomized controlled trial to assess the short-term effectiveness of tailored web- and text-based facilitation of smoking cessation in primary care (iQuit in Practice). *Addiction* 2014;**109**(7):1184-93. [CENTRAL: 1051137; CRS: 9400129000001642; PUBMED: 24661312]

Sutton S, Smith S, Jamison J, Boase S, Mason D, Prevost AT, et al. Study protocol for iQuit in Practice: a randomised controlled trial to assess the feasibility, acceptability and effectiveness of tailored web- and text-based facilitation of smoking cessation in primary care. *BMC Public Health* 2013;**13**:324. [CENTRAL: 873088; CRS: 940012600000015; PUBMED: 23575031]

**Peiris 2019** {published data only}

Peiris D, Wright L, News M, Rogers K, Redfern J, Chow C, et al. A smartphone app to assist smoking cessation among Aboriginal Australians: findings from a pilot randomized controlled trial. *JMIR MHealth and UHealth* 2019;**7**(4):e12745.

**Rodgers 2005** {published data only}

Bramley D, Riddell T, Whittaker R, Corbett T, Lin R-B, Wills M. Smoking cessation using mobile phone text messaging is as effective in Maori as non-Maori. *New Zealand Medical Journal* 2005;**118**(1216):1494-504.

\* Rodgers A, Corbett T, Bramley D, Riddell T, Wills M, Lin R-B, et al. Do u smoke after txt? Results of a randomised trial of smoking cessation using mobile phone text messaging. *Tobacco Control* 2005;**14**:255-61. [doi: [10.1136/tc.2005.011577](https://doi.org/10.1136/tc.2005.011577)]

**Squiers 2017** {published data only}

Coa K, Augustson E, Kaufman A. The impact of weight and weight-related perceptions on smoking status among young adults in a text-messaging cessation program. *Nicotine & Tobacco Research* 2018;**20**(5):614-9. [DOI: [10.1093/ntr/ntx053](https://doi.org/10.1093/ntr/ntx053)]

\* Squiers L, Augustson E, Brown D, Kelly B, Southwell B, Dever J, et al. An experimental comparison of mobile texting

programs to help young adults quit smoking. *Health Systems* 2017;**6**:1-14.

Squiers L, Brown D, Parvanta S, Dolina S, Kelly B, Dever J, et al. The smokefreeTXT (SFTXT) study: web and mobile data collection to evaluate smoking cessation for young adults. *JMIR Research Protocols* 2016;**5**(2):e134. [DOI: [10.2196/resprot.5653](https://doi.org/10.2196/resprot.5653)]

**Tseng 2017** {published and unpublished data}

NCT01898195. Improving adherence to smoking cessation medication among PLWHA (HIV). [clinicaltrials.gov/show/NCT01898195](http://clinicaltrials.gov/show/NCT01898195) (first received 12 July 2013). [CRS: 9400129000001221]

Shelley D, Krebs P, Schoenthaler A, Urbina A, Gonzalez M, Tseng T-Y, et al. Correlates of adherence to varenicline among HIV+ smokers: a path analysis. Society for Research on Nicotine and Tobacco 21st Annual Meeting; 2015 Feb 25-28; Philadelphia. 2015. [CRS: 9400131000001079; PA14-2]

\* Tseng TY, Krebs P, Schoenthaler A, Wong S, Sherman S, Gonzalez M, et al. Combining text messaging and telephone counselling to increase varenicline adherence and smoking abstinence among cigarette smokers living with HIV: a randomized controlled trial. *AIDS Behavior* 2017;**21**(7):1964-74.

**Whittaker 2011** {published and unpublished data}

Whittaker R, Dorey E, Bramley D, Bullen C, Denny S, Elley C, et al. A theory-based video messaging mobile phone intervention for smoking cessation: randomised controlled trial. *Journal of Medical Internet Research* 2011;**13**(1):e10.

**Wilson 2016** {published data only}

NCT01723163. Abstinence Reinforcement Therapy (ART) for rural veteran smokers. [www.clinicaltrials.gov/ct2/show/NCT01723163](http://www.clinicaltrials.gov/ct2/show/NCT01723163) (first received 7th November 2012).

Wilson S, Hair L, Hertzberg J, Kirby A, Olsen M, Lindquist J, et al. Abstinence reinforcement therapy (ART) for rural veterans: methodology for an mHealth smoking cessation intervention. *Contemporary Clinical Trials* 2016;**50**:157-65.

**Yu 2017** {published data only}

ChiCTR-OIC-17010803. Changchun Smoke-Free Home mHealth Program [Creating smoke-free homes for infants and increasing quitting among fathers through education and mHealth interventions in Changchun, China]. [chictr.org.cn/showprojen.aspx?proj=18143](http://chictr.org.cn/showprojen.aspx?proj=18143) (first received 06 March 2017).

\* Yu S, Duan Z, Redmon PB, Eriksen MP, Koplan JP, Huang C. MHealth intervention is effective in creating smoke-free homes for newborns: a randomized controlled trial study in China. *Scientific Reports* 2017;**7**(1):9276. [DOI: [10.1038/s41598-017-08922-x](https://doi.org/10.1038/s41598-017-08922-x)]

**References to studies excluded from this review**

**ACTRN12617000491369** {published data only}

ACTRN12617000491369. Quittr: a game that wants smokers to quit [Quittr: a smoking cessation serious game. Determining if game features influence smoker engagement and retention

during a quit attempt]. [anzctr.org.au/Trial/Registration/TrialReview.aspx?id=372661](http://anzctr.org.au/Trial/Registration/TrialReview.aspx?id=372661) (first received 29 March 2017).

**Aigner 2017** {published data only}

Aigner C, Gritz E, Tami-Maury I, Baum G, Arduino R, Vidrine D. The role of pain in quitting among human immunodeficiency virus (HIV)-positive smokers enrolled in a smoking cessation trial. *Substance Abuse* 2017;**38**(3):249-52. [DOI: [10.1080/08897077.2017.1291466](https://doi.org/10.1080/08897077.2017.1291466)]

**Applegate 2007** {published data only}

Applegate B, Raymond C, Collado-Rodriguez A, Riley W, Schneider N. Improving adherence to nicotine gum by SMS text messaging: a pilot study. Society for Research on Nicotine and Tobacco 13th Annual Meeting; 2007 Feb 21-24; Austin, Texas. 2007. [RPOS3-57]

**Bamidis 2017** {published data only}

Bamidis P, Paraskevopoulos E, Konstantinidis E, Spachos D, Billis A. Multimodal e-health services for smoking cessation and public health: the SmokeFreeBrain project approach. *Studies in Health Technology and Informatics* 2017;**245**:5-9.

**Bernstein 2018** {published data only}

Bernstein S, Dziura J, Weiss J, Miller T, Vickerman K, Grau L, et al. Tobacco dependence treatment in the emergency department: a randomized trial using the Multiphase Optimization Strategy. *Contemporary Clinical Trials* 2018;**66**:1-8. [DOI: [10.1016/j.cct.2017.12.016](https://doi.org/10.1016/j.cct.2017.12.016)]

**Blasco 2012** {published data only}

Blasco A, Carmona M, Fernandez-Lozano I, Salvador CH, Pascual M, Sagredo PG, et al. Evaluation of a telemedicine service for the secondary prevention of coronary artery disease. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2012;**32**(1):25-31. [CENTRAL: 814645; CRS: 9400123000012404; 6831; PUBMED: 22113368]

**Brendryen 2008** {published data only}

Brendryen H, Drozd F, Kraft P. A digital smoking cessation program delivered through internet and cell phone without nicotine replacement (Happy Ending): randomized controlled trial. *Journal of Medical Internet Research* 2008;**10**(5):e51.

\* Brendryen H, Kraft P. Happy Ending: a randomized controlled trial of a digital multi-media smoking cessation intervention. *Addiction* 2008;**103**:478-84. [doi:10.1111/j.1360-0443.2007.02119.x]

**Bricker 2014** {published data only}

\* Bricker J, Mull K, Kientz J, Vilardaga R, Mercer L, Akioka K, et al. Randomized, controlled pilot trial of a smartphone app for smoking cessation using acceptance and commitment therapy. *Drug and Alcohol Dependence* 2014;**143**(1):87-94. [CENTRAL: 1047686; CRS: 9400050000000151; EMBASE: 2014721228; PUBMED: 25085225]

Heffner JL, Vilardaga R, Mercer LD, Kientz JA, Bricker JB. Feature-level analysis of a novel smartphone application for smoking cessation. *American Journal of Drug and Alcohol Abuse* 2015;**41**(1):68-73. [CENTRAL: 1036931; CRS: 9400129000003871; EMBASE: 2014973943]

**Brinker 2016** {published data only}

Brinker T, Holzapfel J, Baudson T, Sies K, Jakob L, Baumert H, et al. Photoaging smartphone app promoting poster campaign to reduce smoking prevalence in secondary schools: the Smokerface randomized trial: design and baseline characteristics. *BMJ Open* 2016;**6**(11):e014288. [DOI: [10.1136/bmjopen-2016-014288](https://doi.org/10.1136/bmjopen-2016-014288)]

**Bronshtein 2016** {published data only}

Bronshtein E, Ondersma S, Sokol R. Optimizing ehealth for smoking in pregnancy: a pilot factorial evaluation of providing single vs multiple options. *Reproductive Sciences* 2016;**S1**:195A.

**Buller 2014** {published data only}

Buller DB, Borland R, Bettinghaus EP, Shane JH, Zimmerman DE. Randomized trial of a smartphone mobile application compared to text messaging to support smoking cessation. *Telemedicine Journal and E-health* 2014;**20**(3):206-14. [CENTRAL: 1053488; CRS: 9400129000003463]

**Chow 2012** {published data only}

Chow CK, Redfern J, Thiagalingam A, Jan S, Whittaker R, Hackett M, et al. Design and rationale of the tobacco, exercise and diet messages (TEXT ME) trial of a text message-based intervention for ongoing prevention of cardiovascular disease in people with coronary disease: a randomised controlled trial protocol. *BMJ Open* 2012;**2**(1):e000606.

**Dale 2014** {published data only}

Dale L, Whittaker R, Jiang Y, Stewart R, Rolleston A, Maddison R. Improving coronary heart disease self-management using mobile technologies (Text4Heart): a randomised controlled trial protocol. *Trials* 2014;**15**:71. [CENTRAL: 984614; CRS: 9400129000001546; EMBASE: 2014230967; PUBMED: 24588893]

**Fingrut 2014** {published data only}

Fingrut W, Stewart L, Cheung K. Choice of smoking cessation counselling via phone, text, or email in emergency department patients. *Canadian Journal of Emergency Medicine* 2014;**16**:S86. [CENTRAL: 1009852; CRS: 9400129000003122; EMBASE: 75007022]

**Fraser 2014** {published data only}

Fraser D, Kobinsky K, Smith SS, Kramer J, Theobald WE, Baker TB. Five population-based interventions for smoking cessation: a MOST trial. *Translational Behavioral Medicine* 2014;**4**(4):382-90. [CENTRAL: 1040081; CRS: 9400129000003838; EMBASE: 2015657527]

**Gritz 2013** {published data only}

\* Gritz E, Danysh H, Fletcher F, Tami-Maury I, Fingeret M, King R, et al. Long-term outcomes of a cell phone-delivered intervention for smokers living with HIV/AIDS. *Clinical Infectious Diseases* 2013;**57**(4):608-15.

Gritz E, Vidrine D, Marks R, Arduino R. A randomized trial of an innovative cell phone intervention for smokers living with HIV/AIDS. Society for Research on Nicotine & Tobacco 17th Annual Meeting; 2011 Feb 16-19; Toronto, Canada. 2011. [SYM 10A]



NCT00502827. Smoking Cessation for HIV/AIDS Patients. [clinicaltrials.gov/ct2/show/NCT00502827](https://clinicaltrials.gov/ct2/show/NCT00502827) (first received 26 February 2016).

Tami-Maury I, Vidrine D, Fletcher F, Danysh H, Arduino R, Gritz E. Poly-tobacco use among HIV-positive smokers: implications for smoking cessation efforts. *Nicotine & Tobacco Research* 2013;**15**(12):2100-6.

Vidrine D, Kypriotakis G, Li L, Arduino R, Fletcher F, Tami-Maury I, et al. Mediators of a smoking cessation intervention for persons living with HIV/AIDS. *Drug and Alcohol Dependence* 2015;**147**:76-80. [DOI: [10.1016/j.drugalcdep.2014.12.003](https://doi.org/10.1016/j.drugalcdep.2014.12.003)]

Vidrine D, Marks R, Arduino R, Gritz E. Efficacy of cell phone-delivered smoking cessation counseling for persons living with HIV/AIDS: 3-month outcomes. *Nicotine & Tobacco Research* 2012;**14**(1):106-10.

#### Halpern 2018 {published data only}

\* Halpern S, Harhay M, Saulsgiver K, Brophy C, Troxel A, Volpp K. A pragmatic trial of e-cigarettes, incentives, and drugs for smoking cessation. *New England Journal of Medicine* 2018;**378**(24):2302-10.

NCT02328794. Randomized clinical trial to reduce harm from tobacco. [clinicaltrials.gov/ct2/show/NCT02328794](https://clinicaltrials.gov/ct2/show/NCT02328794) (first received 31 December 2014).

#### Hammett 2018 {published data only}

Hammett E, Veldheer S, Hrabovsky S, Yingst J, Berg A, Poole E, et al. TXT2STAYQUIT: pilot randomized trial of brief automated smoking cessation texting intervention for inpatient smokers discharged from the hospital. *Journal of Hospital Medicine* 2018;**13**(7):488-9. [DOI: [10.12788/jhm.2907](https://doi.org/10.12788/jhm.2907)]

#### Hassandra 2017 {published data only}

Hassandra M, Lintunen T, Hagger MS, Heikkinen R, Vanhala M, Kettunen T. An mHealth App for supporting quitters to manage cigarette cravings with short bouts of physical activity: a randomized pilot feasibility and acceptability study. *JMIR Mhealth and Uhealth* 2017;**5**(5):e74. [DOI: [10.2196/mhealth.6252](https://doi.org/10.2196/mhealth.6252)]

#### Haug 2008 {published data only}

Haug S, Meyer C, Gross B, Schorr G, Thyrian JR, Kordy H, et al. Continuous individual support of smoking cessation in socially deprived young adults via mobile phones - results of a pilot study. *Gesundheitswesen* 2008;**70**(6):364-71.

#### Haug 2009 {published data only}

Haug S, Meyer C, John U. Individual support of smoking cessation using text messaging: results of two pilot studies. *Sucht* 2008;**54**(5):316. [CENTRAL: 794115; CRS: 940012300006096]

\* Haug S, Meyer C, Schorr G, Bauer S, John U. Continuous individual support of smoking cessation using text messaging: a pilot experimental study. *Nicotine & Tobacco Research* 2009;**11**(8):915-23.

#### Haug 2014 {published data only}

Haug S, Paz Castro R, Filler A, Kowatsch T, Fleisch E, Schaub MP. Efficacy of an internet and SMS-based integrated smoking cessation and alcohol intervention for smoking cessation in young people: study protocol of a two-arm cluster randomised controlled trial. *BMC Public Health* 2014;**14**:1140. [CENTRAL: 1053487; CRS: 9400129000003374]

#### Kiselev 2011 {published data only}

Kiselev AR, Shvarts VA, Posnenkova OM, Gridnev VI, Dovgalevskii P, Oshchepkova EV, et al. Outpatient prophylaxis and treatment of arterial hypertension with application of mobile telephone systems and Internet techniques. *Terapevticheskiĭ Arkhiv* 2011;**83**(4):46-52.

#### Lazev 2004 {published data only}

Lazev A, Vidrine D, Arduino R, Gritz E. Increasing access to smoking cessation treatment in a low-income, HIV-positive population: the feasibility of using cellular telephones. *Nicotine & Tobacco Research* 2004;**6**(2):281-6. [doi: [10.1080/4622200410001676314](https://doi.org/10.1080/4622200410001676314)]

#### Mason 2016 {published data only}

Mason M, Campbell L, Way T, Keyser-Marcus L, Benetsch E, Mennis J, et al. Development and outcomes of a text messaging tobacco cessation intervention with urban adolescents. *Substance Abuse* 2015;**36**(4):500-6. [DOI: [10.1080/08897077.2014.987946](https://doi.org/10.1080/08897077.2014.987946)]

Mason M, Mennis J, Way T, Floyd Campbell L. Real-time readiness to quit and peer smoking within a text message intervention for adolescent smokers: modeling mechanisms of change. *Journal of Substance Abuse Treatment* 2015;**59**:67-73. [DOI: [10.1016/j.jsat.2015.07.009](https://doi.org/10.1016/j.jsat.2015.07.009)]

\* Mason M, Mennis J, Way T, Zaharakis N, Campbell LF, Benetsch E, et al. Text message delivered peer network counseling for adolescent smokers: a randomized controlled trial. *Journal of Primary Prevention* 2016;**37**(5):403-20. [DOI: [10.1007/s10935-016-0439-2](https://doi.org/10.1007/s10935-016-0439-2)]

Mason M, Mennis J, Zaharakis N, Way T. The dynamic role of urban neighborhood effects in a text-messaging adolescent smoking intervention. *Nicotine & Tobacco Research* 2016;**18**(5):1039-45. [DOI: [10.1093/ntr/ntv254](https://doi.org/10.1093/ntr/ntv254)]

#### Mehring 2014 {published data only}

Mehring M, Haag M, Linde K, Wagenpfeil S, Schneider A. Effects of a guided web-based smoking cessation program with telephone counseling: a cluster randomized controlled trial. *Journal of Medical Internet Research* 2014;**16**(9):e218. [CENTRAL: 1051138; CRS: 9400129000003393]

#### Naughton 2012 {published data only}

Naughton F, Prevost A, Gilbert H, Sutton S. Randomized controlled trial evaluation of a tailored leaflet and SMS text message self-help intervention for pregnant smokers (MiQuit). *Nicotine & Tobacco Research* 2012;**14**(5):569-77.

#### Naughton 2017 {published data only}

Cooper S, Foster K, Naughton F, Leonardi-Bee J, Sutton S, Ussher M, et al. Pilot study to evaluate a tailored text message

intervention for pregnant smokers (MiQuit): study protocol for a randomised controlled trial. *Trials* 2015;**16**(1):29. [DOI: <http://dx.doi.org/10.1186/s13063-014-0546-4>]

Naughton F, Cooper S, Foster K, Emery J, Leonardi-Bee J, Sutton S, et al. Large multi-centre pilot randomized controlled trial testing a low-cost, tailored, self-help smoking cessation text message intervention for pregnant smokers (MiQuit). *Addiction* 2017;**112**(7):1238-49. [DOI: [10.1111/add.13802](https://doi.org/10.1111/add.13802)]

**NCT01454999** {unpublished data only}

Jordan P. Tailored texting to enhance a stage-based online tailored program for veterans who smoke: a randomized breakthrough and benchmark trial [personal communication]. Email 2015.

\* NCT01454999. Cell phone-based expert systems for smoking cessation (TXT). [clinicaltrials.gov/ct2/show/NCT01454999](https://clinicaltrials.gov/ct2/show/NCT01454999) (first received 19 October 2011).

**NCT02245308** {published data only}

NCT02245308. Abstinence reinforcement therapy (ART) for homeless veteran smokers. [clinicaltrials.gov/show/NCT02245308](https://clinicaltrials.gov/show/NCT02245308) (first received 19 September 2014). [CRS: 9400131000001050]

**NCT02844296** {published data only}

NCT02844296. Short-bout handgrip exercise for smoking cessation (SHESC). [clinicaltrials.gov/ct2/show/NCT02844296](https://clinicaltrials.gov/ct2/show/NCT02844296) (first received 26 July 2016).

**NCT03177265** {published data only}

NCT03177265. Text messaging to engage and retain veterans in smoking cessation counseling (TiMES). [clinicaltrials.gov/ct2/show/NCT03177265](https://clinicaltrials.gov/ct2/show/NCT03177265) (first received 6 June 2017).

**Obermayer 2004** {published data only}

Obermayer JL, Riley WT, Asif O, Jean-Mary J. College smoking cessation using cell phone text messaging. *Journal of American College Health* 2004;**53**(2):71-8.

**Pechmann 2015** {published data only}

\* Pechmann C, Pan L, Delucchi K, Lakon CM, Prochaska JJ. Development of a Twitter-based intervention for smoking cessation that encourages high-quality social media interactions via automessages. *Journal of Medical Internet Research* 2015;**17**(2):e50. [CRS: 9400131000001029; PUBMED: 25707037]

Prochaska JJ, Pechmann C, Lakon C, Delucchi K. Randomized controlled trial of tweet2quit for smoking cessation. Society for Research on Nicotine and Tobacco 21st Annual Meeting; 2015 Feb 25-28; Philadelphia. 2015. [CRS: 9400131000001026; PA21-2]

**Peng 2013** {published data only}

Peng WB, Schoech D. Evaluation of a web-phone intervention system in changing smoking behavior - a randomized controlled trial. *Journal of Technology in Human Services* 2013;**31**(3):248-68. [CENTRAL: 1053486; CRS: 9400129000000455]

**Pollak 2013** {published data only}

Pollak KI, Lyna P, Bilheimer A, Farrell D, Gao X, Swamy GK, et al. A pilot study testing SMS text delivered scheduled gradual reduction to pregnant smokers. *Nicotine & Tobacco Research* 2013;**15**(10):1773-6. [CENTRAL: 982524; CRS: 9400129000000412; EMBASE: 2014131762; PUBMED: 23569007]

**Riley 2008** {published data only}

Riley W, Obermayer J, Jean-Mary J. Internet and mobile phone text messaging intervention for college smokers. *Journal of American College Health* 2008;**57**(2):245-8.

**Shi 2013** {published data only}

Shi HJ, Jiang XX, Yu CY, Zhang Y. Use of mobile phone text messaging to deliver an individualized smoking behaviour intervention in Chinese adolescents. *Journal of Telemedicine and Telecare* 2013;**19**(5):282-7. [CENTRAL: 984362; CRS: 9400126000000380; PUBMED: 24163238]

**Skov-Ettrup 2013** {published data only}

Skov-Ettrup LS, Dalum P, Ekholm O, Tolstrup JS. Reach and uptake of Internet- and phone-based smoking cessation interventions: results from a randomized controlled trial. *Preventive Medicine* 2014;**62**:38-43. [CENTRAL: 1001289; CRS: 9400129000000774; EMBASE: 2014140700]

\* Skov-Ettrup LS, Dalum P, Tolstrup JS. Internet-based intervention and telephone counselling for smoking cessation: results from a 4-arm randomised controlled trial. *European Journal of Epidemiology* 2013;**28**(S1):S48-9. [CENTRAL: 1011884; CRS: 9400130000000339; EMBASE: 71300831]

**Skov-Ettrup 2014** {published data only}

Skov-Ettrup LS, Ringgaard LW, Dalum P, Flensburg-Madsen T, Thygesen LC, Tolstrup JS. Comparing tailored and untailored text messages for smoking cessation: a randomized controlled trial among adolescent and young adult smokers. *Health Education Research* 2014;**29**(2):195-205. [CRS: 94001290000003462; PUBMED: 24399268]

**Skov-Ettrup 2016** {published data only}

Skov-Ettrup LS, Dalum P, Bech M, Tolstrup JS. The effectiveness of telephone counselling and Internet- and text-message-based support for smoking cessation: results from a randomized controlled trial. *Addiction* 2016;**111**(7):1257-66. [DOI: [10.1111/add.13302](https://doi.org/10.1111/add.13302)]

**Snider 2011** {published data only}

Snider J. Cell phone text messaging may boost smoking quit rates. *Journal of the American Dental Association* 2011;**142**(8):901-2.

**Stanczyk 2014** {published data only}

\* Stanczyk N, Bolman C, Van Adrichem M, Candel M, Muris J, De Vries H. Comparison of text and video computer-tailored interventions for smoking cessation: randomized controlled trial. *Journal of Medical Internet Research* 2014;**16**(3):e69. [CENTRAL: 1046762; CRS: 94001290000002402]

Stanczyk NE, Bolman C, Muris JW, De Vries H. Study protocol of a Dutch smoking cessation e-health program. *BMC Public Health*

2011;**11**:847. [CENTRAL: 814250; CRS: 940010000000130; PUBMED: 22059446]

Stanczyk NE, Crutzen R, Bolman C, Muris J, De Vries H. Influence of delivery strategy on message-processing mechanisms and future adherence to a Dutch computer-tailored smoking cessation intervention. *Journal of Medical Internet Research* 2013;**15**(2):e28. [CENTRAL: 873085; CRS: 9400107000001436; PUBMED: 23388554]

#### **Vidrine 2006** {published data only}

\* Vidrine D, Arduino R, Gritz E. Impact of a cell phone intervention on mediating mechanisms of smoking cessation in individuals living with HIV/AIDS. *Nicotine & Tobacco Research* 2006;**8**(S1):S103-8.

Vidrine D, Arduino R, Lazev A, Gritz E. A randomized trial of a proactive cellular telephone intervention for smokers living with HIV/AIDS. *AIDS* 2006;**20**:253-60. [ISSN 0269-9370]

#### **Vilaplana 2014** {published data only}

Vilaplana J, Solsona F, Abella F, Cuadrado J, Alves R, Mateo J. S-PC: an e-treatment application for management of smoke-quitting patients. *Computer Methods and Programs in Biomedicine* 2014;**115**(1):33-45. [CENTRAL: 988111; CRS: 9400050000000047; EMBASE: 2014296941; PUBMED: 24742965]

#### **Wizner 2009** {published data only}

Wizner B, Gaciong Z, Narkiewicz K, Grodzicki T. Education using SMS increases efficacy of treatment of hypertensive patients [Zwiększenie skuteczności terapii hipotensyjnej u pacjentów z nadciśnieniem tętniczym dzięki edukacji przez SMS]. *Nadciśnienie Tętnicze* 2009;**13**(3):147-57.

#### **Ybarra 2012** {published and unpublished data}

\* Ybarra M, Bagci Bosi AT, Korchmaros J, Emri S. A text messaging-based smoking cessation program for adult smokers: randomized controlled trial. *Journal of Medical Internet Research* 2012;**14**(6):e172. [CENTRAL: 863878; CRS: 9400107000000182; PUBMED: 23271159]

Ybarra ML, Holtrop JS, Bagci Bosi AT, Emri S. Design considerations in developing a text messaging program aimed at smoking cessation. *Journal of Medical Internet Research* 2012;**14**(4):e103. [CRS: 9400123000016823; PUBMED: 22832182]

Ybarra ML, Holtrop JS, Bosi AT, Bilir N, Korchmaros JD, Emri AK. Feasibility and acceptability of a text messaging-based smoking cessation program in Ankara, Turkey. *Journal of Health Communication* 2013;**18**(8):960-73. [CRS: 9400126000000575; PUBMED: 23627304]

#### **Ybarra 2013** {published and unpublished data}

\* Ybarra M, Holtrop J, Prescott T, Rahbar M, Strong D. Pilot RCT results of Stop My Smoking (SMS) USA: a text messaging-based smoking cessation program for young adults. *Nicotine & Tobacco Research* 2013;**15**(8):1388-99.

Ybarra ML, Holtrop JS, Prescott TL, Strong D. Process evaluation of a mHealth program: lessons learned from Stop my Smoking USA, a text messaging-based smoking cessation program for young adults. *Patient Education and Counseling*

2014;**97**(2):236-43. [CRS: 9400131000000800; PUBMED: 25103183]

#### **Yuhongxia 2011** {published data only}

Yuhongxia L. The compliance of varenicline usage and the smoking abstinence rate via mobile phone text messaging combine with varenicline: a single-blind, randomised control trial. *Respirology* 2011;**16**:46-7.

### References to ongoing studies

#### **Cambon 2017** {published data only}

Cambon L, Bergman P, Le Faou A, Vincent I, Le Maitre B, Pasquereau A, et al. Study protocol for a pragmatic randomised controlled trial evaluating efficacy of a smoking cessation e-'Tabac Info Service': ee-TIS trial. *BMJ Open* 2017;**7**(2):e013604. [DOI: [10.1136/bmjopen-2016-013604](https://doi.org/10.1136/bmjopen-2016-013604)]

#### **Collins 2017** {published data only}

Collins BN, Lepore SJ. Babies living safe & smokefree: randomized controlled trial of a multilevel multimodal behavioral intervention to reduce low-income children's tobacco smoke exposure. *BMC Public Health* 2017;**17**(1):249. [DOI: [10.1186/s12889-017-4145-7](https://doi.org/10.1186/s12889-017-4145-7)]

#### **CTRI201801011643** {published data only}

CTRI/2018/01/011643. Tobacco cessation at non communicable disease clinics [Development of tobacco cessation training package and assessing its impact on tobacco use behavior of patients attending non communicable diseases clinics of Punjab]. [ctri.nic.in/Clinicaltrials/showallp.php?mid1=20950&EncHid=&userName=Tobacco%20Cessation%20at%20Non%20Communicable%20Disease%20Clinics](http://ctri.nic.in/Clinicaltrials/showallp.php?mid1=20950&EncHid=&userName=Tobacco%20Cessation%20at%20Non%20Communicable%20Disease%20Clinics) (first received 31 January 2018).

#### **CTRI201803012401** {published data only}

CTRI/2018/03/012401. A clinical trial to study the effect of WhatsApp and pamphlet based quit smoking interference among software professionals in Bengaluru City [Effectiveness of WhatsApp based smoking cessation intervention among software professional in Bengaluru City: a randomised controlled trial]. [ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=21448&EncHid=&userName=CTRI/2018/03/012401](http://ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=21448&EncHid=&userName=CTRI/2018/03/012401) (first received 07 March 2018).

#### **Graham 2016** {published data only}

Graham AL, Jacobs MA, Cohn AM, Cha S, Abrams LC, Papandonatos GD, et al. Optimising text messaging to improve adherence to web-based smoking cessation treatment: a randomised control trial protocol. *BMJ Open* 2016;**6**(3):e010687. [DOI: [10.1136/bmjopen-2015-010687](https://doi.org/10.1136/bmjopen-2015-010687)]

#### **Graham 2017** {published data only}

Graham A, Burke M, Jacobs M, Cha S, Croghan I, Schroeder D, et al. An integrated digital/clinical approach to smoking cessation in lung cancer screening: study protocol for a randomized controlled trial. *Trials* 2017;**18**(1):568. [DOI: [10.1186/s13063-017-2312-x](https://doi.org/10.1186/s13063-017-2312-x)]

**ISRCTN1154315** *{published data only}*

ISRCTN1154315. Efficacy of a smoking cessation intervention using smartphones. [isrctn.com/ISRCTN1154315](http://isrctn.com/ISRCTN1154315) (first received 04 April 2018).

**ISRCTN11318024** *{published data only}*

ISRCTN11318024. Impact of a smartphone application on smoking cessation: a randomized controlled trial. [isrctn.com/ISRCTN11318024](http://isrctn.com/ISRCTN11318024) (first received 26 April 2018).

**ISRCTN15396225** *{published data only}*

ISRCTN15396225. Evaluation of the effectiveness of a text-based mHealth smoking cessation intervention among high school students in Sweden. [isrctn.com/ISRCTN15396225](http://isrctn.com/ISRCTN15396225) (first received 10 October 2017).

**ISRCTN16022919** *{published data only}*

ISRCTN16022919. Mobile health interventions for smoking cessation services uptake and smoking cessation: a factorial randomised trial in Thailand. [isrctn.com/ISRCTN16022919](http://isrctn.com/ISRCTN16022919) (first received 11 November 2016).

**ISRCTN17964518** *{published data only}*

ISRCTN17964518. Evaluation of the "Stop Tabac" Android phone application. [isrctn.com/ISRCTN17964518](http://isrctn.com/ISRCTN17964518) (first received 24 June 2015).

**ISRCTN33869008** *{published data only}*

ISRCTN33869008. Mobile phone-based smoking cessation intervention for patients with elective surgery. [isrctn.com/ISRCTN33869008](http://isrctn.com/ISRCTN33869008) (first received 17 August 2018).

**NCT01982110** *{published data only}*

NCT01982110. A mindfulness based application for smoking cessation. [clinicaltrials.gov/show/NCT01982110](http://clinicaltrials.gov/show/NCT01982110) (first received 13 November 2013). [CRS: 940013100001042]

**NCT01990079** *{published data only}*

NCT01990079. Use of technological advances to prevent smoking relapse among smokers with PTSD. [clinicaltrials.gov/show/NCT01990079](http://clinicaltrials.gov/show/NCT01990079) (first received 21 November 2013). [CRS: 940013100001036]

**NCT01995097** *{published data only}*

NCT01995097. BABY STEPS II: SMS scheduled gradual reduction text messages to help pregnant smokers quit. [clinicaltrials.gov/ct2/show/NCT01995097](http://clinicaltrials.gov/ct2/show/NCT01995097) (first received 26 November 2013).

**NCT02037360** *{published data only}*

NCT02037360. Mobile mindfulness training for smoking cessation. [clinicaltrials.gov/show/NCT02037360](http://clinicaltrials.gov/show/NCT02037360) (first received 15 January 2014). [CRS: 940013100001048]

**NCT02218281** *{published data only}*

NCT02218281. Developing a smartphone app with mindfulness training for teen smoking cessation. [clinicaltrials.gov/show/NCT02218281](http://clinicaltrials.gov/show/NCT02218281) (first received 15 January 2014). [CRS: 940013100001087]

**NCT02218944** *{published data only}*

NCT02218944. Response inhibition training in smoking cessation. [clinicaltrials.gov/ct2/show/NCT02037360](http://clinicaltrials.gov/ct2/show/NCT02037360) (first received 15 January 2014).

**NCT02237898** *{published data only}*

NCT02237898. Harnessing the power of technology: MOMBA for postpartum smoking. [clinicaltrials.gov/show/NCT02237898](http://clinicaltrials.gov/show/NCT02237898) (first received 11 September 2014). [CRS: 940013100001085]

**NCT02420015** *{published data only}*

NCT02420015. Mobile health technology to enhance abstinence in smokers with schizophrenia. [clinicaltrials.gov/ct2/show/NCT02420015](http://clinicaltrials.gov/ct2/show/NCT02420015) (first received 17 April 2015).

**NCT02665208** *{published data only}*

NCT02665208. A pilot text messaging intervention to reduce smoking in office-based buprenorphine and inpatient detoxification patients. [clinicaltrials.gov/ct2/show/NCT02665208](http://clinicaltrials.gov/ct2/show/NCT02665208) (first received 27 January 2016).

**NCT02724462** *{published data only}*

NCT02724462. Trial of an innovative smartphone intervention for smoking cessation. [clinicaltrials.gov/ct2/show/NCT02665208](http://clinicaltrials.gov/ct2/show/NCT02665208) (first received 27 January 2016).

**NCT02840513** *{published data only}*

NCT02840513. Smartphone app and CO self-monitoring for smoking cessation. [clinicaltrials.gov/ct2/show/NCT02840513](http://clinicaltrials.gov/ct2/show/NCT02840513) (first received 21 July 2016).

**NCT02901171** *{published data only}*

NCT02901171. The contribution of a smartphone application to acceptance and commitment therapy group treatment for smoking cessation. [clinicaltrials.gov/show/NCT02901171](http://clinicaltrials.gov/show/NCT02901171) (first received 21 July 2016).

**NCT03021655** *{published data only}*

NCT03021655. A pilot randomized control trial to help youth smokers to quit smoking. [clinicaltrials.gov/ct2/show/NCT03021655](http://clinicaltrials.gov/ct2/show/NCT03021655) (first received 16 January 2017).

**NCT03038542** *{published data only}*

NCT03038542. Quit4hlth: enhancing tobacco and cancer control through framed text messages. [clinicaltrials.gov/ct2/show/NCT03021655](http://clinicaltrials.gov/ct2/show/NCT03021655) (first received 16 January 2017).

**NCT03191019** *{published data only}*

NCT03191019. A mobile-phone based intervention to support smoking cessation among Chilean women. [clinicaltrials.gov/ct2/show/NCT03191019](http://clinicaltrials.gov/ct2/show/NCT03191019) (first received 19 June 2017).

**NCT03445507** *{published data only}*

NCT03445507. Effectiveness of a chat bot for smoking cessation: a pragmatic trial in primary care. [clinicaltrials.gov/show/NCT03445507](http://clinicaltrials.gov/show/NCT03445507) (first received 19 June 2017).

**NCT03495622** *{published data only}*

NCT03495622. Effectiveness of a combined CHW and text messaging-based tobacco intervention in India.



clinicaltrials.gov/show/NCT03495622 (first received 12 April 2018).

**NCT03538938** {published data only}

NCT03538938. Improving quitline support study (IQS). clinicaltrials.gov/show/NCT03538938 (first received 28 May 2018).

**NCT03552978** {published data only}

NCT03552978. Tech and telephone smoking cessation treatment for young veterans with PTSD. clinicaltrials.gov/show/NCT03552978 (first received 12 June 2018).

**NCT03553173** {published data only}

NCT03553173. So-Lo-Mo intervention applied to the smoking cessation process. clinicaltrials.gov/ct2/show/NCT03553173 (first received 12 June 2018).

**Valdivieso-Lopez 2013** {published data only}

Valdivieso-López E, Flores-Mateo G, Molina-Gómez JD, Rey-Reñones C, Uriarte ML, Duch J, et al. Efficacy of a mobile application for smoking cessation in young people: study protocol for a clustered, randomized trial. *BMC Public Health* 2013;**13**(1):704.

**Weng 2018** {published data only}

Weng X, Wang M, Suen Y, Li W, Wu Y, Cheung D, et al. Comparing different intensities of active referral to smoking cessation services in promoting smoking cessation among community smokers: a study protocol of a cluster randomized controlled trial. *BMC Public Health* 2018;**18**(1):830.

**Additional references**

**Abroms 2011**

Abroms LC, Padmanabhan N, Thaweethai L, Phillips T. iPhone apps for smoking cessation: a content analysis. *American Journal of Preventive Medicine* 2011;**40**(3):279-85.

**Armanasco 2017**

Armanasco AA, Miller YD, Fjeldsoe BS, Marshall AL. Preventive health behavior change text message interventions: a meta-analysis. *American Journal of Preventive Medicine* 2017;**52**(3):391-402.

**Boland 2017**

Boland V, Mattick R, McRobbie H, Siahpush M, Courtney R. "I'm not strong enough; I'm not good enough. I can't do this, I'm failing": a qualitative study of low-socioeconomic status smokers' experiences with accessing cessation support and the role for alternative technology-based support. *International Journal for Equity in Health* 2017;**16**:196.

**Byambasuren 2018**

Byambasuren O, Sanders S, Beller E, Glasziou P. Prescribable mHealth apps identified from an overview of systematic reviews. *npj Digital Medicine* 2018;**1**(1):12.

**Covidence [Computer program]**

Veritas Health Innovation. Covidence. Version accessed 29 October 2018. Melbourne, Australia: Veritas Health Innovation.

**Dirieto 2016**

Dirieto A, Carraça E, Rawstorn J, Whittaker R, Maddison R. mHealth technologies to influence physical activity and sedentary behaviors: behavior change techniques, systematic review and meta-analysis of randomized controlled trials. *Annals of Behavioral Medicine* 2016;**51**(2):226-39.

**Guerrero 2013**

Guerrero C, Cairns J, Roberts I, Rodgers A, Whittaker R, Free C. The cost-effectiveness of smoking cessation support delivered by mobile phone text messaging: Txt2stop. *European Journal of Health Economics* 2013;**14**:789-97.

**Higgins 2003**

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60.

**Higgins 2017**

Higgins JP, Altman DG, Sterne JA (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Churchill R, Chandler J, Cumpston MS (editors), *Cochrane Handbook for Systematic Reviews of Interventions* version 5.2.0 (updated June 2017), Cochrane, 2017. Available from [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook).

**IHME 2018**

Institute for Health Metrics and Evaluation. Findings from the Global Burden of Disease Study 2017. Seattle, WA: IHME, 2018.

**ITU 2018**

International Telecommunications Union. Measuring the Information Society Report 2018 - Volume 1. [www.itu.int/en/ITU-D/Statistics/Documents/publications/misr2018/MISR-2018-Vol-1-E.pdf](http://www.itu.int/en/ITU-D/Statistics/Documents/publications/misr2018/MISR-2018-Vol-1-E.pdf) (accessed 29 March 2019).

**Lunde 2018**

Lunde P, Nilsson BB, Bergland A, Kværner KJ, Bye A. The effectiveness of smartphone apps for lifestyle improvement in noncommunicable diseases: systematic review and meta-analyses. *Journal of Medical Internet Research* 2018;**20**(5):e162.

**Schoeppe 2016**

Schoeppe S, Alley S, Van Lippevelde W, Bray NA, Williams SL, Duncan MJ, et al. Efficacy of interventions that use apps to improve diet, physical activity and sedentary behaviour: a systematic review. *International Journal of Behavioral Nutrition and Physical Activity* 2016;**13**(1):127.

**Schünemann 2017**

Schünemann HJ, Oxman AD, Higgins JP, Vist GE, Glasziou P, Akl E, et al. on behalf of the Cochrane GRADEing Methods Group and the Cochrane Statistical Methods Group. Chapter 11: Completing 'Summary of findings' tables and grading the confidence in or quality of the evidence. In: Higgins JPT, Churchill R, Chandler J, Cumpston MS (editors), *Cochrane Handbook for Systematic Reviews of Interventions* version

5.2.0 (updated June 2017). Cochrane, 2017. Available from [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook).

#### Scott-Sheldon 2016

Scott-Sheldon LA, Lantini RC, Jennings EG, Thind H, Rosen RK, Salmoirago-Blotcher E, et al. Text messaging-based interventions for smoking cessation: a systematic review and meta-analysis. *JMIR MHealth and UHealth* 2016;**4**(2):e49.

#### Thakkar 2016

Thakkar J, Kurup R, Laba TL, Santo K, Thiagalingam A, Rodgers A, et al. Mobile telephone text messaging for medication adherence in chronic disease: a meta-analysis. *JAMA Internal Medicine* 2016;**176**(3):340-9.

#### Zhao 2016

Zhao J, Freeman B, Li M. Can mobile phone apps influence people's health behavior change? An evidence review. *Journal of Medical Internet Research* 2016;**18**(11):e287.

### References to other published versions of this review

#### Whittaker 2009

Whittaker R, Borland R, Bullen C, Lin RB, McRobbie H, Rodgers A. Mobile phone-based interventions for smoking cessation. *Cochrane Database of Systematic Reviews* 2009, Issue 4. [DOI: [10.1002/14651858.CD006611.pub2](https://doi.org/10.1002/14651858.CD006611.pub2)]

#### Whittaker 2012

Whittaker R, McRobbie H, Bullen C, Borland R, Rodgers A, Gu Y. Mobile phone-based interventions for smoking cessation. *Cochrane Database of Systematic Reviews* 2012, Issue 11. [DOI: [10.1002/14651858.CD006611.pub3](https://doi.org/10.1002/14651858.CD006611.pub3)]

#### Whittaker 2016

Whittaker R, McRobbie H, Bullen C, Rodgers A, Gu Y. Mobile phone-based interventions for smoking cessation. *Cochrane Database of Systematic Reviews* 2016, Issue 4. [DOI: [10.1002/14651858.CD006611.pub4](https://doi.org/10.1002/14651858.CD006611.pub4)]

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Abroms 2014

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> Internet. Individuals who were searching on Google with keywords related to quitting smoking saw study ads in conjunction with their search results.</p> <p><b>Dates of study:</b> 2011-13</p>
Participants	<p><b>Baseline characteristics (n = 503)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 35.7 years</li> <li>• Female: 65.6%</li> <li>• High school or lower education: 21.9%</li> <li>• FTND: 5.33</li> <li>• White: 78.5%</li> </ul> <p><b>Inclusion criteria:</b> to be eligible for the study, participants were required to (1) be <math>\geq 18</math> years of age; (2) smoke <math>\geq 5</math> cigarettes/day; (3) have a US mailing address; (4) have a working e-mail address; (5) have a cell phone number with an unlimited SMS (i.e. text messaging) plan; (6) express an interest in quitting smoking within the next month; and (7) not be pregnant.</p> <p><b>Exclusion criteria:</b> pregnant</p>
Interventions	<p><b>Text2Quit:</b> automated, tailored, interactive and bidirectional text messaging programme that was supported by email and web portal. Based on social cognitive theory and practice guidelines. Duration 6 months with decreasing frequency of messages.</p> <p><b>Control:</b> weblink to smokefree.gov website, weblink to a guidebook, and study related reminder text messages</p>
Outcomes	<p><b>Definition of abstinence:</b> 6-month biochemically confirmed repeat point prevalence</p>

**Abroms 2014** (Continued)

Funding source	This research was supported by grant no. 5K07 CA124579-02 and the American Recovery and Reinvestment Act supplement to Dr. Lorien Abroms, from the National Cancer Institute of the NIH. Support also came from an award from the Department of Prevention and Community Health at the George Washington University School of Public Health and Health Services to Dr. Lorien Abrom	
Conflicts of interest	The George Washington University/Lorien Abroms has licensed the Text@Quit program to Voxiva Inc; Dr Abroms has stock options in Voxiva Inc	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-generated
Allocation concealment (selection bias)	Low risk	Central randomisation online
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically verified
Incomplete outcome data (attrition bias) All outcomes	Low risk	At 6 months, 52 lost to follow-up in control group (21.6%) and 70 lost in intervention group (26.7%)
Other bias	Unclear risk	Some issues with fraudulent enrolment at outset of study, corrected process once detected

**Abroms 2017**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> participants were recruited from Text4Baby (national text message health information programme for pregnant women) subscribers</p> <p><b>Study dates:</b> 2015-16</p>
Participants	<p><b>Baseline characteristics (n = 497)</b></p> <ul style="list-style-type: none"> <li>• Average age: 26.31 years</li> <li>• Female: 100%</li> <li>• High school or less: 59.96%</li> <li>• White: 63.18%</li> <li>• FTCD: 2.48</li> </ul> <p><b>Inclusion criteria:</b> Text4baby subscribers were eligible if they had a due date 8 weeks in the future at the time of sending. Subscribers were eligible for the Quit4baby study if they had a cell phone for their personal use, were willing to receive text messages on their mobile phone, were aged <math>\geq 14</math> years, were currently pregnant, and had smoked at least 1 puff of a cigarette in the past 2 weeks.</p>

**Abroms 2017** (Continued)

**Exclusion criteria:** Text4baby subscribers from California, Oklahoma, Ohio, and Louisiana were excluded because Quit4baby was already available in those states.

Interventions	<p><b>Quit4Baby:</b> Text4baby plus Quit4baby. Quit4baby: 1-8 text messages/day based on social cognitive theory guidelines for SC in pregnancy, that lasts 3 months</p> <p><b>Control:</b> Text4Baby: 3 health information text messages each week for pregnant women and mothers</p>
Outcomes	<b>Definition of abstinence:</b> self-reported 30-day abstinence at 6 months (only 3-month abstinence was biochemically verified)
Funding source	This research was supported by the National Institute on Drug Abuse of NIH. Support also came from an award from the Department of Prevention and Community Health at the Milken Institute School of Public Health at George Washington University
Conflicts of interest	Dr Abroms has stock in Wellpass Inc (formerly Voxiva Inc) and has licensed Text2Quit and Quit4Baby to WellPass Inc. Dr Johnson is employed by Wellpass Inc, the company that operates Text4Baby and Quit4Baby. Ms Bushar is employed by ZERO TO THREE, a partner operating the Text4Baby service. Dr Brandon has served as a paid consultant to Voxiva In and has received research support from Pfizer Inc.
Notes	

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were recruited online, sequence was generated by REDCap application
Allocation concealment (selection bias)	Low risk	Consented and baseline survey completed then computer allocation using REDCap computerised allocation module
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Contact with investigators was minimal
Incomplete outcome data (attrition bias) All outcomes	Low risk	Follow-up rates at 6 months were 71% and 72% and ITT analysis and imputation conducted

**Alessi 2017**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Study grouping:</b> parallel group</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> through E-mail, flyers, and print advertisements</p> <p><b>Dates of study:</b> 2012-2014</p>
Participants	<p><b>Baseline characteristics (n = 90)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 45</li> <li>• Female: 59% (N = 53)</li> <li>• High school or lower education: not stated</li> </ul>



**Alessi 2017** (Continued)

- FTND: approx 3
- White: 74% (N = 67)

**Inclusion criteria:** inclusion criteria were (1)  $\geq 10$  cigarettes daily verified by  $\text{CO} \geq 8$  ppm, (2) no past-year abstinence  $> 3$  months, (3) intent to quit within 3 weeks (score  $\geq 7$  out of 10, "How much do you want to quit smoking within the next 3 weeks?"<sup>15</sup>), (4) aged  $\geq 18$  years, and (5) mailing address and valid photo ID

**Exclusion criteria:** Exclusion criteria were (1) past month behavioral or pharmacotherapy for smoking, (2) serious and unstable psychiatric illness (e.g. schizophrenia, non-nicotine substance use disorder) or medical disease, or contraindication for transdermal nicotine, (3) pregnant, nursing a child, or not using effective contraceptive if female, (4) ongoing use of monoamine oxidase inhibitors, antipsychotics, mood stabilisers, bupropion, or naltrexone, and (5) not English-speaking

**Interventions**

**mHealth monitoring:** for all participants, brief counselling (~10 min) was scheduled to occur twice weekly for 4 weeks by phone. Discussion included personal reasons for quitting, skills-based items, and craving control strategies. Self-reported smoking status was documented. The study also provided 8 weeks of transdermal nicotine (typically 21 mg patches for 4 weeks, 14 mg for 2 weeks, and 7 mg for 2 weeks). All participants were instructed that an IVR system would send prompts to conduct CO self-tests up to 3 times daily between 7 a.m. and 10 p.m. for the next 4 weeks, with the exact number and timing not disclosed. When prompted, participants used the video-record function on their study cell phone (with a front-facing lens) to record the CO self-test process, and sent the date and time-stamped video to research staff using multimedia messaging. Participants also reported the CO results and number of cigarettes smoked using the IVR. Video test results were compared against IVR reports to confirm accuracy (confirmed in all but 2 instances).

**mHealth reinforcement:** as above, plus mHealth reinforcement participants earned chances for prizes contingent on on-time and smoking-negative breath tests ( $\text{CO} \leq 6$  ppm). Earnings were determined immediately via computer algorithm during IVR calls, and were available for redemption after IVR reports were confirmed against video clips.

**Outcomes**

**Definition of abstinence:** 6-month biochemically confirmed repeat point prevalence

**Funding source**

National Institutes of Health grants R21-DA029215 and R01-DA01344

**Conflicts of interest**

None declared

**Notes**
**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "On the target quit date, participants (N = 90) were randomly assigned (allocated 1:1) to one of two treatment conditions using an urn procedure 34 and stratified on at least one smoking-negative CO during baseline"
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically verified.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Follow-up rates did not differ between conditions ( $P > 0.05$ ). Reinforcement group (n = 45) follow-up questionnaires: 38, samples: 33; usual care group (n = 45) questionnaires 43, samples 39. Used ITT analysis

**Augustson 2017**

Methods	<b>Study design:</b> RCT  <b>Country:</b> China  <b>Recruitment:</b> recruited from subscribers to Nokia Life Tools on Nokia phones in both urban and rural areas of China's Zhejiang, Heilongjiang, and Shaanxi provinces  <b>Date of study:</b> 2013	
Participants	<b>Baseline characteristics (n = 8000)</b> <ul style="list-style-type: none"> <li>• Female %: not stated</li> <li>• Mean age: not stated</li> <li>• High school or lower education: not stated</li> <li>• FTND: not stated</li> <li>• White: not stated</li> </ul> <b>Inclusion criteria:</b> Nokia Life Tools users; adult smokers  <b>Exclusion criteria:</b> none specifically stated	
Interventions	<b>High-frequency text contact (HFTC):</b> 91 messages during the 6 weeks; 3 messages/day for weeks 1 and 2, 2/day for weeks 3-5, and 1/day for week 6. At the end of each text message, participants in both groups were offered the opportunity to cancel the service via text. The text messages provided encouragement, practical advice to help maintain cessation, and information on the health effects of smoking.  <b>Low-frequency text contact (LFTC):</b> 1 text message/week, for a total of 6 text messages during the 6-week intervention period; a subset of text messages on smoking's health effects	
Outcomes	<b>Definition of abstinence:</b> 7-day point prevalence abstinence self-reported via text message at 6-month follow-up	
Funding source	National Cancer Institute, National Institutes of Health, U.S. Dept of Health Human Services	
Conflicts of interest	None declared	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Quote: "Intervention participants who opted into the phase 3 SC trial were randomly assigned to the intervention or comparison group (n ¼ 4000 in each group)"  Therefore, participants were randomly assigned but it was not stated how
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessed by text message with no face-to-face contact  Quote: "participants were not aware of the separate intervention arms and therefore did not know what group they were assigned to"
Incomplete outcome data (attrition bias)	High risk	Loss to follow-up clearly stated but > 50%; 58.4% of intervention group and 56.1% of control group lost to follow-up at 6-month assessment. ITT analysis

**Augustson 2017** (Continued)  
 All outcomes

**Baskerville 2018**

Methods	<b>Study design:</b> RCT  <b>Country:</b> Canada  <b>Setting:</b> recruited through web-based media including Facebook, Google, and other sources  <b>Study dates:</b> 2014-15	
Participants	<b>Baseline characteristics (n = 1599)</b> <ul style="list-style-type: none"> <li>• Age: 49.1% aged 19-23 years</li> <li>• Female: 45.6%</li> <li>• High school or lower education: 44.5%</li> <li>• Moderate to high nicotine dependence: 26.5%</li> <li>• White: 73%</li> <li>• Smokes at least a packet a day: 25.6%</li> </ul> <p><b>Inclusion criteria:</b> aged 19-29 years, smoked cigarettes daily, resided in Canada, were considering quitting smoking in the next 30 days, had an Android (version 2.0-5.0) or iPhone (version 4.0-7.0) smartphone, were able to provide informed consent, were able to comprehend English, and were not referred to the study by an existing study participant.</p> <p><b>Exclusion criteria:</b> not explicitly stated</p>	
Interventions	<p><b>Crush the Crave:</b> a comprehensive and evidence-informed SC smartphone app; enabled users to customise a quit plan by choosing a QD and whether to quit or reduce every week; reminders of money saved and health improvements; contingency reinforcement with milestones tracked as rewards, choose to share to Facebook/Twitter; Facebook community for additional support; supportive messages and inspirational photos; recording smoking; feedback; web-based distractions; evidence-informed information for relapses and cravings; access to cessation services</p> <p><b>Control:</b> standard print-based self-help guide 'On the Road to Quitting' for young adult smokers</p>	
Outcomes	<b>Definition of abstinence:</b> self-reported continuous 6-month abstinence	
Funding source	Health Canada, Federal Tobacco Control Strategy and a grant from the Canadian Institutes of Health Research	
Conflicts of interest	NBB received salary support from the Canadian Cancer Society Research Institute	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation procedure
Allocation concealment (selection bias)	Low risk	Participants were blinded to group allocation and were not aware of which was the control and intervention condition. Investigators were blinded to group allocation until completion of the trial after initial analysis of the primary and secondary outcomes.

**Baskerville 2018** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Contact with investigators was minimal
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT analysis was conducted. High rates of attrition, but no significant difference between groups. Follow-up at 6 months was 60.48%, however complete case follow-up was considered at both 3 and 6 months for primary outcomes data; 43.2% intervention and 47.6% control groups (no significant difference)

**BinDhim 2018**

Methods	<b>Study design:</b> RCT  <b>Countries:</b> USA, Australia, UK and Singapore  <b>Recruitment:</b> users of the Apple App Store in the 4 countries were recruited passively via the app's download page in the Apple App Store  <b>Study date:</b> 2014	
Participants	<b>Baseline characteristics (n = 684)</b> <ul style="list-style-type: none"> <li>• Mean age: 28.3 (SD 10.0)</li> <li>• Female: 55% (N = 376)</li> <li>• &lt; graduate level education: 46.3% (N = 317)</li> <li>• FTND: proportion 6-10 (high-very high): 36.7% (N = 251)</li> <li>• White: not stated</li> </ul> <b>Inclusion criteria:</b> the eligibility criteria were daily smokers of cigarettes, $\geq 18$ years and from USA, UK, Singapore, Australia  <b>Exclusion criteria:</b> occasional smokers and users of other tobacco products	
Interventions	<b>SSC app:</b> decision aid app that included 4 main components that made optimal use of smartphone features: (1) mandatory information about quitting options, with their benefits and harms; (2) daily motivational messages using push notifications sent from the study server, (3) a quitting diary and (4) a quitting benefits tracker. The decision-aid app allowed smokers to freely choose a quit method through a structured process of weighing up the available options and their benefits and harms.  <b>Control App:</b> both groups encouraged to set QD. App with information about quitting only	
Outcomes	<b>Definition of abstinence:</b> self-reported continuous abstinence at 6 months	
Funding source	The app was developed by NFB as part of a PhD degree, advertisement was covered by a small fund from the PhD sponsor (Ministry of Education, Saudi Arabia)	
Conflicts of interest	None declared	

## Notes

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The study app automatically randomised eligible participants (daily cigarette smokers, aged 18 years and above and from the four countries) to either the intervention or the control sub-app using stratified block (age, gen-

**BinDhim 2018** (Continued)

		der, country) randomisation. The strata were defined by age, country and gender."
Allocation concealment (selection bias)	Low risk	Quote: "The study app automatically randomised eligible participants (daily cigarette smokers, aged 18 years and above and from the four countries) to either the intervention or the control sub-app using stratified block (age, gender, country) randomisation. The strata were defined by age, country and gender."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All participant involvement was remote, through the apps, in both study arms
Incomplete outcome data (attrition bias) All outcomes	Low risk	Numbers reported, similar in both groups (289/342 completed follow-up in control group, 294/342 in intervention group). ITT analysis, imputation of missing data including sensitivity analysis and all missing as smoking

**Bock 2013**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> advertisements in local media outlets, Internet sites, radio programmes, asking interested individuals to call or text</p> <p><b>Study date:</b> 2011</p>
Participants	<p>Baseline characteristics (<b>n = 60</b>)</p> <ul style="list-style-type: none"> <li>• Mean age: 30.7 (9.0)</li> <li>• Female: 57% (N = 34)</li> <li>• High school or less: 30% (N = 18)</li> <li>• FTND: not stated</li> <li>• White: 66% (N = 40)</li> </ul> <p><b>Inclusion criteria:</b> current daily smoker, interested in quitting smoking in the next 30 days, have a mobile phone with SMS text messaging capability, use SMS messaging at least once monthly</p> <p><b>Exclusion criteria:</b> not explicitly stated</p>
Interventions	<p>All participants received a single, individual 30-min SC counselling session.</p> <p><b>TXT-2-Quit:</b> an 8-week programme with 1-4 text messages/day (depending on quit stage). SC messages were tailored to the participant's stage of SC, with specialised messages provided on demand, based on user requests for additional support, and an optional peer-to-peer social support network</p> <p><b>Control:</b> an 8-week programme of daily non-smoking-related text messages</p>
Outcomes	Primary outcome: self-reported 7-day point-prevalence abstinence at 6 months
Funding source	National Institute on Drug Abuse
Conflicts of interest	None declared
Notes	

**Bock 2013** (Continued)

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Simple randomisation via computerised random number generator
Allocation concealment (selection bias)	Low risk	Assignments in a sealed envelope delivered after completion of the baseline data collection
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Questionnaires were filled in online with minimal investigator contact
Incomplete outcome data (attrition bias) All outcomes	Low risk	Only 2 participants in control group appeared to be missing at 6 months; ITT analysis presented

**Borland 2013**

Methods	<p><b>Study design:</b> RCT. Participants were not pre-committed to consider using the interventions they were offered</p> <p><b>Country:</b> Australia</p> <p><b>Recruitment:</b> from those having recently sought cessation assistance (mainly Quitline callers who were not seeking help from a counsellor) via the study website, and from a cold-contacted sample taken from two Internet survey panels</p> <p><b>Study dates:</b> 2008-09</p>
Participants	<p><b>Baseline characteristics (n = 3530)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 42.1</li> <li>• Female: 60% (N = 2118)</li> <li>• High school or less: not stated</li> <li>• Smoking: 16.9 cigarettes/day</li> <li>• White: not stated</li> </ul> <p><b>Inclusion criteria:</b> none stated</p> <p><b>Exclusion criteria:</b> none stated</p>
Interventions	<p><b>onQ programme:</b> provides a stream of SMS messages that mix snippets of advice on strategy and motivational messages. The user can interact with it by indicating their stage of quitting so that appropriate stage-specific messages are sent, and once quit can also call up messages in crisis situations.</p> <p><b>QuitCoach:</b> a personalised, automated tailored cessation programme delivered via the Internet. It generates letters of advice based on answers to an assessment questionnaire, including suggestions about strategy and motivational messages. It also provides further untailored supplementary resources.</p> <p><b>Control:</b> brief information on Internet- and phone-based assistance available in Australia</p>
Outcomes	<p><b>Definition of abstinence:</b> self-reported 6-months, sustained abstinence at 7-month follow-up</p> <p>Intention-to-quit analysis and sensitivity analysis around treatment of missing data</p>

**Borland 2013** (Continued)

Funding source	National Health and Medical Research Council (Australia)
Conflicts of interest	JB is currently employed part-time through the University of Freiburg, Germany, on a project funded by Pfizer Global Health Partnership
Notes	OnQ and control arms used in comparison of text messaging with minimal SC. OnQ and QuitCoach arms used in comparison of text messaging with other SC support

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerised random number generator embedded within the baseline survey
Allocation concealment (selection bias)	Low risk	Participant allocation was embedded into the baseline survey, which appeared to be carried out online
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome data assessed through online surveys with no difference in contact between study arms
Incomplete outcome data (attrition bias) All outcomes	Low risk	Loss to follow-up 475 (13% total) with similar numbers across groups (control = 66, onQ = 89, QuitCoach = 104, both = 121, participant choice = 95); 2 excluded as reported to have died at 7-month follow-up

**Chan 2015**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> China</p> <p><b>Recruitment:</b> recruitment activities for the Quit to Win Contest took place at shopping malls or public areas in 16 out of the 18 districts in Hong Kong during May-July 2009. Participants who expressed an interest in joining the contest were screened for eligibility and tested on their exhaled CO to ascertain their smoking status</p> <p>Study year: 2009</p>
Participants	<p><b>Baseline characteristics (n = 1003)</b></p> <ul style="list-style-type: none"> <li>Female: 8.2%</li> <li>Education level primary or below: 21.6%</li> <li>White: not stated</li> <li>Age group: 49.1% aged 40-59 years</li> <li>Nicotine dependency (Heaviness of Smoking Index): 32.6% Heavy</li> </ul> <p><b>Inclusion criteria:</b> eligible participants were (1) Hong Kong residents aged <math>\geq 18</math> years; (2) daily smokers who smoked at least 1 cigarette/day in the past 6 months; (3) exhaled CO of <math>\geq 4</math> ppm; (4) able to communicate in Cantonese and read Chinese and (5) had a mobile phone to receive SMS.</p> <p><b>Exclusion criteria:</b> smokers who were physically or mentally unable to communicate or currently following other forms of SC programme were excluded from this RCT.</p>
Interventions	<p>All participants were given an 8-page self-help SC booklet.</p> <p><b>TEL Group:</b> 5-min telephone SC counselling by a trained nurse within 7 days of enrolment</p>

**Chan 2015** (Continued)

**SMS Group:** 8 mobile telephone text messages that were constructed with reference to the 8-page SC booklet

**Control Group:** self-help booklet and the contact information of the SC services at the enrolment

Outcomes	<b>Definition of abstinence:</b> biochemically validated 7-day point prevalence at 12-month follow-up
Funding source	Hong Kong Council on Smoking and Health (chairman and executive director are co-authors)
Conflicts of interest	TH Lam is the principal investigator of the FAMILY project, which was funded by the Hong Kong Jockey Club Charities Trust.
Notes	The control and SMS group were used in the comparison of text messaging with minimal SC support, and the SMS group and TEL group were used in the comparison of text messaging with other SC support

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "To achieve balanced number of subjects in each arm, the allocation sequence was sequentially generated by the author based on block randomization (with each recruitment session as a block) using the web site <a href="http://www.random.org">http://www.random.org</a> ."
Allocation concealment (selection bias)	Unclear risk	Quote: "The randomization and allocation were conducted by the author who did not participate in subject recruitment to ensure allocation concealment."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Follow-up calls were made to all the participants at 2, 6 and 12 months after the enrolment with standardized questionnaires by trained interviewers who were blinded to the group assignment."  Quote: "The RCT was single-blinded that all recruitment staff and assessors were not aware of the group allocation at the follow-up assessment."  Biochemical validation of abstinence was used.
Incomplete outcome data (attrition bias) All outcomes	Low risk	ITT analysis, at 6 months 66.9%, 73.1% and 70.6% of the 3 groups were available at follow-up

**Cobos-Campos 2017**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> Spain</p> <p><b>Recruitment:</b> participants recruited from 2 health centres, identified through their electronic health record and sent a letter of invitation</p> <p><b>Study dates:</b> 2013-2014</p>
Participants	<p><b>Baseline characteristics (n = 320)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 45.0</li> <li>• Female: 44.0%</li> <li>• High school or lower education: not stated</li> <li>• FTND: 17.5% high dependence</li> </ul>



**Cobos-Campos 2017** (Continued)

- White: not stated

**Inclusion criteria:** smokers aged  $\geq 18$  years, had a mobile phone, were able to receive and send text messages, and were motivated to start a SC programme (based on a score of  $\geq 5$  on the Richmond test)

**Exclusion criteria:** people who were on drug treatment for SC or had a history of mental or behavioral disorders or a diagnosis of depression (using the Goldberg scale; 23), as well as women who were pregnant

Interventions	<p><b>Health advice:</b> usual clinical practice (health advice provided by a doctor or nurse). Both groups followed the usual protocol (health advice) with its 4 visits (protocol according to recommendations of Spanish Society of Family and Community Medicine)</p> <p><b>Text messaging + health advice:</b> as above, plus reinforcement text messages to their mobile phones. 2 automatically generated text messages/day (1 in the morning and 1 in the evening) for the first 5 weeks and 3 messages/week from weeks 6 to 26. Messages were motivational in intent, to encourage participants in their efforts to stop smoking, and also provided information about the health-related risks of smoking. (SMSalud®)</p>
Outcomes	<p><b>Definition of abstinence:</b> biochemically confirmed prolonged abstinence (participant reporting not having smoked &gt; 5 cigarettes since the start of the follow-up period) at 6 months</p>
Funding source	<p>Departamento de Industria del Gobierno Vasco of the Basque Country under the 2012 Saiotek funding round (reference number SAI012-OA12BF001)</p> <p>Departamento de Educación, Política Lingüística y Cultura del Gobierno Vasco (IT620-13)</p>
Conflicts of interest	None declared

## Notes

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomly assigned to groups using computer-generated sequence
Allocation concealment (selection bias)	Low risk	Researchers involved were blind to the computer-generated sequence used for randomisation until the moment of group allocation
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Biochemical confirmation of abstinence
Incomplete outcome data (attrition bias) All outcomes	High risk	Dropout rate was > 50%: 48.75% intervention and 43.75% provided 6-month follow-up data, similar in both groups, all numbers reported, ITT analysis

**Danaher 2019**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> multifaceted nationwide online marketing campaign using Google AdWords and Reddit ads combined with listings on Smokefree.gov and ORI.org</p>
---------	---

**Danaher 2019** (Continued)

**Study dates:** 2015-2017

Participants	<b>Baseline characteristics (n = 1271)</b> <ul style="list-style-type: none"> <li>• Mean age: 44.9</li> <li>• Female: 78% (N = 991)</li> <li>• High school or lower education: 27.9% (N = 355)</li> <li>• FTND (FTND-6): mean: 5.5</li> <li>• White: 76.6% (N = 974)</li> </ul> <p><b>Inclusion criteria:</b> (1) <math>\geq 18</math> years of age; (2) smoked cigarettes as the primary tobacco product they used; (3) smoked <math>\geq 5</math> cigarettes/day for the previous 6 months; (4) wanted to quit smoking in next 14 days; (5) active use of a smartphone (iPhone or Android) and a personal computer or tablet; (6) willing to receive up to 150 text messages over 6 months of the programme; (7) access to the Internet; (8) have a valid personal email address; (9) US resident.</p> <p><b>Exclusion criteria:</b> none explicitly stated</p>	
Interventions	<p>Both interventions presented very similar best-practices SC recommendations and incorporated many of the same interactive and multimedia features (e.g. pictures, audios and videos). Both interventions shared a similar cognitive behavior therapy (CBT) theoretical foundation that was tied to specific programme features. Programme content was framed according to the multiple phases of quitting – Preparing to Quit, Quitting, Maintaining Abstinence, and Retooling – if a lapse/relapse was reported. In addition, the interventions used a series of online engagement activities in order to get the participant actively involved.</p> <p><b>QuitOnline:</b> a web-based intervention that presented best practice SC content using interactive and multimedia features. The content and structure of the programme was similar to the efficacious My-LastDip smokeless tobacco cessation programme.</p> <p><b>MobileQuit:</b> smartphone condition using an integrated web app and text messaging intervention designed for smartphones. The mobile web app used the smartphone’s Web browser and it had an appearance and functionality characteristic of a native app. There were 290 text messages delivered over the 180-day (6-month) study period. Adhoc text messages sent if the participant missed certain programme content, did not quit on QD, reported a lapse, reset the programme quit clock, replied to smoking status texts, and was scheduled to complete an online follow-up assessment</p>	
Outcomes	<b>Definition of abstinence:</b> self-reported abstinence, 7-day repeated point prevalence at 6 months	
Funding source	National Cancer Institute (US National Institutes of Health) - R01CA172205	
Conflicts of interest	None declared	
Notes		
<b>Risk of bias</b>		
	<b>Bias</b>	<b>Authors' judgement</b>
Random sequence generation (selection bias)	Low risk	Randomisation used a computer-generated randomisation sequence.
Allocation concealment (selection bias)	Low risk	No specific detail provided but recruitment, enrolment and randomisation all automated online so can assume allocation concealment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Online assessments. No difference in contact between arms

**Danaher 2019** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	54% completed follow-up, rates similar for both groups. Reasons for loss to follow-up not given, but ITT analysis performed
--	----------	---

**Ferguson 2015**

Methods	<b>Study design:</b> RCT  <b>Country:</b> Australia  <b>Recruitment:</b> advertisements in papers, radio, social media, Facebook, in Tasmania  <b>Study dates:</b> not stated	
Participants	<b>Baseline characteristics (n = 284)</b> <ul style="list-style-type: none"> <li>• Mean age: 42.1 (SD 13.2)</li> <li>• Female: 51.1% (N = 145)</li> <li>• Household income &lt; AUD 45,000: 66.7% (N = 189)</li> <li>• FTND: mean 4.8 (SD 2.0)</li> <li>• White: 93.7% (N = 266)</li> </ul> <b>Inclusion criteria:</b> daily smokers of > 10 cigarettes/day for past 3 years  <b>Exclusion criteria:</b> not explicitly stated	
Interventions	<b>Control:</b> self-help quit booklet containing tips for quitting and cognitive and behavioural coping mechanisms  <b>Intervention:</b> as above, plus 4 or 5 randomly timed text messages/day containing quit smoking advice and encouragement tailored to participants' current quit status (preparing to quit, first week of the quit attempt, second week of attempt etc.). Participants could request additional text messages	
Outcomes	<b>Definition of abstinence:</b> 7-day point prevalence abstinence verified by expired CO at 6 months	
Funding source	National Health and Medical Research Council	
Conflicts of interest	SF has consulted for GlaxoSmithKline Consumer Healthcare on matters relating to SC and has received researcher-initiated project grant funding from Pfizer (through the GRAND initiative)	
Notes		

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomisation schedules for sequential allocation. No further information provided
Allocation concealment (selection bias)	Unclear risk	Quote: "The group allocation procedure was designed to blind study staff with direct participant contact from knowing group assignment"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically verified

**Ferguson 2015** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Loss to follow-up not stated
--	--------------	------------------------------

**Free 2009**

Methods	<b>Study design:</b> RCT  <b>Country:</b> UK  <b>Setting:</b> advertisements on radio, bus billboards, websites, newspapers, primary care centres, pharmacies, SC services. Participants registered their interest by text message or online.  <b>Study dates:</b> 2007	
Participants	<b>Baseline characteristics (n = 200)</b> <ul style="list-style-type: none"> <li>• Mean age: 36 years (SD 9)</li> <li>• Female: 47% (N = 94)</li> <li>• Education: not stated</li> <li>• Cigarettes per day: mean 20</li> <li>• White: not stated</li> </ul> <b>Inclusion criteria:</b> aged $\geq 16$ years; smoking daily and interested in quitting; current owner of mobile phone  <b>Exclusion criteria:</b> not explicitly stated	
Interventions	<b>Intervention:</b> 6-month text messaging programme delivered solely over mobile phone based on programme in <a href="#">Rodgers 2005</a> but messages adapted for UK population. Participant nominated QD and received regular personalised text messages with advice, support and distraction, with a countdown to QD, intensive 4 weeks of 5 or 6 messages/day then maintenance phase of 1 message/2 weeks. Messages selected from database matched to participant characteristics. Free month of text messaging from QD. Optional Quit Buddy, and Text Crave (messages on demand). Interactive polls and quizzes <b>Control:</b> 1 generic text message every 2 weeks about study participation, with no SC content	
Outcomes	<b>Definition of abstinence:</b> salivary cotinine verified continuous abstinence (< 5 cigarettes) at 6 months	
Funding source	UK Medical Research Council, Primary Care Research Networks	
Conflicts of interest	None declared	
Notes	Pilot study carried out as a precursor to <a href="#">Free 2011</a>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Central computerised randomisation
Allocation concealment (selection bias)	Low risk	Participants sent consent via text message, RA entered data into web-based form, system automatically generated intervention or control group texts according to the computer-generated allocation
Blinding of outcome assessment (detection bias)	Low risk	Verified with salivary cotinine

**Free 2009** (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	Lost to follow-up: 8/98 (control) and 8/102 (intervention) at 6 months (92% follow-up)
--	----------	--

**Free 2011**

Methods

**Study design:** RCT

**Country:** UK

**Recruitment:** advertisements on radio, bus billboards, websites, newspapers, primary care centres, pharmacies, SC services. Participants registered their interest by text message or online

**Study dates:** 2007-09

Participants

**Baseline characteristics (n = 5800)**

- Mean age: 37
- Female: 45% (N = 2610)
- Education up to 16 years only: 44% (N = 2552)
- FTND > 5: 40% (N = 2320)
- White: 88.5% (N = 5133)

**Inclusion criteria:** aged ≥ 16 years, willing to make an attempt to quit smoking in the next month and owned a mobile phone.

**Exclusion criteria:** not explicitly stated

Interventions

All participants were free to participate in any other SC service or support that they wished to use, and were offered the QUIT and National Health Service (NHS) SC help line numbers.

**Intervention:** delivered solely over mobile phone based on programme in [Rodgers 2005](#). Participants asked to set a QD within 2 weeks of randomisation. They received 5 text messages/day for the first 5 weeks and then 3/week for the next 26 weeks. Intervention included motivational messages and behaviour-change techniques. The programme was also personalised with an algorithm based on demographic and other information gathered at baseline, such as smoker's concerns about weight gain after quitting. The core programme consisted of 186 messages and the personalised messages were selected from a database of 713 messages. For instance, by texting the word "lapse", participants received a series of 3 text messages that encouraged them to continue with their quit attempt. Participants could also request the mobile phone number of another trial participant so that they could text each other for support. Participants in the intervention group using pay-as-you-go mobile phone schemes were given a £20 top-up voucher to provide sufficient credit to participate in the intervention

**Control:** 2-weekly, simple, short, text messages related to the importance of trial participation (not SC-focused)

Outcomes

**Definition of abstinence:** biochemically validated prolonged abstinence at 6 months of follow-up (no more than 5 cigarettes smoked since the start of the abstinence period)

Funding source

Cancer Research UK

Conflicts of interest

None declared

Notes

**Risk of bias**

**Free 2011** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	An independent telephone randomisation system
Allocation concealment (selection bias)	Low risk	The system automatically generated intervention or control group texts according to the allocation.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically validated.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Primary outcome data were available for 94% of participants in the intervention group and 97% in the control group.

**Garrison 2018**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> online via advertisements</p> <p><b>Study dates:</b> 2014-2015</p>
Participants	<p><b>Baseline characteristics (n = 325)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 41 years</li> <li>• Female: 72% (233/325)</li> <li>• High school or lower education: 16% (52/325)</li> <li>• FTND: not stated</li> <li>• White: 81% (262/325)</li> <li>• Mean cigarettes per day: 16</li> </ul> <p><b>Inclusion criteria:</b> age 18–65 years, smoked <math>\geq 5</math> cigarettes/day, had <math>\leq 3</math> months past-year abstinence, owned an iPhone/Android, and were motivated to quit, indicated by <math>\geq 8/10</math> on the Contemplation Ladder and <math>\geq 4/5</math> on an Action item of the Readiness to Change Questionnaire: “I am trying to smoke less than I used to,” 1 = strongly disagree, 5 = strongly agree</p> <p><b>Exclusion criteria:</b> none specifically stated</p>
Interventions	<p><b>Mobile mindfulness training with experience sampling ((MMT-ES) Craving to quit app):</b> 22 days of training modules (5–15 min/day) teaching mindfulness for SC. The app teaches mindfulness and 3 standard meditation practices: body scan, loving kindness, and breath awareness. Body scan is practiced by bringing awareness to different parts of the body, to foster awareness of body sensations that constitute cravings and affective states. Loving kindness is practiced by directed well-wishing by repeating phrases such as “may X be happy,” to foster acceptance of oneself and others. Breath awareness is practiced by paying attention to the breath wherever one feels it most strongly in the body, to help retrain the mind away from habitual self-related thinking toward a more present-centred awareness. The app also teaches an informal practice to work mindfully with cravings, RAIN: Recognize, Accept, Investigate, and Note what cravings feel like. ES is another feature to measure smoking, craving, and other factors</p> <p><b>Experience sampling (ES) only (Control app):</b> a smartphone app with the same look and feel as MMT-ES, delivering only ES for 22 days, to control for potential effects of ES, expectancy effects and nonspecific effects of using a smartphone for SC.</p>



**Garrison 2018** (Continued)

Outcomes	<b>Definition of abstinence:</b> biochemically verified 1-week point prevalence abstinence at 6 months	
Funding source	American Heart Association (14CRP18200010) and the National Institute on Drug Abuse (K12DA00167)	
Conflicts of interest	JB and PP own stock in Claritas Mindsciences, the company that developed the apps used in this study.	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-generated (reported in protocol)
Allocation concealment (selection bias)	Unclear risk	Detail not stated
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "The primary outcome is one-week point-prevalence abstinence from tobacco smoking at 6-months, verified by carbon monoxide monitoring"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "Retention among full ITT was 72.6% (MMT-ES, 78.4%; ES, 74.2%; $\chi^2$ (1) = 1.2, p = .28) and among modified ITT was 83.7% (MMT-ES, 87.4%; ES, 80.8%; $\chi^2$ (1) = 2.6, p = .11)."  Retention 72.6%, no between-group differences in number of check-ins or days checked-in. ITT analysis presented

**Haug 2013**

Methods	<b>Study design:</b> cluster-RCT  <b>Country:</b> Switzerland  <b>Recruitment:</b> from students in vocational schools  <b>Study dates:</b> 2011-12
Participants	<b>Baseline characteristics (n = 755)</b> <ul style="list-style-type: none"> <li>• Mean age: 18.2 years (SD 2.3)</li> <li>• Female: 51.9% (N = 392)</li> <li>• Secondary school or less: 81.6% (N = 616)</li> <li>• Cigarettes per day mean: 10.6 (SD 7.6)</li> <li>• No immigration background: 53.2% (N = 399)</li> </ul> <b>Inclusion criteria:</b> daily or occasional cigarette smoking (at least 4 cigarettes in the preceding month and at least 1 cigarette during the preceding week), ownership of a mobile phone  <b>Exclusion criteria:</b> not explicitly stated
Interventions	<b>SMS-COACH:</b> a 3-month programme including a weekly SMS text message assessment of smoking-related target behaviour, 2 weekly text messages tailored to baseline data and responses to the SMS text message assessments, and an optional further integrated QD preparation and relapse prevention SMS programme. Participants who did not use the integrated programme for QD preparation and relapse prevention received a total of 37 text messages (1 welcome message, 11 assessment messages, 24 tai-

**Haug 2013** (Continued)

lored feedback messages, 1 goodbye message). Participants, who used the QD preparation and relapse-prevention programme for the whole period from 1 week before the scheduled QD until 3 weeks afterwards, received an additional 42 text messages

**Control:** all students in participating classes were invited to participate in an online health screening survey during a regular school lesson reserved for health education. The control group did not receive anything else

Outcomes	<b>Definition of abstinence:</b> self-reported 4-week point prevalence abstinence at 6-month follow-up
Funding source	Swiss Tobacco Prevention Fund
Conflicts of interest	SH and CM were involved in the development of the intervention
Notes	

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Block randomisation with computer-generated randomly permuted blocks of 4 cases
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Minimal contact with study investigators in both trial arms
Incomplete outcome data (attrition bias) All outcomes	Low risk	111/383 in control and 85/372 in intervention were lost to follow-up at 6 months. ITT analysis conducted
Other bias	Unclear risk	Although clustering is adjusted for in this study's analysis the authors do not report the clustering effect, making it impossible to adjust for this in our analysis. Therefore, it is not clear how much the clustering adjustment influences the result from this study.

**Herbec 2019**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> UK</p> <p><b>Recruitment:</b> online via Twitter and Facebook, supplemented by emails and posters within Bupa and the University College London. The app could also be found through online searches and UK app stores. Interested participants were directed to the project website.</p> <p><b>Study dates:</b> 2015-16</p>
Participants	<p><b>Baseline characteristics (n = 425)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 32.9 (11.19)</li> <li>• Female: 45.5% (N = 193)</li> <li>• High school or lower education: 31.3% (N = 133)</li> <li>• FTND: 21.4% smoke within 5 min of waking</li> </ul>

**Herbec 2019** (Continued)

- White: not stated

**Inclusion criteria:** (1) UK-based, (2)  $\geq 18$  years, (3) smoked daily, (4) wanted to make a serious quit attempt, (5) completed registration, (6) were willing to set a QD within 2 weeks of registration, (7) agreed to follow-up, (8) agreed to, if invited, confirm abstinence with a personal CO monitor posted to them for free, (9) consented and agreed to Bupa's End User License Agreement (EULA). Criteria (1)-(5) were assessed through a baseline questionnaire

**Exclusion criteria:** not explicitly stated

Interventions	<p><b>BupaQuit:</b> including SF28 ('SmokeFree28') app components, including advice, gamification elements + control app functionality</p> <p><b>Control:</b> smartphone app with 'minimum credible intervention' that provided users with brief advice tools for monitoring of the quit progress sharing of progress (number of smoke-free days) on social media or e-mail</p>	
Outcomes	<p><b>Definition of abstinence:</b> self-reported continuous 6-month abstinence</p>	
Funding source	<p>The costs of app development and of conducting the study (including participant recruitment, data collection, the cost of purchasing CO monitors) were covered by Bupa. AH was leading the trial as part of her PhD funded by British Heart Foundation 4-year PhD Studentship at UCL (FS/13/59/30649). JB EB salaries are funded by a programme grant from Cancer Research UK (CRUK; C1417/A22962). EB also receives funding from the NIHR SPHR. RW's salary was funded by CRUK for part of the preparation of this manuscript.</p>	
Conflicts of interest	<p>AH led the BupaQuit trial as part of her PhD funded by the British Heart Foundation and has been employed by Bupa in a casual role. AH has received unrestricted funds as Global Bridges at Mayo Clinic and Pfizer Independent Grants for Learning and Change RFP: European Program. LS has received honoraria for talks, an unrestricted research grant and travel expenses from Pfizer and Johnson &amp; Johnson, and has acted as a paid reviewer for grant awarding bodies and a consultant for health care companies. JB and EB have received an unrestricted grant from Pfizer. AM worked as Digital Manager at Bupa. RW undertakes research and consultancy and receives fees for speaking from companies that develop and manufacture SC medications. JB and RW are unpaid members of the scientific steering group of the Smoke Free mobile application</p>	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Participants were randomised automatically within the app (after registration) in 1:1 ratio to either the intervention or control app (random numbers generated using a standard JavaScript library)
Allocation concealment (selection bias)	Low risk	Participants were randomised automatically within the app (after registration) in 1:1 ratio to either the intervention or control app (random numbers generated using a standard JavaScript library)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Participant involvement was remote in both trial arms
Incomplete outcome data (attrition bias) All outcomes	High risk	High loss to follow-up with 40.2% of participants followed up at 6.5 months, no significant differences between groups

**Liao 2018**

**Methods** **Study design:** RCT

**Country:** China

**Recruitment:** via radio, bus billboards, online (e.g. websites, QQ, WeChat), newspapers, hospitals, and pharmacies in China

**Study dates:** 2016-17

**Participants** **Baseline characteristics (n = 1369)**

- Mean age: 38.1 years (SD 9.79)
- Female: 5.4% (N = 74)
- High school or lower education: 25.5% (N = 349)
- FTND: 4.6 (SD 2.16)
- White: not stated

**Inclusion criteria:** daily Chinese cigarette smokers.  $\geq 18$  years. Being able to read and write in Chinese. Owning a text-capable cell phone and knowing how to text. Willing to make an attempt to quit smoking in the next month. Willing to provide informed consent to participate in the study.

**Exclusion criteria:** non-smokers.  $< 18$  years. Unable to read and write in Chinese.

**Interventions** **High-frequency text messaging (HFM):** "Happy Quit" mobile phone-based HFM for 12 weeks (3-5 messages/day)

**Low-frequency text messaging (LFM):** "Happy Quit" mobile phone-based LFM for 12 weeks (3-5 messages/week)

**Control:** 1 text message every week, thanking them for being in the study

**Outcomes** **Definition of abstinence:** biochemically confirmed continuous abstinence at 6 months

**Funding source** China Medical Board (CMB) Open Competition Program (Grant Number 15-226)

**Conflicts of interest** None declared

**Notes** We compared HFM and LFM to control in the text messaging vs minimal SC support analysis, and compared HFM to LFM in the comparison of higher vs lower frequency text messaging

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Participants were randomly allocated using an independent telephone randomisation system that included a minimisation algorithm balancing for sex (male, female), age (18–34 years, > 34 years), educational level (years of education: $\leq 12$ years, $> 12$ years), and FTND score ( $\leq 5$ , $> 5$ )
Allocation concealment (selection bias)	Low risk	Quote: "Participants, investigators, and research personnel were masked to treatment allocation"  Participants were randomly allocated using an independent telephone randomisation system
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically verified

**Liao 2018** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Quote: "By the end of the 24-week trial period, the trial was completed by 83.2%, 74.6%, and 87.1% of participants in the HFM group, LFM group, and control group, respectively"
--	----------	--

**Naughton 2014**

Methods	<b>Study design:</b> RCT  <b>Country:</b> UK  <b>Recruitment:</b> participants were recruited from 32 participating primary care practices opportunistically, through self-referral or referred by a health professional.  <b>Study dates:</b> 2009-11	
Participants	<b>Baseline characteristics (n = 317)</b> <ul style="list-style-type: none"> <li>• Mean age: 41.8 years (SD 13.0)</li> <li>• Female: 52.7% (N = 167)</li> <li>• High school or lower education: not stated</li> <li>• First cigarette in 30 min: 67.9% (N = 215)</li> <li>• White: 98.0% (311)</li> </ul> <b>Inclusion criteria:</b> participants aged 18-75 years and current smokers ( $\geq 1$ cigarette/day and smoked within previous 7 days) who were willing to quit within 14 days of randomisation, recruited in primary care. Participants were self-referred or referred by a health professional, able to read English and provide written informed consent, with a mobile phone and familiar with sending and receiving text messages  <b>Exclusion criteria:</b> enrolled in another formal SC study or other cessation programme	
Interventions	<b>Control:</b> 'usual care' consisting of routine 'level 2' SC advice delivered by SC adviser. This included a brief discussion about smoking habits and history, measurement of expired-air CO, brief advice to quit, setting a QD within the next 14 days, options for pharmacotherapy, a prescription and arranging a follow-up visit. Usually the opportunity for multiple follow-up visits was offered  <b>Intervention:</b> usual care as above, plus a tailored advice report and a 90-day programme of tailored text messages generated by the iQuit system (number of messages sent each day varied from 0 to 2, mean/day over 90 days 1.2). The messages were designed to advise smokers on their quit attempt, provide information about the consequences of smoking and expectations for quitting, provide encouragement, boost self-efficacy, maintain motivation to quit and remind smokers how to cope with difficult situations	
Outcomes	<b>Definition of abstinence:</b> self-reported continuous abstinence at 6-month follow-up	
Funding source	National Institute for Health Research School for Primary Care Research. GP practice costs (NHS Service Support Costs) were provided by the Comprehensive Local Research Network. ATP was supported by the NIHR Biomedical Research Centre at Guy's and St Thomas's NHS Foundation Trust and King's College London.	
Conflicts of interest	None declared	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>

**Naughton 2014** (Continued)

Random sequence generation (selection bias)	Low risk	Randomised by online programme
Allocation concealment (selection bias)	Low risk	Randomised by online programme once baseline data collected
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcomes collected via postal questionnaire, with the same amount of investigator contact between groups
Incomplete outcome data (attrition bias) All outcomes	Low risk	65/303 in control and 70/299 in intervention lost to follow-up at 6 months. ITT analyses presented

**Peiris 2019**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> Australia</p> <p><b>Recruitment:</b> via an Aboriginal Community Controlled Health Service, a regional community event, and the NSW government telephone coaching service</p> <p><b>Study dates:</b> 2016-17</p>
Participants	<p><b>Baseline characteristics (n = 49)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 42 years</li> <li>• Female: 78% (N = 38)</li> <li>• High school or lower education: 44% (N = 22)</li> <li>• FTND: low 33% (N = 16)</li> <li>• White: 0%</li> </ul> <p><b>Inclusion criteria:</b> participants were eligible if they could provide informed consent and met all of the following criteria: (1) current smokers aged <math>\geq 16</math> years, (2) self-identification as an Aboriginal and/or Torres Strait Islander person, (3) willing to make an attempt to quit smoking in the next month, and (4) had access to an iPhone or Android smartphone. Only 1 person per household was invited to participate in the study</p> <p><b>Exclusion criteria:</b> not explicitly stated</p>
Interventions	<p><b>Intervention:</b> Android or iOS app comprising a personalised profile and quit plan, text and in-app motivational messages, and a challenge feature allowing users to 'compete' with others. Support worker could facilitate and give a tutorial if wanted.</p> <p><b>Control:</b> participants were encouraged to make use of all SC support services available to them.</p>
Outcomes	<p><b>Definition of abstinence:</b> biochemically confirmed continuous abstinence at 6-month follow-up</p>
Funding source	Carried out by the George Institute for Global Health, commissioned by NSW Health
Conflicts of interest	None declared
Notes	
<b>Risk of bias</b>	



**Peiris 2019** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation was conducted via a central computer-based randomisation service. Allocation was 1:1 intervention versus control using a minimisation algorithm to balance for sex, age (< 30 years vs ≥ 30 years), and heaviness of smoking index score (low (score ≤ 2) vs moderate or high addiction (score > 2)) for nicotine dependence.
Allocation concealment (selection bias)	Low risk	Central computer-based randomisation
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically validated
Incomplete outcome data (attrition bias) All outcomes	Low risk	3 lost to follow-up, all in intervention group (3/25). None lost in control group (0/24)

**Rodgers 2005**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> New Zealand</p> <p><b>Recruitment:</b> advertisements on websites, media articles, email and text message mailing lists, and posters at tertiary institutions</p> <p><b>Study date:</b> 2001</p>
Participants	<p><b>Baseline characteristics (n = 1705)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 22 years</li> <li>• Female: 58.5% (N = 997)</li> <li>• High school or lower education: not stated</li> <li>• FTND: 5</li> <li>• Maori (indigenous population): 20.8% (N = 355)</li> </ul> <p><b>Inclusion criteria:</b> aged ≥ 16 years, smoking daily, wanting to quit within the next month, and were able to send and receive text messages on their own mobile phone</p> <p><b>Exclusion criteria:</b> not explicitly stated</p>
Interventions	<p>All participants were informed at their baseline interview of the smoking Quitline and the government subsidy for nicotine replacement therapy that was available.</p> <p><b>Intervention:</b> 6-month programme delivered solely over mobile phone. Participant nominated QD and received regular personalised text messages with advice, support and distraction, with a countdown to QD, intensive 4 weeks of 5 or 6 messages/day then maintenance phase of 1 message/2 weeks. Messages selected from database matched to participant characteristics. Free month of text messaging from QD. Optional Quit Buddy and Text Crave (messages on demand). Interactive polls and quizzes</p> <p><b>Control:</b> 1 text message/2 weeks thanking participants for taking part (text messages had no SC content)</p>
Outcomes	<p><b>Definition of abstinence:</b> self-reported continuous abstinence at 26 weeks</p>

### Rodgers 2005 (Continued)

**Funding source** National Heart Foundation of New Zealand, the Cancer Society of New Zealand, Vodafone NZ, Alcatel and Auckland UniServices.

AR held a Senior Fellowship from the National Heart Foundation.

**Conflicts of interest** None declared

**Notes**

#### **Risk of bias**

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Central computerised randomisation
Allocation concealment (selection bias)	Low risk	Central computerised randomisation, concealed until after assignment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Minimal contact with investigators across groups
Incomplete outcome data (attrition bias) All outcomes	Low risk	Loss to follow-up at 6 months was 179/853 in the control arm versus 261/852 in the intervention arm
Other bias	Unclear risk	The authors suggested that some participants in the control group may have thought their incentive at follow-up (month of free text messaging) depended on reporting quitting. This could account for an unexpected increase in control group participants reporting quitting from 6 weeks (109 participants) to 6 months (202 participants reporting no smoking in the past 7 days), which could have led to an underestimation of the effect of the intervention.

### Squiers 2017

**Methods** **Study design:** RCT

**Country:** USA

**Recruitment:** online advertising and search ads via Facebook, Craigslist, Pandora, and Google, Yahoo! and Bing, plus email recruitment via market research panels

**Study dates:** 2013-14

**Participants** **Baseline characteristics (n = 4027)**

- Age 18-21: 20.8% (N = 839)
- Female: 70.2% (N = 2825)
- Less than high school: 5.6% (N = 225)
- Time to first cigarette 5 min: 27.6% (N = 1110)
- White: 73.7% (N = 2967)

**Inclusion criteria:** aged 18-29 years. Reside in the USA. Smoke cigarettes at least 5 days/month. Be interested in quitting cigarette use. Not be involved in a cessation programme. Have an active email account. Be able to receive text messages on their mobile phone. Be the only member of your household

**Squiers 2017** (Continued)

participating in this study. Be willing to share contact information with the study team in order to share information about the study on a timely basis.

**Exclusion criteria:** didn't complete baseline survey. The anti-fraud process included automated duplication checks of phone numbers, email addresses, and IP addresses. If duplicates were detected, the individual was excluded from the study. Failure to provide informed consent. Refused the honesty pledge. Faced technical difficulties (e.g. undelivered text messages). Texted "STOP" at any point before their QD. Opted out of the study entirely by notifying the project team, typically via email

Interventions	<p><b>Arm 1:</b> periodic cessation assessments and QD reminder messages (total 11 messages)</p> <p><b>Arm 2:</b> arm 1 messages above, plus motivational preparatory messages for 2 weeks prior to participants' QD (total 40 messages)</p> <p><b>Arm 3:</b> all of the messages from Arms 1 and 2 above, plus 6 weeks of follow-up post-QD messages (total 127 messages)</p>
Outcomes	<b>Definition of abstinence:</b> self-reported 7-day repeat point prevalence abstinence at 32 weeks
Funding source	National Institutes of Health, National Cancer Institute HHSN261201400002B, HHSN26100006, HHSN26100007
Conflicts of interest	None
Notes	We split Arm 1 and compared it with arms 2 and 3 in the comparison of higher- versus lower-frequency text messaging

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information about the sequence generation process
Allocation concealment (selection bias)	Low risk	Not stated but as all done online it is very unlikely to have any bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Minimal contact with investigators
Incomplete outcome data (attrition bias) All outcomes	Low risk	There was a 64.4% response rate overall. Arm 1: 846/1313; Arm 2: 933/1400; Arm 3: 824/1314. ITT analysis

**Tseng 2017**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> USA</p> <p><b>Recruitment:</b> from large urban HIV clinics</p> <p><b>Study dates:</b> 2013-14</p>
Participants	<p><b>Baseline characteristics (n = 158)</b></p> <ul style="list-style-type: none"> <li>Mean age: 46.79 years</li> </ul>

**Tseng 2017** (Continued)

- Female: 18.4% (N = 29)
- Less than high school: 22.2% (N = 35)
- Time to first cigarette 5 min: 53.8% (N = 85)
- White: 13.3% (N = 21)

**Inclusion criteria:** ≥ 18 years, current patient of the clinics, current or regular smoker (≥ 5 cigarettes/day), CO ≥ 8 ppm, willing to set a QD within the next 2 weeks, willing to use a mobile phone and able to read text messages, and eligible to take varenicline

**Exclusion criteria:** alcohol dependence and active drug abuse, and conditions that would prevent the use of varenicline

Interventions	<p><b>Control:</b> received standard care, which consisted of a self-help information sheet, tailored to HIV-positive and an offer of varenicline for 12 weeks according to the standard dosage schedule. Participants needed to return to the clinic each 4 weeks to receive further medication. All participants were provided with a pre-paid mobile phone - the control group received phones to facilitate their ability to call the quit line and receive text message appointment reminders only</p> <p><b>Intervention 1:</b> participants received standard care as above, and 2 text messages/day for 12 weeks. 1 message reminding them to take their medication and 1 motivational message regarding cessation</p> <p><b>Intervention 2 (not eligible for inclusion as tests the addition of more intensive behavioural support and not the text messaging intervention):</b> the standard care and text message interventions described above, plus behavioural therapy delivered via 7 proactive mobile phone-delivered counselling sessions over a 6-week period. These combined cognitive behavioural therapy and motivational interviewing techniques</p>	
Outcomes	<b>Definition of abstinence:</b> biochemically confirmed point prevalence abstinence at 6 months	
Funding source	National Institutes of Drug Abuse of the National Institutes of Health. The study medication was provided by Pfizer Inc. The research was supported by the Center for Drug Use and HIV Research. Dr Sherman is supported in part by a grant from NIDA.	
Conflicts of interest	None declared	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-generated randomisation schedule, stratified by people smoking 5-10 and people smoking > 10 cigarettes/day
Allocation concealment (selection bias)	Low risk	After consent and baseline data collected, the research assistant called to receive the assignment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinence was biochemically verified
Incomplete outcome data (attrition bias) All outcomes	Low risk	21/53 participants in the control group and 19/54 in the text message group did not complete 24-week follow-up visits. ITT analysis is used in this meta-analysis

**Whittaker 2011**

Methods	<p><b>Study design:</b> RCT</p> <p><b>Country:</b> New Zealand</p> <p><b>Recruitment:</b> targeted at young people through advertising via radio, Internet, mobile phone, paper-based and online magazines, Maori-specific media of all types, local and national newspapers and media releases to national media outlets, tertiary education institutions, primary healthcare services, SC services, large employer health promotion programmes, and posters at cafes/bars/sports/grounds</p> <p><b>Study dates:</b> 2007-09</p>	
Participants	<p><b>Baseline characteristics (n = 226)</b></p> <ul style="list-style-type: none"> <li>• Mean age: 27 years</li> <li>• Female: 47.3%</li> <li>• Less than high school: not stated</li> <li>• Time to first cigarette 5 min: 23% (N = 52)</li> <li>• White: 52.2% NZ European</li> </ul> <p><b>Inclusion criteria:</b> ≥ 16 years, current daily smokers ready to quit, and had a video message-capable phone.</p> <p><b>Exclusion criteria:</b> not explicitly stated</p>	
Interventions	<p><b>Intervention:</b> received an automated package of video and text messages over 6 months that was tailored to self-selected QD, role model and timing of messages. Video messages were video diary-style from a selected 'ordinary' person going through a quit attempt in advance of the participant. Frequency of messages varied from 1/day in the lead up to QD, 2/day from QD for 4 weeks, then reducing to 1 every 2 days for 2 weeks and then 1 every 4 days for about 20 weeks until 6 months after randomisation. Extra messages were available on demand to beat cravings and address lapses. Additional website for intervention group participants to review video messages they had been sent (and rate them if desired), change their selected time periods and change (or add to) their selected role model.</p> <p><b>Control:</b> set a QD and received a general health video message sent to their phone every 2 weeks</p>	
Outcomes	<p><b>Definition of abstinence:</b> self-reported continuous abstinence at 6-month follow-up</p>	
Funding source	<p>Health Research Council of New Zealand. It was supported by Vodafone NZ who provided free access to their mobile phone network but was otherwise uninvolved. The intervention was previously funded by the Digital Strategy Community Partnership Fund, Dept of Internal Affairs, NZ</p>	
Conflicts of interest	<p>None declared</p>	
Notes	<p></p>	
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Central computerised randomisation
Allocation concealment (selection bias)	Low risk	Baseline data collected online, with computer randomisation on submission of form, and programme automatically assigned - no study staff involved
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Contact with investigators was minimal in both groups

**Whittaker 2011** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	32% intervention and 22% control lost to follow-up at 6 months
--	----------	--

**Wilson 2016**

Methods	<b>Study design:</b> RCT  <b>Country:</b> USA  <b>Recruitment:</b> VAMC patients were sent a letter and were called to complete a telephone survey  <b>Study dates:</b> 2017
Participants	<b>Baseline characteristics (n = 310)</b> <ul style="list-style-type: none"> <li>• Mean age: 57 years</li> <li>• Female: 10.9% (N = 34)</li> <li>• High school or lower education: 34.2% (N = 106)</li> <li>• FTND: 4</li> <li>• White: 98.4% (N = 305)</li> </ul> <b>Inclusion criteria:</b> ≥ 18 years of age, enrolled at Durham VAMC for ongoing medical care, current smoker willing to make a quit attempt in the next 30 days, and English speaking  <b>Exclusion criteria:</b> no access to telephone, severely impaired hearing or speech, active diagnosis of a psychotic disorder, extended serious illness, and current hospitalisation
Interventions	<b>Control:</b> cognitive-behavioral telephone counselling and a tele-medicine clinic for access to NRT  <b>Intervention:</b> as above, plus a mobile contingency management app. Participants were required to record CO readings, upload readings via the app, and use the app to check receipt of compensation and abstinence incentives. Incentives were provided for submitting CO readings pre-QD and for abstinence post-QD
Outcomes	<b>Definition of abstinence:</b> self-reported prolonged abstinence at 6 months
Funding source	Department of Veterans Affairs Health Services Research and Development Service
Conflicts of interest	Quote: "The authors have no conflicts of interest to report"
Notes	Only published on clinicaltrials.gov (trials register) at this point

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Permuted block randomisation generated a priori using computerised methods
Allocation concealment (selection bias)	Unclear risk	Randomisation was concealed from study staff for each participant until completion of baseline measures; however details of how this was concealed were not given
Blinding of outcome assessment (detection bias)	Low risk	Outcome measures collected by phone surveys. There was minimal contact with researchers.



**Wilson 2016** (Continued)

All outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	18/156 in intervention group lost to follow-up, 15/154 in control group lost to follow-up.
--	----------	--

**Yu 2017**

Methods

**Study design:** RCT

**Country:** China

**Recruitment:** trained health workers in local maternal-child health centres asked all mothers attending their initial post-delivery visit (1 month after birth) to complete a short health questionnaire with questions related to tobacco use and household SHS exposure.

**Study date:** 2014

Participants

**Baseline characteristics (n = 342)**

- Mean age: 31.8 years (SD 4.5)
- Female: 0% (0/299)
- High school or lower education: 31.1% (93/299)
- Daily smoking: 77.6% (232/299)
- White: 0%

**Inclusion criteria:** families were eligible for inclusion if they met the following criteria: nonsmoking mothers and their newborns were currently exposed to SHS in the home; fathers currently smoked cigarettes in the home; the parents both owned a mobile phone that could receive text messages; and the family was able to provide informed consent.

**Exclusion criteria:** newborn older than 6 month when the intervention began; family refused to participate when the intervention began

Interventions

**Intervention IA:** in-person health counselling and materials on establishing a smoke-free home

**Intervention IB:** as above, plus a text message intervention targeted at both parents. The text message intervention included messages to the mother and her husband on the harms of SHS to the mother and the infant. The husband received additional cessation text messages to encourage him to quit smoking. A total of 9500 messages were sent to participants.

**Control:** standard postnatal care, which did not include any tobacco control and cessation counselling service

Outcomes

**Definition of abstinence:** self-reported SC at 12 months (6 month data also reported)

Funding source

National Cancer Institute and the Bill and Melinda Gates Foundation

Conflicts of interest

None declared

Notes

Framed as a SHS-reduction programme for families with cessation aimed at fathers

We compared IB and control for the text messaging vs minimal behavioural support analysis and compared IA and IB to test text messaging in addition to another form of SC support.

**Risk of bias**
**Bias**
**Authors' judgement**
**Support for judgement**

**Yu 2017** (Continued)

Random sequence generation (selection bias)	Low risk	Randomisation was fully computerised, using no blocks or strata, and each participant was allocated a number: 1 (then assigned to I-A), 2 (then assigned to I-B), or 3 (then assigned to control group) with equal probability
Allocation concealment (selection bias)	Unclear risk	Method of concealment is not described
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Abstinance was not biochemically validated, however contact was balanced between arms
Incomplete outcome data (attrition bias) All outcomes	Low risk	Group A 103/103 followed up at 6 months, Group B 99/100, Group C 96/96

**CO:** carbon monoxide; **FTCD:** Fagerström test for cigarette dependence; **FTND:** Fagerström Test of Nicotine Dependence; **ITT:** intention to treat; **IVR:** interactive voice response; **NIH:** National Institutes of Health; **NRT:** nicotine replacement therapy; **NSW:** New South Wales; **NZ:** New Zealand; **ppm:** parts per million; **QD:** quit day/date; **RCT:** randomised controlled trial; **SC:** smoking cessation; **SD:** standard deviation; **SHS:** second-hand smoke; **SMS:** short messaging service; **VAMC:** Veterans Affairs Medical Center

**Characteristics of excluded studies** [ordered by study ID]

Study	Reason for exclusion
<a href="#">ACTRN12617000491369</a>	Follow-up < 6 months
<a href="#">Aigner 2017</a>	Wrong intervention
<a href="#">Applegate 2007</a>	Follow-up < 6 months
<a href="#">Bamidis 2017</a>	Wrong study design
<a href="#">Bernstein 2018</a>	Follow-up < 6 months
<a href="#">Blasco 2012</a>	Wrong intervention
<a href="#">Brendryen 2008</a>	Wrong intervention
<a href="#">Bricker 2014</a>	Follow-up < 6 months
<a href="#">Brinker 2016</a>	Wrong intervention
<a href="#">Bronstein 2016</a>	Wrong intervention
<a href="#">Buller 2014</a>	Follow-up < 6 months
<a href="#">Chow 2012</a>	Wrong intervention
<a href="#">Dale 2014</a>	Wrong intervention
<a href="#">Fingrut 2014</a>	Wrong intervention
<a href="#">Fraser 2014</a>	Wrong intervention
<a href="#">Gritz 2013</a>	Wrong intervention

Study	Reason for exclusion
Halpern 2018	Wrong intervention
Hammett 2018	Follow-up < 6 months
Hassandra 2017	Wrong patient population
Haug 2008	Follow-up < 6 months
Haug 2009	Follow-up < 6 months
Haug 2014	Wrong intervention
Kiselev 2011	Wrong intervention
Lazev 2004	Wrong study design
Mason 2016	Wrong outcomes
Mehring 2014	Wrong intervention
Naughton 2012	Follow-up < 6 months
Naughton 2017	Follow-up < 6 months
NCT01454999	Follow-up < 6 months
NCT02245308	Wrong intervention
NCT02844296	Wrong intervention
NCT03177265	Wrong intervention
Obermayer 2004	Wrong study design
Pechmann 2015	Wrong study design
Peng 2013	Wrong intervention
Pollak 2013	Wrong intervention
Riley 2008	Wrong study design
Shi 2013	Follow-up < 6 months
Skov-Ettrup 2013	Wrong intervention
Skov-Ettrup 2014	Wrong study design
Skov-Ettrup 2016	Wrong intervention
Snider 2011	Wrong study design
Stanczyk 2014	Wrong intervention
Vidrine 2006	Follow-up < 6 months

Study	Reason for exclusion
<a href="#">Vilaplana 2014</a>	Follow-up < 6 months
<a href="#">Wizner 2009</a>	Wrong intervention
<a href="#">Ybarra 2012</a>	Follow-up < 6 months
<a href="#">Ybarra 2013</a>	Follow-up < 6 months
<a href="#">Yuhongxia 2011</a>	Wrong study design

### Characteristics of ongoing studies *[ordered by study ID]*

#### **Cambon 2017**

Trial name or title	Pragmatic randomised controlled trial evaluating effectiveness of a smoking cessation e- intervention "Tabac Info Service"
Methods	RCT
Participants	Smokers aged 18+ years in France
Interventions	E-intervention, Tabac Info Service (TIS), by website and mobile application Control: current practices of smoking cessation in France
Outcomes	Point prevalence abstinence at 6 months
Starting date	January 2017
Contact information	Dr Linda Cambon; Linda.cambon@ehesp.fr
Notes	Funding: this study is funded by the CNAMTS for the period 2016–2018

#### **Collins 2017**

Trial name or title	Babies living safe and smokefree (BLISS)
Methods	RCT
Participants	Female smokers aged 18+, living with a child < 6 years old, in USA
Interventions	Multimodal behavioural intervention (MBI) treatment: mobile app on cessation + Ask advise refer (AAR) + telephone cessation counselling + NRT gum/lozenge Control: AAR + telephone nutrition counselling + mobile phone nutrition app
Outcomes	Point prevalence abstinence at 12 months
Starting date	February 2016
Contact information	Bradley N. Collins, collinsb@temple.edu

**Collins 2017** (Continued)

Notes Funding: this project was supported by a grant from the National Institutes of Health (CA188813) to Lepore and Collins (multi-PIs).

**CTRI201801011643**

Trial name or title	Tobacco cessation at non communicable disease clinics
Methods	RCT
Participants	Smokers aged 30-80, with 1+ non communicable disease (diabetes, hypertension, CVD, stroke, cancer), in India
Interventions	Mobile messages and calls on tobacco cessation + counselling Control: counselling
Outcomes	Point prevalence abstinence at 6 months
Starting date	May 2018
Contact information	Garima Bhatt, garimabhattach.90@gmail.com; Dr Sonu Goel, sonugoel007@yahoo.co.in
Notes	Source of monetary or material support: Post Graduate Institute of Medical Education and Research Sector 12 Chandigarh

**CTRI201803012401**

Trial name or title	A clinical trial to study the effect of WhatsApp and pamphlet based quit smoking interference among software professionals in Bengaluru City
Methods	RCT
Participants	Smokers aged 20-40, software professionals, in India
Interventions	WhatsApp text on tobacco cessation counselling Control: self-help pamphlet
Outcomes	Smoking abstinence at 6 months
Starting date	November 2017
Contact information	Silpi Chatterjee, dr.silpi510@gmail.com; Archana Krishna Murty, archanakm20@gmail.com
Notes	

**Graham 2016**

Trial name or title	Optimizing text messaging to improve adherence to web-based cessation treatment
Methods	RCT

**Mobile phone text messaging and app-based interventions for smoking cessation (Review)**

### Graham 2016 *(Continued)*

Participants	Smokers aged 18+ in USA
Interventions	<p>Phase 1: mobile text messaging (personalisation, integration with web-based programme, tailoring, varying levels of message intensity). Phase 2: BecomeAnEX.org web-based smoking cessation programme + optimal-adherence text message programme from Phase 1.</p> <p>Phase 1: full factorial design of 4 factors each with 2 levels (16 arms); Phase 2: 2-arm randomised trial. Phase 1 control: standard text message programme. Phase 2 control: BecomeAnEX.org web-based smoking cessation program.</p> <p>Funding: National Institute on Drug Abuse of the National Institutes of Health (#1 R01 DA 038139-01A1; Graham, PI)</p>
Outcomes	Point prevalence abstinence at 9 months
Starting date	March 2018
Contact information	Amanda L Graham, agraham@truthinitiative.org
Notes	

### Graham 2017

Trial name or title	An integrated digital/clinical approach to smoking cessation in lung cancer screening
Methods	RCT
Participants	Smokers aged 18+ in USA
Interventions	<p>Mobile text messaging + access to BecomeAnEx website + consult with a trained tobacco treatment specialist</p> <p>Control: brief cessation counselling</p>
Outcomes	Smoking abstinence at 6 months
Starting date	August 2017
Contact information	Amanda L Graham, agraham@truthinitiative.org
Notes	Funding: National Cancer Institute of the National Institutes of Health under Award Number R01CA207048

### ISRCTN11154315

Trial name or title	Efficacy of a smoking cessation intervention using smartphones
Methods	RCT
Participants	Smokers aged 18+ and buddies non-smoker aged 18+ in Switzerland
Interventions	Smartphone app (SmokeFree Buddy) to support cessation. The participants choose a buddy (self-chosen from the personal social network)



**ISRCTN11154315** (Continued)

	Control: announces a self-set QD, and try to stop smoking on their own
Outcomes	Smoking abstinence at 6 months
Starting date	June 2017
Contact information	Philipp Schwaninger, philipp.schwaninger@uzh.ch
Notes	Funding: University of Zurich (Switzerland)

**ISRCTN11318024**

Trial name or title	Impact of a smartphone application on smoking cessation: a RCT
Methods	RCT
Participants	Smokers aged 18+ in Switzerland or in France
Interventions	Smartphone app (Stop-tabac) with (1) immediate feedback during episodes of craving and tobacco withdrawal symptoms; (2) an interactive 'coach' who provides individually-tailored counselling messages; (3) a discussion forum (The Tribe) where participants receive support from other users; (4) fact sheets; a calculator of cigarettes not smoked, money saved, and years of life gained; (5) a module on NRT  Control: a placebo smartphone app
Outcomes	Smoking abstinence at 6 months
Starting date	September 2018
Contact information	Jean-François Etter. ORCID ID: <a href="https://orcid.org/0000-0002-1426-3157">orcid.org/0000-0002-1426-3157</a> . ISG-Campus Biotech, 9 ch. des Mines, Geneva 1202 Switzerland
Notes	Funder: Swiss National Science Foundation

**ISRCTN15396225**

Trial name or title	Evaluation of the effectiveness of a text-based mHealth smoking cessation intervention among high school students in Sweden
Methods	RCT
Participants	Smokers, as high school students, in Sweden
Interventions	Mobile phone text messages of 13 weeks  Control: treatment as usual
Outcomes	Point prevalence abstinence at 6 months
Starting date	November 2017

**ISRCTN15396225** (Continued)

Contact information	Ulrika Müssener. ORCID ID: <a href="https://orcid.org/0000-0001-5173-5419">orcid.org/0000-0001-5173-5419</a> . Department of Medical and Health Sciences, Faculty of Medicine and Health, Linköping university, Linköping 58183 Sweden
Notes	Funder: Linköping University (Sweden)

**ISRCTN16022919**

Trial name or title	Mobile health interventions for smoking cessation services uptake and smoking cessation: a factorial randomised trial in Thailand
Methods	RCT
Participants	Smokers aged 18+ in Thailand
Interventions	Mobile phone text messages for 30 days
Outcomes	Smoking abstinence at 6 months  Control: mobile messages thanking participants for being part of project (contain no behaviour change)
Starting date	December 2015
Contact information	Pritaporn Kingkaew, <a href="mailto:umpk@leeds.ac.uk">umpk@leeds.ac.uk</a>
Notes	Funder: Health Promotion Economic Evaluation Collaborative Center (Thailand)

**ISRCTN17964518**

Trial name or title	Evaluation of the "Stop Tabac" Android phone application
Methods	RCT
Participants	Smokers aged 18+ in Switzerland
Interventions	Smartphone app (Stop Tabac) to support cessation  Control: placebo (smartphone application with minimal content on stopping smoking)
Outcomes	Smoking abstinence at 6 months
Starting date	June 2015
Contact information	Céline Mavrot, <a href="mailto:celine.mavrot@kpm.unibe.ch">celine.mavrot@kpm.unibe.ch</a>
Notes	Funder: Tobacco Control Fund of the Swiss Federal Office of Public Health

**ISRCTN33869008**

Trial name or title	Mobile phone-based smoking cessation intervention for patients with elective surgery
---------------------	--

**ISRCTN33869008** (Continued)

Methods	RCT
Participants	Smokers, undergoing elective surgery (not children or neonatal), in Sweden
Interventions	Mobile phone text messaging, including interactive component Control: usual care
Outcomes	Point prevalence abstinence at 6 months
Starting date	September 2017
Contact information	Marcus Bendtsen, ORCID ID: <a href="https://orcid.org/0000-0002-8678-1164">orcid.org/0000-0002-8678-1164</a> . Linköping University, Linköping 58183 Sweden
Notes	Funder: Kamprad Family Foundation (Sweden)

**NCT01982110**

Trial name or title	A mindfulness based application for smoking cessation
Methods	RCT
Participants	Smokers aged $\geq 18$ years in the USA
Interventions	A mindfulness-based smartphone app
Outcomes	Number of cigarettes smoked at 6 months
Starting date	September 2013
Contact information	Jennifer Penberthy <a href="mailto:jkp2n@virginia.edu">jkp2n@virginia.edu</a>
Notes	

**NCT01990079**

Trial name or title	Use of technological advances to prevent smoking relapse among smokers with PTSD
Methods	RCT
Participants	Smokers aged 18-70 years in USA
Interventions	Quit4ever combines counselling sessions, bupropion and NRT mobile contingency management and the smartphone application Stay Quit Coach
Outcomes	Point prevalence abstinence at 3 and 6 months
Starting date	December 2013
Contact information	Jean Beckham, Duke University

**NCT01990079** (Continued)

Notes

**NCT01995097**

Trial name or title	BABY STEPS II: SMS scheduled gradual reduction text messages to help pregnant smokers quit
Methods	RCT
Participants	Female smokers aged 18+, 10-28 week pregnant, in USA
Interventions	Scheduled gradual reduction text messages Control: support messages only
Outcomes	Point prevalence smoking abstinence at late third trimester (35 weeks)
Starting date	March 2014
Contact information	Kathryn Pollak, Duke University
Notes	Sponsor: Duke University

**NCT02037360**

Trial name or title	Mobile mindfulness training for smoking cessation
Methods	RCT
Participants	Smokers aged 18-65 years in USA
Interventions	Smartphone-based training programme Control: free smoking cessation smartphone app
Outcomes	Point prevalence abstinence at 6 months
Starting date	August 2015
Contact information	Judson Brewer judson.brewer@yale.edu
Notes	Sponsor: University of Massachusetts, Worcester

**NCT02218281**

Trial name or title	Developing a smartphone app with mindfulness training for teen smoking cessation
Methods	RCT
Participants	Smokers aged 13-19 years in USA
Interventions	Smoking cessation treatment delivered through a smartphone app via mindfulness training

**Mobile phone text messaging and app-based interventions for smoking cessation (Review)**

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley &amp; Sons, Ltd.

**NCT02218281** (Continued)

	Control: written smoking cessation materials only
Outcomes	Point prevalence abstinence at 3 and 6 months
Starting date	September 2014
Contact information	Lori Pbert, University of Massachusetts
Notes	Sponsor: University of Massachusetts, Worcester

**NCT02218944**

Trial name or title	Smoking response inhibition training
Methods	RCT
Participants	Smokers aged 18-45 years in USA
Interventions	Smoking-specific response inhibition training programme in the context of a quit attempt. The task is based on a modified stop-signal task  Control: placebo has 50% no-go trials, with no-go responses spread evenly across images; active comparator: response inhibition training: B, 20% of responses are no-go trials, but with no-go responses spread evenly across the various images.
Outcomes	Smoking relapse at 6 months
Starting date	September 2014
Contact information	Robert D Dvorak, North Dakota State University
Notes	

**NCT02237898**

Trial name or title	Harnessing the power of technology: MoMba for postpartum smoking
Methods	RCT
Participants	Smokers aged 18-50 years in USA
Interventions	MoMba Live Long smartphone application  Control: traditional contingency management with financial incentives
Outcomes	Point prevalence abstinence at 21 months
Starting date	February 2016
Contact information	Ruth M Arnold, ruth.arnold@yale.edu
Notes	Sponsor: Yale University

**NCT02420015**

Trial name or title	Mobile health technology to enhance abstinence in smokers with schizophrenia
Methods	RCT
Participants	Smokers aged 18-70 with schizophrenia in USA
Interventions	Multi-Component Mobile-enhanced Treatment for Smoking Cessation (iCOMMIT) Control: pharmacotherapy + cessation counselling
Outcomes	Prolonged abstinence at 6 month
Starting date	March 2017
Contact information	Jean C Beckham, Duke University
Notes	Sponsor: Duke University

**NCT02665208**

Trial name or title	A pilot text messaging intervention to reduce smoking in office-based buprenorphine and inpatient detoxification patients
Methods	RCT
Participants	Smokers aged 18+ with opiate dependence and/or alcohol dependence in USA
Interventions	Smokefreetxt by the National Cancer Institute + prescriptions for NRT Control: informational pamphlets + prescriptions for NRT
Outcomes	Smoking abstinence at week 1 (time frame: 24 Weeks)
Starting date	March 2015
Contact information	Babak Tofighi, New York University Medical School
Notes	Sponsor: New York University School of Medicine

**NCT02724462**

Trial name or title	Trial of an innovative smartphone intervention for smoking cessation
Methods	RCT
Participants	Smokers aged 18+ in USA
Interventions	Smartphone-delivered Intervention (SmartQuit) Control: standard of care smartphone smoking cessation app

**NCT02724462** (Continued)

Outcomes	Point prevalence abstinence at 12 months
Starting date	May 2017
Contact information	Jonathan Bricker, Fred Hutchinson Cancer Research Center
Notes	Sponsor: Fred Hutchinson Cancer Research Center

**NCT02840513**

Trial name or title	Smartphone app and CO self-monitoring for smoking cessation (SMART-CO)
Methods	RCT
Participants	HIV-infected smokers aged 16+ in Switzerland
Interventions	Smartphone coaching app/CO self-monitoring Control: usual care as regularly provided by their physicians
Outcomes	Smoking abstinence at 6 months
Starting date	June 2017
Contact information	Dmitry Gryaznov, dmitry.gryaznov@usb.ch
Notes	Sponsor: Alain Nordmann, Basel Institute for Clinical Epidemiology and Biostatistics

**NCT02901171**

Trial name or title	The contribution of a smartphone application to acceptance and commitment therapy group treatment for smoking cessation
Methods	RCT
Participants	Smokers aged 18+ in Ireland
Interventions	Acceptance and Commitment Therapy (ACT) group treatment combined with smartphone application Control: group-based behavioural support programme
Outcomes	Point prevalence abstinence at 6 months
Starting date	September 2016
Contact information	Louise McHugh, University College Dublin
Notes	Sponsor: University College Dublin



**NCT03021655**

Trial name or title	A pilot randomised control trial to help youth smokers to quit smoking
Methods	RCT
Participants	Smokers aged 12-25 in Hong Kong
Interventions	Smartphone WhatsApp support group Control: telephone counselling on quitting
Outcomes	Point prevalence abstinence at 6 months
Starting date	December 2016
Contact information	Ho-cheung Li, william3@hku.hk
Notes	Sponsor: University of Hong Kong

**NCT03038542**

Trial name or title	Quit4hlth: enhancing tobacco and cancer control through framed text messages
Methods	RCT
Participants	Smokers aged 18+ in USA
Interventions	Gain-framed text messages + standard quit line treatment Control: standard care text messages + standard quit line treatment
Outcomes	Point prevalence abstinence at 30 weeks
Starting date	May 2016
Contact information	Benjamin Toll, Medical University of South Carolina
Notes	

**NCT03191019**

Trial name or title	A mobile-phone based intervention to support smoking cessation among Chilean women
Methods	RCT
Participants	Female smokers aged 18-24 in Chile
Interventions	Mobile-phone app for smoking cessation Control: mobile-phone app that will send 1 message every 2 weeks thanking participants for taking part in the study
Outcomes	Point prevalence abstinence at 6 months

**NCT03191019** (Continued)

Starting date	November 2017
Contact information	Carolina Lopez, cxlopez@uc.cl
Notes	

**NCT03445507**

Trial name or title	Effectiveness of a chat bot for smoking cessation: a pragmatic trial in primary care. (Dej@lo)
Methods	RCT
Participants	Smokers aged 18+ in Spain
Interventions	Mobile-phone app of an evidence-based chat bot Control: usual care given by their usual general practitioners and nurses of primary care health centres
Outcomes	Continuous abstinence at 6 months
Starting date	October 2018
Contact information	Eduardo Olano-Espinosa, Eduardo.Olano@salud.madrid.org
Notes	Sponsor: Gerencia de Atención Primaria, Madrid

**NCT03495622**

Trial name or title	Effectiveness of a combined CHW and text messaging-based tobacco intervention in India (MUKTI)
Methods	RCT
Participants	Smokers aged 18-70 in India
Interventions	Text messaging + motivational interviewing Control: brief verbal advice
Outcomes	Smoking abstinence at 12 months
Starting date	January 2018
Contact information	Vittal Hejjaji, vitty.hejy@gmail.com
Notes	Sponsor: University Hospitals Cleveland Medical Center

**NCT03538938**

Trial name or title	Improving Quitline support study (IQS)
---------------------	--

**NCT03538938** (Continued)

Methods	RCT
Participants	Smokers aged 18+, covered by MediAid with no more than a high school education in USA
Interventions	SmokefreeTXT + patch (and lozenge) + counselling + financial incentive
Outcomes	Point prevalence abstinence at 6 months
Starting date	June 2018
Contact information	Danielle E. McCarthy, University of Wisconsin
Notes	Sponsor: University of Wisconsin, Madison

**NCT03552978**

Trial name or title	Tech and telephone smoking cessation treatment for young veterans with PTSD
Methods	RCT
Participants	Veteran smokers aged 18-39 with PTSD in USA
Interventions	Stay Quit Coach (SQC) smartphone app and the iCO Smokerlyzer device and app Control: referral to the VA Quitline
Outcomes	Point prevalence abstinence at 24 weeks
Starting date	February 2019
Contact information	Ellen Herbst, Ellen.Herbst@va.gov
Notes	Sponsor: University of California, San Francisco

**NCT03553173**

Trial name or title	So-Lo-Mo intervention applied to the smoking cessation process (So-Lo-Mo)
Methods	RCT
Participants	Smokers aged 18+ in Spain
Interventions	Smartphone So-Lo_Mo app + usual psycho-pharmacological treatment Control: usual psycho-pharmacological treatment
Outcomes	Smoking abstinence rate at 1 year
Starting date	October 2016
Contact information	Francisco Ortega-Ruiz, Virgen del Rocío University Hospital

**NCT03553173** (Continued)

Notes

Sponsor: Fundación Pública Andaluza para la gestión de la Investigación en Sevilla

**Valdivieso-Lopez 2013**

Trial name or title	Efficacy of a mobile application in the smoking cessation among young people (TOBB_STOP)
Methods	Cluster-RCT
Participants	Smokers aged 18-30 smoking 10+ cigarettes a day, in Spain
Interventions	Mobile-phone app assisting 6-month implementation of recommendations of a Clinical Practice Guideline on smoking cessation  Control: usual cal
Outcomes	Point prevalence abstinence at 6 months
Starting date	January 2013
Contact information	Empar Valdivieso-López: tac.tacneg@sci.etrat.osevidlave
Notes	Funding: Jordi Gol i Gurina Foundation. Note: subgroup result paper <a href="http://www.ncbi.nlm.nih.gov/pubmed/30916655">www.ncbi.nlm.nih.gov/pubmed/30916655</a>

**Weng 2018**

Trial name or title	Building capacity and promoting smoking cessation in the community via "Quit to Win" contest 2016
Methods	RCT
Participants	Smokers aged 18+ in Hong Kong
Interventions	Mobile phone text messaging for 1 month for cessation referral (2 experimental arms: high intensity/low intensity)  Control: general very brief advice
Outcomes	Smoking abstinence at 6 months
Starting date	June 2016
Contact information	Man Ping Kelvin Wang, mpwang@hku.hk
Notes	Funding: this study is funded by the Hong Kong Council on Smoking and Health.

**CVD:** cardiovascular disease; **NRT:** nicotine replacement therapy; **QD:** quit day/date; **RCT:** randomised controlled trial; **VA:** Veterans Affairs


**DATA AND ANALYSES**
**Mobile phone text messaging and app-based interventions for smoking cessation (Review)**

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley &amp; Sons, Ltd.


**Comparison 1. Text messaging versus minimal smoking cessation support**


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Long-term abstinence (all randomised))	13	14133	Risk Ratio (M-H, Random, 95% CI)	1.54 [1.19, 2.00]
2 Long-term abstinence (complete case)	13	11969	Risk Ratio (M-H, Random, 95% CI)	1.56 [1.21, 2.02]

**Analysis 1.1. Comparison 1 Text messaging versus minimal smoking cessation support, Outcome 1 Long-term abstinence (all randomised)).**



**Analysis 1.2. Comparison 1 Text messaging versus minimal smoking cessation support, Outcome 2 Long-term abstinence (complete case).**







**Comparison 2. Text messaging versus other smoking cessation intervention**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Long-term abstinence (all randomised)	2	2238	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.61, 1.40]
2 Long-term abstinence (complete case)	2	1813	Risk Ratio (M-H, Random, 95% CI)	0.93 [0.63, 1.36]

**Analysis 2.1. Comparison 2 Text messaging versus other smoking cessation intervention, Outcome 1 Long-term abstinence (all randomised).**




**Analysis 2.2. Comparison 2 Text messaging versus other smoking cessation intervention, Outcome 2 Long-term abstinence (complete case).**



**Comparison 3. Text messaging + other smoking cessation support versus other smoking cessation support alone**


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Long-term abstinence (all randomised)	4	997	Risk Ratio (M-H, Random, 95% CI)	1.59 [1.09, 2.33]
2 Long-term abstinence (complete case)	4	796	Risk Ratio (M-H, Random, 95% CI)	1.63 [1.12, 2.37]

**Analysis 3.1. Comparison 3 Text messaging + other smoking cessation support versus other smoking cessation support alone, Outcome 1 Long-term abstinence (all randomised).**






**Analysis 3.2. Comparison 3 Text messaging + other smoking cessation support versus other smoking cessation support alone, Outcome 2 Long-term abstinence (complete case).**




**Comparison 4. High-frequency versus low-frequency text messaging**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Long-term abstinence (all randomised)	3	12985	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.95, 1.06]
2 Long-term abstinence (complete case)	3	6798	Risk Ratio (M-H, Random, 95% CI)	1.04 [1.00, 1.09]

**Analysis 4.1. Comparison 4 High-frequency versus low-frequency text messaging, Outcome 1 Long-term abstinence (all randomised).**




**Analysis 4.2. Comparison 4 High-frequency versus low-frequency text messaging, Outcome 2 Long-term abstinence (complete case).**




**Comparison 5. Smartphone app versus lower-intensity smoking cessation support**


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
<b>1 Long-term abstinence (all randomised)</b>	5	3079	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.66, 1.52]
1.1 Comparison: minimal non-app SC support	2	1645	Risk Ratio (M-H, Random, 95% CI)	0.82 [0.56, 1.18]
1.2 Comparator: less intensive smartphone app	3	1434	Risk Ratio (M-H, Random, 95% CI)	1.12 [0.60, 2.09]
<b>2 Long-term abstinence (complete case)</b>	5	1774	Risk Ratio (M-H, Random, 95% CI)	1.06 [0.72, 1.54]
2.1 Comparison: minimal non-app SC support	2	771	Risk Ratio (M-H, Random, 95% CI)	0.87 [0.62, 1.23]
2.2 Comparator: less intensive smartphone app	3	1003	Risk Ratio (M-H, Random, 95% CI)	1.18 [0.67, 2.09]

**Analysis 5.1. Comparison 5 Smartphone app versus lower-intensity smoking cessation support, Outcome 1 Long-term abstinence (all randomised).**






**Analysis 5.2. Comparison 5 Smartphone app versus lower-intensity smoking cessation support, Outcome 2 Long-term abstinence (complete case).**




**Comparison 6. CO monitoring + contingency management versus smoking cessation support**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Long-term abstinence (all randomised)	2		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
1.1 Comparison: counselling, nicotine replacement therapy and monitoring only app	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
1.2 Comparison: counselling and nicotine replacement therapy	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
2 Long-term abstinence (complete case)	2		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
2.1 Comparison: counselling, nicotine replacement therapy and monitoring only app	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
2.2 Comparison: counselling and nicotine replacement therapy	1		Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]

**Analysis 6.1. Comparison 6 CO monitoring + contingency management versus smoking cessation support, Outcome 1 Long-term abstinence (all randomised).**



**Analysis 6.2. Comparison 6 CO monitoring + contingency management versus smoking cessation support, Outcome 2 Long-term abstinence (complete case).**



**Comparison 7. Smartphone app + text messaging versus web-based intervention**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Long-term abstinence (all randomised)	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected
2 Long-term abstinence (complete case)	1		Risk Ratio (M-H, Random, 95% CI)	Totals not selected

**Analysis 7.1. Comparison 7 Smartphone app + text messaging versus web-based intervention, Outcome 1 Long-term abstinence (all randomised).**

Study or subgroup	Smartphone app + TM n/N	Web-based intervention n/N	Risk Ratio M-H, Random, 95% CI	Risk Ratio M-H, Random, 95% CI
Danaher 2019	100/633	56/638	+	1.8[1.32,2.45]

Favours web-based int      Favours app + TM

**Analysis 7.2. Comparison 7 Smartphone app + text messaging versus web-based intervention, Outcome 2 Long-term abstinence (complete case).**

Study or subgroup	Smartphone app + TM n/N	Web-based intervention n/N	Risk Ratio M-H, Random, 95% CI	Risk Ratio M-H, Random, 95% CI
Danaher 2019	100/247	56/216	+	1.56[1.19,2.05]

Favours web-based int      Favours app + TM

**APPENDICES**

**Appendix 1. Tobacco Addiction Group Specialised Register search strategy**

Searched in Cochrane Register of Studies

- #1 Cellular Phone:MH
- #2 Cell Phones:MH
- #3 MeSH DESCRIPTOR Cellular Phone
- #4 MESH DESCRIPTOR Cell Phones
- #5 MeSH DESCRIPTOR Text Messaging
- #6 (mobile NEAR2 (phone\* OR telephon\*)):TI,AB,MH,EMT,XKY,KY,KW
- #7 (cell\* NEAR2 (phone\* OR telephon\*)):TI,AB,MH,EMT,XKY,KY,KW
- #8 smartphone\*:TI,AB,MH,EMT,XKY,KY,KW
- #9 text messag\*:TI,AB,MH,EMT,XKY,KY,KW
- #10 (txt OR pxt OR mms OR sms):TI,AB,MH,EMT,XKY,KY,KW
- #11 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10

**WHAT'S NEW**

Date	Event	Description
23 September 2019	New citation required but conclusions have not changed	Search updated and 13 new studies added
19 June 2019	New search has been performed	Search updated 2018, new studies added and text updated

## HISTORY

Protocol first published: Issue 3, 2007

Review first published: Issue 4, 2009

Date	Event	Description
1 October 2015	New search has been performed	Updated 2015, seven new studies added and text updated
1 October 2012	New citation required and conclusions have changed	Three new included studies added, meta-analysis conducted, conclusions changed (pooled effect statistically significant)
1 October 2012	New search has been performed	Updated 2012, three new studies added and text updated
15 July 2008	Amended	Converted to new review format.
5 September 2006	New citation required and conclusions have changed	Substantive amendment

## CONTRIBUTIONS OF AUTHORS

RW is the lead author of this review.

For the most recent update:

- HM and YG selected studies for inclusion, with assistance from CB;
- RW, RD, HM, CB and YG independently extracted data from the papers;
- all authors contributed to the writing and editing of the review.

## DECLARATIONS OF INTEREST

RW was co-author of one paper on one of the included studies ([Rodgers 2005](#)). She was a co-investigator on included studies ([Baskerville 2018](#); [Free 2009](#); [Free 2011](#)), and principle investigator of a further included study ([Whittaker 2011](#)). RW's institution (Auckland UniServices Ltd) received grant money to cover the costs of providing the text messaging intervention for the study described in [Free 2011](#). RW's institution licensed the STOMP text messaging cessation intervention in 2008, however no royalties were received. The licence has since been rescinded. This is not deemed to be a conflict of interest.

HM was co-author of [Whittaker 2011](#) and received honoraria from Pfizer for speaking at educational events and attending advisory group meetings.

CB was co-author of [Whittaker 2011](#) and his institution received grant money to cover the costs of providing the text messaging intervention for the study described in [Free 2011](#).

AR was a lead author ([Rodgers 2005](#)), and a co-author ([Free 2009](#); [Free 2011](#); [Whittaker 2011](#)), on included studies.

YG none known.

RD's institution received grant money to cover the costs of providing the text messaging intervention for the study described in [Free 2011](#).

---

## SOURCES OF SUPPORT

### Internal sources

- National Institute for Health Innovation (Auckland Uniservices), New Zealand.

### External sources

- No sources of support supplied

## DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We have followed the change of policy of the Cochrane Tobacco Addiction Group, and now report our findings using Mantel-Haenszel random-effect risk ratios rather than as odds ratios.

## INDEX TERMS

### Medical Subject Headings (MeSH)

\*Cell Phone; \*Smoking Cessation; \*Text Messaging; Counseling [\*methods]; Randomized Controlled Trials as Topic

### MeSH check words

Adult; Humans