

2. Senkus E, Kyriakides S, Ohno S, Penault-Llorca F, Poortmans P, Rutgers E, et al. Primary breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2015;26 Suppl. 5, v8-30.
3. Chase A, Cross NC. Aberrations of EZH2 in cancer. *Clin Cancer Res*. 2011;17:2613–8.
4. Yoo KH, Hennighausen L. EZH2 methyltransferase and H3K27 methylation in breast cancer. *Int J Biol Sci*. 2012;8:59–65.
5. Hayashi A, Yamauchi N, Shibahara J, Kimura H, Morikawa T, Ishikawa S, et al. Concurrent activation of acetylation and tri-methylation of H3K27 in a subset of hepatocellular carcinoma with aggressive behavior. *PLoS ONE*. 2014;9:e91330.

<http://dx.doi.org/10.1016/j.pbj.2017.07.116>

PS212

### Is P-glycoprotein relevant for the release of microvesicles by tumor cells?

I. Castro<sup>1,2,3,\*</sup>, C.P.R. Xavier<sup>1,2</sup>, M.H. Vasconcelos<sup>1,2,4</sup>

<sup>1</sup> i3S – Instituto de Investigação e Inovação em Saúde, Universidade do Porto, 4200-135 Porto, Portugal

<sup>2</sup> Cancer Drug Resistance Group, IPATIMUP – Institute of Molecular Pathology and Immunology of the University of Porto, 4200-465 Porto, Portugal

<sup>3</sup> FMUP – Faculty of Medicine of the University of Porto, 4200-319 Porto, Portugal

<sup>4</sup> FFUP – Faculty of Pharmacy of the University of Porto, 4050-313 Porto, Portugal

E-mail address: [msilva@ipatimup.pt](mailto:msilva@ipatimup.pt)

(I. Castro).

**Aim:** In this study, we aimed to verify if MDR cells without expression of P-gp also produced more microvesicles and less exosomes than their DS counterpart cells.

**Introduction:** Cancer multidrug resistance (MDR) is a major cause of chemotherapy failure and is highly associated with overexpression of drug-efflux pumps such as P-glycoprotein (P-gp). The identification of mechanisms specific of P-gp overexpressing cells may contribute to the identification of biomarkers of MDR.

It was recently discovered that a drug-resistant phenotype may be horizontally transferred from drug-resistant (DR) to drug-sensitive (DS) cells, mediated by the cargo of extracellular vesicles (EVs) released by DR cells and captured by DS cells. These EVs include smaller exosomes and larger microvesicles. Our previous work showed that MDR cells with overexpression of P-gp released more microvesicles than exosomes, unlike their DS counterpart cells. However, it is not known if this phenomenon is restricted to MDR cells with overexpression of P-gp or if it is extensive to all DR cells (with other mechanism of drug resistance).

**Methods:** Drug-response curves of MDR and DS counterpart cells were obtained, using resazurin and trypan blue assays, to confirm the resistant or sensitive phenotype of the cell lines. Confirmation of their P-gp status was possible by Western-Blot. EVs released by both DS and MDR cells were isolated by ultracentrifugation and characterized by transmission electron microscopy, dynamic light scattering, nanoparticle tracking analysis and Western blot analysis.

**Results:** We confirmed that MDR cells without expression of P-gp release EVs with similar sizes to the ones released by their DS counterparts.

**Conclusion:** So, P-gp may be associated with the release of larger EVs by MDR cells. These results will be further confirmed by characterizing the EVs released by P-gp overexpressing MDR cell lines following downregulation of P-gp expression and the EVs released by DS cell lines following transfection of P-gp.<sup>1-4</sup>



### References

1. Chen Z, et al. Mammalian drug efflux transporters of the ATP binding cassette (ABC) family in multidrug resistance: a review of the past decade. *Cancer Lett*. 2016;370:153–64.
2. Sousa D, Lima RT, Vasconcelos MH. Intercellular transfer of cancer drug resistance traits by extracellular vesicles. *Trends Mol Med*. 2015;21:595–608.
3. Lopes-Rodrigues V, et al. Data supporting the shedding of larger extracellular vesicles by multidrug resistant tumour cells. *Data Brief*. 2016;6:1023–7.
4. Lopes-Rodrigues V, et al. Multidrug resistant tumour cells shed more microvesicle-like EVs and less exosomes than their drug-sensitive counterpart cells. *Biochim Biophys Acta*. 2016;1860:618–27.

<http://dx.doi.org/10.1016/j.pbj.2017.07.117>

PS215

### Uterine protein oxidative modifications may condition trophoblast function



S. Mendes<sup>1,2,\*</sup>, A.I. Soares<sup>1,2</sup>, S. Silveira<sup>1</sup>, L. Matos<sup>1,2</sup>, L. Guedes-Martins<sup>2,3</sup>, J. Saraiva<sup>2,3,4</sup>, H. Almeida<sup>1,2,5</sup>, E. Silva<sup>1,2</sup>

<sup>1</sup> I3s; IBMC and FMUP, Portugal

<sup>2</sup> FMDUP, Portugal

<sup>3</sup> Centro Hospitalar do Porto EPE, Portugal

<sup>4</sup> Private Hospital Trofa, Portugal

<sup>5</sup> Obstetrics-Gynecology, Hospital-CUF Porto, Portugal

E-mail address: [saramendes313@gmail.com](mailto:saramendes313@gmail.com)

(S. Mendes).

**Aim:** Evaluate whether protein carbonylation resulting from uterine altered redox imbalance interferes with extravillous trophoblast viability.

**Introduction:** Local redox homeostasis is believed to have a pivot role in uterine transformation necessary for blastocyst implantation and placenta development. By contrast, redox status imbalance plays a role in deficient placentation and the development of pregnancy-related complications (e.g. preeclampsia or gestational diabetes) with increased incidence in older women. Thus, it was hypothesized that at an older reproductive age, loss of redox homeostasis is a contributor to disruption of foetal/placental interactions and the development of such complications.

**Methods:** Uterine human samples were collected at delivery by elective caesarean section. The protocol was approved by the ethical committee of “Centro Materno-Infantil do Porto”, volunteers gave written consent to be included in the study. Total protein carbonylation was detected by oxyblot and protein expression was quantified by western blotting. Specific protein carbonylation was verified by immunoprecipitation. Albumin was carbonylated using hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), followed by dialysis, and western blotting to confirm carbonylated albumin. HST-8SV neo extravillous trophoblasts were treated with carbonylated/non-carbonylated albumin, followed by cell viability assay. A *P* value less than 0.05 was assumed to denote significant difference.

**Results:** At the placental site, carbonylated albumin normalized to total albumin expression showed a positive and significant association with maternal age. ( $r=0.6909$ ,  $P=0.0021$ ) In vitro, carbonylated albumin displayed a cytotoxic effect, at concentrations ranging from 10 to 100 µg/ml. Lower concentrations did not affect trophoblast viability.

**Conclusion:** Uterine aging is accompanied by selective albumin oxidative modifications, which appears to interfere with trophoblast ability to invade and transform the maternal placental site.

<http://dx.doi.org/10.1016/j.pbj.2017.07.118>