

ANIMAL GENETICS AND GENOMICS

Association study reveals Th17, Treg, and Th2 loci related to resistance to *Haemonchus contortus* in Florida Native sheep¹

Zaira Magdalena Estrada-Reyes,^{*,2} Owen Rae,[†] Carol Postley,[‡] Myriam Berenice Jiménez Medrano,[†] Joel David Leal Gutiérrez,^{*} and Raluca Georgiana Mateescu^{*}

^{*}Department of Animal Sciences, University of Florida, Gainesville, FL, [†]Department of Large Animal Clinical Sciences, College of Veterinary Medicine, University of Florida, Gainesville, FL, and [‡]Fairmeadow Farm, Ocala, FL

¹This research was supported Southern SARE (GS17-173). ZMER was partially supported by Consejo Nacional de Ciencia y Tecnología. We gratefully acknowledge Dr. Francisco Peñagaricano, Assistant Professor at the University of Florida, for his guidance and support, and Eduardo Rodriguez and Kaitlyn Sarlo for their help during data analysis and sample processing.

²Corresponding author: zestrada@ufl.edu

ORCID number: [0000-0002-1541-5749](https://orcid.org/0000-0002-1541-5749) (Z. M. Estrada-Reyes).

Abstract

The aim of this study was to identify for the first time single nucleotide polymorphisms (SNPs) associated with *Haemonchus contortus* resistance in Florida Native sheep, using a targeted sequencing approach. One hundred and fifty-three lambs were evaluated in this study. At the start of the trial, phenotypic records for fecal egg count (FEC), FAMACHA score, body condition score (BCS), and weight were recorded and deworming of sheep with levamisole (18 mg/kg of body weight) was performed. Ten days post-deworming (baseline) and 28 d post-baseline, a full hematogram of each sheep was obtained and FEC, FAMACHA score, BCS, and weight were assessed. Average daily gain was calculated at the end of the trial. Out of 153 animals, 100 sheep were selected for genotyping using a targeted sequencing approach. Targeted sequencing panel included 100 candidate genes for immune response against *H. contortus*. SNPs were discarded if call rate <95% and minor allele frequency ≤ 0.05 . A mixed model was used to analyze the response variables and included the identity by state matrix to control for population structure. A contemporary group (age, group, and sex) was included as fixed effect. Bonferroni correction was used to control for multiple testing. Eighteen SNPs on chromosomes 1, 2, 3, 4, 6, 7, 11, 15, 18, 20, 24, and 26 were significant for different traits. Our results suggest that loci related to Th17, Treg, and Th2 responses play an important role in the expression of resistant phenotypes. Several genes including *ITGA4*, *MUC15*, *TLR3*, *PCDH7*, *CFI*, *CXCL10*, *TNF*, *CCL26*, *STAT3*, *GPX2*, *IL2RB*, and *STAT6* were identified as potential markers for resistance to natural *H. contortus* exposure. This is the first study that evaluates potential genetic markers for *H. contortus* resistance in Florida Native sheep.

Key words: Florida Native sheep, genome-wide association study, *Haemonchus contortus*, immune response, targeted sequencing

Introduction

Gastrointestinal infections are the main health problem for grazing sheep in the southern United States. The humid environmental conditions in this region are ideal for survival and growth of *Haemonchus contortus* and other gastrointestinal nematodes (GINs) of sheep (Miller et al., 1998). The predominant strategy for controlling GIN is anthelmintic treatment. However, anthelmintic resistance has reached epidemic proportions in small ruminant populations, and many cases the anthelmintic resistance has been reported to almost all available drugs (Miller and Craig, 1996; Zajac and Gipson, 2000; Kaplan, 2004; Howell et al., 2008; Goolsby et al., 2017). Breeding for parasite resistance offers many benefits to long-term sustainable sheep production. The main benefits of using selection to enhance parasite resistance in sustainable sheep breeding programs are that genetic change is permanent, the animal's performance is improved, and infectivity of pastures is decreased, leading to additional benefits for susceptible and young sheep grazing within the same flock (Bishop, 2012). Traditionally, selection of resistant sheep to GIN has focused on estimated breeding values calculated from phenotypic and pedigree information (Goddard and Hayes, 2009). Breeding values for resistance to GIN based on fecal egg count (FEC) are now available for sheep breeds in Australia (www.sheepgenetics.org.au) and New Zealand (www.sil.co.nz). Phenotypic recording for selection for parasite resistance can be complemented with other indicator traits such as FAMACHA® score, packed cell volume (PCV), or hematocrit (HCT), eosinophil count (EC), and determination of IgA, IgM, and IgG levels (Bishop, 2012). With current technologies, incorporation of genotype information, using genetic markers in sheep breeding programs aimed to select individuals for enhanced resistance to GIN, is a promising faster approach that reduces the generation interval in the selection process and avoids the requirement for animals to be challenged with gastrointestinal parasites.

In the United States, several sheep breeds have been studied and appear to have naturally developed enhanced resistance to GIN based on FEC. These breeds include St. Croix, Katahdin, Barbados Black Belly, Gulf Coast Native, and Florida Native (Zajac et al., 1988; Amarante et al., 1999a,b; Baker and Gray, 2004; Estrada-Reyes et al., 2018). Florida Native sheep is a multipurpose (wool and meat), landrace breed locally adapted to the warm humid environmental conditions from Florida. It is believed that this breed was developed from sheep (Churra) introduced by Spanish when they founded St. Agustin in 1565. Sheep became feral for several decades until Florida ended the open range at the end of World War II (<https://livestockconservancy.org/index.php/heritage/internal/cracker-sheep>). According to the American Livestock Breeds Conservancy, this heritage breed is now on the endangered list of sheep breeds (<http://livestockconservancy.org/>). To ensure genetic diversity and long-term survival of the breed, protection of the current genetic stock is critical and conservation efforts are required to promote its breeding and production. Natural parasite resistance is a common feature of Florida Native sheep, and good health and reproductive performance have persisted for many generations. The genetics underlying resistance to GIN in Florida Native sheep have been studied without identifying the genes or gene variants controlling variation, with selection based on phenotypic measures such as FEC and PCV (Zajac et al., 1988; Amarante et al., 1999a,b).

Genome-wide association studies have successfully identified genomic variants associated with complex traits

and biological pathways responsible for genetic variation with moderate to large effects in some livestock populations (Karim et al., 2000; Grisart et al., 2002). However, resistance to GIN is quantitative in nature with polygenic effects, and several studies aimed to identify the genetic variation controlling gastrointestinal parasite suggested a scattering of effects across the sheep genome (Kemper et al., 2011; McRae et al., 2014; Benavides et al., 2016; Sweeney et al., 2016). Many of the genomic regions identified from these studies are related to the sheep immune response against GIN.

The implementation of genomic markers associated to resistance to GIN could improve the selection of animals early in life and benefit the production costs. To date, results from several studies with other sheep breeds suggest that genetic architecture that underlines resistance to GIN relies on DNA polymorphisms within many genes. These studies suggest that resistance to GIN is quantitative in nature and polygenic (controlled by many genes with small effects) (McRae et al., 2014; Benavides et al., 2016; Sweeney et al., 2016). Exploration of genetic variation for resistance to GIN can be performed either with SNP-based genome scans or more specifically in candidate regions involved with innate and adaptive immune pathways related to GIN. Identification of variants such as SNPs in these candidate regions may help to identify a set of genetic markers significantly associated with parasite resistance. Thus, the objectives of this study were: 1) to use a targeted sequencing approach to identify SNPs in 100 genes related to the immune response during natural infections with *H. contortus* in Florida Native sheep; 2) to perform a SNP-based association analysis for FEC, FAMACHA score, HCT, red blood cell count (RBC), hemoglobin level (HGB), white blood cell count (WBC), neutrophil count (NEU), lymphocyte count (LC), monocyte count (MC), basophil count (BC), and EC; and 3) to identify the potential candidate genes associated to resistance to natural *H. contortus* infections in Florida Native sheep.

Materials and Methods

Sheep Population and Phenotypic Data

The research protocol was approved by the University of Florida Institutional Animal Care and Use Committee (approval number 201810108). All animals used in this study were from a commercial Florida Native sheep farm from Ocala, Florida. The selection flock was established in 1998 from an initial group of 80 rams ranked for *H. contortus* FEC, with high and low FEC rams mated with approximately 100 foundation dams. Since then, animals were selected for resistance to GIN solely on the basis of *H. contortus* FEC phenotyping and parentage recording. One hundred and fifty-three female and male sheep were tested for GINs under natural grazing conditions. Animals were born between December 2017 and February 2018 and naturally exposed to parasites since birth. They were approximately 3 and 5 mo old on average and were grouped based on age. At the beginning of the study, initial FEC, FAMACHA score, weight, and body condition score (BCS) were measured and animals were drenched with levamisole (18 mg/kg of body weight) the same day of initial screening. Then, animals returned to grazing conditions. Ten days post-deworming, FEC reduction was confirmed and FAMACHA score, weight, BCS, and a full hematogram of each sheep were performed. This date was used as baseline (0 d) for future phenotype collection dates. Finally, 28 d post-baseline (28 d), the same parasitological and hematological parameters were measured and ADG was determined based on weight and

days under study. Identification of GIN genus was carried out from fresh fecal samples collected directly from the rectum of evaluated sheep as described by Roberts and O'Sullivan (1950) at start of the trial (initial FEC) and during baseline (0 d) and 28 d post-baseline (28 d). However, due to low and relatively constant numbers for the other species, only eggs of *Haemonchus* spp. were recorded and analyzed.

The Shapiro-Wilk test was used to test all variables for normality. Box-Cox transformation was performed for all the traits recorded to obtain a normal distribution of values. The R "car" library was used to estimate the power parameter λ and carry out square root transformation for FEC (Estrada-Reyes et al., 2018) and logarithmic transformation (\log_{10}) for RBC, HGB, WBC, NEU, LC, MC, BC, and EC. For all the traits, the different time points evaluated during the study (initial screening, 0 d, and 28 d) and the difference between 28 d post-baseline and baseline (28 d minus 0 d) were included in the analysis. Spearman phenotypic correlations among all the parameters evaluated were obtained using R software (<https://cran.r-project.org>).

Genotyping Using Capture Sequencing

Genomic DNA from blood samples collected from the jugular vein using vacutainer tubes with anticoagulant EDTA at day 0 post-infection was isolated using DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA) according to the manufacturer's instructions and stored at -20°C . The DNA yield was calculated from a spectrophotometric measurement at 260 nm (NanoDrop-1000, Thermo Scientific), and the purity was assessed using a ratio 260/280 nm.

DNA samples (250 ng/ μL per sample) from 100 selected sheep samples with the highest ($>1,000$ eggs/g of feces, $n = 52$) and lowest (<900 eggs/g of feces, $n = 48$) FEC (Supplementary Figure 1) were sent to RAPiD Genomics (Gainesville, FL) for sequencing of 100 genes using DNA Capture sequencing. Briefly, this approach uses a hybrid capture-based enrichment sequencing in which regions of interest are captured by sequence-specific hybridization probes. A focused panel that contained a set of 100 genes was used and custom designed to include genomic regions of interest. Selection of the gene panel was performed based on results from previous studies in sheep (Araujo et al., 2009; MacKinnon et al., 2009; Silva et al., 2012; Venturina et al., 2013; McRae et al., 2014; Benavides et al., 2015, 2016; Sweeney et al., 2016; Berton et al., 2017; Estrada-Reyes et al., 2018). Also, genes related to the immune response against *H. contortus* and other GIN were considered as candidates for sequencing. The sheep genome assembly from *Ovis aries* (Oar_v4.0) available at the National Center for Biotechnology Information (NCBI) genome browser was used to design the biotinylated 120-mer probes used for sequencing. Probes were designed to capture sequences at each target gene.

For library preparation, Illumina's guidelines were followed and Nextera tagmentation, which converts DNA into adapter-tagged libraries, was used. Libraries were denatured and biotin-labeled probes were used for hybridization. Biotinylated probes were captured by streptavidin-coated beads. Then, DNA fragments bound to the streptavidin-coated beads were magnetically pulled down and enriched DNA fragments were eluted. After library preparation, libraries were captured by surface-bound oligos complementary to library adapters. Then, each library fragment was amplified into distinct, clonal clusters through bridge amplification. After cluster generation, templates were ready for sequencing. Illumina sequencing by synthesis chemistry was used for DNA Capture sequencing and

samples were then sequenced using a depth of 500–1,000 \times . Data from sequencing was demultiplexed, cleaned, and trimmed. The 3' ends were trimmed by removing low quality bases with <20 quality score reads. Clean reads were aligned to genome with MOSAIK software (Lee et al., 2014). For SNP calling, Freebayes software was used and VCFtools (Danecek et al., 2011) was used to generate VCF files. Supplementary Table 1 presents a summary of the genes evaluated in our study.

Quality Control and Association Analysis

Quality control of SNP data set was performed in JMP Genomics 6.0 software (SAS Institute Inc., Cary, NC). A total of 1,546 SNPs were available for analysis. Markers were removed based on genotype call rates per marker ($<95\%$) and minor allele frequency ≤ 0.05 , and the final marker dataset consisted of 1,277 SNPs. For the association study, a mixed model was used to evaluate the *H. contortus* FEC, FAMACHA score, HCT, RBC, HGB, WBC, NEU, LC, MC, BC, and EC. The GenABEL package (Aulchenko et al., 2007) from R software was used to perform the association analysis. Fixed effects included SNP and a contemporary group conformed by age of the animal, sex, and group. To control for population structure, the genomic relationship matrix was included in the analysis as random effect and it was calculated from SNP information. The mixed model used for the association analysis was as follows:

$$y = Xb + Zu + Wa + e$$

where y represents the vector of the phenotypic observation for each animal; b is the vector of fixed effects, u is the vector of random animal polygenic effect, $u \sim N(0, G\sigma_u^2)$, being G the additive genomic relationship matrix; a represents the fixed SNP additive effect, and e represents the residual vector, $e \sim N(0, I\sigma_e^2)$. The matrices X , Z , and W represent the incidence matrices for b , u , and a , respectively. SNP markers were tested one at a time, and a Bonferroni correction was used to control for multiple testing as follows:

$$\alpha = \frac{0.05}{(\text{number of candidate genes})} = 0.0005$$

Bioinformatics Analysis of Significant SNPs

Information regarding the identity of the genes underlying the significant SNPs associated with different traits was obtained from the sheep genome assembly (Oar_v4.0) available at the NCBI genome browser. Then, significant associated SNPs were screened for putative function to evaluate the possible changes in the free energy and the secondary structure of the ovine mRNA. The mRNA sequences from Oar_v4.0 available at the NCBI genome browser were downloaded for the genes with significant SNPs. For each mRNA sequence, a segment of 500 base pairs harboring the SNP in the middle of the sequence was used for prediction of mRNA secondary structures and molecular stability (free energy, G) using the RNA fold web server (Lorenz et al., 2011).

From each association analysis, genes with significant SNPs associated with the evaluated traits were analyzed using Gene Ontology software (Ashburner et al., 2000) to identify molecular function and immunological pathways. These genes were also used in the construction of a gene network related to *H. contortus* infection. The Cytoscape v3.7.1 software was used for gene network graphics (Shannon et al., 2003).

Results and Discussion

Phenotypic Data

The GIN genus identified from eggs were the following: *Haemonchus* spp, *Moniezia* spp, *Coccidia* spp, and *Strongyloides* spp. Table 1 shows the descriptive statistics for parasitological and hematological traits available for this study. Table 2 presents the phenotypic correlations between *H. contortus* FEC and other traits. Significant moderate negative correlations were observed between *H. contortus* FEC (initial and 28 d) and RBC, and HGB ($P < 0.01$). For HCT at 28 d, a significant moderate negative correlation was observed with initial *H. contortus* FEC ($P = 0.00005$), and with *H. contortus* FEC at 28 d ($P = 0.00009$). Significant moderate positive correlated responses were observed between initial *H. contortus* FEC and WBC (0 and 28 d) ($P = 0.0001$ and $P = 0.0008$, respectively). Also, *H. contortus* FEC at 28 d was positively correlated with WBC at 0 and 28 d ($P = 0.0003$ and $P = 0.0002$, respectively). Finally, significant moderate positive correlations were observed between ADG and *H. contortus* FEC (initial and 28 d) ($P = 0.03$ and $P = 0.013$, respectively). For FAMACHA, BC, and EC, no significant correlations were observed with *H. contortus* FEC (initial and 28 d). A significant strong positive correlation (0.89) ($P = 2.2 \times 10^{-16}$) between HGB at 28 d and RBC at 28 d was observed. Supplementary Table 1 shows the phenotypic correlated responses between all the traits under study.

Genomic Regions Identified by the Association Analysis

A total of 23 SNPs covering 14 genes were associated with *H. contortus* FEC, FAMACHA, RBC, HGB, WBC, NEU, BC, and EC.

No significant variants were detected for HCT, LC, and MC. A summary of the significant SNPs associated with these traits is presented in Table 3. Overall, associated SNPs identified in this study were located on intergenic and untranslated regions (UTRs) of 14 genes in OAR1, 2, 3, 4, 6, 7, 11, 15, 18, 19, 20, 24, and 26 (Fig. 1). The proportion of variance explained by majority of the significant SNPs associated with FEC, FAMACHA score, WBC, RBC, BC, and EC was less than 0.2. Thus, based on the current and previous studies, it is possible that many genes from the immune response (with small effects) are contributing to parasite resistance/susceptibility in Florida Native sheep populations.

For initial *H. contortus* FEC, SNPs located in exon 24 of *ITGA4* (OAR2:127159353), intron 13 of *STAT3* (OAR11:41860577), and the 5' UTR of *MUC15* (OAR15:55310482) genes were significant. These proportions of variance explained by these SNPs was 0.14, 0.10 and 0.11, respectively, for initial *H. contortus* FEC. The OAR2:127159353 is a G to A substitution which codes for a synonymous codon (Val), where the GG genotype ($1,179.4 \pm 938.37$ eggs/g of feces) had lower FEC when compared with the AG ($3,239.94 \pm 2,094.891$ eggs/g of feces) and AA ($6,890.1 \pm 4,099.9$ eggs/g of feces) genotypes. For this polymorphism, moderate changes in mRNA stability were observed for the "G" allele (-118.4 kcal/mol) when compared with the "A" allele (-119.1 kcal/mol), with great changes in mRNA secondary structure.

The OAR11:41860577 generates a G to C substitution, and OAR15:55310482 confers a base pair change from T to C. For OAR15:55310482, only two genotypes were observed, and the TT genotype (945.6 ± 804.1 eggs/g of feces) had lower FEC when compared to the CT genotype ($4,444.4 \pm 2,300.3$ eggs/g of feces).

Table 1. Descriptive statistics for *H. contortus* fecal egg count (FEC, eggs/gram) FAMACHA score, average daily gain (ADG), body condition score (BCS), hematocrit (HCT, %), red blood cell count (RBC, M/ μ L), hemoglobin level (HGB, g/dL), white blood cell count (WBC, K/ μ L), neutrophil count (NEU, K/ μ L), lymphocyte count (LC, K/ μ L), monocyte count (MC, K/ μ L), basophil count (BC, K/ μ L), and eosinophil count (EC, K/ μ L) at -10 d baseline, 0 d (baseline), and 28 d post-baseline in Florida Native sheep

Trait	Time	n	Mean	SD	Min	Max
FEC, eggs/gram	Initial FEC (-10 d)	100	1,926.00	312.6	0.00	15,800.00
	FEC (0 d)	100	21.00	7.10	0.00	500.0
	FEC (28 d)	100	1,593.00	270.5	0.00	19,300.00
FAMACHA, score	Initial FAMACHA (-10 d)	100	2.93	0.72	2.00	4.00
	FAMACHA (0 d)	100	2.87	0.63	2.00	5.00
	FAMACHA (28 d)	100	2.75	0.68	1.00	4.00
ADG, g		100	167.83	58.97	49.89	344.73
BCS, score	Initial BCS (-10 d)	100	2.32	0.26	1.75	3.00
	BCS (0 d)	100	2.42	0.28	2.00	3.25
	BCS (28 d)	100	2.37	0.26	1.75	3.00
HCT, %	HCT (0 d)	100	41.00	0.40	24.00	52.00
	HCT (28 d)	100	40.00	0.50	22.00	50.00
RBC, M/ μ L	RBC (0 d)	100	13.11	1.22	10.64	16.40
	RBC (28 d)	100	13.42	1.80	7.63	18.17
HGB, g/dL	HGB (0 d)	100	11.82	1.01	8.60	14.00
	HGB (28 d)	100	11.99	1.37	7.10	15.10
WBC, K/ μ L	WBC (0 d)	100	10.30	2.78	5.22	21.68
	WBC (28 d)	100	10.24	2.24	4.60	15.06
NEU, K/ μ L	NEU (0 d)	100	2.73	1.66	0.55	12.26
	NEU (28 d)	100	2.40	0.79	1.00	4.36
LC, K/ μ L	LC (0 d)	100	5.48	1.90	1.63	9.62
	LC (28 d)	100	5.86	1.97	1.28	10.27
MC, K/ μ L	MC (0 d)	100	1.42	0.72	0.21	3.33
	MC (28 d)	100	1.65	0.74	0.24	4.20
BC, K/ μ L	BC (0 d)	100	0.07	0.10	0.01	0.61
	BC (28 d)	100	0.05	0.06	0.01	0.43
EC, K/ μ L	EC (0 d)	100	0.48	0.47	0.01	3.15
	EC (28 d)	100	0.36	0.24	0.02	1.19

Table 2. Phenotypic correlations between initial (–10 d) or 28 d *H. contortus* fecal egg count (FEC) and neutrophil count (NEU) FAMACHA (28 d), hematocrit (HCT), red blood cell count (RBC, 28 d), hemoglobin level (HGB, 28 d), white blood cell count (WBC, 0 and 28 d), basophil count (BC 0 d), and eosinophil count (EC 28 d)¹

		FAMACHA	HCT	RBC	HGB	WBC		NEU	BC	EC
		28 d	28 d	28 d	28 d	0 d	28 d	0 d	0 d	28 d
FEC	Initial (–10 d)	–0.004	–0.4	–0.36	–0.36	0.37	0.32	0.29	0.03	0.02
	28 d	–0.14	–0.38	–0.27	–0.27	0.34	0.14	0.2	0.04	0.17

¹Values in bold indicate significance at $P < 0.05$.

For this SNP, the “T” allele had greater mRNA stability (–135.6 kcal/mol) when compared with the “C” allele (–145.27 kcal/mol) and strong changes in mRNA structure were observed (Fig. 2).

Integrin subunit alpha 4 (ITGA4) gene codes for a cell adhesion molecule related to eosinophil mobilization (Rothenberg et al., 2001). Therefore, the protein complex co-formed by ITGA4 protein and beta 1 or beta 7 subunits regulates cell motility and migration. Thus, it is possible that eosinophils use this protein complex to be mobilized to the area of infection through IL-5 independent or dependent mechanisms. Eosinophils are produced in the bone marrow and are commonly found in almost all the sheep gastrointestinal tract. Sheep eosinophilia is related to reduction in *H. contortus* larvae establishment (Balic et al., 2006; Terefe et al., 2007). In calves exposed to *Ostertagia*, *Cooperia*, and *Nematodirus* spp., higher levels of ITGA4 expression is observed in the mesenteric lymph nodes of resistant animals. In peripheral blood from Chinese indigenous white goats exposed to *H. contortus*, ITGA4 gene was identified as upregulated in resistant individuals with lower FEC when compared with susceptible animals (Bhuiyan et al., 2017).

The STAT3 gene belongs to the STAT family genes known as signal transducer and activator of transcription. STAT3 protein can be activated by phosphorylation in response to several key activators such as IL-6, IL-10, IL-23, IL-21, and IL-11 cytokines to initiate Th17 signaling and IL17 production. Stronger Th17 response is observed in resistant Caribbean hair sheep at 3 d post-challenge with *H. contortus* (MacKinnon et al., 2009). Therefore, excretory and secretory proteins from *H. contortus* larvae are linked to increased Th17 activation and IL17 production (Gadahi et al., 2016). Thus, after initial mechanisms of parasite rejection and complement activation, Th17 response is activated to regulate pathogen clearance and tissue inflammation.

Mucin 15 (MUC15) gene produces a high-molecular-weight membrane-bound glycoprotein consisting of a mucin peptide backbone and O-linked oligosaccharides. Epithelial surfaces of the gastrointestinal tract are composed of these types of proteins and other mucins to protect against GINs and represent the first line of defense against pathogens (Moncada et al., 2003). It is possible that some of these types of mucins are ligands for GIN adhesins (McGuckin et al., 2011). In Dorper × Red Maasai sheep, MUC15 was identified as potential candidate marker for FEC (Benavides et al., 2015). Thus, ITGA4 and MUC15 could play an important role during the initial establishment and rejection of *H. contortus* in Florida Native sheep.

For *H. contortus* FEC at 28 d and the difference between 28 and 0 d, the same significant SNPs were identified within the 5'UTR of IL2RB (OAR3:180166632 and OAR3: 180167554) and exon 6 of CFI (OAR6: 15681195) genes. The proportion of variance explained by these polymorphisms was 0.14, 0.13, and 0.15, respectively, for FEC at 28 d. These SNPs are substitutions from G to A, C to G, and G to A, respectively. No significant changes were observed in the stability and secondary structure of the mRNA

for OAR3:180166632 in the IL2RB gene. For OAR3: 180167554, the CC genotype ($3,451.9 \pm 1,725.2$ eggs/g of feces) presented more FEC when compared with the GC ($2,397.3 \pm 1,793.3$ eggs/g of feces) and GG (677.2 ± 539.3 eggs/g of feces) genotypes, with no significant changes in stability and secondary structure of the mRNA (Fig. 2).

The OAR6: 15681195 is a synonymous mutation with a minor change in mRNA stability of 0.6 kcal/mol but strong impact on the mRNA secondary structure (Fig. 2). For this polymorphism, the GG genotype (249.9 ± 124.9 eggs/g of feces) had lower FEC when compared with the AG ($2,370.7 \pm 1,775.2$ eggs/g of feces) and AA ($1,142.1 \pm 909.8$ eggs/g of feces) genotypes.

Complement factor I (CFI) gene encodes a serine protease implicated in regulation of the complement cascade. In hair sheep from the Canary Islands, high expression of CFI gene was observed in resistant animals (Guo et al., 2016). In our study, a genetic variant within this gene was associated with FEC at 28 d suggesting that complement activation is one of the earliest events in the Florida Native sheep immune responses to protect against *H. contortus*.

For FAMACHA at 28 d, a significant SNP (OAR3:180148777) in exon 12 of IL2RB associated with this trait. The proportion of variance explained by OAR3:180148777 was 0.13. This SNP (G/A) is part of a synonymous codon and no significant changes in stability and secondary structure of the mRNA were observed. For this locus, the GG genotype (2.11 ± 0.64) had lower FAMACHA score when compared with AG (2.85 ± 0.67) and AA (2.95 ± 0.58) genotypes. Also, for FAMACHA at 28 d, one SNP (OAR7:73930804) in the 3'UTR of GPX2 gene was significantly associated with this trait and explained 0.03 of the proportion of variance observed. The OAR7:73930804 generates an A to G substitution which results in minor changes in mRNA stability (a difference of –0.4 kcal/mol between variants) and mRNA secondary structure (Fig. 3).

Similar findings have been observed in Santa Inês sheep exposed to natural *H. contortus* infections, where genetic variants within IL2RB gene are associated to HCT (Berton et al., 2017). IL2RB protein is part of the IL-2 receptor complex (IL2R) co-formed by IL2RA, IL2RB, and IL2R γ subunits, and it is activated by IL2 cytokine. This cytokine activates Treg cells, a subpopulation of T cells responsible for maintenance of immunological self-tolerance and homeostasis through immune suppression. Long-term infection with GINs is linked to activation of Treg populations (Smits and Yazdanbakhsh, 2007). The exact role of Treg response during *H. contortus* infections in Florida Native sheep remains unclear but a faster switch from Th1 to Th2/Treg response is observed in resistant Suffolk sheep exposed to *Teladorsagia circumcincta* (Hassan et al., 2011).

The GPX2 gene encodes the glutathione peroxidase 2, which catalyzes the reduction of peroxides and protects against oxidative damage in the epithelium of the gastrointestinal tract. The generation of host oxidants such as reactive oxygen and

Table 3. Significant SNPs associated with *H. contortus* fecal egg count (FEC, eggs/gram), FAMACHA score, red blood cell count (RBC, M/ μ L), hemoglobin level (HGB, g/dL), white blood cell count (WBC, K/ μ L), neutrophil count (NEU, K/ μ L), lymphocyte count (LC, K/ μ L), monocyte count (MC, K/ μ L), basophil count (BC, K/ μ L), and eosinophil count (EC, K/ μ L) in Florida Native sheep¹

Trait	OAR	Position (bp)	Gene name	SNP	$-\log_{10}$ P-values	Estimate	Proportion of variance explained	MAF	Mutation
FEC (initial -10 d)	2	127,159,353	Exon 24 ITGA4	G/A	4.108	21.58	0.14	0.16	Synonymous
FEC (initial -10 d)	11	41,860,577	Intron 13 STAT3	G/C	4.052	15.94	0.10	0.11	Intronic
FEC (initial -10 d)	15	55,310,482	5'UTR MUC15	T/C	3.773	24.22	0.11	0.12	UTR
FEC (28 d)	3	180,166,632	5'UTR IL12RB2	G/A	4.065	21.58	0.14	0.11	UTR
FEC (28 d)	3	180,167,554	5'UTR IL12RB2	C/G	3.531	15.94	0.13	0.12	UTR
FEC (28 d)	6	15,681,195	Exon 6 CFI	G/A	4.009	24.22	0.15	0.11	Synonymous
FAMACHA (28 d)	3	180,148,777	Exon 12 IL12RB2	G/A	3.616	-0.33	0.13	0.40	UTR
FAMACHA (28 d)	7	73,930,804	5'UTR GPX2	A/G	3.531	-0.36	0.03	0.49	Synonymous
RBC (28 d)	18	26,116,868	Exon 18 IL16	C/T	4.558	-0.08	0.16	0.11	Synonymous
HGB (28 d)	18	26,116,868	Exon 18 IL16	C/T	3.563	-1.60	0.42	0.11	Synonymous
WBC (0 d)	3	162,039,038	3'UTR STAT6	G/A	3.42	0.06	0.02	0.46	UTR
WBC (28 d)	6	49,768,053	5'UTR PCDH7	G/T	4.781	-0.008	0.17	0.11	UTR
WBC (28 d)	6	49,768,057	5'UTR PCDH7	G/T	3.61	-0.006	0.12	0.11	UTR
WBC (28 d)	15	55,310,748	5'UTR MUC15	C/T	3.999	0.0005	0.14	0.38	UTR
NEU (0 d)	6	90,454,870	3'UTR CXCL10	A/G	3.809	0.46	0.14	0.12	UTR
NEU (0 d)	20	26,766,451	3'UTR TNF	C/T	3.668	0.67	0.13	0.30	UTR
BC (0 d)	24	33,935,613	3'UTR CCL26	C/T	4.335	-1.65	0.16	0.46	UTR
EC (28 d)	2	127,200,458	Intron 10 ITGA4	A/G	3.687	0.25	0.13	0.12	Intronic
EC (28 d)	2	14,799,731	Exon 2 TLR3	T/C	4.094	0.26	0.15	0.11	Synonymous

¹For each SNP, the chromosome (OAR), chromosomal position (bp), gene name and region, base pair substitution (SNP), estimate of SNP effect, proportion of variance explained by SNP, minor allele frequency (MAF), and mutation type (UTR, synonymous or intronic) are presented.




Figure 1. Manhattan plots for natural *H. contortus* fecal egg count (A, FEC), FAMACHA score (B), red blood cell count (C, RBC), hemoglobin level (C, HGB), white blood cell count (D, WBC), neutrophil count (E, NEU), basophil count (E, BC), and eosinophil count (E, EC). The $3.3 \cdot 10^{-6}$ P-value threshold was indicated with the red dotted line. The red dots indicate the significant SNPs identified above the threshold for each trait.

nitrogen species is important for parasite control (Ingham et al., 2008; Patel et al., 2009). In Perendale sheep, expression of *GPX2* was upregulated in the duodenum of susceptible individuals when compared to resistant animals (Keane et al., 2006). In contrast, expression of *GPX2* gene was upregulated in resistant

sheep infected with *H. contortus* (Menzies et al., 2010) during initial days of infection. Similar findings were observed in resistant Australian sheep, where an increase in the expression of *GPX2* gene is related to *H. contortus* challenge during the first days of infection (Lees et al., 2011). Results from these studies




Figure 2. Change in the free energy of the thermodynamic ensemble for the different mRNA variants observed for *H. contortus* fecal egg count (FEC) in Florida Native sheep. Predicted secondary structure of mRNA of 500 bp segment of *ITGA4* (A: G allele and B: A allele), *MUC15* (C: T allele and D: C allele), and *CFI* (E: A allele and F: G allele) genes.

suggest that increase of *GPX2* expression is essential during larval expulsion in the first days of infection.

For RBC and HGB at 28 d, the same SNP located in *IL16* gene (OAR18: 26116868, C/T) was associated with both traits.

The proportion of variance explained by OAR18: 26116868 was 0.16 and 0.12 for these traits, respectively. This SNP codes for a synonymous codon (Ser) which did not have significant changes in stability and secondary structure of the mRNA (Fig. 3). For




Figure 3. Change in the free energy of the thermodynamic ensemble for the different mRNA variants observed for FAMACHA score and white blood cell count (WBC, 0 and 28 d) in Florida Native sheep. Predicted secondary structure of mRNA of 500 bp segment of *GPX2* (A: G allele and B: A allele), *STAT6* (C: G allele and D: A allele), and *MUC15* (E: C allele and F: T allele) genes.




Figure 4. Change in the free energy of the thermodynamic ensemble for the different mRNA variants observed for neutrophil count (NEU, 0 d) and basophil count (BC, 0 d) in Florida Native sheep. Predicted secondary structure of mRNA of 500 bp segment of *CXCL10* (A: A allele and B: G allele), *TNF* (C: C allele and D: T allele), and *CCL26* (E: C allele and F: T allele) genes.

OAR18: 26116868 only two genotypes were observed, and the CC genotype presented higher RBC (13.66 ± 1.58 M/ μ L) and HGB (12.17 ± 1.21 g/dL) when compared with the CT genotype (11.48 ± 1.75 M/ μ L and 10.57 ± 1.67 g/dL, respectively).

IL16 cytokine is related to inflammatory response and its synthesis is induced by T lymphocytes upon antigen exposure. Also, IL-16 promotes CD4⁺ T-cell production and it is commonly known as lymphocyte chemoattractant factor (O'Shea et al., 2013). In susceptible Perendale sheep selected for high FEC, expression of *IL16* gene was downregulated in duodenum tissue (Diez-Tascón et al., 2005). In contrast, in Angus yearlings infected with gastrointestinal parasites, *IL16* gene was highly expressed in the mesenteric lymph nodes of susceptible individuals (Araujo et al., 2009).

For WBC at 0 d, one SNP (OAR3: 162039038) in the 3'UTR of *STAT6* gene was associated with this trait and explained 0.12 of the proportion of variance observed. For OAR3: 162039038, the G to A substitution results in moderate changes in mRNA stability (-1.3 kcal/mol) but great impact on mRNA secondary structure (Fig. 4).

Several studies have suggested that mechanisms related to *STAT6* activation via IL4 receptor are required for the expulsion of gastrointestinal parasites in sheep (Urban et al., 1998; Venturina et al., 2013). Signal transducer and activator of transcription 6 (*STAT6*) gene encodes a protein that mediates Th2 signaling. Th2 response is linked to resistance to GIN in sheep (Pernthaner et al., 2005). Commonly features of Th2 response include IL4, IL5, and IL13 cytokine production, eosinophil infiltration, and activation mediated by IL5, and IgE production. IgE antibodies bind to antigens expressed in the membrane of gastrointestinal parasites. Then, Fc receptors (FcεRI and II) located on the surface of eosinophils recognize IgE antibodies, leading to degranulation and releasing of cytotoxic proteins that damage the parasite membrane (Meeusen and Balic, 2000; McRae et al., 2015). In wool sheep infected with *H. contortus*, stronger Th2 response is observed at 27 d post-infection, with upregulation of *IL4* and downregulation of *IFNγ* expression.

Two SNPs (OAR6:49768053 and OAR6:49768057) in the 5'UTR region of *PCDH7* gene were associated to WBC at 28 d. The




Figure 5. Change in the free energy of the thermodynamic ensemble for the different mRNA variants observed for eosinophil count (EC, 28 d) in Florida Native sheep. Predicted secondary structure of mRNA of 500 bp segment of *ITGA4* (A: A allele and B: G allele), and *TLR3* (C: T allele and D: C allele) genes.

proportion of variance explained by both SNPs was 0.17 and 0.12, respectively. Both polymorphisms generate G to T substitutions with no significant changes in stability and secondary structure of the mRNA. For OAR6:49768053, the GG genotype (10.59 ± 2.34 K/ μ L) presented higher WBC at 28 d when compared with the GT (10.06 ± 2.17 K/ μ L) and the TT (6.97 ± 0.40 K/ μ L) genotypes. For OAR6:49768057, the GT genotype (10.96 ± 2.36 K/ μ L) presented higher WBC when compared with the GG (10.59 ± 2.34 K/ μ L) and the TT (9.65 ± 2.22 K/ μ L) genotypes.

Protocadherin 7 (PCDH7) gene encodes a protein from the cadherin family. Cadherins are cell adhesion molecules responsible for cell interactions and maintenance of tissue structure and morphogenesis (Li and Gasbarre, 2010). Studies in Angus yearlings observed upregulation of PCDH7 expression in susceptible animals infected with GINs. However, the exact role of PCDH7 protein during *H. contortus* infections in Florida Native sheep remains unknown.

One SNP (OAR15: 55310748) in the 5'UTR of MUC15 gene was also associated with WBC at 28 d, which results in a base pair change from C to T. The OAR15: 55310748 explained 0.14 of the proportion of variance observed for this trait. This SNP is a putative functional SNP with great changes in mRNA stability (5 kcal/mol) and mRNA secondary structure (Fig. 4). For this locus, the CC genotype (10.72 ± 2.50 K/ μ L) presented higher number of WBC when compared with the CT (10.01 ± 2.09 K/ μ L) and TT (10.37 ± 2.25 K/ μ L) genotypes.

For NEU at 0 d, two SNPs (OAR6:90454870 and OAR20:26766451) in the 3' UTR of CXCL10 and TNF genes were associated with this trait. These SNPs explained 0.14 and 0.13 of the proportion of variance observed. The OAR6:90454870 is an A to G substitution, and OAR20: 26766451 is a C to T substitution. Both polymorphisms are putative functional SNPs with moderate changes in mRNA stability (a difference of 2.6 kcal/mol and -2.1 kcal/mol between variants, respectively) but great impact on mRNA structure. For the OAR6:90454870, the "A" allele presented higher mRNA stability (-166.5 kcal/mol) than the "G" allele (-169.1 kcal/mol) (Fig. 5). The AA genotype (2.58 ± 1.62 K/ μ L) had lower NEU when compared with the AG (3.19 ± 0.68 K/ μ L) and the GG (6.82 ± 0.72 K/ μ L) genotypes.

For OAR20:26766451 located on TNF gene, the CC genotype (3.20 ± 1.98 K/ μ L) had higher NEU when compared to the CT (2.27 ± 0.91 K/ μ L) and the TT (1.94 ± 0.61 K/ μ L) genotypes, with minor changes in mRNA stability but moderate impact on mRNA secondary structure (Fig. 5).

The TNF and CXCL10 genes encode proteins related to inflammatory response. TNF protein is commonly secreted by macrophages. CXCL10 protein is produced by monocytes and its production is induced by interferon gamma (IFN γ). In Santa Inés sheep, polymorphisms within CXCL10 gene have been associated with platelet count. Thus, it is possible that TNF and CXCL10 genes are related to inflammatory responses in Florida Native sheep exposed to natural *H. contortus* infections.




Figure 6. Gene network for the genes with significant SNPs associated with *H. contortus* fecal egg count (FEC), FAMACHA score (FAM), red blood cell count (RBC), hemoglobin level (HGB), white blood cell count (WBC), neutrophil count (NEU), basophil count (BC), and eosinophil count (EC). The traits FEC, FAMACHA, WBC, and EC shared more genes in common with significant SNPs than RBC, HGB, NEU, and BC.

For BC at 0 d, one SNP (OAR24:33935613) in the 3'UTR region of CCL26 gene was associated with this trait. This SNP explained 0.16 of the proportion of variance observed. The C to T substitution from OAR24: 33935613 results in a great change in mRNA stability of -4.8 kcal/mol and has a great impact on mRNA secondary structure. The "T" allele (-201.9 kcal/mol) had greater mRNA stability than the "C" allele (-206.7 kcal/mol) (Fig. 5). The TT genotype (0.14 ± 0.13 K/ μ L) presented higher BC when compared to the CT (0.07 ± 0.06 K/ μ L) and the CC (0.05 ± 0.04 K/ μ L) genotypes.

The CCL26 protein is a member of the eotaxin family. Mature eosinophils are activated and migrate to the site of infection in response to various chemoattractants, such as IL-5, CCL11, CCL24, and CCL26 (Rosenberg et al., 2013). Upregulation of CCL26 gene has been observed in draining lymph node tissue from Martinik \times Romane back-cross resistant sheep exposed to *H. contortus* (Sallé et al., 2014). Based on these results, eosinophil migration could play an important role to control *H. contortus* in Florida Native sheep.

Two SNPs (OAR2:127200458 and OAR26: 14799731) in ITGA4 gene and exon 2 of TLR3 were associated with EC at 28 d. These polymorphisms explained 0.13 and 0.15 of the proportion of variance observed for EC at 28 d. The A to G substitution from

OAR2:127199427 results in a minor change of -0.4 kcal/mol in mRNA stability but a moderate change in mRNA secondary structure (Fig. 6). Minor changes in mRNA stability were observed for the "A" allele (-127.9 kcal/mol) when compared to the "G" allele (-128.3 kcal/mol). For this SNP, the GG genotype (0.39 ± 0.24 K/ μ L) had greater EC when compared with GA (0.27 ± 0.23 K/ μ L) and AA (0.19 ± 0.18 K/ μ L) genotypes.

The T to C substitution from OAR26: 14799731 results in a minor change of 0.9 kcal/mol but has a strong impact on mRNA secondary structure (Fig. 6). The "T" allele (-91.9 kcal/mol) presented greater mRNA stability than the "C" allele (-92.8 kcal/mol), and the TT genotype (0.39 ± 0.37 K/ μ L) had greater EC than the CT (0.16 ± 0.15 K/ μ L) and CC (0.24 ± 0.23 K/ μ L) genotypes.

Our results could suggest that genetic variation within ITGA4 gene could be essential for controlling *H. contortus* infections in Florida Native sheep, due to its association with FEC and EC. Toll-like receptors (TLRs) are commonly expressed in antigen-presenting cells and other immune cells. TLR genes are found upregulated in the gut mucosa of resistant sheep infected with *H. contortus* and *Trichostrongylus colubriformis* (Ingham et al., 2008). TLR3 signaling could be one of the first immune response mechanisms activated by *H. contortus* in Florida Native sheep.

Table 4. Immunological pathways related to *H. contortus* FEC, FAMACHA score, red blood cell count (RBC), hemoglobin level (HGB), white blood cell count (WBC), neutrophil count (NEU), basophil count (BC), and eosinophil count (EC) in Florida Native sheep; Gene = gene name; GO term = gene ontology pathway name; Number of genes in gene list = number of genes from list identified in the pathway; Number of genes in GO term = total number of genes included in the GO term

Trait	Gene	GO term	GO term name	Number of genes in gene list	Number of genes in GO term
FEC	ITGA4, STAT3, MUC15, IL2RB, CFI	GO:0060397	JAK-STAT cascade involved in growth hormone signaling pathway	2	6
		GO:0043373	CD4-positive, alpha-beta T-cell lineage commitment	2	8
		GO:0002295	T-helper cell lineage commitment	2	8
		GO:0070102	Interleukin-6-mediated signaling pathway	2	9
		GO:0072540	T-helper 17 cell lineage commitment	2	6
		GO:0072539	T-helper 17 cell differentiation	2	6
		GO:0072538	T-helper 17 type immune response	2	6
		GO:1905562	Regulation of vascular endothelial cell proliferation	2	11
		GO:1905564	Positive regulation of vascular endothelial cell proliferation	2	7
		GO:0060259	Regulation of feeding behavior	2	
FAMACHA score	IL2RB, GPX2	GO:0098869	Cellular oxidant detoxification	1	86
		GO:0019221	Cytokine-mediated signaling pathway	1	91
		GO:0005125	Cytokine activity	1	173
		GO:0042093	T-helper cell differentiation	1	17
		GO:0002295	T-helper cell lineage commitment	1	8
RBC, HGB	IL16	GO:0002294	CD4-positive, alpha-beta T-cell differentiation involved in immune response	1	17
		GO:0035771	Interleukin-4-mediated signaling pathway	1	2
		GO:0048295	Positive regulation of isotype switching to IgE isotypes	1	2
		GO:0002439	Chronic inflammatory response to antigenic stimulus	1	1
		GO:1901740	Negative regulation of myoblast fusion	1	1
WBC	STAT6, PCDH7, MUC15	GO:0031728	CCR3 chemokine receptor binding	1	4
		GO:0034138	Toll-like receptor 3 signaling pathway	2	4
NEU	CXCL10, TNF	GO:0002439	Chronic inflammatory response to antigenic stimulus	1	1
		GO:1901740	Negative regulation of myoblast fusion	1	1
BC	CCL26	GO:0031728	CCR3 chemokine receptor binding	1	4
EC	ITGA4, TLR3	GO:0034138	Toll-like receptor 3 signaling pathway	2	4

Overall, significant SNPs associated with FEC, FAMACHA score, WBC, NEU, BC, and EC in Florida Native sheep exposed to natural *H. contortus* infection are putative functional SNPs that affect the mRNA secondary structure and stability. Some of the genes (*ITGA4*, *STAT3*, *MUC15*, *IL2RB*, and *CFI*) containing the significant SNPs for FEC were previously highlighted as potential candidate markers for gastrointestinal parasite resistance in sheep populations (Periasamy et al., 2014, Benavides et al., 2015), and identified as differentially expressed in Chinese goats and Angus cattle during *H. contortus* infection and GIN (Araujo et al., 2009; Bhuiyan et al., 2017).

Results presented in this study confirm that stability of mRNA and conformational changes in the secondary structure of mRNA are important and possible causal mechanisms of gene expression variability. The 5' UTR region is known to influence mRNA translation efficiency and secondary structures within this region could inhibit translational mechanisms (Leppek et al., 2018). Also, polymorphisms within the UTR regions confer more deleterious effects than polymorphisms in coding regions (Johnson et al., 2011).

Immunological Pathways Related to *H. contortus* Infection

Five pathways were related to significant SNPs associated with *H. contortus* FEC. These pathways include JAK-STAT signaling pathway, CD4⁺ T-cell commitment, Th17 signaling, endothelial cell proliferation, and leptin signaling and feeding behavior (Table 4). Leptin is commonly secreted by adipocytes, the placenta, and gastric epithelial cells. High concentrations of leptin in the stomach showed to induce inappetence in rats (Bado et al., 1998). Some studies suggested that increase in leptin concentration is responsible for immune mediated reduction in feed intake in sheep infected with *T. colubriformis* and *T. circumcincta* (Stear

et al., 2003; Greer et al., 2009). Thus, inappetence could be used by Florida Native sheep as a protective response to minimize feed intake while the damaged mucosa by *H. contortus* is being repaired.

For FAMACHA, two pathways related to cellular oxidant detoxification and cytokine-mediated signaling were identified. For ADG, and RBC/HGB, pathways related to positive regulation of T-cell proliferation and cytokine activity were observed. The related pathways to WBC included T-helper differentiation and commitment, CD4⁺ T-cell differentiation, IL4-mediated signaling, and positive regulation of isotype switching to IgE. For NEU, pathways related to inflammatory response were identified. Finally, for BC and EC, pathways such as CCR3 chemokine receptor signaling and toll-like receptor 3 signaling were associated with these traits.

Immune response pathways associated with genes containing significant SNPs for WBC were related to CD4⁺ T-cell differentiation and Th2 response (IL4 signaling and isotype switching to favor IgE antibody production). From these genes, polymorphisms in *MUC15* gene were also associated with FEC in our study. Although different SNPs were associated with both traits, the 5'UTR of *MUC15* gene seems to play an essential role for *H. contortus* protection. Thus, initial establishment and rejection of *H. contortus* could be mediated by *MUC15* gene and other molecules in Florida Native sheep.

From our gene network diagram, FEC, FAMACHA, WBC, and EC had more shared genes with significant SNPs when compared with NEU, HGB, RBC, and BC (Fig. 6). Also, FEC was the trait with more associated SNPs in our population. Some of these traits such as FEC and FAMACHA could be easily incorporated into future breeding programs to control GIN in Florida Native sheep.

Overall, significant polymorphisms associated with these traits highlighted potential immune response mechanisms




Figure 7. Immune response mechanisms associated with natural *H. contortus* infection. Significant polymorphisms in *ITGA4* and *MUC15* genes are related to initial rejection of the larvae. For innate immune response, significant SNPs were identified in *TLR3* and *PCDH7* genes. For the inflammatory response, significant polymorphisms in *CXCL10* and *TNF* genes were observed. Activation of mature eosinophils, antigen recognition by T cells, T-cell proliferation, Th17 signaling, protection against oxidative damage, Treg signaling, and Th2 response were related to significant SNPs within *CCL26*, *IL16*, *STAT3*, *GPX2*, *IL2RB*, and *STAT6* genes, respectively.

related to: initial nematode rejection, innate immune response (inflammatory response and complement activation), activation of eosinophils (IL5 independent), antigen recognition by T cells, T-cell proliferation and cytokine production, Th17 signaling, protection against oxidative damage, Treg signaling, and Th2 signaling and IgE production. These mechanisms could be representative of the potential immune mechanisms used by Florida Native sheep to control natural *H. contortus* infections (Fig. 7).

Implications

This study is the first report of potential candidate markers for parasite resistance in Florida Native sheep. However, future studies using monospecific infection with *H. contortus*, genome-wide scans, and additional animals would be required to validate our findings before these markers could be used with confidence in selection programs.

SNPs within 14 genes were significantly associated with FEC, FAMACHA score, WBC, NEU, BC, and EC. These genes were related to JAK-STAT signaling, CD4⁺ T-cell differentiation and commitment, Th17 signaling, leptin signaling and regulation of feeding behavior, cytokine-mediated signaling, Th2 response, TLR3 signaling, and CCR3 chemokine receptor binding. It is possible that immune response against natural *H. contortus* infections is mediated by these pathways in Florida Native sheep. Our results indicate that *ITGA4*, *MUC15*, *TLR3*, *PCDH7*, *CFI*, *CXCL10*, *TNF*, *CCL26*, *IL16*, *STAT3*, *GPX2*, *IL2RB*, and *STAT6* genes could be potential markers for resistance to *H. contortus* exposure in Florida Native sheep. Future studies will need to validate these markers in other Florida Native sheep populations before they can be applied.

Supplementary Data

Supplementary data are available at *Journal of Animal Science* online.

Literature Cited

- Amarante, A. F. T., T. M. Craig, W. S. Ramsey, S. K. Davis, and F. W. Bazer. 1999a. Nematode burdens and cellular responses in the abomasal mucosa and blood of Florida Native, Rambouillet and crossbred lambs. *Vet. Parasitol.* **80**:311–324. doi:10.1016/S0304-4017(98)00229-5
- Amarante, A. F., T. M. Craig, W. S. Ramsey, N. M. El-Sayed, A. Y. Desouki, and F. W. Bazer. 1999b. Comparison of naturally acquired parasite burdens among Florida native, Rambouillet and crossbred ewes. *Vet. Parasitol.* **85**:61–69. doi:10.1016/S0304-4017(99)00103-x
- Araujo, R. N., T. Padilha, D. Zarlenga, T. Sonstegard, E. E. Connor, C. Van Tassel, W. S. Lima, E. Nascimento, and L. C. Gasbarre. 2009. Use of a candidate gene array to delineate gene expression patterns in cattle selected for resistance or susceptibility to intestinal nematodes. *Vet. Parasitol.* **162**:106–115. doi:10.1016/j.vetpar.2008.12.017
- Ashburner, M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matise, J. E. Richardson, M. Ringwald, G. M. Rubin, and G. Sherlock. 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **25**:25–29. doi:10.1038/75556
- Aulchenko, Y. S., S. Ripke, A. Isaacs, and C. M. van Duijn. 2007. GenABEL: an R library for genome-wide association analysis. *Bioinformatics* **23**:1294–1296. doi:10.1093/bioinformatics/btm108
- Bado, A., S. Levasseur, S. Attoub, S. Kermorgant, J. P. Laigneau, M. N. Bortoluzzi, L. Moizo, T. Lehy, M. Guerre-Millo, Y. Le Marchand-Brustel, and M. J. M. Lewin. 1998. The stomach is a source of leptin. *Nature* **394**:790–793. doi:10.1038/29547
- Baker, R. L., and G. D. Gray. 2004. Appropriate breeds and breeding schemes for sheep and goats in the tropics. In: R. A. Sani, G. D. Gray, and R. L. Baker, editors, *Worm control for small ruminants in tropical Asia*. Australian Center for International Agricultural Research, Canberra, Australia. p. 63–95.
- Balic, A., C. P. Cunningham, and E. N. Meeusen. 2006. Eosinophil interactions with *Haemonchus contortus* larvae in the ovine gastrointestinal tract. *Parasite Immunol.* **28**:107–115. doi:10.1111/j.1365-3024.2006.00816.x
- Benavides, M. V., T. S. Sonstegard, S. Kemp, J. M. Mugambi, J. P. Gibson, R. L. Baker, O. Hanotte, K. Marshall, and C. Van Tassel. 2015. Identification of novel loci associated with gastrointestinal parasite resistance in a Red Maasai x Dorper backcross population. *PLoS ONE* **10**:e0122797. doi:10.1371/journal.pone.0122797
- Benavides, M. V., T. S. Sonstegard, and C. Van Tassel. 2016. Genomic regions associated with sheep resistance to gastrointestinal nematodes. *Trends Parasitol.* **32**:470–480. doi:10.1016/j.pt.2016.03.007
- Berton, M. P., R. M. de Oliveira Silva, E. Peripolli, N. B. Stafuzza, J. F. Martin, M. S. Álvarez, B. V. Gavina, M. A. Toro, G. Banchemo, P. S. Oliveira, J. Pereira Eler, F. Baldi, and J. B. Sterman Ferraz. 2017. Genomic regions and pathways associated with gastrointestinal parasites resistance in Santa Inês breed adapted to tropical climate. *J. Anim. Sci. Biotechnol.* **8**:73. doi:10.1186/s40104-017-0190-4
- Bhuiyan, A. A., J. Li, Z. Wu, P. Ni, A. A. Adetula, H. Wang, C. Zhang, X. Tang, A. A. Bhuyan, S. Zhao, and X. Du. 2017. Exploring the genetic resistance to gastrointestinal nematodes infection in goat using RNA-sequencing. *Int J Mol Sci.* **18**:751. doi:10.3390/ijms18040751
- Bishop, S. C. 2012. Possibilities to breed for resistance to nematode parasite infections in small ruminants in tropical production systems. *Animal* **6**:741–747. doi:10.1017/S1751731111000681
- Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. Mcvean, and R. Durbin; 1000 Genomes Project Analysis Group. 2011. The variant call format and VCFtools. *Bioinformatics* **27**:2156–2158. doi:10.1093/bioinformatics/btr330
- Diez-Tascón, C., O. M. Keane, T. Wilson, A. Zadissa, D. L. Hyndman, D. B. Baird, J. C. McEwan, and A. M. Crawford. 2005. Microarray analysis of selection lines from outbred populations to identify genes involved with nematode parasite resistance in sheep. *Physiol. Genomics* **21**:59–69. doi:10.1152/physiolgenomics.00257.2004
- Estrada-Reyes, Z. M., Y. Tsukahara, A. L. Goetsch, T. A. Gipson, T. Sahlh, R. Puchala, Z. Wang, S. P. Hart, and R. G. Mateescu. 2018. Effect of Ovar-DRA and Ovar-DRB1 genotype in small ruminants with haemonchosis. *Parasite Immunol.* **40**:e12534. doi:10.1111/pim.12534
- Gadahi, J. A., B. Yongqian, M. Ehsan, Z. C. Zhang, S. Wang, R. F. Yan, X. K. Song, L. X. Xu, and X. R. Li. 2016. *Haemonchus contortus* excretory and secretory proteins (HcESPs) suppress functions of goat PBMCs in vitro. *Oncotarget* **7**:35670–35679. doi:10.18632/oncotarget.9589
- Goddard, M. E., and B. J. Hayes. 2009. Mapping genes for complex traits in domestic animals and their use in breeding programmes. *Nat. Rev. Genet.* **10**:381–391. doi:10.1038/nrg2575
- Goolsby, M. K., M. L. Leite-Browning, and R. Browning, Jr. 2017. Evaluation of parasite resistance to commonly used commercial anthelmintics in meat goats on humid subtropical pasture. *Small Rumin. Res.* **146**:37–40. doi:10.1016/j.smallrumres.2016.11.022
- Greer, A. W., Y. R. Boisclair, M. Stankiewicz, and R. W. McNulty. 2009. Leptin concentrations and the immune-mediated

- reduction of feed intake in sheep infected with the nematode *Trichostrongylus colubriformis*. *Brit. J. Nutr.* **102**:954–957.
- Grisart, B., W. Coppieters, F. Farnir, L. Karim, C. Ford, P. Berzi, N. Cambisano, M. Mni, S. Reid, P. Simon, R. Spelman, M. Georges, and R. Snell. 2002. Positional candidate cloning of a QTL in dairy cattle: identification of a missense mutation in the bovine DGAT1 gene with major effect on milk yield and composition. *Genome Res.* **12**:222–231. doi:10.1101/gr.224202
- Guo, Z., J. F. González, J. N. Hernandez, T. M. McNeilly, Y. Corripio-Miyar, D. Frew, T. Morrison, P. Yu, and R. W. Li. 2016. Possible mechanisms of host resistance to *Haemonchus contortus* infection in sheep breeds native to the Canary Islands. *Sci. Rep.* **6**:1–14. doi:10.1038/srep26200
- Hassan, M., J. P. Hanrahan, B. Good, G. Mulcahy, and T. Sweeney. 2011. A differential interplay between the expression of Th1/Th2/Treg related cytokine genes in *Teladorsagia circumcincta* infected DRB1*1101 carrier lambs. *Vet. Res.* **42**(1):45. doi:10.1186/1297-9716-42-45
- Howell, S. B., J. M. Burke, J. E. Miller, T. H. Terrill, E. Valencia, M. J. Williams, L. H. Williamson, A. M. Zajac, and R. M. Kaplan. 2008. Prevalence of anthelmintic resistance on sheep and goat farms in the southeastern United States. *J. Am. Vet. Med. Assoc.* **233**:1913–1919. doi:10.2460/javma.233.12.1913
- Ingham, A., A. Reverter, R. Windon, P. Hunt, and M. Menzies. 2008. Gastrointestinal nematode challenge induces some conserved gene expression changes in the gut mucosa of genetically resistant sheep. *Int. J. Parasitol.* **38**:431–442. doi:10.1016/j.ijpara.2007.07.012
- Johnson, A., H. Trumbower, and W. Sadee. 2011. RNA structures affected by single nucleotide polymorphisms in transcribed regions of the human genome. *Bioinformatics* **2**:WMC001600.
- Kaplan, R. M. 2004. Drug resistance in nematodes of veterinary importance: a status report. *Trends Parasitol.* **20**:477–481. doi:10.1016/j.pt.2004.08.001
- Karim, L., W. Coppieters, L. Grobet, A. Valentini, and M. Georges. 2000. Convenient genotyping of six myostatin mutations causing double-muscling in cattle using a multiplex oligonucleotide ligation assay. *Anim. Genet.* **31**:396–399. doi:10.1046/j.1365-2052.2000.00684.x
- Keane, O. M., A. Zadissa, T. Wilson, D. L. Hyndman, G. J. Greer, D. B. Baird, A. F. McCulloch, A. M. Crawford, and J. C. McEwan. 2006. Gene expression profiling of naive sheep genetically resistant and susceptible to gastrointestinal nematodes. *BMC Genomics* **7**:42. doi:10.1186/1471-2164-7-42
- Kemper, K. E., D. L. Emery, S. C. Bishop, H. Oddy, B. J. Hayes, S. Dominik, J. M. Henshall, and M. E. Goddard. 2011. The distribution of SNP marker effects for faecal worm egg count in sheep, and the feasibility of using these markers to predict genetic merit for resistance to worm infections. *Genet. Res. (Camb)*. **93**:203–219. doi:10.1017/S0016672311000097
- Lee, W. P., M. P. Stromberg, A. Ward, C. Stewart, E. P. Garrison, and G. T. Marth. 2014. MOSAIK: a hash-based algorithm for accurate next-generation sequencing short-read mapping. *PLoS ONE* **9**:e90581. doi:10.1371/journal.pone.0090581
- Lees, M. S., N. A. Robinson, A. B. Ingham, A. C. Kotze, and D. M. Piedrafita. 2011. Dual oxidase 2 and glutathione peroxidase gene expression are elevated in hyperimmunized sheep challenged with *Haemonchus contortus*. *Vet. Parasitol.* **179**:113–122. doi:10.1016/j.vetpar.2011.02.013
- Leppek, K., R. Das, and M. Barna. 2018. Functional 5' UTR mRNA structures in eukaryotic translation regulation and how to find them. *Nat. Rev. Mol. Cell Biol.* **19**:158–174. doi:10.1038/nrm.2017.103
- Li, R. W., and L. C. Gasbarre. 2010. Gene expression in the bovine gastrointestinal tract during nematode infection. In: G. LaMann, editor. *Vet parasitology*. Nova Biomedical Press, New York. p. 157–178.
- Lorenz, R., S. H. Bernhart, C. Höner Zu Siederdisen, H. Tafer, C. Flamm, P. F. Stadler, and I. L. Hofacker. 2011. ViennaRNA package 2.0. *Algorithms Mol. Biol.* **6**:26. doi:10.1186/1748-7188-6-26
- MacKinnon, K. M., J. L. Burton, A. M. Zajac, and D. R. Notter. 2009. Microarray analysis reveals difference in gene expression profiles of hair and wool sheep infected with *Haemonchus contortus*. *Vet. Immunol. Immunopathol.* **130**:210–220. doi:10.1016/j.vetimm.2009.02.013
- McGuckin, M. A., S. K. Lindén, P. Sutton, and T. H. Florin. 2011. Mucin dynamics and enteric pathogens. *Nat. Rev. Microbiol.* **9**:265–278. doi:10.1038/nrmicro2538
- McRae, K. M., J. C. McEwan, K. G. Dodds, and N. J. Gemmell. 2014. Signatures of selection in sheep bred for resistance or susceptibility to gastrointestinal nematodes. *BMC Genomics* **15**:637. doi:10.1186/1471-2164-15-637
- McRae, K. M., M. J. Stear, B. Good, and O. M. Keane. 2015. The host immune response to gastrointestinal nematode infection in sheep. *Parasite Immunol.* **10**:605–613. doi:10.1111/pim.12290
- Meeusen, E. N., and A. Balic. 2000. Do eosinophils have a role in the killing of helminth parasites? *Parasitol. Today* **16**:95–101. doi:10.1016/s0169-4758(99)01607-5
- Menzies, M., A. Reverter, N. Andronicos, P. Hunt, R. Windon, and A. Ingham. 2010. Nematode challenge induces differential expression of oxidant, antioxidant and mucous genes down the longitudinal axis of the sheep gut. *Parasite Immunol.* **32**:36–46. doi:10.1111/j.1365-3024.2009.01156.x
- Miller, J. E., M. Bahirathan, S. L. Lemarie, F. G. Hembry, M. T. Kearney, and S. R. Barras. 1998. Epidemiology of gastrointestinal nematode parasitism in Suffolk and Gulf Coast Native sheep with special emphasis on relative susceptibility to *Haemonchus contortus* infection. *Vet. Parasitol.* **74**:55–74. doi:10.1016/s0304-4017(97)00094-0
- Miller, D. K., and T. M. Craig. 1996. Use of anthelmintic combinations against multiple resistant *Haemonchus contortus* in Angora goats. *Small Rumin. Res.* **19**:281–283.
- Moncada, D. M., S. J. Kammanadiminti, and K. Chadee. 2003. Mucin and Toll-like receptors in host defense against intestinal parasites. *Trends Parasitol.* **19**:305–311.
- O'Shea, J. J., M. Gadina, and R. Siegel. 2013. Cytokines and cytokine receptors. In: R. R. Rich, T. A. Fleisher, W. T. Shearer, H. W. Schroeder, A. J. Frew, and C. M. Weyand, editors. *Clinical immunology*. 4th ed. London. p. 108–135. doi:10.1016/B978-0-7234-3691-1.00033-7
- Patel, N., T. Kreider, J. F. Urban, Jr, and W. C. Gause. 2009. Characterisation of effector mechanisms at the host:parasite interface during the immune response to tissue-dwelling intestinal nematode parasites. *Int. J. Parasitol.* **39**:13–21. doi:10.1016/j.ijpara.2008.08.003
- Periasamy, K., R. Pichler, M. Poli, S. Cristel, B. Cetrá, D. Medus, M. Basar, A. K. Thiruvankadan, S. Ramasamy, M. B. Ellahi, F. Mohammed, A. Teneva, M. Shamsuddin, M. Garcia Podesta, and A. Diallo. 2014. Candidate gene approach for parasite resistance in sheep—variation in immune pathway genes and association with fecal egg count. *PLoS ONE*. **9**:e88337. doi:10.1371/journal.pone.0088337
- Pernthaner, A., S. A. Cole, L. Morrison, and W. R. Hein. 2005. Increased expression of interleukin-5 (IL-5), IL-13, and tumor necrosis factor alpha genes in intestinal lymph cells of sheep selected for enhanced resistance to nematodes during infection with *Trichostrongylus colubriformis*. *Infect. Immunol.* **73**:2175–2183.
- Roberts, F. H. S., and P. J. O'Sullivan. 1950. Methods for egg counts and larval cultures for strongyles infesting the gastro-intestinal tract of cattle. *Aust J Agric Res.* **1**(1):99–102. doi:10.1071/AR9500099
- Rosenberg, H. F., K. D. Dyer, and P. S. Foster. 2013. Eosinophils: changing perspectives in health and disease. *Nat. Rev. Immunol.* **13**:9–22. doi:10.1038/nri3341
- Rothenberg, M. E., A. Mishra, E. B. Brandt, and S. P. Hogan. 2001. Gastrointestinal eosinophils. *Immunol. Rev.* **179**:139–155. doi:10.1034/j.1600-065x.2001.790114.x
- Sallé, G., C. Moreno, S. Boitard, J. Ruesche, A. Tircazes-Secula, F. Bouvier, M. Aletru, J. L. Weisbecker, F. Prévot, J. P. Bergeaud, C. Trumel, C. Grisez, E. Liénard, and P. Jacquiet. 2014. Functional

- investigation of a QTL affecting resistance to *Haemonchus contortus* in sheep. *Vet. Res.* **45**:68. doi:10.1186/1297-9716-45-68
- Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B. Schwikowski, and T. Ideker. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**:2498–2504. doi:10.1101/gr.1239303
- Silva, M. V., T. S. Sonstegard, O. Hanotte, J. M. Mugambi, J. F. Garcia, S. Nagda, J. P. Gibson, F. A. Iraqi, A. E. McClintock, S. J. Kemp, P. J. Boettcher, M. Malek, C. P. Van Tassell, and R. L. Baker. 2012. Identification of quantitative trait loci affecting resistance to gastrointestinal parasites in a double backcross population of Red Maasai and Dorper sheep. *Anim. Genet.* **43**:63–71. doi:10.1111/j.1365-2052.2011.02202.x
- Smits, H. H., and M. Yazdanbakhsh. 2007. Chronic helminth infections modulate allergen-specific immune responses: protection against development of allergic disorders? *Ann. Med.* **39**:428–439. doi:10.1080/07853890701436765
- Stear, M. J., S. C. Bishop, N. G. Henderson, and I. Scott. 2003. A key mechanism of pathogenesis in sheep infected with the nematode *Teladorsagia circumcincta*. *Anim. Health Res. Rev.* **4**:45–52. doi:10.1079/AHRR200351
- Sweeney, T., J. P. Hanrahan, M. T. Ryan, and B. Good. 2016. Immunogenomics of gastrointestinal nematode infection in ruminants—breeding for resistance to produce food sustainably and safely. *Parasite Immunol.* **38**:569–586. doi:10.1111/pim.12347
- Terefe, G., C. Grisez, F. Prevot, J. P. Bergeaud, P. Dorchies, J. C. Brunel, D. François, I. Fourquaux, and P. Jacquet. 2007. In vitro pre-exposure of *Haemonchus contortus* L3 to blood eosinophils reduces their establishment potential in sheep. *Vet. Res.* **38**:647–654. doi:10.1051/vetres:2007019
- Urban, J. F., Jr, N. Noben-Trauth, D. D. Donaldson, K. B. Madden, S. C. Morris, M. Collins, and F. D. Finkelman. 1998. IL-13, IL-4Ralpha, and Stat6 are required for the expulsion of the gastrointestinal nematode parasite *Nippostrongylus brasiliensis*. *Immunity* **8**:255–264. doi:10.1016/s1074-7613(00)80477-x
- Venturina, V. M., A. G. Gossner, and J. Hopkins. 2013. The immunology and genetics of resistance of sheep to *Teladorsagia circumcincta*. *Vet. Res. Commun.* **37**(2):171–181. doi:10.1007/s11259-013-9559-9
- Zajac, A. M., and T. A. Gipson. 2000. Multiple anthelmintic resistance in a goat herd. *Vet. Parasitol.* **87**:163–172. doi:10.1016/s0304-4017(99)00174-0
- Zajac, A. M., R. P. Herd, and K. E. McClure. 1988. Trichostrongylid parasite populations in pregnant or lactating and unmated Florida Native and Dorset/Rambouillet ewes. *Int. J. Parasitol.* **18**:981–985. doi:10.1016/0020-7519(88)90181-6